

Probability&RV

Assignment-03

U Anuradha-ee21resch01008

download Python code from

https://github.com/Anuradha-Uggi/Assignments-AI5002-Probability-and-Random-Variables/blob/main/Prob_ass04/rvsp_urn_balls.py

Download Latex code from

https://github.com/Anuradha-Uggi/Assignments-AI5002-Probability-and-Random-Variables/blob/main/Prob_ass04/UrnBalls.tex

I. QUESTION(Prob-3,8)

Six balls are drawn successively from an urn containing 7 red and 9 black balls. Tell whether or not the trials of drawing balls are Bernoulli trials when after each draw the ball drawn is

- (i) replaced
- (ii) not replaced in the urn.

II. SOLUTION

Properties to be satisfied if a trial needs to be a bernoulli trial: 1.Number of trials should be finite
2.each trial should have outcomes of success and failure
3.if P is the success probability then failure probability should be 1-P.
4.probability of success should not vary in all trials.

case(i):Replaced

Number of red balls = 7

Number of black balls = 9

let X be the random variable that indicates probability of success and failure.

let $P(x=1)$ is the probability of success lets take it as picking a red ball.

and $P(x=0)$ is the probability of failure lets take it as picking a black ball.

$P(x = 1) = \frac{7}{16}$ and its same for any red ball drawn since the drawn ball is replaced.

$$P(x = 0) = \frac{9}{16} = 1 - P(x = 1)$$

so it satisfies all properties of Bernoulli therefore Trials under Replaced case are Bernoulli Trials.

case(ii):Not Replaced

$P(x = 1) = \frac{7}{16}$ and for 2nd trial $P(x = 1) = \frac{6}{15}$ similarly

$P(x = 0) = \frac{9}{16}$ and for 2nd trial $P(x = 0) = \frac{8}{15}$ so here the probability of success and corresponding failure is varying in each trial therefore under Not Replaced case Trials are not Bernoulli Trials.

III. CONCLUSION

Case(i):Drawn balls are Replaced into the Urn

Trials are Bernoulli Trials

Case(ii):Drawn balls are not Replaced

Trials are not Bernoulli Trials.