GENERAL ASSEMBLY

DATA SCIENCE

11 WEEK PART TIME COURSE

Week 3 - Regression
Monday 4th April

AGENDA

1. Motivation
2. Supervised Vs Unsupervised learning
3. What is Linear Regression?
4. How do Run a Linear Regression Model?
5. Lab
6. Discussion / Review / Homework

WHAT ARE THE GOALS OF STATISTICAL LEARNING?


scikit-learn
algorithm cheat-sheet
START
get more data
NO
>50 samples
YES
classification
kernel approximation
NOT WORKING
SVC
Ensemble Classifiers
NOT WORKING
KNeighbors Classifier
SGD Classifier
NO
Naive Bayes
YES
Text Data
NO
NOT WORKING
Linear SVC
YES
<100K samples
NO
regression
SGD Regressor
Lasso
ElasticNet
SVR(kernel='rbf')
EnsembleRegressors
<100K samples
NO
YES
few features should be important
YES
NOT WORKING
RidgeRegression
NO
SVR(kernel='linear')
YES
predicting a category
YES
do you have labeled data
NO
predicting a quantity
YES
NO
clustering
Spectral Clustering
GMM
NOT WORKING
KMeans
YES
number of categories known
YES
<10K samples
NO
NO
<10K samples
YES
MiniBatch KMeans
MeanShift
VBGMM
NO
just looking
YES
Randomized PCA
NOT WORKING
Isomap
Spectral Embedding
NOT WORKING
LLE
YES
<10K samples
NO
kernel approximation
dimensionality reduction
NO
predicting structure
tough luck
Back
scikit learn

We want to predict some value, let's call it **y**, based on some observed data we have, let's call that **x**.

We will use statistical learning to estimate a function that approximates **y** based on the input, **x**.

y is also called; label, dependent variable, target

x is also called; predictor, independent variable, features

We want to predict the price of a house, let's call it y, based on some observed data we have about the area, number of bedrooms, size of the house, and if it has a pool or not.

The area, number of bedrooms, size of the house, and if it has a pool or not would be our x variables (sometimes you might see this denoted as X)

What we want is $y = f(X)$, a way to describe the house price based on observed data

If the y variable is numeric then we have a regression problem - we are trying to predict a continuous number

If the y variable is a category (for example trying to predict a type of flower) the we have a classification problem - we are trying to classify what group that y belongs to.

We want to find some underlying structure or patterns in the data but in this case we don't have any labeled data.

So for example, if we have a large group of customers but would like to separate them into groups (or clusters) to better target them.

WHAT IS LINEAR REGRESSION?

We want to model a linear relationship (think straight line) between our target variable y and our input variable x.

$$y = X\beta + \epsilon$$

$$y = X\beta + \epsilon$$

- y = target variable
- X = input variable
- β = coefficients
- ϵ = error term

Note, one of our input variables can be 1 so we have an intercept parameter

- Linear relationship in the parameters, β, we can transform the actual values of the inputs if we want
- Variance of the error term, ϵ, is constant. This means there is no systematic pattern in the values of X and the variance of ϵ
- The mean of $\epsilon = 0$
- ϵ has a normal distribution
- No perfect (or near perfect) co-linearity between any of the input variables. Otherwise the fitting procedure will break.

DATA SCIENCE PART TIME COURSE

HOW TO RUN LINEAR REGRESSION?

$$SS_{res} = \sum_{i=1}^{n} (y_i - f(x_i))^2$$

Basically, what we are trying to do is minimise the Residual Sum of Squares. This is the Sum of the squared difference between our observed value and the value from the model

$$SS_{res} = \sum_{i=1}^{n} (y_i - f(x_i))^2$$

Basically, what we are trying to do is minimise the Residual Sum of Squares. This is the Sum of the squared difference between our observed value and the value from the model


Oak Diameter vs. Age
DBH (inch)
Age (years)
0
5
10
15
0
20
40
60
80
100

- Make sure you visualise your data and check the actual model fit !!!
- The fitting a model to the four datasets in the table on the right produce the same fit statistics, model coefficients and standard error
- See anything wrong?

Anscombe's quartet

I		II		III		IV	
x	y	x	y	x	y	x	y
10.0	8.04	10.0	9.14	10.0	7.46	8.0	6.58
8.0	6.95	8.0	8.14	8.0	6.77	8.0	5.76
13.0	7.58	13.0	8.74	13.0	12.74	8.0	7.71
9.0	8.81	9.0	8.77	9.0	7.11	8.0	8.84
11.0	8.33	11.0	9.26	11.0	7.81	8.0	8.47
14.0	9.96	14.0	8.10	14.0	8.84	8.0	7.04
6.0	7.24	6.0	6.13	6.0	6.08	8.0	5.25
4.0	4.26	4.0	3.10	4.0	5.39	19.0	12.50
12.0	10.84	12.0	9.13	12.0	8.15	8.0	5.56
7.0	4.82	7.0	7.26	7.0	6.42	8.0	7.91
5.0	5.68	5.0	4.74	5.0	5.73	8.0	6.89

LINEAR REGRESSION – IMPORTANT STEP



LAB

git remote -v

git remote add upstream https://github.com/ihansel/SYD_DAT_3.git

git remote -v

git fetch upstream

git checkout master

git merge upstream/master

DATA SCIENCE – Week 2 Day 2

DISCUSSION TIME

- Review of last class
- Further Reading for Regression
- Check in with homework/course project

WEEK 2 – Wednesday



Chart Suggestions—A Thought-Starter



DATA SCIENCE – Week 3 Day 1

DISCUSSION TIME

An Introduction to Statistical Learning

- **Chapter 3 – Linear Regression**
- **Chapter 6 – Linear Model Selection and Regularization**



DISCUSSION TIME

Homework/Course Project

- How’s it going ?