


Warning notification!!!!

- The PPTs are prepared for the offline interactive teaching in the class using the materials from different books and web. The author may not have the legal permission for online sharing those materials via social media/web/email or use in business etc in public domain.
- Therefore, students are requested not to share the PPTs outside the class/institute, which can violation the copy-write related issues.

- **Evaluate the response** of the LTI system with impulse response

$$h(t) = e^{-at}u(t), \quad a > 0$$

To the input signal $x(t) = e^{-bt}u(t), \quad b > 0$



From **convolution theorem**, we can write $y(t) = \int_{-\infty}^{\infty} x(\tau) h(t - \tau) d\tau$

(using convolution property)

$$y(t) = x(t) * h(t) \xleftrightarrow{\text{FT}} X(\omega)H(\omega)$$

Fourier transform of $x(t)$ and $h(t)$ as, $X(\omega) = \frac{1}{b + j\omega}$ $H(\omega) = \frac{1}{a + j\omega}$

$$Y(\omega) = X(\omega)H(\omega) = \frac{1}{(b + j\omega)(a + j\omega)} = \frac{A}{(a + j\omega)} + \frac{B}{(b + j\omega)}$$

Partial-fraction expansion

$$Y(\omega) = \frac{1}{(b + j\omega)(a + j\omega)} = \frac{A}{(a + j\omega)} + \frac{B}{(b + j\omega)} = \frac{A}{(a + s)} + \frac{B}{(b + s)} \quad \text{Let } s = j\omega$$

$$= \frac{1}{b - a} \frac{1}{(a + j\omega)} + \frac{1}{a - b} \frac{1}{(b + j\omega)}$$

$$= \frac{1}{a - b} \left[\frac{1}{b + j\omega} \right] - \frac{1}{a - b} \left[\frac{1}{a + j\omega} \right]$$

$$e^{-at}u(t) \stackrel{FT}{\leftrightarrow} \frac{1}{a + j\omega}$$

$$A = [Y(\omega)(a + s)]_{s=-a} = \frac{1}{b - a}$$

$$B = [Y(\omega)(b + s)]_{s=-b} = \frac{1}{a - b}$$

$$y(t) = \frac{1}{a - b} e^{-bt}u(t) - \frac{1}{a - b} e^{-at}u(t)$$

$$X(s) = \frac{1}{(s + 1)(s + 2)} = \frac{A}{s + 1} + \frac{B}{s + 2}.$$

we can evaluate the coefficients A and B by

$$A = [(s + 1)X(s)]_{s=-1} = 1,$$

$$B = [(s + 2)X(s)]_{s=-2} = -1.$$

$$X(s) = \frac{1}{s + 1} - \frac{1}{s + 2}.$$

- The input output of a system is described by $y'(t) + 2y(t) = x(t) + x'(t)$

Compute the impulse response $h(t)$ of the system

Taking Fourier-transform both side of the equation

$$y'(t) + 2y(t) = x(t) + x'(t)$$

$$\rightarrow j\omega Y(\omega) + 2 Y(\omega) = X(\omega) + j\omega X(\omega)$$

$$\rightarrow (j\omega + 2)Y(\omega) = (1 + j\omega)X(\omega)$$

$$H(\omega) = \frac{Y(\omega)}{X(\omega)} = \frac{1+j\omega}{2+j\omega} = \frac{2+j\omega-1}{2+j\omega} = 1 - \frac{1}{2+j\omega}$$

$$\text{Taking Inverse: } h(t) = \delta(t) - e^{-2t}u(t)$$

Duality

If


$$x(t) \xleftrightarrow{FT} X(\omega)$$

$$X(t) \xleftrightarrow{FT} ?$$

$$\int_{-\infty}^{\infty} X(t) e^{-j\omega t} dt = ?$$

Called **duality of time and frequency**.

The duality principle may be compared with a *photograph and its negative*. A photograph can be obtained *from its negative*, and by using an identical procedure, *a negative can be obtained from the photograph*



Proof:

We know, $x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(\omega) e^{j\omega t} d\omega \quad \Rightarrow \quad = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(\tau) e^{j\tau t} d\tau$

$\Rightarrow 2\pi x(t) = \int_{-\infty}^{\infty} X(\tau) e^{j\tau t} d\tau$

Putting, $t = -\omega$


$\Rightarrow 2\pi x(-\omega) = \int_{-\infty}^{\infty} X(\tau) e^{j\tau \cdot -\omega} d\tau$

Putting, $\tau = t$

$$= \int_{-\infty}^{\infty} X(t) e^{-j\omega t} dt$$

$$= FT\{X(t)\}$$

$X(t) \xleftrightarrow{FT} 2\pi x(-\omega) = x(-f)$



$$x(t) = \begin{cases} 1, & |t| < \frac{\tau}{2} \\ 0, & \text{else} \end{cases}$$

$$= \text{rect}\left(\frac{t}{\tau}\right)$$



$$X(\omega) = \tau \frac{\sin\left(\omega \frac{\tau}{2}\right)}{\omega \frac{\tau}{2}}$$

$$= \tau \text{sinc}\left(\omega \frac{\tau}{2}\right)$$

$X(t)$ is the same as $X(\omega)$ with ω replaced by t



$$x(t) = \begin{cases} 1, & |t| < \frac{\tau}{2} \\ 0, & \text{else} \end{cases}$$

$$= \text{rect}\left(\frac{t}{\tau}\right)$$




$$X(\omega) = \tau \frac{\sin\left(\omega \frac{\tau}{2}\right)}{\omega \frac{\tau}{2}}$$

$$= \tau \text{sinc}\left(\frac{\omega\tau}{2}\right)$$



$$2\pi x(-\omega)$$

$x(-\omega)$ is the **same** as $x(t)$
with t **replaced by** $-\omega$.



$$X(t) \xleftrightarrow{FT} 2\pi x(-\omega) = x(-f)$$

$$\underbrace{\tau \text{sinc}\left(\frac{\tau t}{2}\right)}_{X(t)} \iff \underbrace{2\pi \text{rect}\left(\frac{-\omega}{\tau}\right)}_{2\pi x(-\omega)} = 2\pi \text{rect}\left(\frac{\omega}{\tau}\right)$$

- Evaluate Fourier transform of the signal $x(t) = \frac{1}{1+t^2}$

$$y(t) = e^{-a|t|}; a > 0$$

$$y(t) \xleftrightarrow{FT} Y(\omega) = \frac{2a}{a^2 + \omega^2}$$

$$a = 1$$

Using dual property: $Z(t) \xleftrightarrow{FT} 2\pi Z(-\omega)$

$$y(t) = e^{-|t|} \xleftrightarrow{FT} Y(\omega) = \frac{2}{1 + \omega^2}$$

$$Y(t) = \frac{2}{1+t^2} \xleftrightarrow{FT} 2\pi y(-\omega)$$

$$\Rightarrow \frac{1}{1+t^2} \xleftrightarrow{FT} \pi e^{-|-\omega|}$$

$$\Rightarrow \frac{1}{1+t^2} \xleftrightarrow{FT} \pi e^{-|\omega|}$$

Given, $X(\omega) = \frac{1}{(1+\omega^2)} e^{-\frac{2\omega^2}{(1+\omega^2)}}$. **Determine the Fourier transform** of the following signals:

(a) $x(t-2)e^{jt}$

(b) $x(1-t)$

(c) $x\left(\frac{t}{2} - 2\right)$

(a) Time shift property: $x(t-t_0) \xleftrightarrow{FT} e^{-j\omega t_0} X(\omega) \Rightarrow y(t) = x(t-2) \xleftrightarrow{FT} Y(\omega) = e^{-j\omega \cdot 2} X(\omega)$

Frequency shift property: $x(t) e^{j\omega_c t} \xleftrightarrow{FT} X(\omega - \omega_c) \Rightarrow y(t) e^{j\omega_c = 1 \cdot t} \xleftrightarrow{FT} Y(\omega - \omega_c)$

$$y(t) e^{j\omega_c = 1 \cdot t} = x(t-2) e^{j\omega_c = 1 \cdot t} \xleftrightarrow{FT} Y(\omega - \omega_c) = e^{-j2(\omega - \omega_c)} X(\omega - \omega_c)$$

where $\omega_c = 1$

(b) Scaling property:

$$x(at) \xleftrightarrow{FT} \frac{1}{|a|} X\left(\frac{\omega}{a}\right)$$

$$\Rightarrow x(-t) \xleftrightarrow{FT} \frac{1}{|-1|} X\left(\frac{\omega}{-1}\right) = X(-\omega) \quad (\text{for } a=-1) \quad [\text{folding}]$$

$$x(-t + 1) \xleftrightarrow{FT} \int_{-\infty}^{\infty} x(-t + 1) e^{-j\omega t} dt$$

(replacing $-t + 1 = \tau$, $dt = -d\tau$)

$$\xleftrightarrow{FT} \int_{+\infty}^{-\infty} x(\tau) e^{-j\omega(-\tau+1)} \cdot -d\tau$$

$$\xleftrightarrow{FT} e^{-j\omega} \int_{+\infty}^{-\infty} x(\tau) e^{j\omega\tau} \cdot -d\tau$$

$$\xleftrightarrow{FT} e^{-j\omega} \int_{-\infty}^{\infty} x(\tau) e^{j\omega\tau} d\tau = e^{-j\omega} \int_{-\infty}^{\infty} x(\tau) e^{-j \cdot (-\omega) \cdot \tau} d\tau = e^{-j\omega} X(-\omega)$$

Using Time shift property (delay):

$$x(t - t_0) \xleftrightarrow{FT} e^{-j\omega t_0} X(\omega)$$

$$x(-t + 1) \xleftrightarrow{FT} e^{-j\omega \cdot 1} X(-\omega)$$

(c) Using Time shift property (delay): $x(t - t_0) \xleftrightarrow{FT} e^{-j\omega t_0} X(\omega)$

$$y(t) = x(t - 2) \xleftrightarrow{FT} Y(\omega) = e^{-j\omega \cdot 2} X(\omega) \quad [\text{shifting}]$$

Using scaling property $x(at) \xleftrightarrow{FT} \frac{1}{|a|} X\left(\frac{\omega}{a}\right)$ [Scaling]

$$y\left(\frac{t}{2}\right) = x\left(\frac{t}{2} - 2\right) \xleftrightarrow{FT} \frac{1}{1/2} Y\left(\frac{\omega}{1/2}\right) = 2 \cdot Y(2\omega) = 2e^{-j \cdot 2\omega \cdot 2} X(2\omega)$$

Solve the problems



Q1. Evaluate the Fourier transform of the following signals:

(a) $e^{-\alpha t} \cos \omega_0 t u(t), \alpha > 0$

(b) $e^{-3|t|} \sin 2t$

(c) $x(t) = \begin{cases} 1 + \cos \pi t, & |t| \leq 1 \\ 0, & |t| > 1 \end{cases}$

(d) $\sum_{k=0}^{\infty} \alpha^k \delta(t - kt), |\alpha| < 1$



(e) $te^{-2t} \sin 4t u(t)$

(f) $x(t) = \begin{cases} 1 - t^2, & 0 < t < 1 \\ 0, & \text{else} \end{cases}$



(g) $x(t) = \sum_{n=-\infty}^{\infty} e^{-|t-2n|}$

Q2. Determine the signal $x(t)$ from the Fourier transform of the signals,

(a) $X(\omega) = \frac{2 \sin[3(\omega - 2\pi)]}{(\omega - 2\pi)}$

(b) $X(\omega) = \cos(4\omega + \frac{\pi}{3})$

(c) $X(\omega) = 2[\delta(\omega - 1) - \delta(\omega + 1)] + 3[\delta(\omega - 2\pi)] + \delta(\omega + 2\pi)$



Thank you