


Introduction and Background to Multilevel Analysis

Dr. J. Kyle Roberts

Southern Methodist University
Simmons School of Education and Human Development
Department of Teaching and Learning

Background and History of Multilevel Analysis

- Robinson (1950) and the problem of contextual effects
- The “Frog-Pond” Theory



History, cont.

- Statistical models that are not hierarchical sometimes ignore nesting structure and therefore report underestimated standard errors.
- Multilevel techniques are more efficient than other typical OLS techniques
- Multilevel techniques assume a General Linear Model framework and can thus perform all types of analyses.
- Multilevel techniques can go beyond questions of “Do schools differ?” to ask questions of “Why do schools differ?”

Types of Multilevel Structures


- Students nested in classrooms
- Students nested in schools
- Students nested in classrooms nested in schools
- Measurement occasions nested inside individuals
- Students cross-classified by middle and high school

Do We Really Need Multilevel Analysis?

- “All data are multilevel!”
- The problem of independence of observations
 - GPA and the SAT in different high schools
 - SES as it relates to school achievement
- The inefficiency of OLS techniques

Differences Between Multilevel and OLS Methods

- MLA is based on maximum likelihood and empirical Bayesian techniques
- $1 + 1 = 1.5$



Notating the Multilevel ANOVA

- Recall from Analysis of Variance (ANOVA) that we may notate a model testing for differences between groups as:

$$Y_{ij} = \mu + \alpha_j + \epsilon_{ij}, \quad \epsilon_{ij} \sim \mathcal{N}(0, \sigma^2), \quad j = 1, \dots, k, \quad i = 1, \dots, n_j$$

- We can further notate the above model as:

$$Y_{ij} = \gamma_{00} + u_{0j} + e_{ij}$$

- which may be decomposed as a Level-1 model

$$Y_{ij} = \beta_{0j} + e_{ij}$$

- and Level-2 model

$$\beta_{0j} = \gamma_{00} + u_{0j}$$

Notating the Multilevel ANOVA (cont.)


- This means that any individual score within each group may be thought of as

$$Y_{ij} = \beta_j + e_{ij} \left\{ \begin{array}{l} Y_{11} = \beta_1 + e_{11} \\ Y_{21} = \beta_1 + e_{21} \\ Y_{31} = \beta_1 + e_{31} \\ \dots \\ Y_{ij} = \beta_j + e_{ij} \end{array} \right.$$


- And any group score may be modeled as:

$$\beta_j = \gamma_{00} + u_{0j} \left\{ \begin{array}{l} \beta_1 = \gamma_{00} + u_{01} \\ \beta_2 = \gamma_{00} + u_{02} \\ \beta_3 = \gamma_{00} + u_{03} \\ \dots \\ \beta_j = \gamma_{00} + u_{0j} \end{array} \right.$$

Graphical Example of Multilevel ANOVA



Understanding Errors



Heuristic Example of Multilevel ANOVA

- Suppose that we have a dataset in which we have scores from 160 students nested inside 16 different schools.
- The dataset may be found at <http://faculty.smu.edu/kyler/courses/7309/sciach.txt>

```
> sciach <- read.table("sciach.txt", header = T)
> str(sciach)
```

```
'data.frame': 160 obs. of 10 variables:
 $ ID      : int  1 2 3 4 5 6 7 8 9 10 ...
 $ GROUP   : int  1 1 1 1 1 1 1 1 1 1 ...
 $ SCIENCE : int  1 1 2 2 3 3 4 4 5 5 ...
 $ URBAN   : int  8 7 7 6 6 5 5 5 3 2 ...
 $ GENDER  : int  1 1 1 1 1 2 2 2 2 2 ...
 $ CONS    : int  1 1 1 1 1 1 1 1 1 1 ...
 $ URB.MEAN : num  -6.43 -7.43 -7.43 -8.43 -8.43 ...
 $ SCH.RES  : num  5 5 5 5 5 5 5 5 5 5 ...
 $ SCH.RES.MEAN: num  -4.97 -4.97 -4.97 -4.97 -4.97 ...
 $ GEND.FAC : Factor w/ 2 levels "Female","Male": 2 2 2 2 2 1 1 1 1 1
```

```
> sciach$GROUP <- factor(sciach$GROUP)
```

ANOVA of Science Achievement Data

- We can run a simple ANOVA on the science achievement data comparing the means of the 16 schools.

```
> mean(sciach$SCIENCE)
```

```
[1] 10.6875
```

```
> with(sciach, tapply(SCIENCE, GROUP, mean))
```

```
      1      2      3      4      5      6      7      8      9     10     11     12
3.0  4.0  5.0  6.0  7.0  8.0  9.0 10.0 11.0 12.0 13.0 14.0
 13   14   15   16
15.0 16.0 18.5 19.5
```

```
> anova(aov(SCIENCE ~ GROUP, sciach))
```

Analysis of Variance Table

Response: SCIENCE

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
GROUP	15	3859.4	257.292	130	< 2.2e-16
Residuals	144	285.0	1.979		

Running the Same Model as a Multilevel ANOVA

- Recall from the multilevel ANOVA notation that we want to test:

$$Y_{ij} = \gamma_{00} + u_{0j} + e_{ij}$$

```
> m0 <- lme(SCIENCE ~ 1, random = ~1 | GROUP, sciach)
> summary(m0)
```

Linear mixed-effects model fit by REML

Data: sciach

	AIC	BIC	logLik
	643.8561	653.0628	-318.9281

Random effects:

Formula: ~1 | GROUP

(Intercept) Residual

StdDev: 5.052846 1.406829

Fixed effects: SCIENCE ~ 1

	Value	Std.Error	DF	t-value	p-value
(Intercept)	10.6875	1.268098	144	8.427975	0

Standardized Within-Group Residuals:

Comparing OLS and MLA Estimates


```
> cbind(means = with(sciach, tapply(SCIENCE, GROUP,  
+   mean)), coef(m0))
```

	means	(Intercept)
1	3.0	3.059135
2	4.0	4.051442
3	5.0	5.043750
4	6.0	6.036058
5	7.0	7.028365
6	8.0	8.020673
7	9.0	9.012981
8	10.0	10.005288
9	11.0	10.997596
10	12.0	11.989904
11	13.0	12.982212
12	14.0	13.974519
13	15.0	14.966827
14	16.0	15.959135
15	18.5	18.439904
16	19.5	19.432212


Checking Assumptions

```
> par(mfrow = c(1, 2))  
> boxplot(resid(m0) ~ GROUP, sciach, horizontal = T,  
+         main = "Homogeneity of Variance")  
> qqnorm(resid(m0), main = "QQplot for Null Model")
```

Homogeneity of Variance



QQplot for Null Model



Model Fit Indices

- Chi-square or χ^2

$$\chi^2 = -2 * \ell$$


- Akaike Information Criteria (AIC)

$$AIC = -2 * \ell + 2K$$


- Bayesian Information Criteria (BIC)

$$BIC = -2 * \ell + K * \ln(N)$$

Fixed and Random Effects




(a) No Random Effects




(b) Random Intercepts

Fixed and Random Effects



(c) Random Slopes



(d) Random Intercepts and Slopes

Recalling the OLS Linear Model

- Consider the following 1-level regression equation:

$$y = a + bx + \epsilon$$

where:

- y - the response variable
 - a - the y -intercept or the expected value when the covariate is 0
 - b - the expected change in the response variable (y) for every one unit change in the covariate
 - x - the covariate
 - ϵ - the residual term
- This model may also be written as:

$$y = \beta_0 + \beta_1 x + \epsilon$$

Adding a Random Effect for the Intercept

- We may further modify this equation to allow for variation among the intercepts for each pre-identified group such that:

$$y = \beta_0 + \beta_1 x + \epsilon$$

now becomes:

$$y_{ij} = \gamma_{00} + \gamma_{10}x_{ij} + u_{0j} + \epsilon_{ij}$$

where:

- y_{ij} - is the response variable for individual i in group j
- γ_{00} - the y -intercept or the expected value when the covariate is 0
- γ_{10} - the expected change in the response variable (y) for every one unit change in the covariate
- x_{ij} - the covariate term for each individual; the subscripts i and j mean that this variable is measured at the first level
- u_{0j} - the residual term defining the random variation of each of the group intercepts around the grand intercept γ_{00}
- ϵ_{ij} - the residual term defining the random variation of each person around their predicted group regression equation

Breaking Down the Mixed Effects Model into Levels

- From the previous slide, our mixed effects model:

$$y_{ij} = \gamma_{00} + \gamma_{10}x_{ij} + u_{0j} + \epsilon_{ij}$$

may be thought of as a 2-level model where:

- Level 1:

$$y_{ij} = \beta_{0j} + \beta_{1j}x_{ij} + \epsilon_{ij}$$

- Level 2:

$$\beta_{0j} = \gamma_{00} + u_{0j}$$


$$\beta_{1j} = \gamma_{10}$$

- where:

$$u_{0j} \sim \mathcal{N}(0, \sigma_{u_{0j}}^2)$$

$$\epsilon_{ij} \sim \mathcal{N}(0, \sigma_{\epsilon_{ij}}^2)$$

Understanding Errors Again



Running the Linear Model in R

```
> m.lm <- lm(SCIENCE ~ URBAN, sciach)
> summary(m.lm)
```

Call:

```
lm(formula = SCIENCE ~ URBAN, data = sciach)
```

Residuals:

Min	1Q	Median	3Q	Max
-5.3358	-2.1292	0.4919	2.0432	5.0090

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-1.25108	0.59371	-2.107	0.0367
URBAN	0.82763	0.03863	21.425	<2e-16


Residual standard error: 2.592 on 158 degrees of freedom

Multiple R-squared: 0.7439, Adjusted R-squared: 0.7423

F-statistic: 459 on 1 and 158 DF, p-value: < 2.2e-16

Running the Linear Model in R (cont.)

```
> plot(SCIENCE ~ URBAN, sciach)  
> abline(lm(SCIENCE ~ URBAN, sciach))
```



Running the Multilevel Model in R

```
> m1 <- lme(SCIENCE ~ URBAN, random = ~1 | GROUP,  
+          sciach)  
> summary(m1)
```

Linear mixed-effects model fit by REML

Data: sciach

AIC	BIC	logLik
508.094	520.3444	-250.047

Random effects:

Formula: ~1 | GROUP

(Intercept) Residual

StdDev: 9.29817 0.809449

Fixed effects: SCIENCE ~ URBAN

	Value	Std.Error	DF	t-value	p-value
(Intercept)	22.302911	2.4263101	143	9.192111	0
URBAN	-0.805228	0.0479985	143	-16.776087	0

Correlation:

(Intr)

URBAN -0.285

Comparing Models

```
> anova(m0, m1)
```

	Model	df	AIC	BIC	logLik	Test	L.Ratio
m0	1	3	643.8561	653.0628	-318.9281		
m1	2	4	508.0940	520.3444	-250.0470	1 vs 2	137.7621

p-value

m0	
m1	<.0001

- So we can see that by the addition of a single fixed effect to our model, we reduced the AIC by ~ 135 and the BIC by ~ 133 .