
Predicting Depression Using Health Care Data

By Vivienne DiFrancesco

Depression

- 264 million people globally have depression WHO
- 7.1% of US adults had a major depressive episode in 2017 with young adults being the most affected - NIMH



The Problem

- Physicians have become the front line for handling mental health disorders.
- The majority of physician visits are driven by mental disorders and few people follow up with a mental health professional.
- 90% of those who commit suicide have a mental health disorder and 40% had visited their doctor in the last month

The Problem

- Study published in JAMA Patients who receive more holistic care with doctors for mental health reduce healthcare costs and improve patient outcomes.
- This can be unfeasible for many reasons
 Buy-in, logistics, training
- Maybe machine learning could fill the gap to predict patients who are depressed.

Using machine learning to identify individuals with depression could connect patients with the help they need more quickly and easily while reducing healthcare costs and burden on physicians.


The Data

- CDC NHANES data
- 2005 to 2018 comprising 36259 entries total
- Tried to use data that was consistent across years and could reasonably be found in a patient's medical file
- Created labels of who was depressed and not depressed based on the "depression screener" in the data

Methods

- Classification of depressed or not depressed
- Roughly followed OSEMiN method
 - Obtain, scrub, explore, model, interpret
- Modeling was done from simple going to complex
- Wanted to use as few features as necessary
- Imbalanced data means accuracy was not a good metric
- Recall weighted more heavily to reduce false negatives


Best Model Results




Logistic regression is a model that calculates probabilities of entries being in one or the other class then uses a 50% threshold to make the prediction.

Recommendation 1:

Watch for patients with other mental problems





Recommendation 2:

Depression


Not Depressed

Depressed

Watch for patients who have trouble sleeping and sleep too much or too little

Trouble Sleeping History


95.0%




Sleep Hours By Percent Depression

Recommendation 3:

Watch for patients who can't work and/or have low household income





Recommendations Recap

- 1. Watch for patients with other mental problems
- 2. Watch for patients with sleep troubles
- 3. Watch for patients who can't work or have low income

Future Work

- Try different models perhaps neural networks
- Add more data
- Tuning and testing of parameters

Thank you for your time

Any questions?

Appendix

PHQ-9

Take the PHQ-9 online here: https://www.mdcalc.com/phq-9-patient-health-questionnaire-9

Over the <u>last 2 weeks</u> , how often have you been bothered by any of the following problems?	Not at	Several days	More than half the days	Nearly every day
Little interest or pleasure in doing things	0	1	2	3
2. Feeling down, depressed, or hopeless	0	1	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3
Feeling tired or having little energy	0	1	2	3
5. Poor appetite or overeating	0	1	2	3
Feeling bad about yourself — or that you are a failure or have let yourself or your family down	0	1	2	3
Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2	3
Moving or speaking so slowly that other people could have noticed? Or the opposite — being so fidgety or restless that you have been moving around a lot more than usual	0	1	2	3
Thoughts that you would be better off dead or of hurting yourself in some way	0	1	2	3
	For offic	E CODING		
	+	+ + +		

If you checked off <u>any</u> problems, how <u>difficult</u> have these problems made it for you to do your work, take care of things at home, or get along with other people?

Not difficult	Somewhat	Very	Extremely
at all	difficult	difficult	difficult

Copyright © 2010 Pfizer, Inc. All rights reserved.

Most Influential Model Features

