

An Easy Method for Cryopreservation of Zebrafish (*Danio rerio*) Sperm

Caroline Caetano Da Silva,^{1,*} Tímea Kollár,^{2,*} Zsolt Csenki-Bakos,²
Bianca H.V. Fernandes,³ Ákos Horváth,² and Luciani R. Carvalho¹

Abstract

We developed an easy, efficient, and cheap protocol for zebrafish sperm cryopreservation carried out on dry ice (20 min) using simple composition solution (200 mM glucose, 40 mM KCl, 30 mM Tris, pH=8.0). The average efficiency of the present cryopreserve method was between 10% and 20% (expressed as fertilization rate). The experiments were conducted and repeated at two different locations, in different countries, yielding very similar results, showing the reproducibility and applicability of the method.

Keywords: zebrafish, *Danio rerio*, sperm, cryopreservation, dry ice

THE WIDESPREAD USE of zebrafish (*Danio rerio*) as an animal model has led to numerous mutant strains and transgenic lines. This created the need to store these genetic resources, to prevent accidental loss, and to reduce space needed to maintain them. Cryopreservation can be a solution allowing to store genetic resources indefinitely and to recover them without compromising physiological function.^{1–3} There are many zebrafish sperm cryopreservation protocols; however, they vary in basic parameters including the freezing method,^{4,5} number of males to collect sperm,^{5,6} and types of extenders and cryoprotectants. Herewith, we develop an optimized, cheap, and straightforward zebrafish sperm cryopreservation protocol, as an adaptation for zebrafish, of a method previously published using grayling extender.⁷ Our method was consistently reproducible in different laboratories.

Wild-type zebrafish from AB lines were kept at the University of São Paulo Medical School, Brazil, whereas one other stock was maintained in Department of Aquaculture, Szent István University, Gödöllő, Hungary. Zebrafish males were separated from females 1 week before sperm collection. The fish were anesthetized before the experiments with 0.06% of tricaine methanesulfonate (MS-222).⁸ The anesthetized zebrafish were gently dried with paper towel and placed belly up in the slit of a dampened foam holder. The urogenital papilla was further dried just before sperm collection to prevent contamination with water. Five microliters of sperm mixed by pooling (1 μ L per male generally) was collected by stripping in 25 μ L of grayling extender.⁷ Immediately before freezing, 30 μ L of diluted sperm (25 μ L grayling extender +5 μ L sperm) was topped up to the volume of 100 μ L by adding 62 μ L of grayling extender and 8 μ L of methanol, resulting in a dilution ratio of 1:19 (sperm to extender+cryoprotectant) and 8% final concentration of methanol as cryoprotectant. Fifteen milliliters Falcon tubes were inserted into crushed dry ice before the stripping. The samples were loaded into 0.25 mL French straws that were

¹Discipline of Endocrinology, Internal Medicine Department, Hospital das Clínicas HCFMUSP, Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil.

²Department of Aquaculture, Szent István University, Gödöllő, Hungary.

³Diretoria Técnica de Apoio ao Ensino e Pesquisa da Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil.

*These authors contributed equally to this work.

placed in Falcon tubes individually. The tubes containing straws were closed with their screw caps and were held for 20 min⁴ in the crushed dry ice, and then they were plunged directly into liquid nitrogen and kept until fertilization. Prior fertilization, the cryopreserved straws were thawed in a water bath at 40°C for 5 s.

Breeding tanks with a barrier separating males and females were prepared one night before fertilization. The sex ratio was 1:2 (one male to two females). In the early morning, once the light turned on, eggs were removed from at least three anesthetized females by squeezing their bellies into a dry 52 mm Petri dish. Only eggs that had a satisfactory visual appearance (round shape, yellowish color, and strong texture) were used for fertilization. The cryopreserved and thawed sperm were added to the eggs with 1 mL of system water to activate the gametes followed by gentle shaking for 2 min. Five minutes after fertilization, water was replaced on the egg batches and they were incubated at 27°C ± 1°C. The fertilization rate was determined on the following day, as the percentage of fertilized embryos related to the total number of eggs per batch. Two groups of experiments were performed, one in São Paulo while the other in Gödöllő. Twelve replicates were done at both locations.

The average fertilization rate with standard deviation was 11.42% ± 8.06% in Brazil and 19.99% ± 9.68% in Hungary (Supplementary Table S1; Supplementary Data available online at www.libertpub.com/zeb). Our lowest fertilization rate was 2.39% and highest was 35.29%. Even in some cases with low fertilization rate, we have not got 0% in any case. Consequently, all samples had a measurable efficiency, thus, the proportion of successful samples is 100%.

Nowadays, there are several methods for zebrafish sperm cryopreservation (Supplementary Table S2), but some can be challenging to reproduce.⁹ Our method is simpler and cheaper than other available protocols in the literature. We have repeated the experiments 12 times at two different locations. However, we also observed variation among the pools, this disparity is normal in the available protocols and experiments as well.^{10–12} The differences of diet composition may influence physiological processes such as survival, growth, and reproductive performance,¹³ endocrine function,¹⁴ differences of the sperm and egg quality, and even the definition of successful fertilization⁵ can influence the disparity of results. In conclusion, this study showed a simple protocol to cryopreserve zebrafish sperm, with a good efficiency and reproducibility that can be done in any zebrafish holding facility without special equipment or devices.

Materials and Methods

See Supplementary Data for methods.

Acknowledgments

We thank the endocrinology discipline, the medical school foundation, FAPESP (2015/26563-7), and the Animal Research Facility (DTAPEP) in the University of São Paulo Medical School for supporting this project. The work is supported as well by the EFOP-3.6.3-VEKOP-16-2017.00008 project that is cofinanced by the European Union and the European Social Fund.

We thank Dr. Elaine Frade Costa for managing the zebrafish facility in the University of São Paulo.

This research was supported by the Higher Education Institutional Excellence Program (1783-3/2018/FEKUTSRAT) awarded by the Ministry of Human Capacities within the framework of water-related researches of Szent István University.

Disclosure Statement

No competing financial interests exist.

Supplementary Material

Supplementary Data
Supplementary Table1
Supplementary Table2

References

1. Mazur P. Kinetics of water loss from cells at subzero temperatures and the likelihood of intracellular freezing. *J Gen Physiol* 1963;47:347–369.
2. Berghmans S, Morris JP, Kanki JP, Look AT. Zebrafish sperm cryopreservation. *Methods Cell Biol* 2004;77:645–659.
3. Mazur P. Freezing of living cells: mechanisms and implications. *Am J Physiol* 1984;247:C125–C142.
4. Draper BW, Moens CB. A high-throughput method for zebrafish sperm cryopreservation and in vitro fertilization. *J Vis Exp* 2009;29:1395.
5. Carmichael C, Westerfield M, Varga ZM. Cryopreservation and in vitro fertilization at the zebrafish international resource center. *Methods Mol Biol* 2009;546:45–65.
6. Matthews JL, Murphy JM, Carmichael C, Yang H, Tiersch T, Westerfield M, *et al.* Changes to extender, cryoprotective medium, and in vitro fertilization improve zebrafish sperm cryopreservation. *Zebrafish* 2018;15:279–290.
7. Horváth Á, Bokor Z, Bernáth G, Csenki Z, Gorjan A, Herráez MP, *et al.* Very low sperm-egg ratios result in successful fertilization using cryopreserved sperm in the Adriatic grayling (*Thymallus thymallus*). *Aquaculture* 2015;435:75–77.
8. Westerfield M: The zebrafish book. A Guide for the Laboratory Use of Zebrafish (*Danio rerio*). Vol. 4th. University of Oregon, Eugene, 2000.
9. Bai C, Wang X, Lu G, Wei L, Liu K, Gao H, *et al.* Cooling rate optimization for zebrafish sperm cryopreservation using a cryomicroscope coupled with SYBR14/PI dual staining. *Cryobiology* 2013;67:117–123.
10. Yang H, Carmichael C, Varga ZM, Tiersch TR. Development of a simplified and standardized protocol with potential for high-throughput for sperm cryopreservation in zebrafish *Danio rerio*. *Theriogenology* 2007;68:128–136.
11. Morris JPt, Berghmans S, Zahrieh D, Neuberg DS, Kanki JP, Look AT. Zebrafish sperm cryopreservation with N,N-dimethylacetamide. *Biotechniques* 2003;35:956–958.
12. Draper BW, McCallum CM, Stout JL, Slade AJ, Moens CB. A high-throughput method for identifying N-ethyl-N-nitrosourea (ENU)-induced point mutations in zebrafish. *Methods Cell Biol* 2004;77:91–112.
13. Monteiro JF, Martins S, Farias M, Costa T, Certal AC. The impact of two different cold-extruded feeds and feeding regimens on zebrafish survival, growth and reproductive performance. *J Dev Biol* 2018;6:15.
14. Siccardi AJ, 3rd, Garriss HW, Jones WT, Moseley DB, D'Abramo LR, Watts SA. Growth and survival of zebrafish (*Danio rerio*) fed different commercial and laboratory diets. *Zebrafish* 2009;6:275–280.

Address correspondence to:

Ákos Horváth, PhD

Department of Aquaculture

Szent István University

Páter Károly u. 1.

Gödöllő H-2100

Hungary

E-mail: horvath.akos@mkk.szie.hu

Luciani R. Carvalho, MD, PhD

Discipline of Endocrinology

Department of Internal Medicine

Hospital das Clínicas HCFMUSP

Faculdade de Medicina

Universidade de São Paulo

Av Dr Enéas de Carvalho Aguiar, 155

São Paulo 05403-000

Brazil

E-mail: luciani.carvalho@hc.fm.usp.br