

Given a set of rings with radius R at varying z values, the B -field at z' on the z -axis due to a ring at z^* is

$$\int_{z^*}^{z'} \frac{\mu_0 I dl}{4\pi} \frac{R}{(z^2 + R^2)^{\frac{3}{2}}} = \frac{\mu_0}{4\pi} \frac{2\pi R^2 I}{(z^2 + R^2)^{\frac{3}{2}}} \Bigg|_{z^*}^{z'}$$

By the Biot-Savart law.

Thus for a solenoid with N loops, approximated as a series of current carrying loops where each loop is at some value z_i where $0 < i < N$, The net B -field at a point z will be

$$\sum_{i=0}^{i=N} \frac{\mu_0}{4\pi} \frac{2\pi R^2 I}{((z)^2 + R^2)^{3/2}} - \frac{\mu_0}{4\pi} \frac{2\pi R^2 I}{((z^*)^2 + R^2)^{3/2}} = B_z$$

A derivation of the Amperian model for forces between magnetic dipoles in non-uniform magnetic fields can be found here, which states:

$$F_z = \mu \frac{\delta B_z}{\delta z}$$

For point-like dipole. For a very thin, bar magnet with length L , the Force would be:

$$F_z = \mu \frac{\Delta B_z}{\Delta z} = \mu \frac{\Delta B_z}{L}$$

We can determine the overall change in kinetic energy, K , by numerically approximating the following. Note: z is the location of the end of the magnet.

$$\Delta K = -\Delta U = \int_{z_0}^z F_z dz = \int_{z_0}^z \mu \frac{B(z) - B(z - L)}{L}$$

And thus we can determine the speed at which we expect a dipole in our coil gun to travel from its starting position, z_0 to the point z .

To determine the moment, μ , of the magnet, we approximate the magnet as a series of looped coils with moving charge. Given a constant charge density, $J = dI/dx$, we can use the Biot-Savart law to derive that the magnetic field strength at the end of a magnet, B_{end} .

$$dI = J dx$$

$$B_{end} = \int_{x=0}^{x=L} \frac{\mu_0 R^2 J dx}{2(R^2 + x^2)^{3/2}} = \frac{\mu_0 J}{2} \frac{L}{\sqrt{L^2 + R^2}}$$

Solving for the charge density and the total current moving in the magnet gives

$$J = B_{end} \frac{\sqrt{L^2 + R^2}}{L} \frac{2}{\mu_0}$$

$$I = JL$$

$$I = B_{end} \frac{2\sqrt{L^2 + R^2}}{\mu_0}$$

Thus as $\boldsymbol{\mu} = AI_{total}$,

$$\boldsymbol{\mu} = \pi R^2 B_{end} \frac{2\sqrt{L^2 + R^2}}{\mu_0}$$

For a given magnet with a field strength B_{end} .