



PECOS

Predictive Engineering and Computational Sciences

Locally Conservative Discontinuous Petrov-Galerkin for Convection-Diffusion

Truman E. Ellis

Advisors: Leszek Demkowicz, Robert Moser

Collaborators: Jesse Chan, Nate Roberts

The University of Texas at Austin

March 8-9, 2013

A Summary of DPG

Overview of Features

- Robust for singularly perturbed problems
- Stable in the preasymptotic regime
- Designed for adaptive mesh refinement

DPG is a minimum residual method:

$$u_h = \arg \min_{w_h \in U_h} \frac{1}{2} \|Bw_h - l\|_{V'}^2$$
$$\Updownarrow$$

$$b(u_h, R_V^{-1} B \delta u_h) = l(R_V^{-1} B \delta u_h) \quad \forall \delta u_h \in U_h$$

where $v_{\delta u_h} := R_V^{-1} B \delta u_h$ are the **optimal test functions**.

DPG for Convection-Diffusion

Start with the strong-form PDE.

$$\nabla \cdot (\beta u) - \epsilon \Delta u = g$$

Rewrite as a system of first-order equations.

$$\nabla \cdot (\beta u - \sigma) = g$$

$$\frac{1}{\epsilon} \sigma - \nabla u = \mathbf{0}$$

Multiply by test functions and integrate by parts over each element, K .

$$-(\beta u - \sigma, \nabla v)_K + ((\beta u - \sigma) \cdot \mathbf{n}, v)_{\partial K} = (g, v)_K$$

$$\frac{1}{\epsilon} (\sigma, \tau)_K + (u, \nabla \cdot \tau)_K - (u, \tau_n)_{\partial K} = 0$$

Use the ultraweak (DPG) formulation to obtain bilinear form $b(u, v) = l(v)$.

$$\begin{aligned} & -(\beta u - \sigma, \nabla v)_K + (\hat{f}, v)_{\partial K} + \frac{1}{\epsilon} (\sigma, \tau)_K \\ & + (u, \nabla \cdot \tau)_K - (\hat{u}, \tau_n)_{\partial K} = (g, v)_K \end{aligned}$$

Local Conservation

The local conservation law in convection diffusion is

$$\int_{\partial K} \hat{f} = \int_K g,$$

which is equivalent to having $\mathbf{v}_K := \{v, \boldsymbol{\tau}\} = \{1_K, \mathbf{0}\}$ in the test space. In general, this is not satisfied by the optimal test functions. Following Moro et al¹, we can enforce this condition with Lagrange multipliers:

$$L(u_h, \boldsymbol{\lambda}) = \frac{1}{2} \|R_V^{-1}(Bu_h - l)\|_V^2 - \sum_K \lambda_K \underbrace{\langle Bu_h - l, \mathbf{v}_K \rangle}_{\langle \hat{f}, 1_K \rangle_{\partial K} - \langle g, 1_K \rangle_K},$$

where $\boldsymbol{\lambda} = \{\lambda_1, \dots, \lambda_N\}$.

¹D. Moro, N.C. Nguyen, and J. Peraire. A hybridized discontinuous Petrov-Galerkin scheme for scalar conservation laws. *Int.J. Num. Meth. Eng.*, 2011. in print

Local Conservation

Finding the critical points of $L(u, \lambda)$, we get the following equations.

$$\begin{aligned} \frac{\partial L(u_h, \lambda)}{\partial u_h} &= b(u_h, R_V^{-1} B \delta u_h) - l(R_V^{-1} B \delta u_h) \\ &\quad - \sum_K \lambda_K b(\delta u_h, \mathbf{v}_K) = 0 \quad \forall \delta u_h \in U_h \end{aligned}$$

$$\frac{\partial L(u_h, \lambda)}{\partial \lambda_K} = -b(u_h, \mathbf{v}_K) + l(\mathbf{v}_K) = 0 \quad \forall K$$

A few consequences:

- We've turned our minimization problem into a saddlepoint problem.
- Only need to find the optimal test function in the orthogonal complement of constants.

Optimal Test Functions

For each $\mathbf{u} = \{u, \boldsymbol{\sigma}, \hat{u}, \hat{f}\} \in \mathbf{U}_h$, find $\mathbf{v}_\mathbf{u} = \{v_\mathbf{u}, \boldsymbol{\tau}_\mathbf{u}\} \in \mathbf{V}$ such that

$$(\mathbf{v}_\mathbf{u}, \mathbf{w})_\mathbf{V} = b(\mathbf{u}, \mathbf{w}) \quad \forall \mathbf{w} \in \mathbf{V}$$

where \mathbf{V} becomes $\mathbf{V}_{p+\Delta p}$ in order to make this computationally tractable. Chan et al² developed the following robust norm for convection-diffusion.

$$\begin{aligned} \|(v, \boldsymbol{\tau})\|_{\mathbf{V}, \Omega_h}^2 &= \|\nabla \cdot \boldsymbol{\tau}\|^2 + \left\| \min \left\{ \frac{1}{\sqrt{\epsilon}}, \frac{1}{\sqrt{|K|}} \right\} \boldsymbol{\tau} \right\|^2 \\ &\quad + \epsilon \|\nabla v\|^2 + \|\boldsymbol{\beta} \cdot \nabla v\|^2 \quad \underbrace{+ \left\| \min \left\{ \sqrt{\frac{\epsilon}{|K|}}, 1 \right\} v \right\|^2}_{\text{no longer necessary to make this a norm}} \end{aligned}$$

²J. Chan, N. Heuer, T Bui-Thanh, and L. Demkowicz. Robust DPG method for convection-dominated diffusion problems ii: a natural inflow condition. Technical Report 21, ICES, 2012

Optimal Test Functions

For each $\mathbf{u} = \{u, \boldsymbol{\sigma}, \hat{u}, \hat{f}\} \in \mathbf{U}_h$, find $\mathbf{v}_{\mathbf{u}} = \{v_{\mathbf{u}}, \boldsymbol{\tau}_{\mathbf{u}}\} \in \mathbf{V}$ such that

$$(\mathbf{v}_{\mathbf{u}}, \mathbf{w})_{\mathbf{V}} = b(\mathbf{u}, \mathbf{w}) \quad \forall \mathbf{w} \in \mathbf{V}$$


where \mathbf{V} becomes $\mathbf{V}_{p+\Delta p}$ in order to make this computationally tractable. Chan et al² developed the following robust norm for convection-diffusion.

$$\begin{aligned} \|(v, \boldsymbol{\tau})\|_{\mathbf{V}, \Omega_h}^2 &= \|\nabla \cdot \boldsymbol{\tau}\|^2 + \left\| \min \left\{ \frac{1}{\sqrt{\epsilon}}, \frac{1}{\sqrt{|K|}} \right\} \boldsymbol{\tau} \right\|^2 \\ &\quad + \epsilon \|\nabla v\|^2 + \|\boldsymbol{\beta} \cdot \nabla v\|^2 + \underbrace{\left(\int_K v \right)^2}_{\text{zero mean term}} \end{aligned}$$


²J. Chan, N. Heuer, T Bui-Thanh, and L. Demkowicz. Robust DPG method for convection-dominated diffusion problems ii: a natural inflow condition. Technical Report 21, ICES, 2012

Vortex Problem

Does enforcing local conservation make a difference?




With Local Conservation




Without Local Conservation

Vortex Problem

Does enforcing local conservation make a difference?




With Local Conservation




Without Local Conservation

Vortex Problem

Does enforcing local conservation make a difference?




With Local Conservation




Without Local Conservation

Vortex Problem

Does enforcing local conservation make a difference?




With Local Conservation




Without Local Conservation

Vortex Problem

Does enforcing local conservation make a difference?






With Local Conservation




Without Local Conservation

Hemker Problem






With Local Conservation




Without Local Conservation

Hemker Problem






With Local Conservation




Without Local Conservation

Hemker Problem






With Local Conservation




Without Local Conservation

Hemker Problem






With Local Conservation




Without Local Conservation

Hemker Problem





With Local Conservation





Without Local Conservation

High Lift Airfoil





With Local Conservation





Without Local Conservation

High Lift Airfoil





With Local Conservation





Without Local Conservation

High Lift Airfoil





With Local Conservation





Without Local Conservation

High Lift Airfoil





With Local Conservation





Without Local Conservation

High Lift Airfoil





With Local Conservation





Without Local Conservation

High Lift Airfoil





With Local Conservation





Without Local Conservation

High Lift Airfoil




With Local Conservation




Without Local Conservation

Double Glazing Problem




With Local Conservation




Without Local Conservation

Double Glazing Problem




With Local Conservation




Without Local Conservation

Double Glazing Problem




With Local Conservation




Without Local Conservation

Double Glazing Problem




With Local Conservation




Without Local Conservation

Double Glazing Problem




With Local Conservation




Without Local Conservation

Double Glazing Problem




With Local Conservation




Without Local Conservation

Double Glazing Problem




With Local Conservation




Without Local Conservation

Double Glazing Problem




With Local Conservation




Without Local Conservation

Double Glazing Problem




With Local Conservation




Without Local Conservation

Inviscid Burgers' Equation

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = 0 \quad \Leftrightarrow \quad \nabla_{x,t} \cdot \begin{pmatrix} \frac{u^2}{2} \\ u \end{pmatrix} = 0$$




With Local Conservation




Without Local Conservation

Inviscid Burgers' Equation

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = 0 \quad \Leftrightarrow \quad \nabla_{x,t} \cdot \begin{pmatrix} \frac{u^2}{2} \\ u \end{pmatrix} = 0$$




With Local Conservation




Without Local Conservation

Inviscid Burgers' Equation

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = 0 \quad \Leftrightarrow \quad \nabla_{x,t} \cdot \begin{pmatrix} \frac{u^2}{2} \\ u \end{pmatrix} = 0$$




With Local Conservation




Without Local Conservation

Inviscid Burgers' Equation

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = 0 \quad \Leftrightarrow \quad \nabla_{x,t} \cdot \begin{pmatrix} \frac{u^2}{2} \\ u \end{pmatrix} = 0$$




With Local Conservation




Without Local Conservation

Inviscid Burgers' Equation

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = 0 \quad \Leftrightarrow \quad \nabla_{x,t} \cdot \begin{pmatrix} \frac{u^2}{2} \\ u \end{pmatrix} = 0$$




With Local Conservation




Without Local Conservation

Inviscid Burgers' Equation

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = 0 \quad \Leftrightarrow \quad \nabla_{x,t} \cdot \begin{pmatrix} \frac{u^2}{2} \\ u \end{pmatrix} = 0$$




With Local Conservation




Without Local Conservation

Inviscid Burgers' Equation

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = 0 \quad \Leftrightarrow \quad \nabla_{x,t} \cdot \begin{pmatrix} \frac{u^2}{2} \\ u \end{pmatrix} = 0$$




With Local Conservation




Without Local Conservation

Inviscid Burgers' Equation

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = 0 \quad \Leftrightarrow \quad \nabla_{x,t} \cdot \begin{pmatrix} \frac{u^2}{2} \\ u \end{pmatrix} = 0$$




With Local Conservation




Without Local Conservation

Inviscid Burgers' Equation

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = 0 \quad \Leftrightarrow \quad \nabla_{x,t} \cdot \begin{pmatrix} \frac{u^2}{2} \\ u \end{pmatrix} = 0$$



With Local Conservation



Without Local Conservation

Summary

What have we done?

- We've turned our minimization problem into a saddlepoint problem.
- The change is computationally feasible.
- Mathematically, it gets rid of troublesome term.

Does it make a difference?

- Enforcement changes refinement strategy.
- Standard DPG is nearly conservative in practice.

	Local Flux Imbalance		Global Flux Imbalance	
	Before	After	Before	After
Vortex	10^{-8}	10^{-15}	10^{-5}	10^{-16}
Hemker	10^{-6}	10^{-14}	10^{-4}	10^{-14}
High Lift Airfoil	10^{-7}	10^{-14}	10^{-5}	10^{-14}
Double Glazing	10^{-8}	10^{-15}	10^{-9}	10^{-15}
Inviscid Burgers	10^{-7}	10^{-17}	10^{-6}	10^{-16}

Summary

What have we done?

- We've turned our minimization problem into a saddlepoint problem.
- The change is computationally feasible.
- Mathematically, it gets rid of troublesome term.


Does it make a difference?

- Enforcement changes refinement strategy.
- Standard DPG is nearly conservative in practice.

	Local Flux Imbalance		Global Flux Imbalance	
	Before	After	Before	After
Vortex	10^{-8}	10^{-15}	10^{-5}	10^{-16}
Hemker	10^{-6}	10^{-14}	10^{-4}	10^{-14}
High Lift Airfoil	10^{-7}	10^{-14}	10^{-5}	10^{-14}
Double Glazing	10^{-8}	10^{-15}	10^{-9}	10^{-15}
Inviscid Burgers	10^{-7}	10^{-17}	10^{-6}	10^{-16}

We need to study the effect on real fluid dynamics.

Questions?





J. Chan, N. Heuer, T Bui-Thanh, and L. Demkowicz.

Robust DPG method for convection-dominated diffusion problems ii: a natural inflow condition.
Technical Report 21, ICES, 2012.



D. Moro, N.C. Nguyen, and J. Peraire.

A hybridized discontinuous Petrov-Galerkin scheme for scalar conservation laws.

Int.J. Num. Meth. Eng., 2011.

in print.