

Atrial Fibrillation Detection Fishing for An Irregular Heartbeat Before and After Stroke

Vincent Thijs, MD, PhD

See related articles, pp 2653, 2654, 2660, 2665

Atrial fibrillation (AF) is present in $\pm 3\%$ of the general population above age 20, and its prevalence increases substantially in the ≥ 65 -year-olds age group.¹ It is expected that AF prevalence will increase as populations get older. AF is asymptomatic in $\leq 40\%$ of patients. Unfortunately, the absence of symptoms does not suggest a benign course.² The Framingham Heart Study found that stroke is the first manifestation of AF in at least 2% to 5% of AF patients.³ In a hospital-based series, $\leq 20\%$ of stroke patients had previously unidentified or unrecognized AF.⁴ AF also increases the risk of cognitive impairment and dementia.⁵ Identifying AF and reducing stroke risk in patients with AF before stroke occurs is, therefore, an important goal.

Should We Detect AF in the General Population Prior to Stroke?

Whether screening for AF in the general population is warranted is heavily debated. Opportunistic screening refers to screening offered to people as part of a routine medical checkup or when examined for another reason, whereas systematic screening entails screening in the general population (ie, mass screening) or a high-risk, target group (ie, patients with heart failure or diabetes mellitus). Opportunistic screening in patients aged >65 years with pulse palpation or ECG rhythm strip is the current recommended method by the European Society of Cardiology.⁶ No recommendation is given in the US guidelines.⁷ Pulse palpation is sensitive for detection of permanent AF (the sensitivity is 94%, and its specificity is 72%) but is not specific and requires confirmation with ECG. The equipment and time required to perform an ECG examination are a significant barrier to perform screening, but fortunately new methods for screening are being developed rapidly.⁸ Proponents of screening cite the relatively high frequency of AF, the ease of detection and prevention of stroke with AF, and the high risk of stroke in patients with undetected AF as the main reasons for advocating systematic screening in the >65 years age group. The yield of single time-point screening for unknown AF is 1% (1.4% in those >65 years) and depends on the age and

ethnicity and risk factor profile of the target population.⁹ Only a minority of identified AF patients report symptoms. A significant proportion of patients with known AF who are undertreated are also identified.¹⁰ The concerns with screening are the optimal method, frequency and setting of screenings, the lack of randomized evidence of whether screening will lead to measurable reductions in stroke incidence, and the questionable cost-efficacy of screening approaches. One randomized trial found no clear evidence of benefit of mass screening, but found opportunistic screening with pulse palpation, and found confirmation with 12-lead ECG to be probably cost-effective and superior to both mass screening and targeted screening.¹¹ However, this trial was performed in the warfarin era before non-vitamin K antagonists with an improved safety profile and reduced patient burden were available. Only half of the patients identified through mass screening attended a follow-up confirmation ECG.

Devices Used in Screening Programs

Technological advances have made screening without standard 12-lead ECG machines possible, removing at least one barrier to screening in high-risk populations. Low-cost devices attached to cell phones permitting single-lead ECG or dedicated screening devices are now available, which permit easy diagnosis of AF on the spot or allow wireless transmission to a cardiologist or technician for confirmation.^{12–14} These devices provide single-lead ECGs of diagnostic quality. Studies comparing these devices head to head with standard 12-lead ECG report sensitivities of 90% to 100% and specificities between 90% and 97%.⁸ Indirect methods include blood pressure measurement devices which permit simultaneous screening for AF and high BP and devices that analyze plethysmographic waveforms of cardiac activity attached to a smartphone camera with flash or a pulse oximeter.^{15,16} False positives are possible with these indirect methods, and subsequent confirmation with ECG is still required.

These methods allow screening in primary-care doctor's offices, pharmacies, or other settings where screenings are performed, for example, during influenza vaccinations. The costs are also substantially less.

Received May 29, 2017; final revision received July 14, 2017; accepted July 18, 2017.

From the Stroke Division, Florey Institute of Neuroscience and Mental Health, University of Melbourne, and Department of Neurology, Austin Health, Heidelberg, Victoria, Australia.

Correspondence to Vincent Thijs, MD, PhD, Stroke Division, Florey Institute of Neuroscience and Mental Health, University of Melbourne, and Department of Neurology, Austin Health, 245 Burgundy St, Heidelberg, Victoria 3084, Australia. E-mail vincent.thijs@florey.edu.au

(*Stroke*. 2017;48:2671–2677. DOI: 10.1161/STROKEAHA.117.017083.)

© 2017 American Heart Association, Inc.

Stroke is available at <http://stroke.ahajournals.org>

DOI: 10.1161/STROKEAHA.117.017083

Follow-Up After Screening

For screening programs to be effective, adequate follow-up and institution of appropriate anticoagulant treatment is required. Most patients identified through screening seem to have at least a moderate to high risk of stroke and require treatment with an oral anticoagulant based on risk stratification according to guidelines. A systematic review found that $\approx 2/3$ of identified cases were at high risk of stroke and would be eligible for oral anticoagulation.^{9,17}

Single Time-Point Screening Versus Repeated Screening

As AF is not a permanent arrhythmia, one-time screening may miss nonpermanent forms of AF, especially paroxysmal AF. In a screening program of 75/76-year-olds in Sweden, involving twice daily screening over a period of 2 weeks, new AF was identified in 3% of 7173 respondents.¹⁸ Surprisingly, AF was identified only in 17% on the first ECG, strongly suggesting that repeated screening is important to identify more AF. The yield of repeated screening increased to a 7% detection rate when analysis was restricted to patients with at least one stroke risk factor. Further screening programs, combining biomarkers and repeated, intermittent screening, are planned.¹⁹

Relevance of AF Identified Through Screening in Primary Prevention

The relevance of screening-detected AF has been questioned. The natural history of incidentally detected AF through screening has indeed not been studied in detail nor have randomized clinical trials been performed that found a decrease in vascular end points with widespread screening. However, the evidence from studies comparing asymptomatic AF with symptomatic AF does not show a lower risk of cardiovascular outcomes with asymptomatic status, and some studies even reported a worse prognosis.²⁰ There is growing consensus that screening-detected AF in primary prevention is associated with a high risk of stroke, and anticoagulation should be prescribed according to guidelines.⁸

Relevance of AF Identified in Patients With Pacemakers or Implantable Cardioverter-Defibrillator

Traditionally, the risk of stroke was thought to be similar in permanent and nonpermanent forms of AF. Therefore, none of the risk stratification schemes have incorporated the type of AF, and none of the guidelines have adopted a different stance toward recommending anticoagulation in patients with paroxysmal AF. A recent systematic review, however, found that nonpermanent forms of AF may have a 25% lower risk of stroke compared with permanent forms of AF.²¹ Even though the risk of stroke may be reduced, post hoc analyses from the NOAC studies (Non-Vitamin K Oral Anticoagulant) report a similar efficacy and safety compared with warfarin in these patients.²²

The relevance of AF is more contested when short episodes are identified, especially in patients who undergo continuous monitoring with pacemakers or implantable cardioverter-defibrillators.²³ These so-called atrial high-rate episodes may be because of various causes and require review before

being labeled AF. They may occur in $\leq 30\%$ to 60% of monitored patients. There is ongoing debate about the duration of AF required on these devices to increase the risk of stroke independently from the general cardiovascular risk profile. Reported studies cite an increase in risk from anywhere between 5 minutes and 24 hours. Moreover, it is unclear what the relationship is between device-detected AF and the type of AF that was studied in the trials that established superiority of anticoagulants over aspirin and placebo. In the clinical trials, for patients with paroxysmal AF to be enrolled, 2 episodes of AF lasting longer than 30 seconds had to be present on separate occasions, with at least one documented on ECG.

In ASSERT (Asymptomatic Atrial Fibrillation and Stroke Evaluation in Pacemaker Patients and the AF Reduction Atrial Pacing trial), a study that focused on patients without history of AF, the risk of stroke associated with episodes of device-detected atrial high-rate episodes was analyzed in 2580 patients. Episodes longer than 6 minutes were associated with a clinical diagnosis of AF and an increased risk of stroke or systemic embolism.²⁴ A post hoc analysis suggests that only episodes lasting longer than 24 hours are relevant and carry a 3-fold risk of future thromboembolism over the next 2.5 years.²⁵ It is probable that the length of the episode is not the only determinant of risk, and risk scores like the CHA₂D₂-VASC score or echocardiographic features or the burden of AF may have to be taken into account.

The relevance of these findings for patients without preexistent heart disease who are specifically monitored for subclinical AF is unclear. Three prospective studies have assessed the presence of AF in patients at high risk of stroke using an insertable cardiac monitor (ICM). The ASSERT-II study assessed the risk of AF using ICM in 273 high-risk patients with a CHA₂D₂-VASC score of ≥ 3 , large left atria, elevated serum N-terminal pro-brain-type natriuretic peptide levels, or obstructive sleep apnea. Episodes lasting ≥ 5 minutes occurred in 34.4% of patients per year, with a higher risk in older patients and patients with heart failure. Episodes of subclinical AF were not related to stroke occurrence. In the PREDATE AF Study (Predicting Determinants of Atrial Fibrillation or Flutter for Therapy Elucidation in Patients at Risk for Thromboembolic Events), 22.4% of 245 patients with a CHA₂D₂-VASC score of ≥ 2 were found to have AF episodes of >6 minutes on ICM during an average follow-up of 1.2 years.²⁶ In another study with similar inclusion criteria, the rate of AF was 29.3% at 18 months in 394 patients with ICM.²⁷ In none of these studies, an association was found with a prior history of stroke or transient ischemic attack (TIA), but details about the types of stroke patients included in these studies are scant and follow-up probably too short to reveal an association with stroke.

Whether anticoagulation is warranted in patients with device-detected AF is being further explored in the Apixaban for the Reduction of Thrombo-Embolism in Patients With Device-Detected Sub-Clinical Atrial Fibrillation study and the Non-vitamin K Antagonist Oral Anticoagulants in Patients With Atrial High Rate Episodes study.^{28,29}

Other High-Risk Groups for AF

AF is often diagnosed in the context of systemic conditions like pneumonia and thyroid disease or after cardiothoracic surgery.³⁰

In these patients, anticoagulation is not typically considered because this type of AF was typically considered transient and benign. Recent data suggest that these patients are at high risk of recurrent AF and stroke.^{30,31} In one study, the risk of developing AF was 6- to 8-fold with postoperative AF.³² Whether patients with secondary AF will benefit from early anticoagulation or should undergo repeated screening for AF is currently unknown.³³

AF Detection After TIA or Stroke?

The detection of AF after stroke is important because it provides clues to the mechanism of stroke and generally leads to a change in antithrombotic strategy from antiplatelets to anticoagulation. It is estimated that $\approx 25\%$ of patients with stroke or TIA will be diagnosed with AF if systematic, long-term screening is performed.³⁴ After TIA or stroke, most guidelines, therefore, recommend screening patients for the presence of AF, with 12-lead ECG, Holter monitoring, telemetry, or monitoring devices, but the exact timing and duration of screening with these techniques are undefined. Table 1 describes the available methods for prolonged screening.

In the 2008 European Stroke Organisation guidelines, the recommended duration is a 12-lead ECG, continuous ECG monitoring for an undefined duration, and 24-hour Holter monitoring if no cause of stroke can be found.⁴¹ In the 2013 and 2014 American Heart Association/American Stroke Association guidelines, cardiac monitoring is recommended for 24 hours, and prolonged rhythm monitoring for 30 days is considered reasonable in patients without an apparent cause.^{42,43} According to the 2016 guidelines for the management of AF from the European Society of Cardiology more prolonged monitoring is warranted with 72-hour monitoring for all stroke patients and consideration should be made of long-term recordings in some patients.⁶

Four randomized trials have been performed that examined the yield of more intensive monitoring strategies after TIA or stroke. These trials generally show that more intensive monitoring detects more AF, but were underpowered to show a reduction in TIA, stroke, or systemic emboli. In a small trial ($n=100$), a 7-day continuous event-monitoring device detected AF in an additional 16% of patients compared with standard practice.⁴⁴ No difference in vascular events was identified (4 events in both study arms). In the EMBRACE trial (Event Monitor Belt for Recording Atrial Fibrillation After a Cerebral Ischemic Event),

a 30-day external cardiac monitoring device was compared with an extra Holter testing in patients with stroke of unknown cause. AF was detected in 16.1% compared with 3.2% of patients in the control group. The vascular event rate was not reported.⁴⁵ In CRYSTAL AF (Cryptogenic Stroke and Underlying AF trial), an ICM detected AF at 6 months in 8.9% versus 1.4% of patients in the control group in patients who underwent extensive diagnostic testing prior to inclusion.⁴⁶ The control group underwent 6 monthly clinical follow-up that consisted of probing for symptoms of AF and ECG or Holter as per the investigator's discretion. A nonsignificant TIA or stroke event reduction was found at 1 year (5.2% in the ICM arm versus 8.2% in the control group). The FIND AF study randomized 398 patients with recent (<7 days) TIA or stroke who did not have carotid or intracranial artery stenosis to a strategy of enhanced monitoring or standard of care.⁴⁷ The enhanced monitoring strategy consisted of repeated 10-day monitoring at 3 months' intervals for 1 year. At 6 months, the intensive monitoring strategy detected AF in 14% of patients compared with 5% in the control group. At 1 year, the stroke rate was 3.7% with enhanced monitoring compared with 5.4% with standard monitoring.

Who to Screen After TIA or Stroke?

Indirect clues of the presence of AF can often be obtained from clinical information, biomarkers, or echocardiographic findings. These are summarized in Table 2.

No clear algorithm has been developed that reliably identifies patients in whom prolonged monitoring is required. Prolonged screening will lead to higher detection rates in patients with cryptogenic stroke or embolic stroke of unknown source, but the exact yield in other subtypes of stroke is not yet clear.⁶⁷ Patients with large vessel atherosclerotic disease and small vessel disease often have risk factors that contribute to AF development and atherosclerosis, and AF often coexists. A systematic review identified a rate of detection of 2.4% in small vessel disease and 2.2% with large vessel disease. This was generally lower than the 9.2% rate in studies that did not define subtype of the stroke.⁶⁸ No studies were found that performed longer monitoring than 7 days in the small/large vessel disease subtypes, except FIND AF, which did not find a difference in the rate of detection between cryptogenic stroke and noncryptogenic stroke (excluding large vessel disease). The yield

Table 1. Overview of Noninvasive Monitoring Techniques and Insertable Cardiac Monitoring Devices

Methods for Detecting Paroxysmal AF
Noninvasive screening for AF over the medium term is possible with traditional Holter monitors, wearable patches, or belts. ^{35–37} These provide recording periods of ≤ 2 –4 wk. Real-time transmission and analysis is possible with the most sophisticated devices. Holters and some patches store the ECG information and only provide diagnosis in retrospect. Activity monitors (eg, Fitbit devices, smartwatches) used to track sports activities or walking also record heart rates, and are promising for detecting AF when supplemented with deep learning algorithms, but do not produce diagnostic ECG and when abnormalities are found require further confirmation with ECG. The indirect information provided by these devices, for example, sudden increases in heart rates, may, however, provide clues to the presence of AF. Other issues with many of the wearable, consumer-oriented technologies include the need for validation, the cost-effectiveness, data security, and privacy. Yet it is hoped that in the foreseeable future, relatively cheap wearables will become available that reliably track AF.
Insertable cardiac monitoring devices are slightly more invasive devices that store abnormal rhythm episodes in memory, which can be remotely analyzed after wireless transmission. ^{38–40} Confirmation of the nature of the episodes by a specialist is still required. The durability of these devices is restricted by battery life, which is generally ≤ 3 y. The devices are MRI compatible. Dependent on the device's algorithm and settings, the sensitivity and specificity compared with Holter is high. ³⁹ Insertable cardiac monitoring devices cannot detect all types of atrial flutter and may miss brief episodes of AF.

MRI indicates magnetic resonance imaging.

Table 2. Clues From ECG, Echocardiography, and Blood Biomarkers That Suggest a Higher Risk of AF

Characteristic	Method of Detection	Threshold or Value	Comment	Reference
P-wave dispersion	ECG	≥40 ms	Defined as the difference between the longest and shortest P-wave duration from different surface ECG leads; reflects inhomogeneous intraatrial conduction	Dilaveris and Gialafos ⁴⁸
P-wave duration	ECG	Very long P-wave duration in the top fifth percentile associated with doubling of risk of AF		Guidera and Steinberg ⁴⁹ ; Magnani et al ⁵⁰
PR interval	ECG	Each 20 ms increases risk of AF with 20% in primary population	In cryptogenic stroke, 30% increase with 10 ms increase Influenced by cardiac and antihypertensive medication	Thijs et al ⁵¹ ; Cheng et al ⁵²
R-R variability analysis	Continuous cardiac monitoring	Requires special commercial software to postprocess continuous cardiac monitoring devices	Further studies are needed to assess yield of this method in patients with large R-R variability without AF	
Atrial premature beats (ABP)	Holter	Low (7%–9%) probability of AF with <100 ABP/24 h and high probability (>37%) with ≥1000 ABP/24 h	Validation of cutoff needed in separate population	Gladstone et al ⁵⁴
Supraventricular ectopic activity	Holter	Thirty supraventricular ectopic complexes per hour/any episode of run of ≥20 SV complexes	Increases risk of clinical AF diagnosis and AF in primary prevention population over 6 y	Binici et al ⁵⁵
Left atrial volume	Echocardiography	Risk of score increases 2.4× per 10 mm increase in LA volume	Echocardiographic features are not independent predictors in AF risk score after correction for other risk factors ⁵⁶	Benjamin et al ⁵⁷
Left atrial appendage velocity and spontaneous echo contrast	Echocardiography	Low velocities found in patients with paroxysmal AF		Kim et al ⁵⁸ ; Patti et al ⁵⁹
NT-proBNP	Laboratory	Variable	Promising factor, but variable cutoffs	Hijazi et al ⁶⁰ ; Patton et al ⁶¹ ; Shibasaki et al ⁶² ; Samuel et al ⁶³ ; Rodríguez-Yáñez et al ⁶⁴
TSH	Laboratory	30% increase of risk of AF in subclinical hyperthyroidism; high normal thyroid function 12% increase in risk of AF		Selmer et al ⁶⁵
Single nucleotide polymorphisms	Genetic testing	Increased burden of AF predisposing genes	Untested strategy	Lubitz et al ⁶⁶

AF indicates atrial fibrillation; NT-proBNP, N-terminal pro-B-type natriuretic peptide; and TSH, thyroid stimulating hormone.

of AF detection in other stroke subtypes is further investigated in the STROKE-AF study (Stroke of Known Cause and Underlying AF), where 500 patients with stroke associated with large or small vessel disease will be randomized to an ICM or standard of care.⁶⁹ Similarly, TIA patients are often less intensively examined and monitored, and in some settings, even no cardiac monitoring is performed as patients are discharged home. A systematic review of the limited number of studies specifically assessing the yield of screening in TIA found a rate of detection of TIA in 4%.⁷⁰ Detection rates were higher with longer monitoring and in patients with unexplained TIA.

The yield of monitoring is probably the highest early after the cerebral ischemic event, but late detection of AF is also common. The earliest opportunity for detection is in the pre-hospital setting; however, most episodes will be detected during the in-hospital stay.⁷¹

At the time of stroke unit admission or in intensive care unit, telemetry or continuous electrocardiographic monitoring is performed.^{72,73} Cardiac monitoring devices require special software to detect and record AF. In the absence of such dedicated software, irregular heart rhythms must be detected and recorded by stroke unit nurses. Post hoc analysis of recordings of telemetry devices is possible using RR interval analysis performed off-line. This approach has been shown to increase the detection rate compared with standard stroke unit care.⁷² In the FIND AF study, 2/3 of the AF events were detected with the first 10-day Holter monitor. In EMBRACE, half of the AF events were detected in the first week of monitoring, but patients were enrolled 2.5 months after the ischemic event. Similarly, in CRYSTAL AF, half of the events detected within 6 months occurred within 42 days of monitoring. This suggests that the yield of AF detection is highest in the first weeks to months after the cerebral ischemic event. Nevertheless,

30-day monitoring would miss a relevant number of AF episodes. For instance, in CRYSTAL AF, AF was detected by 3 years in $\leq 30\%$ of these patients (compared with 8.9% at 6 months).⁴⁶ Simulations from CRYSTAL AF suggest that any intermittent method of screening AF, be it 30-day monitoring or repeating annual monitoring, would have a limited sensitivity of only $\leq 22.8\%$ and a negative predictive value of 85.7%.⁷⁴

After stroke or TIA, it seems that the highest yield for detecting AF is early after symptom onset, in patients with ischemic stroke (as opposed to TIA), and in patients with cryptogenic stroke. However, significant gaps remain, in our knowledge, on the yield of screening after TIA and stroke, and none of the current screening strategies are fully satisfactory. A commonly performed strategy consists of of stepwise screening with ECG, continuous inpatient electrocardiographic monitoring followed by prolonged monitoring for up to 30 days, after which ICM is considered in patients with unexplained stroke.⁵⁴ Further selection to proceed with AF may be based on indirect markers (see Table 2). Sequential testing and analysis are, however, cumbersome and require multiple visits and close collaboration with cardiologists. Forgoing the prolonged monitoring and immediate ICM use at discharge is a sensitive strategy, but its cost-efficacy and budget impact remain to be determined in various healthcare settings.⁷⁵ The anticoagulation trials in patients with embolic stroke of unknown source will also shed new light on the importance of prolonged screening after stroke and will have to be compared with screening strategies.

Conclusions

AF detection is important in primary prevention and secondary prevention after stroke. A plethora of screening devices is now available that allow reliable detection of AF, but a cheap, noninvasive way of detecting AF over extended periods of time is still missing. The implications of AF that is detected with the new technologies on anticoagulant management are not always clear. Screening for AF in selected populations in primary prevention and prolonged screening after cryptogenic stroke is recommended.

Disclosures

Dr Thijs reports receiving consulting fees from Medtronic. Dr Thijs is on the steering committee of the DIAGNOSE-AF (Detection in Transient Ischemic Attack Patients Using Continuous monitoring vs Standard Evaluation for Atrial Fibrillation) and REACT AF (Rhythm Evaluation for Anticoagulation With Continuous Monitoring of Atrial Fibrillation Clinical Trial) clinical trials sponsored by Medtronic. Dr Thijs reports speaker and consulting fees and travel support by Boehringer Ingelheim, Pfizer/BMS, Daichi Sankyo, and Bayer.

References

- Rahman F, Kwan GF, Benjamin EJ. Global epidemiology of atrial fibrillation. *Nat Rev Cardiol*. 2014;11:639–654. doi: 10.1038/nrcardio.2014.118.
- Flaker GC, Belew K, Beckman K, Vidaillet H, Kron J, Safford R, et al; AFFIRM Investigators. Asymptomatic atrial fibrillation: demographic features and prognostic information from the Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM) study. *Am Heart J*. 2005;149:657–663. doi: 10.1016/j.ahj.2004.06.032.
- Lubitz SA, Yin X, McManus DD, Weng L-C, Aparicio HJ, Walkey AJ, et al. Stroke as the initial manifestation of atrial fibrillation: The

- Framingham Heart Study. *Stroke*. 2017;48:490–492. doi: 10.1161/STROKEAHA.116.015071.
- Borowsky LH, Regan S, Chang Y, Ayres A, Greenberg SM, Singer DE. First diagnosis of atrial fibrillation at the time of stroke. *Cerebrovasc Dis*. 2017;43:192–199. doi: 10.1159/000457809.
- Singh-Manoux A, Fayosse A, Sabia S, Canonic M, Bobak M, Elbaz A, et al. Atrial fibrillation as a risk factor for cognitive decline and dementia. *Eur Heart J*. 2017. doi: 10.1093/eurheartj/ehx208.
- Kirchhof P, Benussi S, Kotecha D, Ahlsson A, Atar D, Casadei B, et al. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Eur Heart J*. 2016;37:2893–2962. doi: 10.1093/eurheartj/ehw210.
- January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC Jr, et al; ACC/AHA Task Force Members. 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines and the Heart Rhythm Society. *Circulation*. 2014;130:e199–e267. doi: 10.1161/CIR.0000000000000041.
- Freedman B, Camm J, Calkins H, Healey JS, Rosenqvist M, Wang J, et al; AF-Screen Collaborators. Screening for atrial fibrillation: a Report of the AF-SCREEN International Collaboration. *Circulation*. 2017;135:1851–1867. doi: 10.1161/CIRCULATIONAHA.116.026693.
- Lowres N, Neubeck L, Redfern J, Freedman SB. Screening to identify unknown atrial fibrillation. A systematic review. *Thromb Haemost*. 2013;110:213–222. doi: 10.1160/TH13-02-0165.
- Engdahl J, Andersson L, Mirskaya M, Rosenqvist M. Stepwise screening of atrial fibrillation in a 75-year-old population: implications for stroke prevention. *Circulation*. 2013;127:930–937. doi: 10.1161/CIRCULATIONAHA.112.126656.
- Hobbs FD, Fitzmaurice DA, Mant J, Murray E, Jowett S, Bryan S, et al. A randomised controlled trial and cost-effectiveness study of systematic screening (targeted and total population screening) versus routine practice for the detection of atrial fibrillation in people aged 65 and over. The SAFE study. *Health Technol Assess*. 2005;9:iii–iv, ix.
- Vaes B, Stalpaert S, Tavernier K, Thaelts D, Lapeire D, Mullens W, et al. The diagnostic accuracy of the MyDiagnostick to detect atrial fibrillation in primary care. *BMC Fam Pract*. 2014;15:113. doi: 10.1186/1471-2296-15-113.
- Lau JK, Lowres N, Neubeck L, Brieger DB, Sy RW, Galloway CD, et al. iPhone ECG application for community screening to detect silent atrial fibrillation: a novel technology to prevent stroke. *Int J Cardiol*. 2013;165:193–194. doi: 10.1016/j.ijcard.2013.01.220.
- Doliwa PS, Frykman V, Rosenqvist M. Short-term ECG for out of hospital detection of silent atrial fibrillation episodes. *Scand Cardiovasc J*. 2009;43:163–168. doi: 10.1080/14017430802593435.
- Wiesel J, Fitzig L, Herschman Y, Messineo FC. Detection of atrial fibrillation using a modified microlife blood pressure monitor. *Am J Hypertens*. 2009;22:848–852. doi: 10.1038/ajh.2009.98.
- Chan PH, Wong CK, Poh YC, Pun L, Leung WWC, Wong YF, et al. Diagnostic performance of a smartphone-based photoplethysmographic application for atrial fibrillation screening in a primary care setting. *J Am Heart Assoc*. 2016;5:e003428. doi: 10.1161/JAHA.116.003428.
- Chan NY, Choy CC. Screening for atrial fibrillation in 13 122 Hong Kong citizens with smartphone electrocardiogram. *Heart*. 2017;103:24–31. doi: 10.1136/heartjnl-2016-309993.
- Svennberg E, Engdahl J, Al-Khalili F, Friberg L, Frykman V, Rosenqvist M. Mass screening for untreated atrial fibrillation: the STROKESTOP Study. *Circulation*. 2015;131:2176–2184. doi: 10.1161/CIRCULATIONAHA.114.014343.
- Engdahl J, Svennberg E, Friberg L, Al-Khalili F, Frykman V, Kemp Gudmundsdottir K, et al. Stepwise mass screening for atrial fibrillation using N-terminal pro b-type natriuretic peptide: the STROKESTOP II study design. *Europace*. 2017;19:297–302. doi: 10.1093/europace/euw319.
- Xiong Q, Proietti M, Senoo K, Lip GY. Asymptomatic versus symptomatic atrial fibrillation: a systematic review of age/gender differences and cardiovascular outcomes. *Int J Cardiol*. 2015;191:172–177. doi: 10.1016/j.ijcard.2015.05.011.
- Ganesan AN, Chew DP, Hartshorne T, Selvanayagam JB, Aylward PE, Sanders P, et al. The impact of atrial fibrillation type on the risk of thromboembolism, mortality, and bleeding: a systematic review and meta-analysis. *Eur Heart J*. 2016;37:1591–1602. doi: 10.1093/eurheartj/ehw007.
- Vanassche T, Lauw MN, Eikelboom JW, Healey JS, Hart RG, Alings M, et al. Risk of ischaemic stroke according to pattern of atrial fibrillation: analysis of 6563 aspirin-treated patients in ACTIVE-A and AVERROES. *Eur Heart J*. 2015;36:281–287a. doi: 10.1093/eurheartj/ehu307.

23. Hess PL, Healey JS, Granger CB, Connolly SJ, Ziegler PD, Alexander JH, et al. The role of cardiovascular implantable electronic devices in the detection and treatment of subclinical atrial fibrillation: a review. *JAMA Cardiol.* 2017;2:324–331. doi: 10.1001/jamacardio.2016.5167.
24. Healey JS, Connolly SJ, Gold MR, Israel CW, Van Gelder IC, Capucci A, et al; ASSERT Investigators. Subclinical atrial fibrillation and the risk of stroke. *N Engl J Med.* 2012;366:120–129. doi: 10.1056/NEJMoa1105575.
25. Van Gelder IC, Healey JS, Crijns HJGM, Wang J, Hohnloser SH, Gold MR, et al. Duration of device-detected subclinical atrial fibrillation and occurrence of stroke in ASSERT. *Eur Heart J.* 2017;38:1339–1344. doi: 10.1093/eurheartj/ehx042.
26. Nasir JM, Pomeroy W, Marler A, Hann M, Baykaner T, Jones R, et al. Predicting Determinants of Atrial Fibrillation or Flutter for Therapy Elucidation in Patients at Risk for Thromboembolic Events (PREDATE AF) Study. *Heart Rhythm.* 2017;14:955–961. doi: 10.1016/j.hrthm.2017.04.026.
27. Reiffel J, Verma A, Halperin JL, Gersh B, Tombul S, Carrithers J, et al. Rationale and design of REVEAL AF: a prospective study of previously undiagnosed atrial fibrillation as documented by an insertable cardiac monitor in high-risk patients. *Am Heart J.* 2014;167:22–27. doi: 10.1016/j.ahj.2013.10.007.
28. Lopes RD, Alings M, Connolly SJ, Beresh H, Granger CB, Mazuecos JB, et al. Rationale and design of the Apixaban for the Reduction of Thrombo-Embolic in Patients With Device-Detected Sub-Clinical Atrial Fibrillation (ARTESiA) trial. *Am Heart J.* 2017;189:137–145. doi: 10.1016/j.ahj.2017.04.008.
29. Kirchhof P, Blank BF, Calvert M, Camm AJ, Chlouverakis G, Diener HC, et al. Probing oral anticoagulation in patients with atrial high rate episodes: rationale and design of the Non-vitamin K antagonist Oral anticoagulants in patients with Atrial High rate episodes (NOAH-AFNET 6) trial. *Am Heart J.* 2017;190:12–18. doi: 10.1016/j.ahj.2017.04.015.
30. Lubitz SA, Yin X, Rienstra M, Schnabel RB, Walkey AJ, Magnani JW, et al. Long-term outcomes of secondary atrial fibrillation in the community: the Framingham Heart Study. *Circulation.* 2015;131:1648–1655. doi: 10.1161/CIRCULATIONAHA.114.014058.
31. Gialdini G, Nearing K, Bhavé PD, Bonuccelli U, Iadecola C, Healey JS, et al. Perioperative atrial fibrillation and the long-term risk of ischemic stroke. *JAMA.* 2014;312:616–622. doi: 10.1001/jama.2014.9143.
32. Ahlsson A, Fengsrud E, Bodin L, Englund A. Postoperative atrial fibrillation in patients undergoing aortocoronary bypass surgery carries an eightfold risk of future atrial fibrillation and a doubled cardiovascular mortality. *Eur J Cardiothorac Surg.* 2010;37:1353–1359. doi: 10.1016/j.ejcts.2009.12.033.
33. Lowres N, Mulcahy G, Gallagher R, Ben Freedman S, Marshman D, Kirkness A, et al. Self-monitoring for atrial fibrillation recurrence in the discharge period post-cardiac surgery using an iPhone electrocardiogram. *Eur J Cardiothorac Surg.* 2016;50:44–51. doi: 10.1093/ejcts/ezv486.
34. Sposato LA, Cipriano LE, Saposnik G, Ruiz Vargas E, Riccio PM, Hachinski V. Diagnosis of atrial fibrillation after stroke and transient ischaemic attack: a systematic review and meta-analysis. *Lancet Neurol.* 2015;14:377–387. doi: 10.1016/S1474-4422(15)70027-X.
35. Rosenberg MA, Samuel M, Thosani A, Zimetbaum PJ. Use of a noninvasive continuous monitoring device in the management of atrial fibrillation: a pilot study. *Pacing Clin Electrophysiol.* 2013;36:328–333. doi: 10.1111/pace.12053.
36. Walsh JA 3rd, Topol EJ, Steinhubl SR. Novel wireless devices for cardiac monitoring. *Circulation.* 2014;130:573–581. doi: 10.1161/CIRCULATIONAHA.114.009024.
37. Barrett PM, Komatireddy R, Haaser S, Topol S, Sheard J, Encinas J, et al. Comparison of 24-hour Holter monitoring with 14-day novel adhesive patch electrocardiographic monitoring. *Am J Med.* 2014;127:95.e11–95.e17. doi: 10.1016/j.amjmed.2013.10.003.
38. Lacour P, Dang PL, Huemer M, Parwani AS, Attanasio P, Pieske B, et al. Performance of the new BioMonitor 2-AF insertable cardiac monitoring system: Can better be worse? *Pacing Clin Electrophysiol.* 2017;40:516–526. doi: 10.1111/pace.13059.
39. Hindricks G, Pokushalov E, Urban L, Taborsky M, Kuck KH, Lebedev D, et al; XPECT Trial Investigators. Performance of a new leadless implantable cardiac monitor in detecting and quantifying atrial fibrillation: results of the XPECT trial. *Circ Arrhythm Electrophysiol.* 2010;3:141–147. doi: 10.1161/CIRCEP.109.877852.
40. Tomson TT, Passman R. The Reveal LINQ insertable cardiac monitor. *Expert Rev Med Devices.* 2015;12:7–18. doi: 10.1586/17434440.2014.953059.
41. European Stroke Organisation (ESO) Executive Committee; ESO Writing Committee. Guidelines for management of ischaemic stroke and transient ischaemic attack 2008. *Cerebrovasc Dis.* 2008;25:457–507. doi: 10.1159/000131083.
42. Jauch EC, Saver JL, Adams HP Jr, Bruno A, Connors JJ, Demaerschalk BM, et al; American Heart Association Stroke Council; Council on Cardiovascular Nursing; Council on Peripheral Vascular Disease; Council on Clinical Cardiology. Guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke.* 2013;44:870–947. doi: 10.1161/STR.0b013e318284056a.
43. Kernan WN, Ovbiagele B, Black HR, Bravata DM, Chimowitz MI, Ezekowitz MD, et al; American Heart Association Stroke Council, Council on Cardiovascular and Stroke Nursing, Council on Clinical Cardiology, and Council on Peripheral Vascular Disease. Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke.* 2014;45:2160–2236. doi: 10.1161/STR.0000000000000024.
44. Higgins P, MacFarlane PW, Dawson J, McInnes GT, Langhorne P, Lees KR. Noninvasive cardiac event monitoring to detect atrial fibrillation after ischemic stroke: a randomized, controlled trial. *Stroke.* 2013;44:2525–2531. doi: 10.1161/STROKEAHA.113.001927.
45. Gladstone DJ, Spring M, Dorian P, Panzov V, Thorpe KE, Hall J, et al; EMBRACE Investigators and Coordinators. Atrial fibrillation in patients with cryptogenic stroke. *N Engl J Med.* 2014;370:2467–2477. doi: 10.1056/NEJMoa1311376.
46. Sanna T, Diener HC, Passman RS, Di Lazzaro V, Bernstein RA, Morillo CA, et al; CRYSTAL AF Investigators. Cryptogenic stroke and underlying atrial fibrillation. *N Engl J Med.* 2014;370:2478–2486. doi: 10.1056/NEJMoa1313600.
47. Wachter R, Gröschel K, Gelbrich G, Hamann GF, Kermer P, Liman J, et al; Find-AF(randomised) Investigators and Coordinators. Holter-electrocardiogram-monitoring in patients with acute ischaemic stroke (Find-AFRANDOMISED): an open-label randomised controlled trial. *Lancet Neurol.* 2017;16:282–290. doi: 10.1016/S1474-4422(17)30002-9.
48. Dilaveris PE, Gialafos JE. P-wave dispersion: a novel predictor of paroxysmal atrial fibrillation. *Ann Noninvasive Electrocardiol.* 2001;6:159–165.
49. Guidera SA, Steinberg JS. The signal-averaged P wave duration: a rapid and noninvasive marker of risk of atrial fibrillation. *J Am Coll Cardiol.* 1993;21:1645–1651.
50. Magnani JW, Johnson VM, Sullivan LM, Gorodeski EZ, Schnabel RB, Lubitz SA, et al. P wave duration and risk of longitudinal atrial fibrillation in persons ≥ 60 years old (from the Framingham Heart Study). *Am J Cardiol.* 2011;107:917–921.e1. doi: 10.1016/j.amjcard.2010.10.075.
51. Thijs VN, Brachmann J, Morillo CA, Passman RS, Sanna T, Bernstein RA, et al. Predictors for atrial fibrillation detection after cryptogenic stroke: results from CRYSTAL AF. *Neurology.* 2016;86:261–269. doi: 10.1212/WNL.0000000000002282.
52. Cheng S, Keyes MJ, Larson MG, McCabe EL, Newton-Cheh C, Levy D, et al. Long-term outcomes in individuals with prolonged PR interval or first-degree atrioventricular block. *JAMA.* 2009;301:2571–2577. doi: 10.1001/jama.2009.888.
53. Duning T, Kirchhof P, Wersching H, Hepp T, Reinhardt R. Extended electrocardiographic poinecare analysis (EPA) for better identification of patients with paroxysmal atrial fibrillation. *J Clin Exp Cardiol.* 2011;2:1–7.
54. Gladstone DJ, Dorian P, Spring M, Panzov V, Mamtani M, Healey JS, et al; EMBRACE Steering Committee and Investigators. Atrial premature beats predict atrial fibrillation in cryptogenic stroke: results from the EMBRACE trial. *Stroke.* 2015;46:936–941. doi: 10.1161/STROKEAHA.115.008714.
55. Binici Z, Intzilakis T, Nielsen OW, Køber L, Sajadieh A. Excessive supraventricular ectopic activity and increased risk of atrial fibrillation and stroke. *Circulation.* 2010;121:1904–1911. doi: 10.1161/CIRCULATIONAHA.109.874982.
56. Schnabel RB, Sullivan LM, Levy D, Pencina MJ, Massaro JM, D'Agostino RB, Sr, et al. Development of a risk score for atrial fibrillation (Framingham Heart Study): a community-based cohort study. *Lancet.* 2009;373:739–745. doi: 10.1016/S0140-6736(09)60443-8.
57. Benjamin EJ, D'Agostino RB, Belanger AJ, Wolf PA, Levy D. Left atrial size and the risk of stroke and death. The Framingham Heart Study. *Circulation.* 1995;92:835–841.
58. Kim D, Shim CY, Hong GR, Kim MH, Seo J, Cho JJ, et al. Clinical implications and determinants of left atrial mechanical dysfunction in patients with stroke. *Stroke.* 2016;47:1444–1451. doi: 10.1161/STROKEAHA.115.011656.

59. Patti G, Pengo V, Marcucci R, Cirillo P, Renda G, Santilli F, et al. The left atrial appendage: from embryology to prevention of thromboembolism. *Eur Heart J*. 2017;38:877–887. doi: 10.1093/eurheartj/ehw159.
60. Hijazi Z, Oldgren J, Siegbahn A, Granger CB, Wallentin L. Biomarkers in atrial fibrillation: a clinical review. *Eur Heart J*. 2013;34:1475–1480. doi: 10.1093/eurheartj/ehv024.
61. Patton KK, Ellinor PT, Heckbert SR, Christenson RH, DeFilippi C, Gottdiener JS, et al. N-terminal pro-B-type natriuretic peptide is a major predictor of the development of atrial fibrillation: the Cardiovascular Health Study. *Circulation*. 2009;120:1768–1774. doi: 10.1161/CIRCULATIONAHA.109.873265.
62. Shibasaki K, Kimura K, Fujii S, Sakai K, Iguchi Y. Brain natriuretic peptide levels as a predictor for new atrial fibrillation during hospitalization in patients with acute ischemic stroke. *Am J Cardiol*. 2012;109:1303–1307. doi: 10.1016/j.amjcard.2011.12.022.
63. Samuel J-L, Wachter R, Lahno R, Haase B, Weber-Krüger M, Seegers J, et al. Natriuretic peptides for the detection of paroxysmal atrial fibrillation in patients with cerebral ischemia – the FIND-AF study. *PLoS ONE*. 2012;7:e34351. doi: 10.1371/journal.pone.0034351.
64. Rodríguez-Yáñez M, Arias-Rivas S, Santamaría-Cadavid M, Sobrino T, Castillo J, Blanco M. High pro-BNP levels predict the occurrence of atrial fibrillation after cryptogenic stroke. *Neurology*. 2013;81:444–447. doi: 10.1212/WNL.0b013e31829d8773.
65. Selmer C, Olesen JB, Hansen ML, Lindhardsen J, Olsen AM, Madsen JC, et al. The spectrum of thyroid disease and risk of new onset atrial fibrillation: a large population cohort study. *BMJ*. 2012;345:e7895. doi: 10.1136/bmj.e7895.
66. Lubitz SA, Parsons OE, Anderson CD, Benjamin EJ, Malik R, Weng L-C, et al; International Stroke Genetics Consortium, and AFGen Consortia. Atrial fibrillation genetic risk and ischemic stroke mechanisms. *Stroke*. 2017;48:1451–1456. doi: 10.1161/STROKEAHA.116.016198.
67. Hart RG, Catanese L, Perera KS, Ntaios G, Connolly SJ. Embolic stroke of undetermined source: a systematic review and clinical update. *Stroke*. 2017;48:867–872. doi: 10.1161/STROKEAHA.116.016414.
68. Demeestere J, Fieuws S, Lansberg MG, Lemmens R. Detection of atrial fibrillation among patients with stroke due to large or small vessel disease: a meta-analysis. *J Am Heart Assoc*. 2016;5:e004151.
69. Bernstein RA, Kamel H, Granger CB, Kowal RC, Ziegler PD, Schwamm LH. Stroke of Known Cause and Underlying Atrial Fibrillation (STROKE-AF) randomized trial: design and rationale. *Am Heart J*. 2017;190:19–24. doi: 10.1016/j.ahj.2017.04.007.
70. Korompoki E, Del Giudice A, Hillmann S, Malzahn U, Gladstone DJ, Heuschmann P, et al. Cardiac monitoring for detection of atrial fibrillation after TIA: a systematic review and meta-analysis. *Int J Stroke*. 2017;12:33–45. doi: 10.1177/1747493016669885.
71. Bobinger T, Kallmünzer B, Kopp M, Kurka N, Arnold M, Heider S, et al. Diagnostic value of prehospital ECG in acute stroke patients. *Neurology*. 2017;88:1894–1898. doi: 10.1212/WNL.0000000000003940.
72. Rizos T, Güntner J, Jenetzky E, Marquardt L, Reichardt C, Becker R, et al. Continuous stroke unit electrocardiographic monitoring versus 24-hour Holter electrocardiography for detection of paroxysmal atrial fibrillation after stroke. *Stroke*. 2012;43:2689–2694. doi: 10.1161/STROKEAHA.112.654954.
73. Rizos T, Rasch C, Jenetzky E, Hametner C, Kathoefer S, Reinhardt R, et al. Detection of paroxysmal atrial fibrillation in acute stroke patients. *Cerebrovasc Dis*. 2010;30:410–417. doi: 10.1159/000316885.
74. Choe WC, Passman RS, Brachmann J, Morillo CA, Sanna T, Bernstein RA, et al; CRYSTAL AF Investigators. A comparison of atrial fibrillation monitoring strategies after cryptogenic stroke (from the Cryptogenic Stroke and Underlying AF Trial). *Am J Cardiol*. 2015;116:889–893. doi: 10.1016/j.amjcard.2015.06.012.
75. Ziegler PD, Rogers JD, Ferreira SW, Nichols AJ, Sarkar S, Koehler JL, et al. Real-world experience with insertable cardiac monitors to find atrial fibrillation in cryptogenic stroke. *Cerebrovasc Dis*. 2015;40:175–181. doi: 10.1159/000439063.

KEY WORDS: atrial fibrillation ■ cardioembolic stroke ■ cryptogenic stroke ■ embolic stroke of unknown source ■ screening ■ stroke