## THE LNM INSTITUTE OF INFORMATION TECHNOLOGY JAIPUR, RAJASTHAN

## End Semester Exam Part-I

MATH-III,, 21st Nov, 2016

Time: 45 minutes, Maximum Marks:10

Name:	Roll No.:	

**Note:** Each question carry 2 marks. Overwriting will be treated as a wrong answer. Use only the last page of main answer sheet for rough work and calculation. Write down the final answer, no marks for formula or incomplete solution.

1. The PDE which characterizes the surfaces

$$F(xyz, x + y + z) = 0,$$

is \_\_\_\_\_

Solution: x(y-z)p+y(z-x)q=(x-y)z (Here  $u=xyz,\ v=x+y+z,\ P=\frac{\partial(u,v)}{\partial(y,z)}=x(z-y),\ Q=\frac{\partial(u,v)}{\partial(z,x)}=y(x-z),\ R=\frac{\partial(u,v)}{\partial(x,y)}=z(y-x))$ 

2. (a) The classification (elliptic, parabolic, or hyperbolic) of PDE

$$xu_{xx} + (x - y)u_{xy} - yu_{yy} = 0, \ x > 0, \ y < 0,$$

is \_\_\_\_\_ type.

(b) Tick Appropriately: The PDE  $x^3u_{xxxx} + xy^3u_{yy} = 0$  is first/second/third/forth order and linear/semi-linear/quasi-linear/non-linear PDE.

Solution: (a) Hyperbolic if  $x + y \neq 0$  and parabolic for x + y = 0 (Here  $b^2 - 4ac = (x - y)^2 + 4xy = (x + y)^2$ )

- (b) Forth order linear PDE
- 3. Solution for following PDE

$$u_{tt} - u_{xx} = 0,$$
  $0 < x < \infty, \ t > 0$   
 $u(x,0) = \cos x,$   $u_t(x,0) = 0, 0 \le x < \infty,$   
 $u(0,t) = 0,$   $t \ge 0.$ 

is 
$$u(x,t) = \begin{cases} \cos x \cos t & x \ge t \\ -\sin x \sin t & x \le t \end{cases}$$

4. Solution of the wave equation (Hint: Duhamel's principle):

$$u_{tt} = u_{xx} + 2t,$$
  $-\infty < x < \infty, t > 0,$   
 $u(x,0) = u_t(x,0) = 0,$   $-\infty < x < \infty.$ 

is 
$$\frac{t^3}{3}$$

5. Solution for the following Laplace equation:

$$u_{xx} + u_{yy} = 0,$$
  $0 < x, y < 2\pi,$   
 $u(x,0) = 2x, \ 0 \le x \le 2\pi,$   $u(x,2\pi) = 0, \ 0 \le x \le 2\pi,$   
 $u(0,y) = u(2\pi,y) = 0,$   $0 \le y \le 2\pi.$ 

is 
$$u(x,y) = \sum_{n=1}^{\infty} \frac{(-1)^n 8 \sin(\frac{nx}{2}) \sinh[\frac{n(y-2\pi)}{2}]}{n \sinh(n\pi)}$$
.