

MA1522 Linear Algebra for Computing

Lecture 10: Applications (of Orthogonality) and Eigenvalues

Yang Yue

Department of Mathematics
National University of Singapore

24 March, 2025

Outline

Exercises and Questions posed in Dr.Teo's Lectures

Challenges posed in Dr.Teo's Lectures

Exercise One in Section 5.4

1. Prove that $\mathbf{Q}^T \mathbf{Q} = \mathbf{I}_n$.
2. Prove that the diagonal entries of \mathbf{R} are positive, $r_{ii} > 0$ for all $i = 1, \dots, n$.

3. Prove that the upper triangular matrix

$$\mathbf{R} = \begin{pmatrix} r_{11} & r_{12} & \cdots & r_{1n} \\ 0 & r_{22} & \cdots & r_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & r_{nn} \end{pmatrix} \text{ is invertible.}$$

(Exercises are part of proofs that Dr. Teo skipped.)

Slide 57: QR Factorization

Theorem (QR Factorization)

Suppose \mathbf{A} is an $m \times n$ matrix with *linearly independent* columns.
Then \mathbf{A} can be written as

$$\mathbf{A} = \mathbf{QR}$$

for some $m \times n$ matrix \mathbf{Q} such that $\mathbf{Q}^T \mathbf{Q} = \mathbf{I}_n$ and invertible upper triangular matrix \mathbf{R} with *positive* diagonal entries.

Definition

The decomposition given in the theorem above is called a QR factorization of \mathbf{A} .

Where do **Q** and **R** come from?

Suppose $\mathbf{A} = (\mathbf{a}_1 \ \mathbf{a}_2 \ \cdots \ \mathbf{a}_n)$ whose columns are **linearly independent**. Applying the Gram-Schmidt process on the columns, we obtain a matrix $\mathbf{Q} = (\mathbf{q}_1 \ \mathbf{q}_2 \ \cdots \ \mathbf{q}_n)$ whose columns are orthonormal.

By Gram-Schmidt, $\text{span}\{\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_i\} = \text{span}\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_i\}$. Thus,

$$\begin{aligned} \mathbf{a}_i &= r_{1i}\mathbf{q}_1 + r_{2i}\mathbf{q}_2 + \cdots + r_{ii}\mathbf{q}_i + 0\mathbf{q}_{i+1} + \cdots + 0\mathbf{q}_n \\ &= \begin{pmatrix} \mathbf{q}_1 & \cdots & \mathbf{q}_i & \cdots & \mathbf{q}_n \end{pmatrix} \begin{pmatrix} r_{1i} \\ \vdots \\ r_{ii} \\ 0 \\ \vdots \\ 0 \end{pmatrix}. \end{aligned}$$

Where do **Q** and **R** come from? (conti.)

Putting things together,

$$\begin{aligned}\mathbf{A} &= \begin{pmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{q}_1 & \mathbf{q}_2 & \cdots & \mathbf{q}_n \end{pmatrix} \begin{pmatrix} r_{11} & r_{12} & \cdots & r_{1n} \\ 0 & r_{22} & \cdots & r_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & r_{nn} \end{pmatrix} \\ &= \mathbf{QR}\end{aligned}$$

for some $m \times n$ matrix **Q** with **orthonormal columns**, and an **upper triangular** $n \times n$ matrix **R**.

Algorithm to QR Factorization

Let \mathbf{A} be an $m \times n$ matrix with **linearly independent** columns.

1. Perform Gram-Schmidt on the columns of $\mathbf{A} = (\mathbf{a}_1 \ \mathbf{a}_2 \ \cdots \ \mathbf{a}_n)$ to obtain an **orthonormal set** $\{\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_n\}$.
2. Set $\mathbf{Q} = (\mathbf{q}_1 \ \mathbf{q}_2 \ \cdots \ \mathbf{q}_n)$.
3. Compute $\mathbf{R} = \mathbf{Q}^T \mathbf{A}$.

Note, item 3 gave us an alternative way to calculate \mathbf{R} .

Answer to Exercise One in Section 5.4 (part 1)

1. Prove that $\mathbf{Q}^T \mathbf{Q} = \mathbf{I}_n$.

Proof. Since \mathbf{Q} is an $m \times n$ matrix, \mathbf{Q}^T is $n \times m$. Thus $\mathbf{Q}^T \mathbf{Q}$ is $n \times n$. By block multiplication,

$$\mathbf{Q}^T \mathbf{Q} = \begin{pmatrix} \mathbf{q}_1^T \\ \vdots \\ \mathbf{q}_n^T \end{pmatrix} (\mathbf{q}_1 \cdots \mathbf{q}_n) = (\mathbf{q}_i^T \mathbf{q}_j)$$

Observe that $\mathbf{q}_i^T \mathbf{q}_j$ (as matrix multiplication) is equal to the dot product $\mathbf{q}_i \cdot \mathbf{q}_j$, and $\{\mathbf{q}_1, \dots, \mathbf{q}_n\}$ is orthonormal, we then have

$$(\mathbf{q}_i^T \mathbf{q}_j) = \mathbf{I}_n.$$

Answer to Exercise One in Section 5.4 (part 2)

2. Prove that the diagonal entries of \mathbf{R} are positive, $r_{ii} > 0$ for all $i = 1, \dots, n$.

Proof. By earlier slides, we have, for each $i \leq n$,

$$\mathbf{a}_i = r_{1i}\mathbf{q}_1 + r_{2i}\mathbf{q}_2 + \cdots + r_{ii}\mathbf{q}_i. \quad (1)$$

Using \mathbf{q}_i to dot multiply both sides of (1), we have

$$r_{ii} = \mathbf{a}_i \cdot \mathbf{q}_i.$$

Now back to Gram-Schmidt, we have

$$\mathbf{v}_i = \mathbf{a}_i - \left(\frac{\mathbf{v}_1 \cdot \mathbf{a}_i}{\|\mathbf{v}_1\|^2} \right) \mathbf{v}_1 - \left(\frac{\mathbf{v}_2 \cdot \mathbf{a}_i}{\|\mathbf{v}_2\|^2} \right) \mathbf{v}_2 - \cdots - \left(\frac{\mathbf{v}_{i-1} \cdot \mathbf{a}_i}{\|\mathbf{v}_{i-1}\|^2} \right) \mathbf{v}_{i-1}. \quad (2)$$

and $\mathbf{q}_i = \frac{\mathbf{v}_i}{\|\mathbf{v}_i\|}$, using \mathbf{q}_i to dot multiply both sides of (2), we have

$$\mathbf{a}_i \cdot \mathbf{q}_i = \mathbf{v}_i \cdot \mathbf{q}_i = \|\mathbf{v}_i\| > 0.$$

Consequently, $r_{ii} > 0$.

Answer to Exercise One in Section 5.4 (part 3)

3. Prove that the upper triangular matrix

$$\mathbf{R} = \begin{pmatrix} r_{11} & r_{12} & \cdots & r_{1n} \\ 0 & r_{22} & \cdots & r_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & r_{nn} \end{pmatrix}$$

is invertible.

Proof. It follows from

$$\det(\mathbf{R}) = r_{11} \cdots r_{nn} > 0.$$

Exercise Two in Section 5.4

Use QR factorization to prove the following

Corollary

Suppose \mathbf{A} is an $m \times n$ matrix with *linearly independent* columns, i.e. $\text{rank}(\mathbf{A}) = n$. Then $\mathbf{A}^T \mathbf{A}$ is invertible, and \mathbf{A} has a *left inverse*; that is, there is a \mathbf{B} such that

$$\mathbf{B}\mathbf{A} = \mathbf{I}_n.$$

Answer to Exercise Two in Section 5.4

Use QR factorization to prove that if \mathbf{A} is an $m \times n$ matrix with linearly independent columns, then $\mathbf{A}^T \mathbf{A}$ is invertible, and \mathbf{A} has a left inverse; that is, there is a \mathbf{B} such that

$$\mathbf{BA} = \mathbf{I}_n.$$

Proof. Since \mathbf{A} has independent columns, \mathbf{A} has a QR-decomposition $\mathbf{A} = \mathbf{QR}$. Then

$$\mathbf{A}^T \mathbf{A} = \mathbf{R}^T \mathbf{Q}^T \mathbf{Q} \mathbf{R} = \mathbf{R}^T \mathbf{R},$$

because $\mathbf{Q}^T \mathbf{Q} = \mathbf{I}_n$ by Exercise above. Since \mathbf{R} is invertible (by the same exercise), \mathbf{R}^T is also invertible, and hence $\mathbf{A}^T \mathbf{A}$ is invertible. Let \mathbf{P} be its inverse, then $\mathbf{B} = \mathbf{PA}^T$ is a left inverse of \mathbf{A} .

Question in Section 5.5

Suppose the system $\mathbf{Ax} = \mathbf{b}$ is consistent.

1. Suppose \mathbf{u} is a solution to $\mathbf{Ax} = \mathbf{b}$. Is \mathbf{u} a least square solution to $\mathbf{Ax} = \mathbf{b}$?
2. Suppose \mathbf{u} is a least square solution to $\mathbf{Ax} = \mathbf{b}$. Is \mathbf{u} a solution to $\mathbf{Ax} = \mathbf{b}$?


Slide 64: Least Square Approximation

Definition

Let \mathbf{A} be an $m \times n$ matrix and \mathbf{b} a vector in \mathbb{R}^m . A vector \mathbf{u} in \mathbb{R}^n is a least square solution of $\mathbf{Ax} = \mathbf{b}$ if for every vector $\mathbf{v} \in \mathbb{R}^n$,

$$\|\mathbf{Au} - \mathbf{b}\| \leq \|\mathbf{Av} - \mathbf{b}\|.$$

Geometrically, by the **best approximation theorem**, the vector $\mathbf{b}' = \mathbf{Au}$ in $\text{Col}(\mathbf{A})$ closest to \mathbf{b} is the **projection** of \mathbf{b} onto $\text{Col}(\mathbf{A})$.



Answer to Question in Section 5.5

Suppose the system $\mathbf{Ax} = \mathbf{b}$ is **consistent**.

1. Suppose \mathbf{u} is a **solution** to $\mathbf{Ax} = \mathbf{b}$. Is \mathbf{u} a **least square solution** to $\mathbf{Ax} = \mathbf{b}$?

Answer: Yes, because for every vector $\mathbf{v} \in \mathbb{R}^n$,

$$\|\mathbf{Au} - \mathbf{b}\| = 0 \leq \|\mathbf{Av} - \mathbf{b}\|.$$

2. Suppose \mathbf{u} is a **least square solution** to $\mathbf{Ax} = \mathbf{b}$. Is \mathbf{u} a **solution** to $\mathbf{Ax} = \mathbf{b}$?

Answer: Yes. By the assumption that $\mathbf{Ax} = \mathbf{b}$ is **consistent**, there is some \mathbf{v} with $\mathbf{Av} = \mathbf{b}$. Since \mathbf{u} is a least square solution,

$$\|\mathbf{Au} - \mathbf{b}\| \leq \|\mathbf{Av} - \mathbf{b}\| = 0.$$

Hence $\mathbf{Au} = \mathbf{b}$.

Exercise in Section 5.5

Suppose \mathbf{A} is an $m \times n$ matrix with linearly independent columns, i.e. $\text{rank}(\mathbf{A}) = n$. QR factorize \mathbf{A} ,

$$\mathbf{A} = \mathbf{QR}.$$

Show that the unique least square solution of $\mathbf{Ax} = \mathbf{b}$ is

$$\mathbf{u} = \mathbf{R}^{-1}\mathbf{Q}^T\mathbf{b}.$$

That is, suffice to solve for

$$\mathbf{Rx} = \mathbf{Q}^T\mathbf{b}.$$

This is easy to solve by hand since \mathbf{R} is an upper triangular matrix (i.e. an REF).

Slide 65: Least Square Approximation

Theorem

Let \mathbf{A} be an $m \times n$ matrix and \mathbf{b} a vector in \mathbb{R}^m . A vector \mathbf{u} in \mathbb{R}^n is a *least square solution* to $\mathbf{Ax} = \mathbf{b}$ if and only if \mathbf{Au} is the *projection* of \mathbf{b} onto the column space of $\text{Col}(\mathbf{A})$.

Theorem

Let \mathbf{A} be an $m \times n$ matrix and \mathbf{b} a vector in \mathbb{R}^m . A vector \mathbf{u} in \mathbb{R}^n is a *least square solution* to $\mathbf{Ax} = \mathbf{b}$ if and only if \mathbf{u} is a *solution* to $\mathbf{A}^T \mathbf{Ax} = \mathbf{A}^T \mathbf{b}$.

Answer to Exercise in Section 5.5

Suppose \mathbf{A} is an $m \times n$ matrix with linearly independent columns, and $\mathbf{A} = \mathbf{QR}$ is a QR factorization. Show that the unique least square solution of $\mathbf{Ax} = \mathbf{b}$ is

$$\mathbf{u} = \mathbf{R}^{-1}\mathbf{Q}^T\mathbf{b}.$$

Proof. By the Theorem on previous slide, the least square solution \mathbf{u} is the solution to

$$\mathbf{A}^T\mathbf{Ax} = \mathbf{A}^T\mathbf{b}.$$

Substituting $\mathbf{A} = \mathbf{QR}$, and using $\mathbf{Q}^T\mathbf{Q} = \mathbf{I}$ and \mathbf{R} and \mathbf{R}^T are invertible, we have

$$\begin{aligned}\mathbf{A}^T\mathbf{Ax} = \mathbf{A}^T\mathbf{b} &\Leftrightarrow \mathbf{R}^T\mathbf{Q}^T\mathbf{QR}\mathbf{u} = \mathbf{R}^T\mathbf{Q}^T\mathbf{b} \\ &\Leftrightarrow \mathbf{R}^T\mathbf{R}\mathbf{u} = \mathbf{R}^T\mathbf{Q}^T\mathbf{b} \\ &\Leftrightarrow \mathbf{u} = \mathbf{R}^{-1}\mathbf{Q}^T\mathbf{b}.\end{aligned}$$

Questions in Section 6.1

1. Let \mathbf{A} and \mathbf{B} be row equivalent order n square matrices.
 - (a) If λ is an eigenvalue of \mathbf{A} , is it an eigenvalue of \mathbf{B} ?
 - (b) If \mathbf{v} is an eigenvector of \mathbf{A} , is it an eigenvector of \mathbf{B} ?
2. Can we compute the characteristic polynomial of a square matrix using row reduction instead of cofactor expansion?

Slide 6: Eigenvalues and Eigenvectors

Definition

Let \mathbf{A} be a **square** matrix of order n . A real number λ is an eigenvalue of \mathbf{A} if there is a **nonzero** vector \mathbf{v} in \mathbb{R}^n , such that

$$\mathbf{A}\mathbf{v} = \lambda\mathbf{v}.$$

In this case, the nonzero vector \mathbf{v} is called an eigenvector **associated** to λ .

Remarks:

- ▶ Think of a matrix as an *action* on \mathbb{R}^n (we will learn linear transformation later). Geometrically, eigenvectors are the vectors that are being scaled (stretch, dilate, or reflect) when acted upon by \mathbf{A} , and eigenvalues are the amount to scale the eigenvectors.
- ▶ We require the eigenvector to be nonzero, $\mathbf{v} \neq \mathbf{0}$, otherwise, the definition becomes uninteresting.

Slides 8 and 9: Characteristic Polynomial

Definition

Let \mathbf{A} be a **square** matrix of order n , the characteristic polynomial of \mathbf{A} , denoted as $\text{char}(\mathbf{A})$, is the **degree n polynomial**

$$\det(x\mathbf{I} - \mathbf{A}).$$

Theorem

Let \mathbf{A} be a **square** matrix of order n . λ is an **eigenvalue** of \mathbf{A} if and only if λ is a **root** of the **characteristic polynomial** $\det(x\mathbf{I} - \mathbf{A})$.

Answer to Questions in Section 6.1 (part 1)

1. Let **A** and **B** be row equivalent order n square matrices.

(a) If λ is an eigenvalue of **A**, is it an eigenvalue of **B**?

(b) If **v** is an eigenvector of **A**, is it an eigenvector of **B**?

Answer: Both (a) and (b) are false. For example, let

$$\mathbf{A} = \begin{pmatrix} 1 & 0 \\ 0 & 4 \end{pmatrix} \quad \mathbf{B} = \begin{pmatrix} 0 & 4 \\ 1 & 0 \end{pmatrix},$$

which are row equivalent. But

$$\det(x\mathbf{I} - \mathbf{A}) = (x - 1)(x - 4) \quad \det(x\mathbf{I} - \mathbf{B}) = (x + 2)(x - 2),$$

which shows that they don't share the same eigenvalues. Moreover

$\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ is an eigenvector of **A** associated with 1, but it is not an eigenvector of **B**.

Answer to Questions in Section 6.1 (part 2)

2. Can we compute the characteristic polynomial of a square matrix using row reduction instead of cofactor expansion?

Answer: Yes, as long as the row operations are on the matrix $x\mathbf{I} - \mathbf{A}$ (not on \mathbf{A}).

Challenge one in Section 5.5

Let \mathbf{A} be an $m \times n$ matrix and \mathbf{b} a vector in \mathbb{R}^m . Prove that for any choice of **least square solution** \mathbf{u} , that is, for any solution \mathbf{u} of $\mathbf{A}^T \mathbf{A} \mathbf{x} = \mathbf{A}^T \mathbf{b}$, the **projection** $\mathbf{A} \mathbf{u}$ is **unique**.

Recall that on Slide 65, we have

Theorem

*Let \mathbf{A} be an $m \times n$ matrix and \mathbf{b} a vector in \mathbb{R}^m . A vector \mathbf{u} in \mathbb{R}^n is a **least square solution** to $\mathbf{A} \mathbf{x} = \mathbf{b}$ if and only if $\mathbf{A} \mathbf{u}$ is the **projection** of \mathbf{b} onto the column space of $\text{Col}(\mathbf{A})$.*

The Challenge is essentially the “only if” direction.

Answer to Challenge one in Section 5.5

Q: Let \mathbf{A} be an $m \times n$ matrix and \mathbf{b} a vector in \mathbb{R}^m . Prove that for any choice of **least square solution** \mathbf{u} , that is, for any solution \mathbf{u} of $\mathbf{A}^T \mathbf{A} \mathbf{x} = \mathbf{A}^T \mathbf{b}$, the **projection** $\mathbf{A} \mathbf{u}$ is **unique**.

Answer: Suppose that $\mathbf{A}^T \mathbf{A} \mathbf{u} = \mathbf{A}^T \mathbf{b}$. Then $\mathbf{A}^T (\mathbf{A} \mathbf{u} - \mathbf{b}) = \mathbf{0}$, in other words, $\mathbf{A} \mathbf{u} - \mathbf{b} \in \text{Null}(\mathbf{A}^T)$.

By Orthogonal to a Subspace algorithm (on Slide 16 in Chapter 5), $\mathbf{A} \mathbf{u} - \mathbf{b}$ is orthogonal to the column space of \mathbf{A} . Hence

$$\mathbf{b} = \mathbf{A} \mathbf{u} + (\mathbf{b} - \mathbf{A} \mathbf{u})$$

is a decomposition as in Orthogonal Projection Theorem. Thus $\mathbf{A} \mathbf{u}$, being the unique orthogonal projection of \mathbf{b} to $\text{Col}(\mathbf{A})$, is unique.

Challenge two in Section 5.5

Let $V \subseteq \mathbb{R}^n$ be a subspace and suppose $\{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_k\}$ is an orthonormal basis of V . Write

$$\mathbf{Q} = \begin{pmatrix} \mathbf{w}_1 & \mathbf{w}_2 & \cdots & \mathbf{w}_k \end{pmatrix}.$$

Then for any $\mathbf{w} \in \mathbb{R}^n$, the projection of \mathbf{w} onto V is

$$\mathbf{Q}\mathbf{Q}^T\mathbf{w}.$$

Answer to Challenge two in Section 5.5

Let $V \subseteq \mathbb{R}^n$ be a subspace and suppose $\{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_k\}$ is an orthonormal basis of V . Write

$$\mathbf{Q} = \begin{pmatrix} \mathbf{w}_1 & \mathbf{w}_2 & \cdots & \mathbf{w}_k \end{pmatrix}.$$

Then for any $\mathbf{w} \in \mathbb{R}^n$, the projection of \mathbf{w} onto V is $\mathbf{Q}\mathbf{Q}^T\mathbf{w}$.

From Challenge one, we know that if $\mathbf{A}^T\mathbf{A}\mathbf{x} = \mathbf{A}^T\mathbf{b}$, then $\mathbf{A}\mathbf{u}$ is the projection of \mathbf{b} onto $\text{Col}(\mathbf{A})$.

In Challenge Two, \mathbf{A} is \mathbf{Q} and \mathbf{b} is \mathbf{w} . It suffices to check that $\mathbf{u} = \mathbf{Q}^T\mathbf{w}$ satisfies

$$\mathbf{Q}^T\mathbf{Q}\mathbf{u} = \mathbf{Q}^T\mathbf{w},$$

which follows immediately from $\mathbf{Q}^T\mathbf{Q} = \mathbf{I}$.

Challenges in Section 6.1

Let \mathbf{A} be an $n \times n$ matrix.

1. Show that the characteristic polynomial of \mathbf{A} is equal to the characteristic polynomial of \mathbf{A}^T . Hence \mathbf{A} and \mathbf{A}^T have the same eigenvalues.
2. Let λ be an eigenvalue of \mathbf{A} . Show that the geometric multiplicity of λ as an eigenvalue of \mathbf{A} is equal to its geometric multiplicity as an eigenvalue of \mathbf{A}^T .

Slide 17: Eigenspace

Recall that **eigenvectors** of **A** associated to eigenvalue λ are **nontrivial** solution to

$$(\lambda \mathbf{I} - \mathbf{A})\mathbf{x} = \mathbf{0}.$$

Since the system is **homogeneous**, the set of all solutions is a subspace. We will call it the **eigenspace** of **A** associated to eigenvalue λ .

Definition

Let **A** be an order n **square** matrix. The **eigenspace** associated to an eigenvalue λ of **A** is

$$E_\lambda = \{ \mathbf{v} \in \mathbb{R}^n \mid \mathbf{A}\mathbf{v} = \lambda\mathbf{v} \} = \text{Null}(\lambda \mathbf{I} - \mathbf{A}).$$

The **geometric multiplicity** of an eigenvalue λ is the **dimension** of its eigenspace,

$$\dim(E_\lambda) = \text{nullity}(\lambda \mathbf{I} - \mathbf{A}).$$

Slide 37 in Chapter 4: Rank-Nullity Theorem

Theorem (Rank-Nullity Theorem)

Let \mathbf{A} be a $m \times n$ matrix. The sum of its rank and nullity is equal to the number of columns,

$$\text{rank}(\mathbf{A}) + \text{nullity}(\mathbf{A}) = n.$$

Sketch of Proof: This follows from the fact that the nullity of \mathbf{A} is equal to the number of non-pivot columns in its reduced row-echelon form, and that the rank of \mathbf{A} is equal to the number of pivot columns of its reduced row-echelon form.

Answer to Challenges in Section 6.1 (part 1)

Let \mathbf{A} be an $n \times n$ matrix.

1. Show that the characteristic polynomial of \mathbf{A} is equal to the characteristic polynomial of \mathbf{A}^T . Hence \mathbf{A} and \mathbf{A}^T have the same eigenvalues.

Proof. The following fact is useful for both parts: For a square matrix \mathbf{A}

$$(\lambda \mathbf{I} - \mathbf{A})^T = (\lambda \mathbf{I})^T - \mathbf{A}^T = \lambda \mathbf{I} - \mathbf{A}^T.$$

For part 1, we have

$$\det(\lambda \mathbf{I} - \mathbf{A}) = \det(\lambda \mathbf{I} - \mathbf{A})^T = \det(\lambda \mathbf{I} - \mathbf{A}^T),$$

where we used the fact in the last equality.

Answer to Challenges in Section 6.1 (part 2)

Let \mathbf{A} be an $n \times n$ matrix.

2. Let λ be an eigenvalue of \mathbf{A} . Show that the geometric multiplicity of λ as an eigenvalue of \mathbf{A} is equal to its geometric multiplicity as an eigenvalue of \mathbf{A}^T .

Proof. For part 2, we have

$$\begin{aligned}\text{nullity}(\lambda \mathbf{I} - \mathbf{A}) &= n - \text{rank}(\lambda \mathbf{I} - \mathbf{A}) \\ &= n - \text{rank}(\lambda \mathbf{I} - \mathbf{A})^T \\ &= n - \text{rank}(\lambda \mathbf{I} - \mathbf{A}^T) \\ &= \text{nullity}(\lambda \mathbf{I} - \mathbf{A}^T).\end{aligned}$$