

A diagnosis of dementia is usually made within memory services. Some care home residents with advanced dementia have never had a formal diagnosis. In these cases a referral to memory services is rarely desirable. It is likely to be distressing for the individual and is usually unnecessary¹.

People with advanced dementia, their families and staff caring for them, still benefit from a formal diagnosis. It enables access to appropriate care to meet individual needs and prompts staff to consider MCA and DOLs issues where appropriate. A diagnosis of dementia can be made with a high degree of certainty if all five criteria listed below are met:



1 Functional impairment

The person is no longer fully independent in relation to basic activities of daily living, washing, dressing, feeding and attending to own continence needs. The requirement of prompting or supervision of staff constitutes a loss of full independence.

☐

2 Cognitive impairment – 6 CIT assessment

Question	Scoring	Score achieved
1.What year is it?	Correct – 0 points; incorrect – 4 points	
2.What month is it?	Correct – 0 points; Incorrect – 3 points	
3.Give an address phase to remember with 5 components e.g. John, Smith, 42, High St, Wakefield		
4.About what time is it (within 1 hour)	Correct – 0 points; Incorrect – 3 points	
5.Count backwards from 20-1	No errors – 0 points; 1 error – 2 points; more than 1 error – 4 points	
6.Say the months of the year in reverse	No errors – 0 points; 1 error – 2 points; more than 1 error – 4 points	
7.Repeat address phase	No errors – 0 points; score 2 points for every component wrong e.g. 3 errors, 6 points	
TOTAL SCORE:		

6 CIT scores: 7 and below normal; 8 and above indicate impairment.

Assessment tool/other than 6CIT can be used. If used does score indicate impairment Y/N?

NB. Scores obtained in this patient group would be expected to be at the severe end of scale and for some patients their cognitive impairment will be of such severity that they cannot undertake the assessment.

Y / N

3 Corroborating History

History of gradual cognitive decline (typically for the last few years) is confirmed by care staff, relatives and medical records. Staff/relatives confirm that in their opinion the patient consistently demonstrates both functional and cognitive impairment.

☐

4 Investigations

Dementia screening **bloods are normal** (where clinically appropriate and patient consents to bloods). If patient lacks capacity to consent to bloods, a best interest decision must be made and documented accordingly. NB. If intracranial pathology (e.g. subdural haematoma, cerebral tumour) is suspected, referral for a brain scan may be appropriate. Otherwise where dementia is advanced, differential diagnosis is unlikely to affect patient management & a brain scan is unnecessary.

☐

5 Exclusion Criteria

There is **no acute underlying cause to explain** confusion i.e. delirium (acute confusional state) has been excluded. Mood disorder or psychosis is also excluded.

☐

A diagnosis of dementia can be made with a high degree of certainty if **all five** criteria listed above are met. If dementia is confirmed, please add this patient to your GP practice dementia register using the recommended codes. Consent should be sought for this from the person themselves or a family carer where the individual lacks capacity.

NB. Where a diagnosis of dementia is confirmed, a copy of the completed DiADeM tool should be saved into the patient's notes as it forms part of their clinical record.

Any enquiries regarding this publication should be sent to: ENGLAND.YHSCN@nhs.net

¹ "Guidance for Commissioners of Dementia Services", published by The Joint Commissioning Panel for Mental Health states patients who present with advanced symptoms of dementia can be diagnosed and managed by primary care with or without CMHT help. www.icpmh.info