

LIS Triangle

Input file: **standard input**
Output file: **standard output**
Time limit: 2 seconds
Memory limit: 1024 megabytes

You are given three positive integers N , K , and L . Your task is to determine whether there exists an integer sequence P of length N that satisfies all of the following conditions. If such a sequence exists, output one such sequence.

- P is a permutation of the sequence $(K, K + 1, \dots, K + N - 1)$.
- The length of the longest increasing subsequence* of P is exactly L .
- For every integer i such that $1 \leq i \leq N - 2$, there exists a non-degenerate triangle[†] with side lengths P_i, P_{i+1} , and P_{i+2} .

You are given T test cases. Answer each test case separately.

Input

The input is given in the following format:

```
T
case1
case2
⋮
caseT
```

Each test case is given in the following format:

```
N K L
```

- All input values are integers.
- $1 \leq T \leq 50000$.
- $3 \leq N \leq 2 \times 10^5$.
- $1 \leq K \leq 2 \times 10^5$.
- $1 \leq L \leq N$.
- Over all test cases in a single input, the sum of N is at most 2×10^5 .

Output

If there is no sequence P that satisfies all the conditions, output:

```
No
```

* A **subsequence** of a sequence P is a sequence formed by removing zero or more elements from P without changing the order of the remaining elements. A **longest increasing subsequence** of P is a strictly increasing subsequence of P that has the maximum possible length.

[†] A non-degenerate triangle is a triangle whose three vertices do not lie on the same straight line.

Otherwise, output:

Yes
 $P_1 P_2 \dots P_N$

Any valid sequence P that satisfies all conditions is accepted.

Example

standard input	standard output
3	Yes
6 3 4	3 6 4 7 5 8
5 5 5	Yes
7 1 2	5 6 7 8 9
	No

Note

In the first example, one valid sequence is $P = (3, 6, 4, 7, 5, 8)$. Other valid sequences may also exist.

In the second example, the only valid sequence is $P = (5, 6, 7, 8, 9)$.

In the third example, no sequence P satisfies all the conditions.