

## Drill 8

1. Birnbaum and Saunders (1969) proposed the following distribution. Given shape parameter  $\alpha > 0$  and scale parameter  $\beta > 0$ , the probability density function (pdf) and the cumulative distribution function (cdf) are given by

$$f(t) = \frac{1}{2\alpha\beta\sqrt{2\pi}} \left[ \sqrt{\frac{\beta}{t}} + \left( \frac{\beta}{t} \right)^{3/2} \right] \exp \left[ -\frac{1}{2\alpha^2} \left( \frac{t}{\beta} - 2 + \frac{\beta}{t} \right) \right]$$

and

$$F(t) = \Phi \left[ \frac{1}{\alpha} \left( \sqrt{\frac{t}{\beta}} - \sqrt{\frac{\beta}{t}} \right) \right], \quad t > 0,$$

respectively, where  $\Phi(\cdot)$  is the standard normal cdf.

- (a) Analogous to the Weibull plot, make a formula for plotting the data from the Birnbaum and Saunders (BS) distribution which draws a straight line.
- (b) Make the plots (Weibull plot and BS plot) using the given data sets in `Data.r` file.
2. Make a R function for Table 2 of Looney and Gullledge, Jr. (1985). The output would be as below:
- ```
> set.seed(1)
> EPP(n=3, levels=c(0.005, 0.010) )
      0.5%      1%
0.8675 0.8685

> EPP(n=3, levels=c(0, 0.005, 0.010) )
      0%      0.5%      1%
0.8661 0.8675 0.8687
```
3. Make the above R function without using `for( )` loop in R.

## References

- Birnbaum, Z. W. and Saunders, S. C. (1969). A new family of life distributions. *Journal of Applied Probability*, 6:319–327.
- Looney, S. W. and Gulledge, Jr., T. R. (1985). Use of the correlation coefficient with normal probability plots. *The American Statistician*, 39:75–79.