

# Several inequalities used for construction of the $\bar{X}$ and $S$ control charts with unequal sample sizes

Chanseok PARK (朴燦奭)

Applied Statistics Laboratory  
Department of Industrial Engineering  
Pusan National University

May 24, 2019

Hosted by [IEICE Technical Committee Conference on Reliability](#)



부산대학교  
PUSAN NATIONAL UNIVERSITY

# Overview

- 1 Basics on the Shewhart-type  $\bar{X}$  and  $S$  charts
- 2 Location parameter estimation
- 3 Scale parameter estimation
- 4 Inequalities and their proofs
- 5 Summary

# Overview

- 1 Basics on the Shewhart-type  $\bar{X}$  and  $S$  charts
- 2 Location parameter estimation
- 3 Scale parameter estimation
- 4 Inequalities and their proofs
- 5 Summary

# Overview

- 1 Basics on the Shewhart-type  $\bar{X}$  and  $S$  charts
- 2 Location parameter estimation
- 3 Scale parameter estimation
- 4 Inequalities and their proofs
- 5 Summary

# Overview

- 1 Basics on the Shewhart-type  $\bar{X}$  and  $S$  charts
- 2 Location parameter estimation
- 3 Scale parameter estimation
- 4 Inequalities and their proofs
- 5 Summary

# Overview

- 1 Basics on the Shewhart-type  $\bar{X}$  and  $S$  charts
- 2 Location parameter estimation
- 3 Scale parameter estimation
- 4 Inequalities and their proofs
- 5 Summary

# Shewhart-type $\bar{X}$ and $S$ charts

## Shewhart-type $\bar{X}$ and $S$ charts

In general, the Shewhart-type control charts are made up of

- **UCL**: the upper control limit
- **CL**: the center line
- **LCL**: the lower control limit

This control charts are widely used and well developed when sample sizes are **equal**.

- However, when sample sizes are **not equal**, there exist only *ad hoc* methods.
- For the location estimator, two estimators are used. Then when one is better?
- Especially for the scale estimator, they are all **biased**.

# Basics on the Shewhart-type charts

## Assumptions

- $m$  samples but each subgroup has a **different sample size**.
- Let  $X_{ij}$  denote the  $i$ th subgroup of size  $n_i$ , where  $i = 1, 2, \dots, m$  and  $j = 1, 2, \dots, n_i$ .
- Assume that  $X_{ij}$  are IID normal normal with mean  $\mu$  and variance  $\sigma^2$ .

## $\bar{X}$ chart

For the  $i$ th ( $i = 1, 2, \dots, m$ ) subgroup, we have

$$\frac{\bar{X}_i - E(\bar{X}_i)}{SE(\bar{X}_i)} \stackrel{\circ}{\sim} N(0, 1),$$

where

$$\bar{X}_i = \frac{1}{n_i} \sum_{j=1}^{n_i} X_{ij}.$$



## $\bar{X}$ chart — continued

Since  $E(\bar{X}_i) = \mu$  and  $SE(\bar{X}_i) = \sigma/n_i$ , the  $CL \pm 3 \cdot SE$  control limits are obtained by setting

$$\frac{\bar{X}_i - \mu}{\sigma/n_i} = \pm 3$$

and solving for  $\bar{X}_i$ .

Note that we can obtain the CI if we solve the above for  $\mu$ .

# Basics on the Shewhart-type charts

## $\bar{X}$ chart — continued

Solving for  $\bar{X}_i$ , the  $\bar{X}$  control chart is constructed as follows

$$\text{UCL} = \mu + \frac{3\sigma}{\sqrt{n_i}}$$

$$\text{CL} = \mu$$

$$\text{LCL} = \mu - \frac{3\sigma}{\sqrt{n_i}}.$$

## Issues

- The population parameters  $\mu$  and  $\sigma$  are **not** known in general.
- Especially, conventional *ad hoc* scale estimators for  $\sigma$  are **biased** when the sample sizes are **not** equal.

## S chart

Similarly, we also have

$$\frac{S_i - E(S_i)}{SE(S_i)} \sim N(0, 1),$$

where  $S_i$  is the sample standard deviation with a sample of size  $n_i$ .  
For the  $CL \pm 3 \cdot SE$  control limits, we can set up

$$\frac{S_i - E(S_i)}{SE(S_i)} = \pm 3,$$

where  $E(S_i) = c_4(n_i)\sigma$ ,  $\text{Var}(S_i) = (1 - c_4(n_i)^2)\sigma^2$ , and

$$c_4(n_i) = \sqrt{\frac{2}{n_i - 1}} \frac{\Gamma(n_i/2)}{\Gamma((n_i - 1)/2)}$$

# Basics on the Shewhart-type charts

## S chart — continued

Using  $E(S_i) = c_4(n_i)\sigma$  and  $\text{Var}(S_i) = (1 - c_4(n_i)^2)\sigma^2$ , we have

$$\frac{S_i - c_4(n_i)\sigma}{\sqrt{1 - c_4(n_i)^2}\sigma} = \pm 3.$$

Solving for  $S_i$ ,

$$\mathbf{UCL} = c_4(n_i)\sigma + 3\sqrt{1 - c_4(n_i)^2}\sigma$$

$$\mathbf{CL} = c_4(n_i)\sigma$$

$$\mathbf{LCL} = c_4(n_i)\sigma - 3\sqrt{1 - c_4(n_i)^2}\sigma.$$

## Issue again

- Especially, conventional *ad hoc* scale estimators for  $\sigma$  are **biased**.

# Location parameter estimation

## Two conventional estimators for $\mu$

Montgomery (2013) provided the following location estimators for  $\mu$  in Equations (6.2) and (6.30) of his book

- $$\bar{\bar{X}}_A = \frac{\bar{X}_1 + \bar{X}_2 + \cdots + \bar{X}_m}{m} = \frac{1}{m} \sum_{i=1}^m \bar{X}_i$$
- $$\bar{\bar{X}}_B = \frac{n_1 \bar{X}_1 + n_2 \bar{X}_2 + \cdots + n_m \bar{X}_m}{N} = \frac{1}{N} \sum_{i=1}^m n_i \bar{X}_i.$$

where  $N = \sum_{i=1}^m n_i$ .

Note that these estimators for  $\mu$  are unbiased while the *ad hoc* estimates for  $\sigma$  are **biased**.

## Issue

Which one of  $\bar{\bar{X}}_A$  or  $\bar{\bar{X}}_B$  is selected?

**The one with the smaller variance!**

## Theorem 1

If  $X_{ij}$  are normal with mean  $\mu$  and variance  $\sigma^2$ , then we have

$$\text{Var}(\bar{\bar{X}}_A) \geq \text{Var}(\bar{\bar{X}}_B).$$

## Proof.

It is immediate from  $\text{Var}(\bar{X}_i) = \sigma^2/n_i$  that we have

- $\text{Var}(\bar{\bar{X}}_A) = \frac{1}{m^2} \sum_{i=1}^m \text{Var}(\bar{X}_i) = \frac{\sigma^2}{m^2} \sum_{i=1}^m \frac{1}{n_i} = \frac{\sigma^2}{m} \frac{1}{\bar{n}_h}$
- $\text{Var}(\bar{\bar{X}}_B) = \frac{1}{N^2} \sum_{i=1}^m n_i^2 \text{Var}(\bar{X}_i) = \frac{\sigma^2}{N^2} \sum_{i=1}^m n_i = \frac{\sigma^2}{N} = \frac{\sigma^2}{m} \frac{1}{\bar{n}_a}$

where

$$\bar{n}_h = \left( \frac{1}{m} \sum_{i=0}^m n_i^{-1} \right)^{-1} \quad \text{and} \quad \bar{n}_a = \frac{1}{m} \sum_{i=1}^m n_i.$$

Then we have

$$\frac{\text{Var}(\bar{\bar{X}}_A)}{\text{Var}(\bar{\bar{X}}_B)} = \frac{\bar{n}_a}{\bar{n}_h}.$$

- $\bar{n}_h$  and  $\bar{n}_a$  are harmonic mean and arithmetic mean, respectively.
- $\bar{n}_a \geq \bar{n}_h$ .
- Thus,  $\text{Var}(\bar{\bar{X}}_A)/\text{Var}(\bar{\bar{X}}_B) \geq 1$  which results in  $\text{Var}(\bar{\bar{X}}_A) \geq \text{Var}(\bar{\bar{X}}_B)$ .



Which one of  $\bar{\bar{X}}_A$  or  $\bar{\bar{X}}_B$  is selected?

**$\bar{\bar{X}}_B$ !**

## $\bar{X}$ chart — Recalled

Substituting  $\mu = \bar{\bar{X}}_B$ , we have

$$\mathbf{UCL} = \bar{\bar{X}}_B + \frac{3\sigma}{\sqrt{n_i}}$$

$$\mathbf{CL} = \bar{\bar{X}}_B$$

$$\mathbf{LCL} = \bar{\bar{X}}_B - \frac{3\sigma}{\sqrt{n_i}}.$$

Thus, we need to estimate  $\sigma$ .



# Scale parameter estimation

Section 6.3.2 of Montgomery (2013) recommended the followings to estimate  $\sigma$ :

- $S_p$   
where  $S_p^2 = \sum_{i=1}^m (n_i - 1) S_i^2 / (N - m)$  (pooled sample variance).
- $\bar{S}^*$   
where  $\bar{S}^* = \bar{S} / c_4(\bar{n})$ ,  $\bar{S} = \sum_{i=1}^m S_i / m$  and  $\bar{n} = \sum_{i=1}^m n_i / m$ .

## Issues

- Are they unbiased?
- Which one is better?

# Scale parameter estimation

Note that  $S_p^2$  is unbiased for  $\sigma^2$  so  $S_p$  should underestimate  $\sigma$ . This is easily proved using the Jensen's inequality as follows.

Since  $\sqrt{T}$  is concave, we have

$$E(\sqrt{T}) < \sqrt{E(T)}.$$

With  $T = S_p^2$ , we have

$$E(\sqrt{S_p^2}) = \mathbf{E}(S_p) < \sqrt{E(S_p^2)} = \sqrt{\sigma^2} = \sigma \implies E(S_p) < \sigma.$$

- $S_p$  always **underestimate**  $\sigma$ ,
- so it is also **biased**.

Note: whenever  $S^2$  is unbiased for  $\sigma^2$ ,  $S$  underestimates  $\sigma$  regardless of an underlying distribution.

# Scale parameter estimation

Then how about  $\bar{S}^*$ ?

Recall  $\bar{S}^*$

$$\bar{S}^* = \frac{\bar{S}}{c_4(\bar{n})},$$

where  $\bar{S} = \sum_{i=1}^m S_i/m$  and  $\bar{n} = \sum_{i=1}^m n_i/m$ .

It **underestimates**  $\sigma$  since

$$E[\bar{S}^*] = \frac{\frac{1}{m} \sum_{i=1}^m E[S_i]}{c_4(\bar{n})\sigma} = \frac{\frac{1}{m} \sum_{i=1}^m c_4(n_i)}{c_4(\bar{n})} \cdot \sigma < \sigma.$$

How to prove the above? The claim is to show (the proof is tricky)

$$\frac{1}{m} \sum_{i=1}^m c_4(n_i) < c_4(\bar{n}) \quad (\text{the proof will be provided later})$$

# Scale parameter estimation

As aforementioned, both  $S_p$  and  $\bar{S}^*$  underestimate  $\sigma$  — **biased!**

We propose

- $$\bar{S}_A = \frac{S_1/c_4(n_1) + S_2/c_4(n_2) + \cdots + S_m/c_4(n_m)}{m} = \frac{1}{m} \sum_{i=1}^m \frac{S_i}{c_4(n_i)}$$
- $$\bar{S}_B = \frac{S_1 + S_2 + \cdots + S_m}{c_4(n_1) + c_4(n_2) + \cdots + c_4(n_m)} = \frac{\sum_{i=1}^m S_i}{\sum_{i=1}^m c_4(n_i)}$$

Using  $E(S_i) = c_4(n_i)\sigma$  under the normality assumption, we see that these are **unbiased** for  $\sigma$ .

## Issue

Which one is used?

That is, which one has a smaller variance?

**Answer:**  $\bar{S}_B$  (the proof will be provided later).

# Scale parameter estimation

It is immediate upon using

$$\text{Var}(S_i) = E(S_i^2) - E(S_i)^2 = \sigma^2 \{1 - c_4(n_i)^2\}$$

that we have

$$\text{Var}(\bar{S}_A) = \frac{\sigma^2}{m^2} \sum_{i=1}^m \left\{ \frac{1}{c_4(n_i)^2} - 1 \right\}$$

and

$$\text{Var}(\bar{S}_B) = \sigma^2 \cdot \frac{\sum_{i=1}^m \{1 - c_4(n_i)^2\}}{\left\{ \sum_{i=1}^m c_4(n_i) \right\}^2}.$$

In the following section, we will provide several inequalities which are needed for proving it.

# Inequalities and their proofs

In the previous sections, the followings are left to prove

- $\text{Var}(\bar{S}_A) \geq \text{Var}(\bar{S}_B)$

- $E[\bar{S}^*] = \frac{\frac{1}{m} \sum_{i=1}^m E[S_i]}{c_4(\bar{n})\sigma} = \frac{\frac{1}{m} \sum_{i=1}^m c_4(n_i)}{c_4(\bar{n})} \cdot \sigma < \sigma$

Note:  $\bar{S}^*$  is recommended for the estimator of  $\sigma$  in Section 6.3.2 of Montgomery (2013).

- To prove the above, it is essential to figure out the behaviors of the  $c_4$  function which is again given by

$$c_4(x) = \sqrt{\frac{2}{(x-1)}} \frac{\Gamma(x/2)}{\Gamma((x-1)/2)}$$

- To this end, we will use the Wallis (1656) and Watson (1959) representations.

# Inequalities and their proofs

Wallis' production formula (Wallis, 1656)

$$\frac{2}{1} \cdot \frac{4}{3} \cdot \frac{6}{5} \cdot \frac{8}{7} \cdot \frac{10}{9} \cdot \frac{12}{11} \cdots \rightarrow \infty \text{ (diverges to } \infty \text{)}$$

$$\times \frac{2}{3} \cdot \frac{4}{5} \cdot \frac{6}{7} \cdot \frac{8}{9} \cdot \frac{10}{11} \cdot \frac{12}{13} \cdots \rightarrow 0 \text{ (trivially goes to } 0 \text{)}$$

---

$$= \frac{2}{1} \cdot \frac{2}{3} \cdot \frac{4}{3} \cdot \frac{4}{5} \cdot \frac{6}{5} \cdot \frac{6}{7} \cdot \frac{8}{7} \cdot \frac{8}{9} \cdot \frac{10}{9} \cdot \frac{10}{11} \cdots \rightarrow \frac{\pi}{2}$$



Another version is also given by

$$\frac{1}{\sqrt{\pi(n + \frac{1}{2})}} < \boxed{\frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdots \frac{2n-1}{2n}} < \frac{1}{\sqrt{\pi n}}$$

# Inequalities and their proofs

The Wallis' production formula can also be rewritten as

$$\boxed{\frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdots \frac{2n-1}{2n}} = \frac{1}{\sqrt{\pi(n+\theta)}},$$

where and  $\boxed{n \geq 1}$  and  $0 < \theta < \frac{1}{2}$ .

This formula will play a pivotal role in proving the inequalities in this talk.

Note: it is provided in his book, *Arithmetica Infinitorum* in the year of 1656, written in Latin. Recently, Stedall (2004) translated into English.

- Extension of the Wallis' production formula by Watson (1959)
- $c_4(x)$  function written by the Watson's  $\theta(x)$  function
- Using the Watson's  $\theta(x)$ , we can provide several useful results



# Inequalities and their proofs

Recall

$$\boxed{\frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdots \frac{2n-1}{2n}} = \frac{1}{\sqrt{\pi(n+\theta)}},$$

where and  $\boxed{n \geq 1}$  and  $0 < \theta < \frac{1}{2}$ . We have

$$\begin{aligned} \frac{\Gamma(n + \frac{1}{2})}{\Gamma(n + 1)} &= \frac{\sqrt{\pi} \cdot \frac{1}{2} \cdot \frac{1 + \frac{1}{2}}{2} \cdot \frac{2 + \frac{1}{2}}{3} \cdots \frac{n - \frac{1}{2}}{n}}{1} \\ &= \sqrt{\pi} \cdot \boxed{\frac{1}{2} \cdot \frac{3}{4} \cdot \frac{5}{6} \cdots \frac{2n-1}{2n}} = \frac{1}{\sqrt{n+\theta}} \end{aligned}$$

Motivated by the above, Watson (1959) extended to

$$\frac{\Gamma(x + \frac{1}{2})}{\Gamma(x + 1)} = \frac{1}{\sqrt{x + \theta(x)}}, \quad (1)$$

where  $\boxed{x > -\frac{1}{2}}$  and  $0 < \theta(x) < \frac{1}{2}$ .

# Inequalities and their proofs

Solving (1) for  $\theta(x)$ , we have

$$\theta(x) = -x + \frac{\Gamma(x+1)^2}{\Gamma(x+\frac{1}{2})^2} = -x + x \frac{\Gamma(x)\Gamma(x+1)}{\Gamma(x+\frac{1}{2})^2}. \quad (2)$$

Since

$$c_4(x)^2 = \frac{1}{(x-1)/2} \frac{\Gamma(x/2)^2}{\Gamma((x-1)/2)^2} = \frac{\Gamma(x/2)^2}{\Gamma((x-1)/2)\Gamma((x+1)/2)},$$

we have

$$c_4(2x+1)^2 = \frac{\Gamma(x+\frac{1}{2})^2}{\Gamma(x)\Gamma(x+1)}. \quad (3)$$

Substituting (3) into (2), we have

$$\boxed{\theta(x) = -x + x \cdot \frac{1}{c_4(2x+1)^2}} \quad \text{or} \quad \boxed{c_4(2x+1) = \sqrt{\frac{x}{\theta(x) + x}}}$$

# Inequalities and their proofs

- We succeeded in rewriting  $c_4$  function using the Watson's  $\theta(x)$  function (extended version of Wallis product).
- Using the nice properties of  $\theta(x)$ , we can get several useful results easily.

## Properties of $\theta(x)$

- $0 < \theta < \frac{1}{2}$ .
- $\theta(x)$  is monotonically decreasing.

# Inequalities and their proofs

## Lemma 1

$$\bullet \sqrt{\frac{x-1}{x}} < c_4(x) < 1$$

## Proof.

Recall  $\theta(x) = -x + x/c_4(2x+1)^2$  and  $0 < \theta(x) < \frac{1}{2}$ . Let  $x = (t-1)/2$ . Then we have

$$0 < -x + \frac{x}{c_4(2x+1)^2} < \frac{1}{2} \implies 0 < -\left(\frac{t-1}{2}\right) + \frac{(t-1)/2}{c_4(t)^2} < \frac{1}{2}.$$

Note:  $c_4(t)$  is defined on  $t > 1$ . Solving the above for  $c_4(t)^2$ , we have

$$\frac{t-1}{t} < c_4(t)^2 < 1.$$



## Lemma 2

- $c_4(x)$  is monotonically increasing
- The function

$$\frac{1}{c_4(x)^2} - 1$$

is monotonically decreasing.

## Proof.

Proofs are omitted. Please refer to the manuscript. □

# Inequalities and their proofs

- Recall: we claim  $\text{Var}(\bar{S}_A) \geq \text{Var}(\bar{S}_B)$ .
- To prove this, we need Lemma 3 below.

## Lemma 3 (Chebyshev's sum inequality)

If  $a_1 \geq a_2 \geq \cdots \geq a_m$  and  $b_1 \geq b_2 \geq \cdots \geq b_m$ , then

$$m \sum_{i=1}^m a_i b_i \geq \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i.$$

Similarly, if  $a_1 \leq a_2 \leq \cdots \leq a_m$  and  $b_1 \geq b_2 \geq \cdots \geq b_m$ , then

$$m \sum_{i=1}^m a_i b_i \leq \sum_{i=1}^m a_i \cdot \sum_{i=1}^m b_i.$$

- The original formula is in an integral form (Chebyshev, 1882).
- Its proof can be found in Chebyshev (1883).

# Inequalities and their proofs

## Proof of Chebyshev's sum inequality.

When  $a_i$  and  $b_i$  are both decreasing,  $(a_i - a_j)$  and  $(b_i - b_j)$  have the **same sign** for any  $i$  and  $j$ . Thus, we have

$$\sum_{i=1}^m \sum_{j=1}^m (a_i - a_j)(b_i - b_j) \geq 0,$$

which results in

$$2m \sum_{i=1}^m a_i b_i - 2 \sum_{i=1}^m a_i \sum_{j=1}^m b_j \geq 0.$$

Next, when  $a_i$  are increasing and  $b_i$  are decreasing,  $(a_i - a_j)$  and  $(b_i - b_j)$  have **different signs** or zero for any  $i$  and  $j$ . Thus, we have

$$\sum_{i=1}^m \sum_{j=1}^m (a_i - a_j)(b_i - b_j) \leq 0.$$



# Inequalities and their proofs

Recall: we propose

- $$\bar{S}_A = \frac{S_1/c_4(n_1) + S_2/c_4(n_2) + \cdots + S_m/c_4(n_m)}{m} = \frac{1}{m} \sum_{i=1}^m \frac{S_i}{c_4(n_i)}$$
- $$\bar{S}_B = \frac{S_1 + S_2 + \cdots + S_m}{c_4(n_1) + c_4(n_2) + \cdots + c_4(n_m)} = \frac{\sum_{i=1}^m S_i}{\sum_{i=1}^m c_4(n_i)}$$

## Theorem 2

We have  $\text{Var}(\bar{S}_A) \geq \text{Var}(\bar{S}_B)$ .

Recall: we showed

$$\text{Var}(\bar{S}_A) = \frac{\sigma^2}{m^2} \sum_{i=1}^m \left\{ \frac{1}{c_4(n_i)^2} - 1 \right\} \quad \text{and} \quad \text{Var}(\bar{S}_B) = \sigma^2 \cdot \frac{\sum_{i=1}^m \{1 - c_4(n_i)^2\}}{\left\{ \sum_{i=1}^m c_4(n_i) \right\}^2}$$



# Inequalities and their proofs

## Sketch proof.

It suffices to show

$$\frac{1}{m^2} \sum_{i=1}^m \left\{ \frac{1}{c_4(n_i)^2} - 1 \right\} \geq \frac{\sum_{i=1}^m \{1 - c_4(n_i)^2\}}{\left\{ \sum_{i=1}^m c_4(n_i) \right\}^2}$$

- First, let  $a_i = c_4(n_i)$  (increasing) and  $b_i = 1/c_4(n_i)^2 - 1$  (decreasing)
- Apply the Chebyshev's sum inequality with  $a_i$  and  $b_i$ . Then we have an intermediate result.
- Next, let  $a_i = c_4(n_i)$  (increasing) and  $b_i = 1/c_4(n_i) - c_4(n_i)$  (decreasing)
- Apply the Chebyshev's sum inequality again to the above intermediate result with  $a_i$  and  $b_i$ . Then we have a result.



# Inequalities and their proofs

## Recall

Section 6.3.2 of Montgomery (2013) uses

$$\bar{S}^* = \frac{\bar{S}}{c_4(\bar{n})},$$

where  $\bar{S} = \sum_{i=1}^m S_i/m$  and  $\bar{n} = \sum_{i=1}^m n_i/m$ .

Claim:  $\bar{S}^*$  **underestimates**  $\sigma$  since

$$E[\bar{S}^*] = \frac{\frac{1}{m} \sum_{i=1}^m E[S_i]}{c_4(\bar{n})\sigma} = \frac{\frac{1}{m} \sum_{i=1}^m c_4(n_i)}{c_4(\bar{n})} \cdot \sigma < \sigma.$$

It suffices to show

$$\frac{1}{m} \sum_{i=1}^m c_4(n_i) < c_4(\bar{n})$$

# Inequalities and their proofs

## Lemma 4

The  $c_4(x)$  function is log-concave.

## Sketch proof.

Recall the  $c_4$  function which is again given by

$$c_4(x) = \sqrt{\frac{2}{(x-1)} \frac{\Gamma(x/2)}{\Gamma((x-1)/2)}}$$

Thus, we have

$$c_4(x)^2 = \frac{1}{(x-1)/2} \frac{\Gamma(x/2)^2}{\Gamma((x-1)/2)^2} = \frac{\Gamma(x/2)^2}{\Gamma((x-1)/2)\Gamma((x+1)/2)}.$$

## Sketch proof – continued.

- Taking the logarithm and dividing by two, we have

$$\log c_4(x) = \log \Gamma\left(\frac{x}{2}\right) - \frac{1}{2} \log \Gamma\left(\frac{x-1}{2}\right) - \frac{1}{2} \log \Gamma\left(\frac{x+1}{2}\right).$$

- From Section 11.14 (iv) of Schilling (2005) and Merkle (1996), the second derivative of  $\log \Gamma(x)$  can be written as the sum of the series so that we have

$$\frac{d^2}{dx^2} \log \Gamma(x) = \sum_{k=0}^{\infty} \frac{1}{(x+k)^2}. \quad (4)$$

## Sketch proof — continued.

After tedious algebra, we have

$$\begin{aligned}\frac{d^2}{dx^2} \log c_4(x) &= \sum_{k=0}^{\infty} \frac{1}{(\frac{x}{2} + k)^2} - \frac{1}{2} \sum_{k=0}^{\infty} \frac{1}{(\frac{x-1}{2} + k)^2} - \frac{1}{2} \sum_{k=0}^{\infty} \frac{1}{(\frac{x+1}{2} + k)^2} \\ &= \sum_{k=0}^{\infty} \frac{-12(x+2k)^2 + 4}{(x+2k)^2(x-1+2k)^2(x+1+2k)^2} < 0.\end{aligned}$$

- Thus, we have  $\frac{d^2}{dx^2} (-\log c_4(x)) > 0$  so  $-\log c_4(x)$  is convex due to Theorem 6.4.6 of Bartle and Sherbert (2011).
- Then  $\log c_4(x)$  is concave and  $c_4(x)$  is thus log-concave, which completes the proof.



# Inequalities and their proofs

## Lemma 5

We have the following inequality

$$\frac{1}{m} \sum_{i=1}^m c_4(n_i) \leq c_4(\bar{n}).$$

## Proof.

The log-concavity of  $c_4(x)$  from Lemma 4 guarantees that  $c_4(x)$  is concave.

Thus, we can apply the Jensen's inequality to  $c_4(x)$  and we have

$$\frac{1}{m} \sum_{i=1}^m c_4(n_i) \leq c_4(\bar{n}),$$

where  $\bar{n} = \sum_{i=1}^m n_i / m$ . This completes the proof. □

## Summary

- The  $c_4$  function plays a pivotal role in constructing the  $\bar{X}$  and  $S$  charts.
- With the useful inequalities, we showed that the conventional *ad hoc* methods are **biased**, which are used for the  $\bar{X}$  and  $S$  charts.
- We also proposed new **unbiased** estimators for  $\sigma$  which can be used for various control charts.
- Thus, we can construct **proper  $\bar{X}$  and  $S$  charts** based on the new **unbiased** estimators.
- The behaviors of the  $c_4$  function are much understood thanks to the connection with Wallis' product and Watson representation.
- Hopefully, these results on  $c_4$  can produce more useful inequalities.

# References

- Bartle, R. G. and Sherbert, D. R. (2011). Introduction to Real Analysis. Wiley, 4th edition.
- Chebyshev, P. L. (1882). Sur les expressions approximatives des intégrales définies par les autres prises entre les même limites. In Markov, A. A. and Sonin, N., editors, Oeuvres de P. L. Tchebychef I–II, Vol. 2, pages 716–719. Imprimerie de l'Academie imperiale des sciences, St. Petersburg.
- Chebyshev, P. L. (1883). Sur une série qui fournit les valeurs extrêmes des intégrales, lorsque la fonction sous le signe est décomposée en deux facteurs. In Markov, A. A. and Sonin, N., editors, Oeuvres de P. L. Tchebychef I–II, Vol. 2, pages 405–419. Imprimerie de l'Academie imperiale des sciences, St. Petersburg.
- Merkle, M. (1996). Logarithmic convexity and inequalities for the gamma function. Journal of Mathematical Analysis and Applications, 203:369–380.



# References

- Montgomery, D. C. (2013). Statistical Quality Control: An Modern Introduction. John Wiley & Sons, Singapore, 7th edition.
- Schilling, R. L. (2005). Measures, Integrals and Martingales. Cambridge University Press, Cambridge, UK.
- Stedall, J. A. (2004). Arithmetica Infinitorum: John Wallis 1656. Springer, New York. (English translation from the original book written in Latin).
- Wallis, J. (1656). Arithmetica Infinitorum. Oxford, England.
- Watson, G. N. (1959). A note on gamma functions. Edinburgh Mathematical Notes, 42:7–9.