

Matrix functions and stability

Optimal uniform growth after inversion of an exponentially stable generator

Difficulty: challenging **Importance:** interesting to the community**Status:** [Lean verified](#)

Rating rationale: Challenging because matching growth bounds require uniform control over arbitrary Hilbert-space generators; community impact includes semigroup stability and rational time discretization.

Provenance: explicit fixed-bound formalization of the source's growth question.

Resolution — 30 September 2026

Part VI of the [submitted manuscript](#), Theorem S.1.1 (printed p. 42; PDF page 43), states the matching fixed-bound asymptotic

$$c_M L(t)^{\alpha(M)} \leq G_M(t) \leq C_M L(t)^{\alpha(M)} \quad (t \geq t_M),$$

where

$$L(t) = \log \log(t + e^e), \quad \alpha(M) = \frac{2}{\pi} \arccos(M^{-1}).$$

For every fixed $M > 1$, the constants $c_M, C_M, t_M > 0$ depend only on M . The upper bound covers all strongly continuous semigroups in the original class, including unbounded generators. The lower bound has finite-dimensional witnesses with the same prescribed bound M ; the witnesses may depend on t . The formalization also proves $G_1(t) = 1$ for all $t \geq 0$.

This determines the asymptotic order requested below, with the exact fixed- M exponent. It does not assert a limiting leading coefficient or uniform constants as $M \downarrow 1$. Part VI, Sections S.2–S.8, supplies the proposed upper and lower arguments; Section S.9 treats the endpoint. The manuscript is retained unchanged, including material outside this submission's sharp-rate claim.

The [accompanying Lean source](#) contains the declarations `ProofProject.sharp_growth`, `ProofProject.finite_dimensional_lower`, `ProofProject.contractive_envelope`, `ProofProject.attainableNorms_bddAbove`, and `ProofProject.stableSemigroup_hasGeneratorInverse`. Its definitions represent the full strong generator graph and quantify over arbitrary complete complex Hilbert spaces in a fixed universe. The supplied project specifies Lean 4.33.1 and Mathlib v4.33.1. The [statement correspondence](#) and [reproduction instructions](#) describe the five contracts.

Lean proof and verification evidence

Lean verified, 30 September 2026. All five declarations above passed a fresh non-root Linux Comparator run, including statement identity, transitive permitted-axiom checks and Lean's default-kernel replay. The complete harness exited successfully after all sandbox and rejection controls and cleanup. The [verification report](#), [successful run receipt](#), [Comparator log](#), and [driver log](#) retain the actual local execution evidence.

The immutable [proof-source snapshot](#) is commit `ac582d01532a914538cbfb74863437b18433038a`. It uses Lean 4.33.1 and Mathlib `0df444a360eaa60ab8c11dca51a86af692955474`, with all other dependencies fixed in `lean/lake-manifest.json`. This submission is a local PR draft: the commit link becomes publicly accessible when that branch is published. No remote repository or hosted CI was modified.

The inverse and envelope-finiteness contracts cover every semigroup in the original class for $M \geq 1$ (and nonnegative time for the envelope). The sharp-growth and finite-witness contracts cover every fixed

$M > 1$; the endpoint contract covers $M = 1$ and every $t \geq 0$. The [statement correspondence](#) expands the definitions, quantifiers and conclusions, and identifies the adapted scalar lower proof and refined constants. There are no unproved extra premises or replaceable definition holes. The [five transitive axiom reports](#) contain exactly *propext*, *Classical.choice* and *Quot.sound*.

Two independent retrospective AI statement reviews approve all five contracts; separate upper and lower source reviews approve their inspected proof paths. The [reports](#) record their coverage and accepted corrections. These are AI source reviews, not external human peer review.

On Linux with the [checker prerequisites](#), run from the repository root at the recorded source revision:

```
python3 tools/lean/harness.py bootstrap /tmp/mf17-tools
python3 tools/lean/harness.py verify \
  matrix-functions-and-stability/MF-17/lean /tmp/mf17-tools
```

The [project instructions](#) also reproduce the ordinary build, independent target-type check and axiom report. The original statement and historical checks remain below; the ratings describe the original question. This date records review and verification of the submitted solution, not a new comprehensive literature search.

Original problem statement

For each fixed real number $M > 1$, let A_M consist of all pairs (H, A) such that H is a complex Hilbert space and A is the generator of a strongly continuous semigroup $T(s)$ on H satisfying

$$\|T(s)\|_{\mathcal{L}(H)} \leq Me^{-s} \quad (s \geq 0).$$

Here A may be unbounded and is understood on its dense domain. Exponential stability implies that 0 belongs to the resolvent set of A , so A^{-1} is bounded on H and its exponential is defined by the operator-norm convergent power series.

Determine, for each fixed $M > 1$, the asymptotic order as $t \rightarrow \infty$ of

$$G_M(t) = \sup_{(H,A) \in A_M} \|\exp(tA^{-1})\|_{\mathcal{L}(H)}.$$

An asymptotic order means a rate $g_M(t)$ with matching positive upper and lower multiplicative bounds for all sufficiently large t ; their constants may depend on M , but must be uniform over $(H, A) \in A_M$.

The source's Problem 1.3 asks for the optimal inverse-semigroup growth for exponentially stable Hilbert-space semigroups. The envelope above makes the dependence on the original semigroup bound explicit. This formalization is supported by uniform upper estimates and examples in the same normalized class, rather than attributed verbatim to the source. The problem concerns operator stability and rational time-stepping analysis, extending finite-dimensional matrix exponential questions.

References

1. E. Lorist, M. Meyries, and M. Veraar, *A solution to the inverse generator problem and related questions*, arXiv:2608.06272v3 (2026), Problem 1.3, Theorem 1.2(2), and Section 1.3 on numerical stability. [Paper](#).
2. C. J. K. Batty, A. Gomilko, and Y. Tomilov, *A Besov algebra calculus for generators of operator semigroups and related norm-estimates*, *Mathematische Annalen* 379 (2021), pp. 23–93, Corollary 5.7: explicit constants depending only on the semigroup bound and decay rate. Their generator notation is $-A$. [Published paper](#).
3. H. Zwart, *Growth Estimates for $\exp(A^{-1}t)$ on a Hilbert Space*, *Semigroup Forum* 74 (2007), pp. 487–494, Theorem 2.2 and Corollary 2.3: the logarithmic upper bound. [Published paper](#).

Status check — 2026-09-08

checked the current arXiv history, with latest version v3 dated 2026-08-26 and no withdrawal, and searched “inverse generator Problem 1.3” and “inverse generator growth 2026 logarithmic”. Corollary 5.7 of reference 2 gives, in the present sign convention and at decay rate one,

$$G_M(t) \leq 2M^2(2 - e^{-1} + e^{-1} \log t), \quad t > 1.$$

Theorem 1.2(2) of reference 1 gives examples with a doubly logarithmic lower bound raised to a positive exponent, while the original semigroup bound is $1 + C\alpha/(1 - \alpha)$ for an absolute C and $0 < \alpha < 1$. Thus its parameters allow examples within each fixed $M > 1$ class. These estimates do not determine the matching asymptotic order. The older yes/no inverse-generator problem was resolved negatively in the 2026 paper and is not listed as open here.

Audit — 2026-09-10

Rechecked [Lorist–Meyries–Veraar, Problem 1.3 and Theorem 1.2](#). The sharp growth question survives their negative inverse-generator theorem; logarithmic upper and doubly logarithmic lower bounds do not match. Title and inverse-semigroup-growth searches found no improvement resolving the fixed- M envelope. Difficulty was raised to reflect the operator-theoretic obstruction.