Exponential ergodicity and the space–time Gibbs states of one-dimensional probabilistic cellular automata

Seth Douglas Nidhal Mghirbi

Version 1.1.0, November 2026

Abstract

A positive-rate probabilistic cellular automaton (PCA) defines a Gibbs specification in one more dimension, its space–time interaction, in which the future is a boundary like the past and the sides. For finite-alphabet, finite-radius, strictly positive PCA on $\mathbb{Z}$ we prove that uniform exponential ergodicity of the dynamics is equivalent to exponential weak spatial mixing of this specification in arbitrary finite volumes of $\mathbb{Z}^2$ under arbitrary exterior conditions, so that the specification has exactly one DLR state, the stationary trajectory law, with exponentially decaying correlations. This answers in one dimension, under exponential ergodicity, the question of Lebowitz, Maes and Speer whether every Gibbs measure of the space–time interaction is a trajectory law of the PCA, previously known for periodic measures. In every dimension and with no rate, the relative-entropy defect of any Gibbs measure from the dynamics is a surface term; hence for an ergodic PCA the stationary trajectory is the only uniformly recurrent Gibbs measure, the translation action on the Gibbs simplex is uniquely ergodic, and all-DLR uniqueness holds exactly when every Gibbs measure is uniformly recurrent. The forward proof removes the far future boundary from Bérard's coupling from the past at an entropy cost that only one-dimensional geometry absorbs. Finite certificates for the hypothesis are exactly computable; for the noisy majority rule they give a unique space–time Gibbs state down to noise 0.485, including noise one half, and Marsan, Sablik and Törmä's fixed radius-one automaton has one space–time Gibbs state at small noise (granting structural estimates from their low-noise proof) and at large noise, and several in between.

AI systems assisted with this work. The authors take responsibility for its claims; see "How this work was done." This preprint has not been peer reviewed.

Programs and finite checks:
https://github.com/Apsiape/exponential-ergodicity-space-time-gibbs.

Contents

1 Introduction **3**
1.1 A dynamics is an equilibrium system in one more dimension 3
1.2 Results 4
1.3 Why the future boundary is the difficulty 5
1.4 Relation to prior work 7
1.5 Reading guide 8

2 Setting **8**
2.1 Probabilistic cellular automata 8
2.2 Random-map representations 9
2.3 The space–time specification 10
2.4 Trajectories are DLR states 10

3 The main theorem **11**

4 Finite certificates for the dynamical hypothesis **11**

5 The coupling input and its finite-volume adapter **13**
5.1 Bérard's local coupling . . . 14
5.2 Clipping a tile sampler to a finite volume . . . 14

6 The cap comparison **15**
6.1 Coarse cells and bad markers . . . 15
6.2 Expanding volumes and the cap identity . . . 16
6.3 The deterministic splice . . . 17
6.4 Resampling at a fixed cluster . . . 18
6.5 Cluster entropy and the harmonic gain . . . 18
6.6 Common extinction and the root comparison . . . 19

7 Every DLR state is the stationary trajectory **20**

8 Weak mixing in arbitrary volumes, and clustering **20**
8.1 Whole-tile roots . . . 21
8.2 The joint coupling . . . 21
8.3 Clustering . . . 23
8.4 Return to arbitrary radius . . . 23

9 Converses in every dimension **24**

10 Every dimension, no rate: the defect bound **25**
10.1 The causal defect and the surface bound . . . 25
10.2 Translation-invariant, periodic and uniformly recurrent states . . . 26
10.3 Ergodic PCA . . . 27

11 Examples **29**
11.1 Any rule at high noise . . . 29
11.2 The noisy shift . . . 30
11.3 The noisy majority rule: exact certificates . . . 30
11.4 Two further sources of the hypothesis . . . 31
11.5 A fixed reentrant family: the Marsan–Sablik–Törmä automaton . . . 32

12 Consequences in equilibrium language **34**

13 Discussion and open problems **37**

Credit **38**

How this work was done **39**

Verification status **39**

A Reproducibility **40**

References **40**

1 Introduction

1.1 A dynamics is an equilibrium system in one more dimension

A probabilistic cellular automaton on $\mathbb{Z}^d$ updates every site of a row independently, each site reading a finite neighbourhood of the preceding row through one fixed transition table $p(a \mid u)$. Run it forever and record every row: the result is a random configuration on $\mathbb{Z}^d \times \mathbb{Z}$. Multiply the transition probabilities of the sites in a finite space–time volume: the product is a Boltzmann weight for the interaction

$$\Phi(\sigma) \;=\; -\log p\big(\sigma_{x,t} \,\big|\, \sigma_{x+[-r,r],\,t-1}\big),$$

translated over the lattice. When every entry of the table is positive this is a finite-range, translation-invariant interaction on $\mathbb{Z}^{d+1}$ in the usual sense, and it comes with the usual specification: the conditional law of a finite volume given its exterior is the normalized product of every translated factor that meets the volume. This specification does not know that time runs upward. It conditions on the future rows above the volume exactly as it conditions on the past rows below it and on the sides.

The correspondence between the dynamics and this equilibrium system is classical in the translation-invariant setting. Goldstein, Kuik, Lebowitz and Maes [9] identify the stationary laws of the PCA that are invariant under spatial translations with the Gibbs measures of Φ that are invariant under every space–time translation; Lebowitz, Maes and Speer [15] develop the statistical mechanics of PCA on this basis. Their identification says nothing about Gibbs measures that are not translation invariant, and in particular nothing about the finite-volume conditional laws with an arbitrary future boundary, which are the objects one must control to prove that a specification has only one Gibbs state.

This paper proves that in one spatial dimension the dynamics controls every boundary, and in every dimension it bounds how far any Gibbs measure can be from the dynamics.

What is settled. Two of the open problems listed by Lebowitz, Maes and Speer [15, Section 6.6] concern the questions treated here, and a third question was raised by Marcovici, Sablik and Taati [19, Section 2.2].

- *Is every Gibbs measure of the space–time interaction a Markov chain measure of the PCA?* (LMS, Problem 2; known for translation-invariant and periodic measures [9].) Yes in one dimension under uniform exponential ergodicity, for every Gibbs measure and every boundary (Theorem 3.1); yes in every dimension for every uniformly recurrent Gibbs measure (Theorem 10.6). What remains open is a Gibbs measure that is not uniformly recurrent: for an ergodic PCA in $d \geq 2$ (Corollary 10.12, Open problem 13.1), for an ergodic PCA on $\mathbb{Z}$ without an exponential rate, if one exists (Remark 10.13, Open problem 13.2), and for a non-ergodic PCA in any dimension.
- *Is there a family of finite-volume conditions, like the Dobrushin–Shlosman conditions C_V but involving only the dependence on spins in the past of V, that characterizes the uniqueness region completely?* (LMS, Problem 4.) On the dynamical side Maes and Shlosman [17] introduced the conditions $C_{N,R}$ on finite space–time trapezoids and proved that they imply exponential ergodicity. For positive PCA in one dimension these conditions are also necessary under (2), by Proposition 12.2, which uses Bérard's uniform triangular coupling [2, Lemma 8]; Bérard's finite-size criterion [2, eq. (5)] is another characterization. What this paper adds is the equilibrium side: in one dimension the region of all-DLR

uniqueness with exponential mixing is exactly the region of exponential ergodicity of the dynamics (Theorem 3.1), so these dynamical conditions characterize it (Corollary 12.3); it contains an open set of positive weights on the fixed supports (Theorem 12.1), and Section 4 gives elementary sufficient conditions that are exactly computable. In $d = 1$ the question left is whether uniqueness can hold with sub-exponential ergodicity (Open problem 13.2); in $d \geq 2$ the forward implication is open (Open problem 13.1), and necessity of $C_{N,R}$ is not established here.

- *Is every ergodic PCA uniformly ergodic?* (Marcovici, Sablik and Taati's conjecture [19, Section 2.2].) For positive rates, yes, by Halawi's Baire-category argument [11], which is not ours; we reproduce it (Proposition 10.7) because it is what lets the dimension-free results rest on bare ergodicity.

What is new beyond that. The equivalence of exponential ergodicity with exponential weak mixing under arbitrary future boundaries, in one dimension (Theorem 3.1); the surface bound on the relative-entropy defect of any Gibbs measure (Theorem 10.3); unique ergodicity of the translation action on the Gibbs simplex of an ergodic PCA, with the stationary trajectory as its only uniformly recurrent point (Theorem 10.10); the characterization of all-DLR uniqueness as ergodicity plus uniform recurrence of every Gibbs measure (Corollary 10.12); the placement of ergodicity of rational positive PCA at Π^0_2 (Remark 10.9); the exact certificates for the noisy majority rule at and below noise one half (Section 11.3); and the reentrant fixed family (Section 11.5), conditional on the structural inputs of [22]. We also show that in one dimension the Dobrushin–Shlosman block conditions for the space–time specification and the past-only conditions $C_{N,R}$ of Maes and Shlosman hold in the same region of positive PCA tables (Corollary 12.3); the direction from the block conditions to $C_{N,R}$ rests on Proposition 12.2, which derives $C_{N,R}$ from uniform exponential ergodicity using Bérard's triangular coupling [2, Lemma 8].

1.2 Results

Throughout, a PCA has a finite alphabet $\mathcal{A}$, a finite radius r, and a strictly positive table $p(a \mid u) > 0$. The precise definitions are in Section 2; the two properties compared are the following.

- *Uniform exponential ergodicity*: there are a stationary row law μ and constants $C, c > 0$ such that, for every finite $I \subset \mathbb{Z}^d$, every time t and every initial row η, $d_{\mathrm{TV}}\big((P^t\eta)_I, \mu_I\big) \leq C|I|e^{-ct}$.
- *Exponential weak spatial mixing* of the space–time specification: there are $C, c > 0$ such that, for every finite volume $\Lambda \subset \mathbb{Z}^{d+1}$, every $E \subset \Lambda$ and all exterior configurations τ, τ', $d_{\mathrm{TV}}\big((\gamma^{\tau}_{\Lambda})_E, (\gamma^{\tau'}_{\Lambda})_E\big) \leq C \sum_{v \in E} e^{-c\ \mathrm{dist}(v, \Lambda^c)}$.

Theorem A (Theorem 3.1). For a finite-alphabet, finite-radius, strictly positive PCA on $\mathbb{Z}$, uniform exponential ergodicity is equivalent to exponential weak spatial mixing of its space–time specification. Under either condition the specification has exactly one DLR state, namely the two-sided stationary trajectory law $\bar{\mu}$; the finite-volume laws converge to $\bar{\mu}$ exponentially fast in the distance to the boundary, uniformly in the exterior condition; and bounded local observables have exponentially decaying covariances under $\bar{\mu}$.

Theorem B (Proposition 9.1, Theorems 9.2, 10.3 and 10.10, and Corollary 10.12). In every spatial dimension $d \geq 1$, exponential weak spatial mixing of the space–time specification implies

uniform exponential ergodicity, and uniqueness of the DLR state alone implies uniform local convergence of the dynamics to its stationary law, without a rate. Moreover, for every DLR state the relative-entropy defect from the forward dynamics, summed over a finite volume, is at most $\log(1/p_-)$ times the number of exterior children of the volume, where p_- is the least table entry. Consequently, if the PCA is ergodic, its stationary trajectory law is the only uniformly recurrent DLR state and the only periodic one, it is extremal, the translation action on the simplex of DLR states is uniquely ergodic, and the specification has a unique DLR state exactly when every DLR state is uniformly recurrent.

Theorem C (Propositions 4.1 to 4.4). Uniform exponential ergodicity follows from a finite certificate in a common random-map representation: either the fixed-pair inequality $(2rL+1)\theta_L < 1$ or the common-map inequality $(2rL+1)q_L < 1$ at some depth L. Both coefficients are polynomials in the table entries of the representation, exactly computable in rational arithmetic, and the two coincide on a binary alphabet when every map of the representation is monotone.

Theorem D (Section 11). Every radius-r rule perturbed by uniform noise $\varepsilon > 1 - 1/(2r+1)$ has a unique space–time Gibbs state with exponential mixing. The noisy shift has one at every positive noise. For the noisy majority rule on three cells, exact certificates give the conclusion at noise 0.667, 0.627, 0.587, 0.554, 0.527, 0.504 and 0.485, at depths one to seven, and in particular at noise $1/2$. Combined with Theorems A and C, the high-noise bound of Lebowitz, Maes and Speer [15, (4.1), (4.13)] gives the same conclusion at every noise above $1/2$, the range in which they note ergodicity [15, Example 1.1]; the depth-seven certificates at $1/2$ and 0.485 go beyond that range.

Theorem E (Section 11.5). Let T be the fixed radius-one automaton of Marsan, Sablik and Törmä [22], and p_ε its perturbation by uniform noise ε. Under the structural estimates of their low-noise proof, stated in Section 11.5, there is $\varepsilon_0 > 0$ such that for every $0 < \varepsilon < \varepsilon_0$ and every $2/3 < \varepsilon \le 1$ the space–time specification of p_ε has a unique DLR state with exponential weak mixing, while at some intermediate noise it has several DLR states. The alphabet and the four-site interaction support are the same throughout.

1.3 Why the future boundary is the difficulty

Fix a finite space–time volume and an exterior configuration. The past and lateral boundaries enter the specification through the parents of interior sites, exactly as in the forward dynamics. The future boundary enters through the *children*: an exterior site whose parents lie inside the volume contributes the factor $p(\tau_{\text{child}} \mid \text{interior parents})$, which tilts the forward law of the interior by the likelihood of the imposed future. For a cap of m fixed future spins this tilt lies between p_-^m and p_+^m, where $p_\pm$ are the extreme table entries, so Bayes' formula gives only

$$\mathbb{P}(E \mid \text{cap}) \le (p_+/p_-)^m \, \mathbb{P}(E)$$

for a past event E. The factor grows with the cap. Even a binary temporal copy chain with flip probability $\varepsilon/2$ has $\mathbb{P}(X_t = -1 \mid X_{t-1} = X_{t+1} = +1)$ of order ε^2, quadratic in the noise, although the forward flip probability is linear. Conditioning on the future changes the conditional laws of the past in a way that no forward contraction estimate accounts for by itself.

The proof removes the cap by a combination of three ideas.

1. *Coalescence.* Bérard [2] proves that an exponentially ergodic positive PCA on $\mathbb{Z}$ admits a local coupling-from-the-past flow, built from independent couplings of complete

Result	Statement	Where
Dynamics controls every boundary	In $d = 1$, exponential ergodicity of a positive PCA $\Leftrightarrow$ exponential weak spatial mixing of its space–time specification, with all-DLR uniqueness and exponential clustering.	Theorem 3.1, Sections 6 to 8
Converses in every dimension	Weak mixing $\Rightarrow$ exponential ergodicity; all-DLR uniqueness $\Rightarrow$ uniform local convergence, no rate.	Proposition 9.1 and Theorem 9.2
The defect bound, every dimension	The relative-entropy defect of any DLR state is a surface term. Ergodic $\Rightarrow$ the stationary trajectory is the only uniformly recurrent DLR state, extremal, with a uniquely ergodic simplex; all-DLR uniqueness $\Leftrightarrow$ ergodic and every DLR state uniformly recurrent.	Section 10
Finite certificates	$(2rL+1)\theta_L < 1$ or $(2rL+1)q_L < 1$ implies the hypothesis; both exactly computable; equal on a binary alphabet when every map of the representation is monotone.	Section 4
Noisy majority	Unique space–time Gibbs state at every noise above $1/2$ (via the high-noise bound of [15]), and at $1/2$ and 0.485 (depth-seven certificates).	Section 11.3
A reentrant fixed family	The MST automaton: one DLR state at small noise (granting structural estimates from the low-noise proof of [22]) and at large noise, several in between, on one four-site support.	Section 11.5
Equilibrium consequences	Dobrushin–Shlosman block certificate; strong mixing on squares; an open uniqueness region in positive weights on fixed supports.	Section 12

Table 1: Results at a glance.

histories on triangular tiles, whose composition over m tiles becomes constant in the initial row except with probability exponentially small in m. A root site influences the terminal row of an expanding volume only through a *cluster* of tiles on which this coalescence failed.

2. *Resampling on a fixed cluster.* For a fixed candidate cluster, replacing the coupling seeds on it by independent copies changes the terminal row only near the cluster's last level, so the cap likelihood changes by at most an exponential in the number of terminal cells of the cluster, while the probability that the candidate is bad is an exponential in its total size. The replacement field has the original law and is independent of the badness event, which is what makes the comparison a bound on the actual, cap-tilted law rather than on the forward law.

3. *Entropy against the cap, with a harmonic gain.* The number of connected candidates of a given size is exponential in the size; the badness probability can be made as small as one likes by taking long tiles; and the cap cost, an exponential in the *terminal* size d of the cluster, is absorbed because in one dimension a cluster that reaches d terminal cells has total size at least $d \sum_{\ell<n} (2R\ell+1)^{-1} \asymp d \log n$. This harmonic sum diverges only in one dimension; in $d \geq 2$ the corresponding sum converges, and this is where the forward proof stops.

Once the cap is removed for a single root, the DLR equations identify every Gibbs state as a two-sided trajectory of the PCA, and exponential ergodicity identifies every such trajectory with the stationary one. Weak mixing in an arbitrary volume then follows by running the same comparison with an entire tile as the root and coupling the actual Gibbs sampling, tile by tile, to a stationary reference evolved with the same coupling tables.

1.4 Relation to prior work

The identification of translation-invariant stationary PCA laws with fully translation-invariant space–time Gibbs measures is due to Goldstein, Kuik, Lebowitz and Maes [9], with the Ising-model reading of the space–time weights going back to Domany and Kinzel [6]; the statistical-mechanical development, including the zero pressure of the space–time interaction, is Lebowitz, Maes and Speer's [15], whose open Problems 2 and 4 [15, Section 6.6] are partially answered here, Problem 2 by Theorem 3.1 and Theorem 10.6, and Problem 4 by Corollary 12.3, which identifies in one dimension the equilibrium block conditions with the past-only conditions of Maes and Shlosman [17], using Proposition 12.2 for their necessity, as set out in Section 1.1. The time-reversal and reversibility aspects go back to Künsch [12]. Miękisz and Taati [20] use the translation-invariant correspondence to build a four-dimensional lattice gas with quasicrystalline Gibbs states and note that a complete description of the two-sided trajectories of their PCA is missing; in the Pirogov–Sinai regime of equilibrium statistical mechanics, Lebowitz and Mazel [14] prove that uniqueness among translation-periodic Gibbs states implies global uniqueness, the analogue in that setting of the passage from the translation-invariant correspondence to all-DLR uniqueness. The Baire-category proof that ergodic positive PCA are uniformly ergodic, reproduced as Proposition 10.7, is Halawi's [11]; it settles for positive rates a question raised in [19, Section 2.2]. Dai Pra, Louis and Roelly [7] study the space–time Gibbs measures of a class of PCA and their phase transition; Louis [16] proves, for attractive PCA, an equivalence between exponential temporal convergence and exponential decay of boundary influence for the invariant measures of the dynamics restricted to finite boxes; Chazottes, Redig and Ugalde [5] obtain uniqueness among translation-invariant stationary measures from a Gaussian concentration bound on the stationary path measure. None of these results controls an arbitrary future boundary of the space–time specification, which is the content of Theorem 3.1.

Maes and Shlosman [17] gave dynamical analogues of the Dobrushin–Shlosman conditions, of the kind that Problem 4 of [15] asks for. Their condition $C_{N,R}$ asks for couplings, on the trapezoids of an admissible covering of a space–time strip of height N, of the conditional laws given two configurations on the base of the trapezoid; it yields exponential ergodicity by contraction of the variational seminorm [17, Section 3, Main Theorem]. Maes and Shlosman also state a converse under exponential convergence with a constant proportional to the total single-site oscillation of the observable [17, Theorem 2], and a criterion involving convergence faster than n^{-d} [17, Corollary 3].[1] For positive PCA on $\mathbb{Z}$, Proposition 12.2 proves necessity of $C_{N,R}$ directly from (2), using Bérard's boundary-uniform coupling [2, Lemma 8]. Maes and Shlosman also observe that the Dobrushin–Shlosman condition for the space–time interaction implies uniqueness of the invariant measure [17, p. 235]; Proposition 9.1 strengthens this to

[1] The proof of [17, Theorem 2, pp. 242–243] extends a coupling of the terminal rows to the whole trapezoid without showing that, for base configurations differing at a single site, the two histories agree outside the forward cone of that site, including below the terminal row; these agreements are needed for their eq. (3-3) with the coefficients chosen below the terminal row, and for the light-cone count in their eq. (3-16). Their Corollary 3 uses the same estimate. Its little-o hypothesis also requires a uniform interpretation: pointwise in the initial pair and the finite cube it is satisfied by deterministic translation, which is not ergodic.

uniform exponential ergodicity under weak mixing alone. The single-site coupling criterion of [18, Theorem C] is a simpler form of the same method: its influence coefficients vanish outside the light cone and are bounded by θ_L inside it, so the finite-alphabet certificate below is an elementary form of their method, and we claim no novelty for it. Their application to equilibrium [17, Section 4, Example 3] concerns a spatial Gibbs measure made invariant by a reversible, sublattice-updating dynamics, for which $C_{N,R}$ can hold where C_V fails for every V; that measure is a different object from an arbitrary DLR state of the normalized space–time interaction, which [17] treats only through the fact, cited from [9], that the space–time measure of a stationary law is Gibbs [17, p. 234]. The implication proved here takes bare exponential ergodicity as its hypothesis, uses no constructive criterion, and controls every DLR state in every finite volume under arbitrary future and lateral conditions. Relative to [17, 18] we make no priority claim beyond that statement.

Bérard's construction [2] is the probabilistic input of the forward proof, and we use it exactly as stated in Proposition 5.1: the local flow, the couplings of complete histories on tiles, and the exponential coalescence of their composition. The finite-volume adapter, the cap comparison, the identification of all DLR states and the whole-tile chain-rule coupling are proved here. Bérard's later example [1], an ergodic positive-rate PCA that admits no successful basic i.i.d. single-site coupling, is the reason the general proof works with whole-history tiles and not with the basic single-site maps: exponential ergodicity must not be replaced by coalescence of a given representation.

On the equilibrium side we use the Dobrushin–Shlosman constructive criterion [8], its block-coupling presentation as developed by Weitz [24], and, in two dimensions, the theorem of Martinelli, Olivieri and Schonmann that weak mixing implies strong mixing [23], which turns the conclusion of Theorem 3.1 into an open uniqueness region in the space of positive weights on the fixed supports. The reentrant automaton is Marsan, Sablik and Törmä's [22]; the noisy-nilpotency and rapid-forgetting results we mention are in [19, 21].

1.5 Reading guide

Section 2 fixes the definitions and proves that PCA trajectories are DLR states in every dimension. Section 3 states the main theorem. Section 4 gives the finite certificates. Sections 5 to 8 prove the forward implication of Theorem 3.1: the coupling input and its clipping (Section 5), the cap comparison (Section 6), the identification of all DLR states (Section 7), and weak mixing with clustering (Section 8). Section 9 proves the converses in every dimension, and Section 10 proves the defect bound and what follows from it in every dimension without a rate. Section 11 contains the examples, including the exact computations and the reentrant family. Section 12 draws the classical equilibrium consequences, Section 13 discusses the boundaries of the result and the open problems, and the closing sections record credit, how the work was done, its verification status and the reproducibility package. A reader who wants only the mechanism can read Section 1.3, Section 2 and Section 6.

2 Setting

2.1 Probabilistic cellular automata

Let $\mathcal{A}$ be a finite alphabet with at least two symbols, $d \geq 1$ the spatial dimension, and $N \subset \mathbb{Z}^d$ a finite neighbourhood containing 0; in dimension one we take $N = [-r, r]$ and call $r \geq 1$ the

radius. A PCA is a transition table

$$p(a \mid u) \;>\; 0, \qquad \sum_{a \in \mathcal{A}} p(a \mid u) = 1, \qquad u \in \mathcal{A}^N. \tag{1}$$

Conditional on a row $\eta \in \mathcal{A}^{\mathbb{Z}^d}$, the next row has independent coordinates, the coordinate at x having law $p(\,\cdot \mid \eta_{x+N})$. We write P for this row kernel, $P^t\eta$ for the law of the row after t updates from the deterministic row η, and νP^t for the same from a random row of law ν. A stationary law is a probability measure μ with $\mu P = \mu$; one always exists, because the configuration space is compact, P is Feller, and a weak limit of Cesàro averages of iterates of any law is stationary. Strict positivity in (1) means positivity of every entry of one finite table; it does not mean a lower bound uniform over a family of tables.

For a finite set $I \subset \mathbb{Z}^d$ and a law ν on $\mathcal{A}^{\mathbb{Z}^d}$, ν_I is its marginal on I. The total variation distance of two laws on a finite set is $d_{\mathrm{TV}}(\nu, \nu') = \frac{1}{2}\sum_a |\nu(a) - \nu'(a)| \in [0, 1]$. All distances on $\mathbb{Z}^d$ and $\mathbb{Z}^{d+1}$ are sup-norm distances, and the distance between two sets is the least distance between a point of one and a point of the other.

Definition 2.1 (Uniform exponential ergodicity)**.** The PCA is *uniformly exponentially ergodic* if it has a stationary law μ and constants $C, c > 0$ such that for every finite $I \subset \mathbb{Z}^d$, every $t \geq 0$ and every deterministic row η,

$$d_{\mathrm{TV}}\big((P^t\eta)_I, \mu_I\big) \;\leq\; C\,|I|\,e^{-ct}. \tag{2}$$

Integrating (2) over the initial row extends it to every initial law, and it implies that μ is the only stationary law. This is the hypothesis of Bérard's theorem [2, eq. (4)]. Ergodicity in this paper always includes convergence; uniqueness of the stationary law is never silently substituted for it. The same name is used in [17, Definition 3], where the constant may depend on the observable; (2) asks for a constant proportional to $|I|$, and the form $\|P^n f - \mu(f)\|_\infty \leq c\,e^{-\lambda n}\sum_x \delta_x f$ used in [17, Theorem 2] implies it.

2.2 Random-map representations

A *local-map representation* is a random function $M\colon \mathcal{A}^N \to \mathcal{A}$ with $\mathbb{P}(M(u) = a) = p(a \mid u)$ for every u and a, used independently at every space–time update. One exists for every table: for instance, draw the values $M(u)$ independently over u. The joint law of the different values $M(u)$ is information beyond the table, and the choice matters. Given one i.i.d. field of such maps, every initial row η is evolved by the same maps; write $F_t(\eta)$ for the row after t updates and define

$$\theta_t = \sup_{\eta,\zeta} \mathbb{P}\big(F_t(\eta)(0) \neq F_t(\zeta)(0)\big), \tag{3}$$

$$q_t = \mathbb{P}\big(\eta \mapsto F_t(\eta)(0) \text{ is not constant}\big). \tag{4}$$

In (3) the two rows are fixed before the innovations are drawn; in (4) one realization of the innovations is tested against all rows. Thus $\theta_t \leq q_t$, and the supremum and the probability may not be interchanged. We call θ_t the *fixed-pair coefficient* and q_t the *common-map coefficient*.

Remark 2.2 (Ergodicity is not coalescence of a given representation). Bérard [1] exhibits a positive-rate PCA on $\mathbb{Z}$ that is ergodic but for which, at suitable parameters, no basic i.i.d. nearest-neighbour local-map representation coalesces. For such a PCA every q_t of such a representation stays bounded below. Exponential ergodicity is therefore a property of the table, not of a representation, and the forward proof of Theorem 3.1 never assumes that the single-site maps coalesce.

2.3 The space–time specification

Points of $\mathbb{Z}^{d+1} = \mathbb{Z}^d \times \mathbb{Z}$ are written $v = (x, t)$. The *parents* of v are $\mathrm{U}(v) = \{(x + y, t - 1) : y \in N\}$, its *children* are the sites whose parent set contains v, and $S_v = \{v\} \cup \mathrm{U}(v)$. The normalized space–time interaction assigns to every v the factor

$$e^{-\Phi_{S_v}(\sigma)} \;=\; p\big(\sigma_v \,\big|\, \sigma_{\mathrm{U}(v)}\big), \tag{5}$$

where $\sigma_{\mathrm{U}(v)}$ is read in the order fixed by N. For a finite $\Lambda \subset \mathbb{Z}^{d+1}$ and an exterior configuration $\tau \in \mathcal{A}^{\Lambda^c}$, the specification is

$$\gamma^{\tau}_{\Lambda}(\omega) \;=\; \frac{1}{Z^{\tau}_{\Lambda}} \prod_{v:\, S_v \cap \Lambda \neq \varnothing} p\big((\omega\tau)_v \,\big|\, (\omega\tau)_{\mathrm{U}(v)}\big), \qquad \omega \in \mathcal{A}^{\Lambda}, \tag{6}$$

where $\omega\tau$ is the configuration equal to ω on Λ and to τ outside. The product is finite and includes the factors of exterior children of interior sites. Removing those outgoing factors would define a different, directed, finite-volume law; the equilibrium problem keeps them. A *DLR state* is a probability measure ν on $\mathcal{A}^{\mathbb{Z}^{d+1}}$ such that for every finite Λ the conditional law of σ_Λ given σ_{Λ^c} is $\gamma^{\sigma_{\Lambda^c}}_{\Lambda}$, ν-almost surely. No invariance is part of this definition.

Definition 2.3 (Arbitrary-volume exponential weak spatial mixing)**.** The specification has *exponential weak spatial mixing* if there are $C, c > 0$ such that

$$d_{\mathrm{TV}}\big((\gamma^{\tau}_{\Lambda})_E, (\gamma^{\tau'}_{\Lambda})_E\big) \;\le\; C \sum_{v \in E} e^{-c\ \mathrm{dist}(v, \Lambda^c)} \tag{7}$$

for every finite $\Lambda \subset \mathbb{Z}^{d+1}$, every $E \subset \Lambda$ and all exterior configurations τ, τ'. Volumes need not be connected, temporally convex, or free of holes.

2.4 Trajectories are DLR states

The following lemma holds in every dimension and is used throughout.

Lemma 2.4 (PCA histories satisfy the DLR equations)**.** *Fix a row η at time 0 and run the PCA at times $1, 2, \dots$. Let ν^{η} be the law of the history on $\mathbb{Z}^d \times \{1, 2, \dots\}$. For every finite $G \subset \mathbb{Z}^d \times \{1, 2, \dots\}$, the conditional law under ν^{η} of σ_G given all other spins of the history, with row 0 equal to η, is $\gamma^{\cdot}_{G}$ of (6). Consequently every two-sided stationary trajectory law of the PCA is a DLR state for the normalized interaction.*

Proof. Let $\mathrm{Ch}(G)$ be the set of children of sites of G and let F be a finite set of sites of rows $1, 2, \dots$, disjoint from G, containing $\mathrm{Ch}(G) \setminus G$ and all parents of $G \cup \mathrm{Ch}(G)$ that lie in rows $1, 2, \dots$ and outside G. Put $D = G \cup F$. For a configuration ζ on D consider the process Y^{ζ} obtained by running the PCA from η and, at every row, overwriting the sites of D by their values in ζ before the next row is generated. Because the sites of a row are conditionally independent given the preceding row, requiring the D-sites of row s to carry their values in ζ multiplies the probability by the product of their transition factors read off the actual row $s - 1$, and leaves the conditional law of the other sites of row s unchanged; the actual row $s - 1$, on this event, is the row $s - 1$ of Y^{ζ}. By induction over rows,

$$\nu^{\eta}(\sigma_D = \zeta) \;=\; \mathbb{E} \prod_{v \in D} p\big(\zeta_v \,\big|\, Y^{\zeta}_{\mathrm{U}(v)}\big),$$

where the expectation is over the non-D coordinates of Y^{ζ}, generated as described. For $v \in G \cup \mathrm{Ch}(G)$ the parents lie in D or in row 0, so these factors are the deterministic numbers $p(\zeta_v \mid \zeta_{\mathrm{U}(v)})$, with η in row 0. A non-D site has no parent in G: such a site would be a child of G, hence in F. Therefore the law of the non-D coordinates of Y^{ζ} depends on ζ only through ζ_F, and the factors of $v \in F \setminus \mathrm{Ch}(G)$ do not involve ζ_G. Writing $\zeta = \omega\xi$ with $\omega \in \mathcal{A}^G$ and $\xi \in \mathcal{A}^F$,

$$\nu^{\eta}(\sigma_G = \omega,\ \sigma_F = \xi) \;=\; \Big[\prod_{v:\, S_v \cap G \neq \varnothing} p\big((\omega\xi\eta)_v \,\big|\, (\omega\xi\eta)_{\mathrm{U}(v)}\big)\Big] \cdot K(\xi),$$

with $K(\xi)$ independent of ω. Dividing by the sum over ω gives $\nu^{\eta}(\sigma_G = \omega \mid \sigma_F = \xi) = \gamma_G^{\xi\eta}(\omega)$ for every such F. Letting F increase to all sites of rows $1, 2, \ldots$ outside G and applying the martingale convergence theorem gives the conditional law given the whole history outside G, which is $\gamma_G^{\cdot}$ because the right side depends only on the finitely many exterior spins in $\mathrm{U}(G \cup \mathrm{Ch}(G)) \cup \mathrm{Ch}(G)$.

For a two-sided stationary trajectory and a finite G, condition first on a row at a time t_0 strictly below the earliest time of G minus one. Given that row, the later history is the PCA started from it, and the argument above applies with that row in place of η. The conditional kernel does not depend on the choice of t_0, which proves the DLR equation for G. □

3 The main theorem

Theorem 3.1 (One-dimensional dynamical–equilibrium equivalence)**.** *Let p be a finite-alphabet, finite-radius, strictly positive PCA on $\mathbb{Z}$, and let γ be the specification (6) of its normalized space–time interaction on $\mathbb{Z}^2$. The following are equivalent:*

(i) the PCA is uniformly exponentially ergodic, (2);

(ii) γ has arbitrary-volume exponential weak spatial mixing, (7).

Under either condition γ has exactly one DLR state, $\bar{\mu}$, the two-sided stationary trajectory law of the unique stationary row law μ; there are $C_*, c_* > 0$ such that for every finite Λ, every $E \subset \Lambda$ and every exterior τ,

$$d_{\mathrm{TV}}\big((\gamma_\Lambda^{\tau})_E, \bar{\mu}_E\big) \;\le\; C_* \sum_{v \in E} e^{-c_*\, \mathrm{dist}(v, \Lambda^c)}; \tag{8}$$

and for bounded observables f, g supported on finite sets E, F,

$$\big|\mathrm{Cov}_{\bar{\mu}}(f, g)\big| \;\le\; C_* \|f\|_\infty \|g\|_\infty \,(|E| + |F|)\, e^{-c_*\, \mathrm{dist}(E,F)}, \tag{9}$$

after enlarging the constants if necessary.

The implication (ii)⇒(i) holds in every spatial dimension (Proposition 9.1). The proof of (i)⇒(ii) occupies Sections 5 to 8; it uses one spatial dimension twice, in the imported triangular coupling and in a divergent harmonic sum, and it does not establish the forward implication in higher dimensions (Open problem 13.1).

4 Finite certificates for the dynamical hypothesis

A reader who wants to apply Theorem 3.1 to a given table needs a way to verify (2). The two certificates below are finite, and exactly computable when the table is rational.

Proposition 4.1 (Fixed-pair certificate). *Let θ_L be the fixed-pair coefficient (3) of some local-map representation of a radius-r PCA on $\mathbb{Z}$. If*

$$\alpha \;:=\; (2rL+1)\,\theta_L \;<\; 1 \tag{10}$$

for some finite depth L, then the PCA is uniformly exponentially ergodic, and all conclusions of Theorem 3.1 hold.

Proof. Couple two trajectories with the same maps and look at them at times $0, L, 2L, \ldots$. Conditional on the two rows η, ζ at a block boundary, the next L layers of innovations are fresh, so the output at x after the block is the depth-L map applied to the two input words on $[x-rL, x+rL]$. If the words agree the outputs agree; otherwise, by the definition of θ_L, they differ with conditional probability at most θ_L. Hence the conditional disagreement probability at x is at most $\theta_L \sum_{|y-x|\le rL} \mathbf{1}\{\eta(y) \neq \zeta(y)\}$. Taking expectations and the supremum over x gives, for $d_n = \sup_x \mathbb{P}(X_{nL}(x) \neq Y_{nL}(x))$, the recursion $d_{n+1} \le \alpha d_n$, so $d_n \le \alpha^n$.

Let μ be a stationary row law and couple a deterministic row η to a row of law μ drawn independently of the maps. A union bound over I gives $d_{\mathrm{TV}}((P^{nL}\eta)_I, \mu_I) \le |I|\alpha^n$. For $t = nL + s$ with $0 \le s < L$, the output at x at time t is a function of the row at time nL on $[x-rs, x+rs]$ and of the last s layers, so its disagreement probability is at most $(2rL+1)\alpha^n$. This is (2) with explicit constants; it also shows that μ is the only stationary law. □

The contraction step is the established constructive method of Maes and Shlosman: the single-site influence coefficients of [18, Theorem C] vanish outside the depth-L light cone and are bounded by θ_L inside it, and (10) is their condition $\gamma_L < 1$ in that case. What the next sections add is not this forward contraction but the control of an arbitrary future boundary.

Proposition 4.2 (Common-map certificate). *For a fixed local-map representation of a radius-r PCA on $\mathbb{Z}$, the following are equivalent:*

(a) $(2rL+1)q_L < 1$ *for some finite L;*

(b) $q_t \le Ce^{-ct}$ *for some $C, c > 0$;*

(c) $\liminf_{L\to\infty} L\, q_L = 0$.

Each implies (10) at the same depth, hence uniform exponential ergodicity.

Proof. Put $\rho = (2rL+1)q_L$. The composite map of $(n+1)L$ steps is $g_L \circ$ (a map of nL steps) at each of the $2rL+1$ input sites of the last block. It can be nonconstant only if the last-block map g_L is nonconstant and at least one of its $2rL+1$ input-site maps from the preceding nL steps is nonconstant. The last block is independent of the earlier maps, so a union bound gives

$$q_{(n+1)L} \;\le\; (2rL+1)\, q_L\, q_{nL} \;=\; \rho\, q_{nL}, \qquad q_{nL} \le \rho^n.$$

Holding the most recent maps fixed and moving the initial time further into the past cannot turn a constant composite map into a nonconstant one, so q_t is nonincreasing and $q_t \le \rho^{\lfloor t/L \rfloor}$. This gives (a)⇒(b); (b)⇒(c) is immediate; and (c) supplies (a) at a sufficiently large L. Finally $\theta_L \le q_L$. □

Proposition 4.3 (Exact computability). *If the local-map representation has finitely many atoms with rational probabilities, then θ_L and q_L are rational numbers computable by finite enumeration: the depth-L output reads a finite cone of $\sum_{a=0}^{L-1}(2ra+1) = L + rL(L-1)$ updates, and one sums the probabilities of the innovation assignments on that cone over which the composite map is nonconstant, or disagrees on a given pair of input words. In particular, for a rational table, the search over L for a*

certificate (10) *or* $(2rL+1)q_L < 1$ *is a terminating semidecision of the corresponding property. Its nontermination is not evidence of non-ergodicity.*

Proof. The statements about enumeration are immediate from the finite cone. Nontermination cannot be read as a converse because, by Remark 2.2, a fixed representation may fail to coalesce for an ergodic table. □

Proposition 4.4 (Monotone rules)**.** *Let* $\mathcal{A} = \{0,1\}$ *with* $0 < 1$*, and suppose every map in the support of the representation is monotone with respect to the coordinatewise order (for instance a monotone deterministic rule perturbed by noise that outputs a constant). Then every composite map is monotone, and for every* t*,*

$$\theta_t \;=\; q_t \;=\; \mathbb{P}\big(F_t(\mathbf{1})(0) = 1\big) - \mathbb{P}\big(F_t(\mathbf{0})(0) = 1\big),$$

where $\mathbf{0}, \mathbf{1}$ *are the constant rows.*

Proof. Compositions of monotone maps are monotone. A monotone map g on $\{0,1\}^I$ is nonconstant exactly when $g(\mathbf{0}) = 0$ and $g(\mathbf{1}) = 1$, and since $g(\mathbf{0}) \le g(\mathbf{1})$ always, $\mathbb{P}(g \text{ nonconstant}) = \mathbb{P}(g(\mathbf{1}) = 1) - \mathbb{P}(g(\mathbf{0}) = 1)$. For any two rows η, ζ, a disagreement $g(\eta) \neq g(\zeta)$ forces g to be nonconstant, so the pair $(\mathbf{0}, \mathbf{1})$ attains the supremum in (3) and $\theta_t = q_t$. □

For monotone binary rules the fixed-pair certificate therefore gains nothing over the common-map certificate, and both reduce to two probabilities, each computable by a row-by-row transfer over the cone without enumerating the innovation field. Section 11.3 uses this.

Proposition 4.5 (Openness within normalized kernels)**.** *Suppose* $\rho = (2rL+1)q_L < 1$ *for a representation of* p*, let* $v_L = L + rL(L-1)$ *be the number of updates in the depth-*L *cone,* $m = |\mathcal{A}|^{2r+1}$ *the number of input words, and* $\delta(p, p') = \max_u d_{\mathrm{TV}}(p(\cdot \mid u), p'(\cdot \mid u))$*. If*

$$\delta(p, p') \;<\; \frac{1-\rho}{2\,(2rL+1)\,v_L\,m},$$

then p' *admits a representation with a common-map certificate at the same depth.*

Proof. For each of the m input words couple the old output to one with the new distribution using additional local randomness; the coupled values define one complete table for p'. A union bound gives probability at most $m\,\delta(p, p')$ that any entry of the table changed. The depth-L composite maps of the two tables then coincide except with probability at most $v_L\, m\, \delta(p, p')$, so $q'_L \le q_L + v_L m\, \delta(p, p')$ and $(2rL+1)q'_L < (1+\rho)/2 < 1$. □

This openness is within normalized kernels. A perturbation of the Boltzmann weights on the fixed supports that is not PCA-normalized is handled in Section 12 through the classical block criterion, not by renormalizing: normalization inserts a parent-dependent factor at every update, which changes the space–time interaction. Openness of the region defined by a finite criterion is already noted in [17, p. 242], and robustness of exponential ergodicity in one dimension is Bérard's [2, Corollary 1].

5 The coupling input and its finite-volume adapter

From here to the end of Section 8 the PCA is on $\mathbb{Z}$ and we assume (2). We first take $r = 1$; Section 8.4 returns to arbitrary radius.

5.1 Bérard's local coupling

The following is the form of Bérard's construction that we use. Its proof is in [2, Theorem 1 and its proof, Section 3.2 and Lemma 9]; the tiles there are called downward and upward trapezoids, and the flow is built from independent couplings of complete histories on them.

Proposition 5.1 (Imported coupling input)**.** *Assume* (2) *and* $r = 1$*. There are a slab height* $T \geq 1$*, the spatial period* $2T$*, and a decomposition of every slab* $\mathbb{Z} \times \{s+1, \dots, s+T\}$ *into shrinking tiles* D_j *and expanding gaps* G_j*,* $j \in \mathbb{Z}$*, together with independent random finite tables, the* seeds*, one attached to every tile and every gap of every slab, with the following properties.*

(B1) Exact law. *For every fixed initial row of a slab, the slab history sampled from the tables has the law of the PCA run for* T *steps from that row.*

(B2) Inputs. *The table of a shrinking tile takes as input the tile's initial word; the table of a gap takes its initial word and the histories of the two adjacent shrinking tiles on its lateral sides, the* lateral record*. Each table returns the complete history of its tile or gap.*

(B3) Locality. *All inputs, outputs and seed dependencies have spatial radius at most a fixed* A*, and different slabs use independent seeds.*

(B4) Exponential coalescence. *Let* Ψ_{mT} *be the composition of the terminal-row maps of* m *consecutive slabs, as a function of the bottom row. Uniformly in* x*,*

$$\mathbb{P}\big(\eta \mapsto \Psi_{mT}(\eta)(x) \text{ is not constant}\big) \;\leq\; C_B e^{-d_B m}. \tag{11}$$

The history information in (B2) is what the proof needs; an endpoint perfect sampler would not suffice. A seed is a random finite table because its input and output sets are finite. Spatial periodicity costs nothing: every constant below is maximized over the finitely many residue classes.

A concrete indexing of the geometry is convenient (Figure 1). In a slab with initial time 0 put $c_j = 2Tj$ and $m_j = c_j + T$. For $t = 1, \dots, T$ the update sites of the shrinking tile D_j and of the gap G_j at time t are

$$D_j(t) = [c_j - T + t,\ c_j + T - t] \cap \mathbb{Z}, \qquad G_j(t) = [m_j - t + 1,\ m_j + t - 1] \cap \mathbb{Z}. \tag{12}$$

These intervals partition each positive-time row of the slab. A shrinking tile is closed under taking parents except at time 0; every parent of a gap site lies in that gap, in an adjacent shrinking tile, or in the initial row; no shrinking site reads a gap site of its slab, and no gap reads another gap. These facts give an exact sampling order: sample all shrinking histories conditional on the initial row, then all gap histories conditional on the lateral records already sampled. Multiplying the conditional probabilities in that order gives exactly the product of the one-site transition factors, which is (B1). The tables need not be chronological internally; only their complete-history laws for every fixed input matter.

5.2 Clipping a tile sampler to a finite volume

Lemma 5.2 (Clipped sampler)**.** *Fix a finite space–time region* V*, an exterior history* τ*, and a row below* V *from which the forward process in* V *starts. Sample the sites of* V *in increasing time with the ordinary transition factors, exterior parents fixed to* τ*, as follows: within each slab use an original table on every tile or gap wholly contained in* V*, and independent basic local maps, drawn chronologically, on every tile or gap cut by the boundary of* V*; generate the cut shrinking parts before the cut gap parts. This samples the exact forward law in* V *with exterior* τ*, and all seeds keep their original independent product law.*




Figure 1: One slab of height T. Shrinking tiles D_j (triangles with base on the initial row) and expanding gaps G_j (triangles with apex on the initial row) partition the slab. A tile is sampled from its initial word; a gap from its initial word and the histories of its two lateral tiles. Different tiles, gaps and slabs use independent seeds.

Proof. Clipping removes update sites and replaces their values, where needed, by the fixed exterior values; it creates no new dependency. A cut shrinking part depends only on the initial row and its own fresh maps; a complete shrinking table has the correct history law for its supplied initial word by (B1), and a cut part has the correct law by chronological generation. When a gap is reached, every lateral input has already been generated or is fixed outside V, and is independent of the gap's fresh seed; a complete gap table has the correct conditional history law for that record by (B2), and a cut gap again has the correct law chronologically. The conditional densities multiply to the product of the incoming factors of the sites of V, which chronological summation shows to be normalized. The procedure evaluates independent seeds at inputs and never conditions a seed on an imposed output, so the product law of the seeds is unchanged. □

The last sentence is the point. Sampling an unrestricted history and then conditioning its outer shell to equal τ would in general destroy the independence of the seeds; the lemma constructs the finite law directly. Only finitely many shapes of cut parts occur, so their dependence radii are uniformly bounded. Enlarge A once and for all so that it controls every clipped sampler as well as the unrestricted tables, and so that $A \geq T$.

6 The cap comparison

This section proves the estimate that removes an arbitrary future boundary. It is stated for a single root site and, in Lemma 6.2 below, for a bounded initial footprint; the second form is used for whole tiles in Section 8.

6.1 Coarse cells and bad markers

Group m small slabs into a large slab of height $H = mT$. Choose a cell width B, a multiple of the period $2T$, with $B \geq (m+2)A$, and take the smallest such; thus $B = O(m)$ as m grows, with T and A fixed. Let $C_k = [kB, (k+1)B - 1] \cap \mathbb{Z}$. In large slab j call the *marker* (j,k) *bad* if the terminal word of the large slab on C_k, under the unrestricted table flow, is not constant as a function of the input row of that large slab. By (11) and a union bound over the B sites of the cell,

$$\mathbb{P}\big((j,k) \text{ is bad}\big) \;\leq\; p_m \;:=\; B\, C_B e^{-d_B m}. \tag{13}$$

Finite locality supplies an integer $R_0 \geq 1$, *independent of* m, such that the input support and the seed support of the terminal word on C_k lie in the cells $k - R_0, \ldots, k + R_0$: the word is a composition of m slab maps of radius at most A, which span at most $mA \leq B$ sites




Figure 2: The expanding volume V_n and the cap. The root influences the terminal row only through the reached cluster $S_1, \dots, S_n$ of bad markers (shaded), which starts at the root cell and spreads by at most R cells per large slab. The cap factors (red) tilt the forward law by at most e^{β} per changed terminal spin, so the tilt costs at most $e^{b|S_n|}$, while the probability that a candidate cluster is bad is at most $p_m^{M/D}$ with M its total size. In one dimension $M \geq |S_n| H_n/(2R+1)$, and the logarithm wins.

horizontally. Let $U_{j,k}$ be a deterministic set of seeds containing the whole dependence support of that word, chosen so that the same bound holds for clipped samplers, and so that changing a seed of $U_{j,k}$ can directly affect only terminal cells within $2R_0$ of k. Within one large slab, $U_{j,k}$ and $U_{j,k'}$ are disjoint when $|k-k'| > 2R_0$; different large slabs use disjoint seeds. Set

$$D = 2R_0 + 1, \qquad R = 3R_0, \qquad \Delta = 2(2R+1), \qquad \widehat{\Delta} = \Delta + 2. \tag{14}$$

The number D is a spacing at which markers of one slab have disjoint seeds; R bounds the horizontal spread of influence from one large slab to the next, counted in cells; Δ bounds the degree of the graph on markers used in Section 6.5; the $+2$ in $\widehat{\Delta}$ allows horizontal edges between consecutive root cells.

6.2 Expanding volumes and the cap identity

Consider first the root $v = (0,0)$ and, for $n \geq 1$, the region

$$V_n \;=\; \{v\} \cup \big\{(x,t) : 1 \leq t \leq nH,\ |x| \leq W + ct\big\}, \tag{15}$$

with outward integer rounding, where the buffer W and the speed $c \geq 1$ are fixed once m is fixed.

The first positive row of V_n contains every child of v. We choose W and c large enough that every marker cone used below lies in complete tiles: if the root cells have indices in $[-s_*, s_*]$, every cell reached at level j by the clusters of Section 6.3 has $|k| \leq s_* + Rj$, and all its relevant seed footprints, input supports and splice neighbourhoods lie horizontally within $|x| \leq (s_* + Rj + 4R_0 + 4)B + 4A$ during that large slab. It suffices, for instance, to take integers

$$W \;\geq\; (s_* + R + 4R_0 + 10)B + 10A, \qquad c \;\geq\; 2 + \frac{(R + 4R_0 + 10)B}{H} + \frac{10A}{T}. \tag{16}$$

At the bottom of large slab $j \geq 2$ the half-width $W + c(j-1)H$ already exceeds the displayed support bound; the first large slab reads its arbitrary complete initial row. The margins contain entire tiles, not only their anchors, so only complete interior tiles are ever resampled.

Because the sides of V_n expand at speed at least one, an exterior site at a time $1, \dots, nH$ has no interior parent: its parents are at horizontal distance at most one, hence beyond the half-width of the preceding row. The only exterior children of interior sites are therefore in the *cap*, the row $nH + 1$. Fix the exterior τ. For $a \in \mathcal{A}$ let X^a be the forward process in $V_n \setminus \{v\}$ with $\sigma_v = a$, started from the row 0 that equals τ off v, sampled by Lemma 5.2, and let X^a_{nH} be its terminal row. Direct factorization of (6) gives the exact identity

$$\gamma^{\tau}_{V_n}(\sigma_v = a) \;=\; \frac{p(a \mid \tau_{\mathrm{U}(v)})\, Z_a}{\sum_b p(b \mid \tau_{\mathrm{U}(v)})\, Z_b}, \qquad Z_a = \mathbb{E}\, h\big(X^a_{nH}\big), \tag{17}$$

where h is the product of the transition factors of the cap sites, evaluated with their cap values from τ and their parents in the terminal row. Every Z_a is positive. There is no omitted boundary-dependent normalization in (17): the interior incoming factors form a normalized forward law, and h contains all remaining factors touching V_n.

Put

$$K = \max_{b,u,u'} \frac{p(b \mid u)}{p(b \mid u')}, \qquad \beta = 3 \log K. \tag{18}$$

Each terminal spin is a parent of at most three cap sites, so for terminal rows z, z' agreeing outside the volume,

$$h(z) \;\leq\; h(z')\, \exp\big\{\beta\, d_{\mathrm{H}}(z, z')\big\}, \tag{19}$$

with d_{H} the Hamming distance. This charges for the number of changed terminal spins, not for the width of the cap.

6.3 The deterministic splice

Let S_0 be a fixed nonempty interval of root cells; the singleton application has $S_0 = \{0\}$, and $|S_0|$ stays bounded by a fixed constant throughout. For a realization x of the seed field define the *reached cluster*

$$S_j(x) \;=\; \big\{k :\ \mathrm{dist}(k, S_{j-1}(x)) \leq R,\ (j,k) \text{ is bad in } x\big\}, \qquad 1 \leq j \leq n. \tag{20}$$

Once a level is empty, all later levels are empty. The law of the cluster depends only on the seeds, not on the sampled spins or on the exterior history.

Lemma 6.1 (Seed splice)**.** *Fix x and its reached cluster $S = (S_0, \dots, S_n)$. Replace the seeds in*

$$U(S) = \bigcup_{j=1}^{n} \bigcup_{k \in S_j} U_{j,k}$$

arbitrarily, obtaining y. Run the forward samplers of Lemma 5.2 with x and with y from the same initial row and exterior. Their terminal rows at level j differ only in cells within distance $2R_0$ of S_j. In particular, with $d = |S_n|$,

$$d_{\mathrm{H}}\big(X_{nH}(x), X_{nH}(y)\big) \;\leq\; B(4R_0 + 1)\, d. \tag{21}$$

With the same seeds x, two initial rows that differ only in S_0 have identical cap rows whenever $S_n = \varnothing$.

Proof. Induct over large slabs. A terminal word at level j that reads a replaced seed directly has index within $2R_0$ of some cell of S_j, by the support bounds. Otherwise its seed cone is unchanged. If its word nevertheless changes, some input cell at level $j-1$ changed; by the inductive hypothesis that input lies within $2R_0$ of S_{j-1}, so the terminal cell is within $R_0 + 2R_0 = R$ of S_{j-1} and is a candidate in (20). Its cone lies in complete tiles by (16). If its marker were good, the whole terminal word would be constant over all input rows, which is incompatible with a changed word under unchanged cone seeds; so the marker is bad and the cell belongs to S_j. These two cases prove the induction and (21), since each of the d terminal cells of S_n accounts for at most $4R_0 + 1$ cells of width B. For two initial rows supported on S_0 there are no seed replacements; the same induction starts with the disagreements in S_0, and if S_n is empty there is no terminal disagreement. □

No property of the replacement seeds is asserted. A replacement can change a neighbouring good marker, but all its direct effects are inside the paid $2R_0$ neighbourhood.

6.4 Resampling at a fixed cluster

Fix a deterministic candidate cluster $S = (S_0, \dots, S_n)$ and let x' be an independent copy of the product seed field. Define the hybrid field

$$y_z = \begin{cases} x'_z, & z \in U(S), \\ x_z, & z \notin U(S). \end{cases} \tag{22}$$

Let $\mathcal{B}_S$ be the event that every nonroot marker listed in S is bad in x. It depends only on $x|_{U(S)}$. The hybrid y depends only on $x|_{U(S)^c}$ and $x'|_{U(S)}$; it therefore has the original product law and is independent of $\mathcal{B}_S$. The exact-cluster event $\{S(x) = S\}$ is contained in $\mathcal{B}_S$ but need not be independent of y.

Let $M = \sum_{j=1}^{n} |S_j|$, $d = |S_n|$ and $b = \beta B(4R_0 + 1)$. For a fixed root value a the splice bound (21) together with (19) holds for every x' on $\{S(x) = S\}$, so

$$\begin{aligned} \mathbb{E}_x\big[\mathbf{1}_{\{S(x)=S\}}\, h(X^a_{nH}(x))\big] &\le e^{bd}\, \mathbb{E}_{x,x'}\big[\mathbf{1}_{\{S(x)=S\}}\, h(X^a_{nH}(y))\big] \\ &\le e^{bd}\, \mathbb{E}_{x,x'}\big[\mathbf{1}_{\mathcal{B}_S}\, h(X^a_{nH}(y))\big] \;=\; e^{bd}\, \mathbb{P}(\mathcal{B}_S)\, Z_a. \end{aligned} \tag{23}$$

The last equality uses the original law of y and its independence from $\mathcal{B}_S$; division by the same Z_a that appears in (17) is thus justified. Choose a residue class of cell indices modulo D containing at least M/D of the listed nonroot vertices; their seed sets are pairwise disjoint within each slab and different slabs are independent, so $\mathbb{P}(\mathcal{B}_S) \le p_m^{M/D}$ and

$$\frac{\mathbb{E}\big[\mathbf{1}_{\{S(x)=S\}}\, h(X^a_{nH})\big]}{Z_a} \;\le\; e^{bd}\, p_m^{M/D}. \tag{24}$$

This calculation is for a fixed candidate. Resampling a random set and asserting independence without the fixed-set calculation would not establish (24).

6.5 Cluster entropy and the harmonic gain

Consider the graph on vertices (j,k) with an edge between (j,k) and $(j+1,k')$ whenever $|k-k'| \le R$; its maximum degree is Δ. Add horizontal edges between consecutive cells of the root interval S_0; the degree is then at most $\widehat{\Delta}$, and the root interval together with any reached cluster is connected. A deterministic spanning tree and a depth-first walk starting

at the leftmost root cell recover a candidate with M nonroot vertices by a walk of length $2(M + |S_0| - 1)$, so the number of candidates with M nonroot vertices is at most

$$C_{\mathrm{root}}\,\widehat{\Delta}^{\,2M}, \qquad C_{\mathrm{root}} = \widehat{\Delta}^{\,2(|S_0|-1)}. \tag{25}$$

Counting all connected sets overcounts the reached clusters, which is harmless. A uniformly bounded root interval changes only the prefactor.

Now suppose $d = |S_n| > 0$. Then there is a vertex at every positive level, so $M \geq n$. For $\ell = 0, \ldots, n-1$ every terminal vertex has an ancestor at level $n - \ell$, and an ancestor can reach at most $2R\ell + 1$ distinct terminal cells, whatever the splitting and merging of paths. Hence

$$|S_{n-\ell}| \;\geq\; \frac{d}{2R\ell + 1}, \qquad M \;\geq\; d \sum_{\ell=0}^{n-1} \frac{1}{2R\ell + 1} \;\geq\; \frac{d\,H_n}{2R+1}, \qquad H_n = \sum_{j=1}^{n} \frac{1}{j}. \tag{26}$$

This is the one-dimensional gain: it counts possible terminal endpoints, not paths, and in $d \geq 2$ spatial dimensions the corresponding sum $\sum_\ell (2R\ell + 1)^{-d}$ would be bounded.

Since p_m decays exponentially in m while R_0 and hence $D, \Delta, \widehat{\Delta}$ are fixed, choose m once and for all so that

$$\tau_* \;:=\; e\,\widehat{\Delta}^{\,2}\,p_m^{1/D} \;<\; 1. \tag{27}$$

With that m fixed, take n_0 such that $H_{n_0} \geq b(2R+1)$. For $n \geq n_0$ and every candidate with $S_n \neq \varnothing$, (26) gives $e^{bd} \leq e^M$, and summing (24) over candidates with (25) and $M \geq n$ gives

$$\frac{\mathbb{E}\big[\mathbf{1}_{\{S_n \neq \varnothing\}}\, h(X^a_{nH})\big]}{Z_a} \;\leq\; C_{\mathrm{root}} \sum_{M \geq n} \tau_*^M \;=\; \frac{C_{\mathrm{root}}\,\tau_*^n}{1 - \tau_*} \;=:\; \delta_n. \tag{28}$$

The reached cluster is thus exponentially unlikely under the actual, cap-tilted law. We have not claimed that the tilted law has independent seeds; the comparison went through the fixed-cluster resampling.

6.6 Common extinction and the root comparison

On $\{S_n = \varnothing\}$ Lemma 6.1 makes the cap row identical for every root value, so the common contribution $Z_0 = \mathbb{E}[\mathbf{1}_{\{S_n=\varnothing\}} h(X^a_{nH})]$ does not depend on a. Write $Z_a = Z_0 + R_a$ with $R_a \geq 0$; (28) says $R_a \leq \delta_n Z_a$ for $n \geq n_0$. More generally, let w be any probability mass function on $\mathcal{A}$ and $\bar{Z} = \sum_a w(a) Z_a$, $\bar{R} = \sum_a w(a) R_a$. Then

$$\frac{w(a)Z_a}{\bar{Z}} \;=\; \frac{Z_0}{\bar{Z}}\,w(a) \;+\; \frac{\bar{R}}{\bar{Z}} \cdot \frac{w(a)R_a}{\bar{R}}, \qquad \frac{\bar{R}}{\bar{Z}} \leq \delta_n, \tag{29}$$

with the second term omitted when $\bar{R} = 0$: the law $w(\cdot)Z_\cdot/\bar{Z}$ is a mixture of w and another law with mixing weight at most δ_n, so its total variation distance from w is at most δ_n. This also shows directly why a very small partition function causes no loss.

The argument remains valid when, conditional on a, the initial row inside a fixed finite footprint has any normalized law π_a and the row outside the footprint is fixed: sample that initial row independently of the future seeds and include its expectation in Z_a. For a fixed sampled initial row the splice calculation is unchanged, and averaging preserves (23). On extinction the cap row is the same for every assignment of the footprint, so the law π_a integrates to one and the common contribution remains independent of a. This is the form used for whole tiles. Absorbing the finitely many $n < n_0$ into a constant, we have proved:

Lemma 6.2 (Uniform root comparison). *There are $C_0 < \infty$ and $s \in (0,1)$ such that for every $n \geq 1$ and every exterior history τ,*

$$d_{\mathrm{TV}}\big((\gamma^{\tau}_{V_n})_v,\ p(\,\cdot \mid \tau_{\mathrm{U}(v)})\big) \ \leq\ \min\{1,\, C_0 s^n\}. \tag{30}$$

The same bound, with constants enlarged uniformly over finitely many footprint types, holds for a root consisting of a uniformly bounded initial footprint carrying any normalized law, in the sense of (29).

7 Every DLR state is the stationary trajectory

Let ν be an arbitrary DLR state. For $v = (x,t)$ let $\mathcal{F}_v$ be the σ-field generated by all spins at times strictly below t and by the other spins of row t. The translated volume $v + V_n$ of (15) contains only v in row t and only future spins otherwise, so $\mathcal{F}_v$ is contained in the exterior σ-field of $v + V_n$. The DLR equation and the tower property give

$$\nu(\sigma_v = a \mid \mathcal{F}_v) \ =\ \mathbb{E}_\nu\big[\gamma^{\sigma}_{v+V_n}(\sigma_v = a) \,\big|\, \mathcal{F}_v\big].$$

The target $p(a \mid \sigma_{\mathrm{U}(v)})$ is $\mathcal{F}_v$-measurable, and (30) is uniform in the exterior. Letting $n \to \infty$,

$$\nu(\sigma_v = a \mid \mathcal{F}_v) \ =\ p(a \mid \sigma_{\mathrm{U}(v)}) \qquad \nu\text{-almost surely.} \tag{31}$$

No limiting exterior σ-field needs to be identified.

Let $\mathcal{P}_t$ be the σ-field of the strict past of row t. For distinct sites $x_1, \dots, x_k$ of row t, condition the product of their indicators first on $\mathcal{F}_{(x_1,t)}$: the other indicators are measurable there, and the first is replaced by its prescribed factor, which is $\mathcal{P}_t$-measurable. Repeating for each site,

$$\nu\big(\sigma_{x_j,t} = a_j,\ 1 \leq j \leq k \,\big|\, \mathcal{P}_t\big) \ =\ \prod_{j=1}^{k} p\big(a_j \,\big|\, \sigma_{\mathrm{U}(x_j,t)}\big). \tag{32}$$

Finite cylinders determine the row law, so under ν every row is obtained from the preceding one by the PCA kernel: ν is a two-sided PCA trajectory, at this stage possibly an arbitrary entrance law rather than a stationary one.

Uniform ergodicity identifies the entrance law. Fix a finite space–time cylinder E with earliest time t_0, and let I be the finite set of sites of row t_0 from which all spins of E are generated, by finite propagation. Under any entrance law the row at time $-N$ is some probability law, and (2) makes its marginal at row t_0 on I converge to μ_I uniformly in that earlier law. Applying the same forward kernel from I to E cannot increase total variation, so letting $N \to \infty$ identifies the E-marginal of ν with that of the stationary trajectory. Hence every DLR state equals $\bar{\mu}$. Existence follows from compactness of $\mathcal{A}^{\mathbb{Z}^2}$ and the continuity of the positive finite-range specification, or directly from Lemma 2.4. Uniqueness implies invariance under every space–time translation, because translates obey the same specification. This proves the uniqueness assertion of Theorem 3.1.

8 Weak mixing in arbitrary volumes, and clustering

The singleton comparison proves uniqueness, but weak mixing from bare exponential ergodicity needs one more step, because there may be no successful basic-map representation (Remark 2.2). We therefore compare an entire tile with its forward history law and couple with the independent history tables themselves.

8.1 Whole-tile roots

For a tile or gap Q of a small slab, write I_Q for its input record in the sense of (B2), and $\mathcal{G}_Q(\cdot \mid I_Q)$ for the law of its complete history under the normalized forward dynamics given that record.

Lemma 8.1 (Whole-tile comparison). *There are $C_1 < \infty$ and $s_1 \in (0,1)$ such that for every tile or gap Q and every $n \geq 1$ there is a finite comparison volume $V_{Q,n} \supset Q$, lying in the slab of Q and the slabs above it, with*

$$d_{\mathrm{TV}}\big((\gamma^{\tau}_{V_{Q,n}})_Q,\ \mathcal{G}_Q(\cdot \mid I_Q(\tau))\big) \ \leq\ C_1 s_1^n \tag{33}$$

uniformly in the exterior history τ. The diameter of $V_{Q,n}$ and its displacement from Q are at most $a_0 + a_1 n$ for fixed constants. In the slab of Q, $V_{Q,n}$ consists of Q and its two adjacent gaps when Q is a shrinking tile, and of Q alone when Q is a gap.

Proof. Suppose first $Q = D_j$ is a shrinking tile, and put $W_Q = D_j \cup G_{j-1} \cup G_j$ in its slab. The neighbouring shrinking tiles are parent-closed, so no update in them reads W_Q; every outgoing arrow from D_j within the slab enters one of the two included gaps; a gap has no outgoing arrow to another tile at a later time of the same slab. Hence every outgoing factor from W_Q lies beyond the slab's terminal row.

Fix a complete history q on D_j. Its incoming product is $\mathcal{G}_Q(q \mid I_Q)$, with I_Q fixed outside the volume. Conditional on q, the two gaps carry their normalized forward laws, reading q on one lateral side and the fixed exterior record on the other; this holds for arbitrary exterior records, whether or not they could occur in a sampled trajectory, because the normalization is an algebraic consequence of the stochastic factors. Append above the slab a buffered expanding tail of n large slabs, as in (16), containing every child of the terminal footprint of W_Q at its first time. The exact finite-volume marginal is

$$(\gamma^{\tau}_{V_{Q,n}})_Q(q) \ =\ \frac{\mathcal{G}_Q(q \mid I_Q)\, Z_q}{\sum_{q'} \mathcal{G}_Q(q' \mid I_Q)\, Z_{q'}}, \tag{34}$$

where Z_q averages the final cap likelihood over the normalized gap histories and the normalized tail. At the tail's starting row all dependence on q and on the gap histories is confined to the terminal footprint $W_Q(T)$, which has width at most $4T - 1$ by (12) and, since $B \geq T$, is covered by at most five consecutive coarse cells. The footprint form of Lemma 6.2 applies with these cells as S_0: the initial law of the footprint may depend on q but is normalized and independent of the future seeds, and on extinction the cap row is identical for every assignment of the footprint, so the weighted common contribution to Z_q is independent of q. Equations (28)–(29) with $w(q) = \mathcal{G}_Q(q \mid I_Q)$ prove (33).

If $Q = G_j$ is a gap, take $W_Q = G_j$. Its entire initial and lateral record is exterior, and no arrow leaves it for a different tile of the slab; (34) holds without auxiliary gaps, and the terminal footprint is again bounded. Finitely many tile types and residue classes give common constants; the initial pieces are fixed finite sets and the tails have width and height linear in n. □

8.2 The joint coupling

Proposition 8.2 (Arbitrary-volume convergence). *Under (2) and $r = 1$, there are $C_*, c_* > 0$ such that (8) and (7) hold.*

Proof. Let Λ be a finite volume and τ an exterior. Sample γ_Λ^τ by a chain rule in this order: slabs in increasing time; within each slab, all intersections of Λ with shrinking tiles, then all intersections with gaps, in any fixed order within each family. These intersections partition the sites of Λ. A cut tile needs no comparison estimate.

For a tile Q wholly contained in Λ, let K_Q be the exact conditional law of its history given the previously exposed spins. If $V_{Q,n} \subset \Lambda$, all previously exposed spins lie outside $V_{Q,n}$: for a shrinking tile, the only other sites of its slab in $V_{Q,n}$ are the two adjacent gaps, which have not been exposed; for a gap, $V_{Q,n}$ contains nothing else of its slab; and the rest of $V_{Q,n}$ lies in later slabs. Its input record I_Q is exposed when Q is processed. Specification consistency and the tower property give

$$d_{\mathrm{TV}}\big(K_Q,\ \mathcal{G}_Q(\cdot \mid I_Q^{\mathrm{actual}})\big) \ \le\ C_1 s_1^n, \tag{35}$$

a comparison of complete histories, not only of one-site marginals.

Construct a joint coupling as follows. Draw the history table $\mathcal{V}_Q$ of Q independently of the whole joint past, evaluate it at I_Q^{actual} to obtain Y_Q, of law $\mathcal{G}_Q(\cdot \mid I_Q^{\mathrm{actual}})$, choose a maximal coupling of that law with K_Q, and sample the actual Gibbs history of Q from its conditional distribution given Y_Q. Averaging over the fresh seed gives exactly K_Q, so the Gibbs marginal is correct; the seed is neither altered nor reweighted, so the seeds retain their product law. The construction is conditional on the full joint past, not only on the actual Gibbs history, and its local mismatch probability is bounded by (35); different mismatches need not be independent.

Use these same tables to evolve a *reference* PCA trajectory from its stationary row law at a time preceding all the finite dependence cones needed, that row drawn independently of all tables; draw further reference tables outside Λ independently, and at a cut tile keep the fresh reference table while sampling the actual Gibbs part by any exact conditional sampler. Working through a finite dependence hull avoids any sampling order on an infinite row. The reference has law $\bar{\mu}$, because the tables have the original history laws and the product seed law.

Fix a target $x \in \Lambda$ at distance $\delta = \mathrm{dist}(x, \Lambda^c)$. Its backward tile-dependence cone through n large slabs contains at most $b_0(n+1)^2$ small tiles, each within distance $b_1 + b_2 n$ of x, and Lemma 8.1 puts every comparison template $V_{Q,n}$ of such a tile within distance $b_3 + b_4 n$ of x. Choose

$$n(\delta) = \max\Big\{0,\ \Big\lfloor \frac{\delta - b_3 - 1}{b_4} \Big\rfloor\Big\}. \tag{36}$$

For $n(\delta) \ge 1$ all these templates lie in Λ and all these tiles are complete; this uses only the distance to the complement, so it is valid in volumes with holes. By a union bound,

$$\mathbb{P}(\text{some mismatch in the cone}) \ \le\ C_2\,(n(\delta)+1)^2\, s_1^{n(\delta)}. \tag{37}$$

If there is no mismatch, the actual target is the unrestricted table-flow output applied to the actual bottom row of the cone, and the reference is the same flow applied to its own bottom row. For a target on a terminal row, (11) bounds the failure to coalesce by an exponential in $n(\delta)$; for a target inside the last small slab, its value is a function of a bounded number of sites of the preceding terminal row and of the last slab's tables, and a union bound over those sites gives the same with changed constants. Hence

$$\mathbb{P}\big(\sigma_x^{\mathrm{actual}} \neq \sigma_x^{\mathrm{reference}}\big) \ \le\ C_2(n(\delta)+1)^2 s_1^{n(\delta)} + C_3 e^{-c_3 n(\delta)}, \tag{38}$$

and the trivial bound one when $n(\delta) = 0$. The coalescence estimate has not been conditioned on successful local matches: the two failure events are union-bounded under the unaltered

product law of the tables. The construction is one joint coupling for all targets, and $n(\delta)$ grows linearly in δ, so the polynomial factor is absorbed into an exponential. A union over $x \in E$ proves (8), and the triangle inequality between two exteriors proves (7). □

Remark 8.3 (A simpler final coupling under a fixed-pair certificate). If (10) is available, the whole-tile step can be replaced by a site-by-site argument after Lemma 6.2. Sample the Gibbs spins in increasing time with a fixed order inside each row; when $z + V_n \subset \Lambda$, every previously exposed spin lies outside $z + V_n$, so the chain-rule conditional K_z satisfies $d_{\mathrm{TV}}(K_z, p(\cdot \mid \text{actual parents})) \le C_0 s^n$. Draw a fresh basic map independently of the joint past, evaluate it at the actual parents, couple maximally with K_z, and run a stationary reference with the same maps; the Gibbs marginal and the product law of the maps are both preserved. For a target at distance δ, choose n proportional to δ so that its depth-nL cone and all comparison templates for the $O(n^2)$ updates in it lie in Λ; the local mismatch probability is at most $Cn^2 s^n$, and at the bottom of the cone the two exposed input rows are independent of the future maps, so Proposition 4.1 bounds the native disagreement of the two outputs by α^n. There is no conditioning on successful matches, which is why a fixed-pair certificate suffices here although the cap argument used all-input coalescence of the independent tile flow.

8.3 Clustering

Proposition 8.4 (Exponential clustering)**.** *Under* (2) *and* $r = 1$*, the bound* (9) *holds.*

Proof. Approximate the stationary trajectory at each target by the tile construction started from a fixed deterministic row a finite number m' of small slabs earlier. At terminal-row sites (11) bounds the approximation error by $C_B e^{-d_B m'}$; at interior-slab sites the preceding-terminal-row argument of Proposition 8.2 changes only constants. The approximation at a target is a function of an independent-seed cone of diameter at most $a + bm'$. For finite supports E, F use one common seed field and choose m' with $2(a + bm') < \operatorname{dist}(E, F)$: the seed cones of the two collections are disjoint, so the approximating observables $f^{(m')}, g^{(m')}$ are independent. If $\varepsilon_E, \varepsilon_F$ are the probabilities that some spin of the respective support is incorrectly approximated, then $\varepsilon_E \le C|E|e^{-cm'}$, $\varepsilon_F \le C|F|e^{-cm'}$, and since $|f - f^{(m')}| \le 2\|f\|_\infty$ on the mismatch event and zero otherwise,

$$|\operatorname{Cov}(f, g)| \;\le\; 4\,\|f\|_\infty \|g\|_\infty\,(\varepsilon_E + \varepsilon_F).$$

Take m' proportional to the separation and enlarge the constant for the finitely many small separations. The stationary process and all approximations are coupled by starting the reference before the finitely many cones from an independent stationary row; alternatively the exponentially coalescing flow defines them directly from the infinite-past seed field. □

8.4 Return to arbitrary radius

For a radius-r PCA, group r consecutive sites into one block. The blocked alphabet is $\mathcal{A}^r$ and the blocked radius is one; the single-block transition probability is the product of the r original transition probabilities, strictly positive and normalized, and conditional independence between block updates is preserved. Estimate (2) transfers with a factor r. The blocked space–time potential is the original one regrouped, not a different interaction, and DLR states correspond under the invertible blocking.

To transfer the arbitrary-volume estimates, let Λ be an original finite volume and $\hat{\Lambda}$ the blocked volume of its completely included spatial blocks. Condition on the spins of partially included blocks; the conditional law on $\hat{\Lambda}$ is the blocked Gibbs kernel with the induced

exterior. Apply its uniform estimate and integrate over the conditioned spins. A site at original distance δ from Λ^c has blocked distance at least $\delta/r - 1$ from the complement of $\hat{\Lambda}$; sites in partially included blocks have bounded original distance and are covered by enlarging the prefactor. A union bound over the blocks meeting E proves (8) in original coordinates, and clustering transfers by the same fixed rescaling.

Proof of Theorem 3.1, forward direction. Assume (2). Block to radius one by Section 8.4. Section 7 proves that $\bar{\mu}$ is the only DLR state; Proposition 8.2 proves (8) and (7); Proposition 8.4 proves (9). The converse direction is Proposition 9.1 below. □

9 Converses in every dimension

In this section the spatial lattice is $\mathbb{Z}^d$ for any $d \geq 1$, with a finite alphabet, a finite neighbourhood and a strictly positive table.

Proposition 9.1 (Weak mixing implies exponential ergodicity)**.** *In every spatial dimension,* (7) *implies* (2)*. For this implication it suffices to assume* (7) *on rectangular space–time volumes.*

Proof. Choose a stationary row law μ and its two-sided stationary trajectory $\bar{\mu}$. Fix a finite $I \subset \mathbb{Z}^d$, a deterministic row η and $t \geq 1$, and put

$$\Lambda_\varrho = [-\varrho, \varrho]^d \times \{1, \dots, 2t\}, \qquad E = I \times \{t\}, \qquad M = \max_{x \in I} \|x\|_\infty.$$

By Lemma 2.4, both $(\nu^\eta)_E$ and $\bar{\mu}_E$ are mixtures of the finite-volume kernels $(\gamma^\tau_{\Lambda_\varrho})_E$ over their respective exterior distributions. Integrating (7) against any coupling of these exterior laws, and using convexity of total variation,

$$d_{\mathrm{TV}}\big((P^t\eta)_I, \mu_I\big) \;\leq\; C\,|I|\,e^{-c\min\{t,\,\varrho - M + 1\}}.$$

Let $\varrho \to \infty$. This is (2), uniformly in η, and it implies uniqueness of the stationary law. □

Theorem 9.2 (Uniqueness implies uniform local convergence)**.** *In every spatial dimension, if the normalized space–time interaction has exactly one DLR state, then for every finite $I \subset \mathbb{Z}^d$,*

$$\sup_\eta \; d_{\mathrm{TV}}\big((P^t\eta)_I, \mu_I\big) \;\longrightarrow\; 0 \qquad (t \to \infty), \tag{39}$$

where μ is the unique stationary row law. No rate is asserted.

Proof. Start with any stationary row law μ; its trajectory law is DLR by Lemma 2.4. If (39) fails, there are a finite I, some $\delta > 0$, times $t_n \to \infty$ and rows η_n with $d_{\mathrm{TV}}((P^{t_n}\eta_n)_I, \mu_I) \geq \delta$. Run the PCA from η_n at time $-t_n$, fill the earlier past arbitrarily, and extract a weakly convergent subsequence of these measures on the compact space $\mathcal{A}^{\mathbb{Z}^{d+1}}$, with limit ν. Every fixed finite region eventually lies strictly above the initial row, so Lemma 2.4 gives its DLR identity for all large n; for a finite-range positive specification, a finite-volume kernel sends cylinder functions to continuous cylinder functions, so the integrated DLR identities, including multiplication by exterior cylinder functions, pass to the weak limit, and a monotone-class argument extends them to bounded measurable functions. Thus ν is DLR. Finite-alphabet marginal total variation is continuous under weak convergence, so the row-zero I-marginal of ν is at distance at least δ from μ_I and $\nu \neq \bar{\mu}$, contradicting uniqueness. Uniform local convergence follows, and any other stationary law equals μ by integrating (39). □

Corollary 9.3. *In every dimension, a positive PCA that is not ergodic has a normalized space–time specification with more than one DLR state.*

The second Gibbs state furnished by the compactness argument need not be stationary; the corollary does not produce two stationary row laws from non-ergodicity alone. Whether a non-ergodic positive PCA must have two stationary laws is itself open [4].

10 Every dimension, no rate: the defect bound

The converses of Section 9 hold in every dimension but say nothing about which DLR states can exist beside the stationary trajectory when $d \geq 2$. This section proves, in every dimension and with no rate, that the distance of any DLR state from the forward dynamics, measured by relative entropy, is a surface term. Most of the section is a few lines from that bound: the classical identification of translation-invariant Gibbs states with stationary laws, the fact that every *uniformly recurrent* Gibbs state of an ergodic PCA is the stationary trajectory, and the unique ergodicity of the translation action on the simplex of Gibbs states. The only other input is Halawi's Baire-category argument that ergodic positive PCA are uniformly ergodic. The section depends only on Section 2, on the row equation (32), and on Theorem 9.2.

Throughout, the spatial lattice is $\mathbb{Z}^d$ with a finite neighbourhood N, and $p_- = \min_{a,u} p(a \mid u) > 0$ is the least table entry. Order $\mathbb{Z}^{d+1}$ by time first and lexicographically within a row, and write $w \prec v$ for this translation-invariant order. The *strict past* of v is $H_v = \{w : w \prec v\}$ and $\mathcal{P}_v$ is the σ-field generated by σ_{H_v}, extending the notation $\mathcal{P}_t$ of Section 7 from rows to sites; the parents $\mathrm{U}(v)$ lie in H_v. For probability vectors α, β on $\mathcal{A}$ with β strictly positive, $\mathrm{KL}(\alpha \,\|\, \beta) = \sum_a \alpha(a) \log\big(\alpha(a)/\beta(a)\big) \geq 0$, with equality only when $\alpha = \beta$; the map $\alpha \mapsto \mathrm{KL}(\alpha \,\|\, \beta)$ is convex and continuous.

10.1 The causal defect and the surface bound

Definition 10.1 (Causal defect)**.** For a probability measure ν on $\mathcal{A}^{\mathbb{Z}^{d+1}}$ and a site v,

$$\kappa_v(\nu) \;=\; \mathbb{E}_\nu \,\mathrm{KL}\Big(\nu\big(\sigma_v \in \cdot \,\big|\, \mathcal{P}_v\big) \,\Big\|\, p\big(\cdot \,\big|\, \sigma_{\mathrm{U}(v)}\big)\Big) \;\in\; [0, \log(1/p_-)]. \tag{40}$$

The defect measures how far the conditional law of a spin given its whole past is from the transition law read off its parents. It vanishes everywhere exactly for the trajectories of the PCA.

Lemma 10.2 (Zero defect means trajectory)**.** $\kappa_v(\nu) = 0$ *for every v if and only if ν is a two-sided trajectory law of the PCA, that is,* (32) *holds for every row.*

Proof. $\kappa_v(\nu) = 0$ means $\nu(\sigma_v = a \mid \mathcal{P}_v) = p(a \mid \sigma_{\mathrm{U}(v)})$ almost surely. Suppose this holds at every site and let $x_1 \prec \cdots \prec x_k$ be sites of row t. The σ-field generated by $\mathcal{P}_t$ and $\sigma_{x_1,t}, \ldots, \sigma_{x_{j-1},t}$ is contained in $\mathcal{P}_{(x_j,t)}$, so the tower property replaces the indicator of $\{\sigma_{x_j,t} = a_j\}$ by $p(a_j \mid \sigma_{\mathrm{U}(x_j,t)})$, which is $\mathcal{P}_t$-measurable; doing this for $j = k, k-1, \ldots, 1$ gives (32). Conversely, under a trajectory law (32) gives, for $v = (x,t)$ and every finite set F of sites of row t preceding v, $\nu(\sigma_v = a \mid \mathcal{P}_t, \sigma_F) = p(a \mid \sigma_{\mathrm{U}(v)})$; letting F increase to all such sites and applying martingale convergence gives the same conditional law given $\mathcal{P}_v$, so $\kappa_v(\nu) = 0$. □

Theorem 10.3 (Surface defect bound)**.** *Let ν be any DLR state of the normalized space–time interaction of a positive PCA on $\mathbb{Z}^d$, and let $\Lambda \subset \mathbb{Z}^{d+1}$ be finite. Write $\partial_+\Lambda = \{v \notin \Lambda : \mathrm{U}(v) \cap \Lambda \neq \varnothing\}$ for the exterior children of Λ. Then*

$$\sum_{v\in\Lambda} \kappa_v(\nu) \;\le\; |\partial_+\Lambda| \,\log(1/p_-). \tag{41}$$

No invariance is assumed.

Proof. Fix an exterior configuration ξ. The product in (6) runs over the sites of $\Lambda \cup \partial_+\Lambda$, so

$$\gamma_\Lambda^\xi \;=\; \frac{\lambda\, B}{\lambda(B)}, \qquad \lambda(\omega) = \prod_{v\in\Lambda} p\big((\omega\xi)_v \,\big|\, (\omega\xi)_{\mathrm{U}(v)}\big), \qquad B(\omega) = \prod_{v\in\partial_+\Lambda} p\big(\xi_v \,\big|\, (\omega\xi)_{\mathrm{U}(v)}\big).$$

Here λ is a probability measure on $\mathcal{A}^\Lambda$: summing out the $\prec$-last site of Λ removes one factor, which sums to one because that site is a parent of no other site of Λ, and induction does the rest. The same telescoping shows that under λ the conditional law of ω_v given the sites of Λ preceding v is $p(\cdot \mid (\omega\xi)_{\mathrm{U}(v)})$. Since $p_-^{|\partial_+\Lambda|} \le B \le 1$,

$$\mathrm{KL}(\gamma_\Lambda^\xi \,\|\, \lambda) \;=\; \int \log B \, d\gamma_\Lambda^\xi - \log\lambda(B) \;\le\; |\partial_+\Lambda| \log(1/p_-).$$

Expand the left side by the chain rule along $\prec$: it is the sum over $v \in \Lambda$ of the expected relative entropy of the conditional law of ω_v given the preceding sites of Λ under γ_Λ^ξ with respect to the same conditional law under λ, which is $p(\cdot \mid (\omega\xi)_{\mathrm{U}(v)})$. Now let $\xi = \sigma_{\Lambda^c}$ be distributed under ν. By the DLR equation $\gamma_\Lambda^{\sigma_{\Lambda^c}}$ is the conditional law of σ_Λ given σ_{Λ^c}, so the v-th term of the chain rule, integrated over the exterior, becomes $\mathbb{E}_\nu \,\mathrm{KL}\big(\nu(\sigma_v \in \cdot \mid \mathcal{A}_v) \,\|\, p(\cdot \mid \sigma_{\mathrm{U}(v)})\big)$ with $\mathcal{A}_v$ generated by σ_{Λ^c} and the spins of Λ preceding v. Since H_v is contained in Λ^c together with the sites of Λ preceding v, we have $\mathcal{P}_v \subset \mathcal{A}_v$, and $p(\cdot \mid \sigma_{\mathrm{U}(v)})$ is $\mathcal{P}_v$-measurable. Conditional Jensen for the convex map $\alpha \mapsto \mathrm{KL}(\alpha \,\|\, \beta)$, with $\mathbb{E}[\nu(\sigma_v \in \cdot \mid \mathcal{A}_v) \mid \mathcal{P}_v] = \nu(\sigma_v \in \cdot \mid \mathcal{P}_v)$, bounds the v-th term below by $\kappa_v(\nu)$. □

For the box $\Lambda_n = [-n, n]^{d+1}$ the exterior children lie within the range of N of its lateral faces or just above its top row, so $|\partial_+\Lambda_n| \le C_N n^d$ while $|\Lambda_n| \asymp n^{d+1}$:

$$\sup_{\nu \text{ DLR}} \sup_{w\in\mathbb{Z}^{d+1}} \frac{1}{|\Lambda_n|} \sum_{v\in w+\Lambda_n} \kappa_v(\nu) \;\le\; \frac{C_N \log(1/p_-)}{n}. \tag{42}$$

The defect of a Gibbs state is therefore a boundary effect: it can be large at some sites, but not on a set of positive density.

10.2 Translation-invariant, periodic and uniformly recurrent states

Translations τ_w, $w \in \mathbb{Z}^{d+1}$, act on configurations and on measures; since the order $\prec$ is translation invariant, $\kappa_{v+w}(\nu) = \kappa_v(\tau_w\nu)$ with the appropriate sign convention, which plays no role below. The specification is translation invariant, so translations act on the set $\mathcal{G}$ of DLR states, which is compact and convex in the weak topology.

Corollary 10.4 (Periodic states are stationary trajectories)**.** *A DLR state invariant under a finite-index subgroup of $\mathbb{Z}^{d+1}$, in particular a translation-invariant one, is a two-sided trajectory law; if it is invariant under the time translation its row law is stationary. In particular the translation-invariant DLR states are exactly the trajectory laws of the translation-invariant stationary row laws, which is the correspondence of [9].*

Proof. A subgroup of index L contains $L\mathbb{Z}^{d+1}$. If ν is invariant under $L\mathbb{Z}^{d+1}$ then $v \mapsto \kappa_v(\nu)$ is L-periodic, so the averages in (42) converge to the average over one period cell, which is therefore 0; as every term is nonnegative, $\kappa_v(\nu) = 0$ for all v, and Lemma 10.2 applies. Time invariance of a trajectory law says that its row law is stationary. The converse direction is Lemma 2.4. □

A measure $\nu \in \mathcal{G}$ is *uniformly recurrent* if for every weak neighbourhood W of ν the set $\{w \in \mathbb{Z}^{d+1} : \tau_w \nu \in W\}$ is syndetic, that is, meets every translate of some finite box depending on W. Periodic states are uniformly recurrent, and so is every point of a minimal orbit closure; in fact uniform recurrence of ν is equivalent to minimality of its orbit closure, by the usual argument, and the property is shared by ν and all its translates.

Lemma 10.5 (The defect is lower semicontinuous)**.** *On $\mathcal{G}$ with the weak topology, $\nu \mapsto \kappa_0(\nu)$ is a supremum of continuous functions, hence lower semicontinuous and Borel.*

Proof. For n large enough that $\mathrm{U}(0) \subset [-n,n]^{d+1}$, let $D_n = H_0 \cap [-n,n]^{d+1}$ and $\kappa^{(n)}(\nu) = \mathbb{E}_\nu \mathrm{KL}\big(\nu(\sigma_0 \in \cdot \mid \sigma_{D_n}) \,\|\, p(\cdot \mid \sigma_{\mathrm{U}(0)})\big)$. Conditional Jensen, as in the proof of Theorem 10.3, makes $\kappa^{(n)}$ nondecreasing in n, and martingale convergence together with the boundedness and continuity of $\mathrm{KL}(\cdot \,\|\, \beta)$ for $\beta \geq p_-$ gives $\kappa^{(n)} \uparrow \kappa_0$. Finally $\kappa^{(n)}(\nu) = H_\nu(\sigma_{D_n}) - H_\nu(\sigma_0, \sigma_{D_n}) - \mathbb{E}_\nu \log p(\sigma_0 \mid \sigma_{\mathrm{U}(0)})$, where H_ν is the Shannon entropy of a finite marginal: a continuous function of the marginal of ν on $\{0\} \cup D_n$. □

Theorem 10.6 (Uniformly recurrent Gibbs states are trajectories)**.** *Every uniformly recurrent DLR state of a positive PCA on $\mathbb{Z}^d$ is a two-sided trajectory law.*

Proof. Suppose $\kappa_0(\nu) > \varepsilon > 0$. By Lemma 10.5 the set $W = \{\nu' \in \mathcal{G} : \kappa_0(\nu') > \varepsilon\}$ is open, so the set S of w with $\kappa_w(\nu) > \varepsilon$ is syndetic: it meets every translate of some finite box Q_0. A box Λ_n contains at least $c\, n^{d+1}$ disjoint translates of Q_0, so $\sum_{v \in \Lambda_n} \kappa_v(\nu) \geq \varepsilon\, c\, n^{d+1}$, which contradicts (42) for large n. Hence $\kappa_0(\nu) = 0$. Every translate of ν is uniformly recurrent, so $\kappa_v(\nu) = 0$ for every v, and Lemma 10.2 concludes. □

10.3 Ergodic PCA

The PCA is *ergodic* if it has a stationary law μ with $\nu P^t \to \mu$ weakly for every initial law ν, and *uniformly ergodic* if moreover (39) holds, that is, the convergence is uniform in the initial row on every finite set of sites. Uniform exponential ergodicity implies uniform ergodicity, which implies ergodicity. For positive rates the last two notions coincide; the following argument is Halawi's [11, Theorem 4.2.3 and Proposition 4.2.4], and we reproduce it because the thesis is not widely available.

Proposition 10.7 (Ergodic positive PCA are uniformly ergodic)**.** *A strictly positive PCA on $\mathbb{Z}^d$ that is ergodic is uniformly ergodic.*

Proof. Fix a finite I, a pattern $c \in \mathcal{A}^I$ and $\varepsilon > 0$, and for $m \geq 1$ let E_m be the set of rows η with $|(P^t\eta)_I(c) - \mu_I(c)| \leq \varepsilon/2$ for all $t \geq m$. Each $\eta \mapsto (P^t\eta)_I(c)$ depends on finitely many coordinates of η, so E_m is closed, and ergodicity applied to the deterministic rows says that the E_m cover the compact space $\mathcal{A}^{\mathbb{Z}^d}$. By Baire's theorem some E_m has nonempty interior, hence contains a cylinder $[\zeta_J] = \{\eta : \eta_J = \zeta_J\}$ with J finite. By positivity, from any row the next row lies in $[\zeta_J]$ with probability at least $p_-^{|J|}$, so the first entrance time T of the chain (X_t) into $[\zeta_J]$ satisfies $\mathbb{P}(T > s) \leq (1 - p_-^{|J|})^s$ uniformly in the initial row. Fix s and $t \geq m + s$. By the strong Markov property at the stopping time $T \wedge s$, $(P^t\eta)_I(c) = \mathbb{E}\big[(P^{t-T\wedge s} X_{T\wedge s})_I(c)\big]$. On

$\{T \le s\}$ the row X_T lies in $[\zeta_J] \subset E_m$ and $t - T \ge m$, so the integrand is within $\varepsilon/2$ of $\mu_I(c)$; on $\{T > s\}$ it is within 1. Hence, for every η,

$$\big|(P^t\eta)_I(c) - \mu_I(c)\big| \;\le\; \frac{\varepsilon}{2} + \mathbb{P}(T > s) \;\le\; \frac{\varepsilon}{2} + (1 - p_-^{|J|})^s.$$

Choosing s with $(1 - p_-^{|J|})^s \le \varepsilon/2$ and taking the finitely many patterns c gives (39). □

Lemma 10.8 (Uniform ergodicity is uniqueness of the entrance law)**.** *For a positive PCA on $\mathbb{Z}^d$, with $\Delta(I,t) = \sup_{\eta,\zeta} d_{\mathrm{TV}}\big((P^t\eta)_I, (P^t\zeta)_I\big)$, the following are equivalent:*

(i) *the PCA is uniformly ergodic;*

(ii) *$\Delta(I,t) \to 0$ as $t \to \infty$ for every finite I;*

(iii) *the PCA has exactly one two-sided trajectory law.*

Proof. (i)⇔(ii) is the triangle inequality in one direction and integration of ζ against μ in the other. (ii)⇒(iii): two two-sided trajectory laws have row-0 laws of the form $\nu_{-s}P^s$ and $\nu'_{-s}P^s$ for every s, so their I-marginals at row 0 are within $\Delta(I,s) \to 0$ of each other; the same holds at every row, and a trajectory law is determined by its row laws and the kernel. (iii)⇒(ii): if $\Delta(I,t_n) \ge \delta$ along $t_n \to \infty$, with rows η_n, ζ_n, run the PCA from η_n and from ζ_n at time $-t_n$, fill the earlier rows deterministically, and extract weak limits along a common subsequence. Every fixed finite region eventually lies above the initial row, so the identities (32), integrated against finitely many past cylinders, hold for all large n and pass to the limit, and a monotone-class argument gives (32) for both limits: they are two-sided trajectory laws. Their row-0 marginals on I are at distance at least δ, contradicting (iii). □

Remark 10.9 (Ergodicity of a rational positive PCA is Π_2^0). $(P^t\eta)_I$ depends on η only through the sites within distance $t \max_{y\in N} \|y\|_\infty$ of I, and is a polynomial in the table entries, so $\Delta(I,t)$ is computable from a rational table. It is nonincreasing in t, because $(P^{t+1}\eta)_I$ is a mixture of the laws $(P^t\zeta)_I$ and total variation is convex. By Lemma 10.8, uniform ergodicity is the statement $\forall I\, \forall k\, \exists t\ \Delta(I,t) < 1/k$, and by Proposition 10.7 so is ergodicity: a Π_2^0 property of the table. We have not found this placement stated elsewhere.

Theorem 10.10 (The Gibbs simplex of an ergodic PCA)**.** *Let a strictly positive PCA on $\mathbb{Z}^d$ be ergodic, with stationary law μ and stationary trajectory law $\bar\mu$, and let $\mathcal{G}$ be the set of DLR states of its normalized space–time interaction. Then:*

(i) $\bar\mu$ is the only two-sided trajectory law and the only uniformly recurrent DLR state; in particular it is the only periodic DLR state;

(ii) $\delta_{\bar\mu}$ is the only translation-invariant probability measure on $\mathcal{G}$; consequently, for every continuous f on $\mathcal{G}$, $|\Lambda_n|^{-1}\sum_{w\in\Lambda_n} f(\tau_w\nu) \to f(\bar\mu)$ uniformly in $\nu \in \mathcal{G}$;

(iii) $\bar\mu$ is an extreme point of $\mathcal{G}$.

Every DLR state other than $\bar\mu$ therefore fails uniform recurrence: it has a neighbourhood whose return set under translation has arbitrarily large gaps.

Proof. (i) Proposition 10.7 and Lemma 10.8 give the uniqueness of the trajectory law, and Theorem 10.6 identifies every uniformly recurrent DLR state with it.

(ii) Let ϱ be a translation-invariant Borel probability measure on $\mathcal{G}$. By Lemma 10.5 the integrals below make sense, and invariance gives $\int \kappa_v\, d\varrho = \int \kappa_0\, d\varrho$ for every v. Integrating (42) against ϱ yields $\int \kappa_0\, d\varrho \le C_N \log(1/p_-)/n$ for every n, so $\kappa_0 = 0$ ϱ-almost surely, hence

$\kappa_v = 0$ ϱ-almost surely for each of the countably many v, and Lemma 10.2 with (i) shows that ϱ-almost every ν equals $\bar{\mu}$. Thus $\varrho = \delta_{\bar{\mu}}$. If the averages of some continuous f failed to converge uniformly, a sequence of empirical measures $|\Lambda_{n_k}|^{-1} \sum_{w \in \Lambda_{n_k}} \delta_{\tau_w \nu_k}$ would have a weak limit point that is translation invariant and different from $\delta_{\bar{\mu}}$.

(iii) By [10, Theorem 7.26] every DLR state has a unique representation $\nu = \int \nu' \, \varrho_\nu(d\nu')$ with ϱ_ν carried by the extreme points of $\mathcal{G}$. Translations map extreme points to extreme points and fix $\bar{\mu}$, so uniqueness of the representation makes $\varrho_{\bar{\mu}}$ translation invariant, whence $\varrho_{\bar{\mu}} = \delta_{\bar{\mu}}$ by (ii), that is, $\bar{\mu}$ is extreme. □

Corollary 10.11 (Finitely many phases)**.** *If a strictly positive PCA on $\mathbb{Z}^d$ is ergodic and $\mathcal{G}$ has finitely many extreme points, then $\mathcal{G} = \{\bar{\mu}\}$.*

Proof. Let K be the number of extreme points. By the Krein–Milman theorem $\mathcal{G}$ is their convex hull, and translations act on $\mathcal{G}$ by affine homeomorphisms, so they permute the extreme points. The translations in $L\mathbb{Z}^{d+1}$ with $L = \operatorname{lcm}(1, \dots, K)$ fix every extreme point and, being affine, every DLR state. By Corollary 10.4 every DLR state is then a trajectory law whose row laws ν_t satisfy $\nu_{t+1} = \nu_t P$ and $\nu_{t+L} = \nu_t$; ergodicity gives $\nu_t = \lim_k \nu_t P^{kL} = \mu$, so every DLR state is $\bar{\mu}$. □

Corollary 10.12 (All-DLR uniqueness in every dimension)**.** *For a strictly positive PCA on $\mathbb{Z}^d$ the following are equivalent:*

(a) the normalized space–time interaction has exactly one DLR state;

(b) the PCA is ergodic and every DLR state is uniformly recurrent.

Proof. (a)⇒(b): Theorem 9.2 gives uniform local convergence, hence ergodicity, and the unique DLR state is $\bar{\mu}$, which is translation invariant and so uniformly recurrent. (b)⇒(a) is Theorem 10.10(i). □

Remark 10.13 (What the defect bound does not exclude). Corollary 10.12 locates the gap in Open problem 13.1 precisely: for an ergodic PCA in $d \geq 2$, a second DLR state would have to be a boundary-selected state that is not uniformly recurrent, returning to a neighbourhood of itself only along a set of translations with arbitrarily large gaps. Equilibrium statistical mechanics knows such objects: for four-dimensional lattice gauge theories with a finite abelian group at weak coupling, Borgs [3] proves that the translation-invariant equilibrium state is unique and constructs a second, non-invariant one, carried by a localized magnetic flux tube that starts and ends at infinity. In the opposite direction, Lebowitz and Mazel [14] prove that in the Pirogov–Sinai regime uniqueness among translation-periodic Gibbs states implies global uniqueness. The space–time interaction of a positive PCA has zero pressure and no low-temperature structure, so neither the example nor the theorem transfers directly. In $d = 1$ under exponential ergodicity such states are excluded by Theorem 3.1; for an ergodic PCA on $\mathbb{Z}$ that is not exponentially ergodic, if one exists (Open problem 13.2), the question is open even in one dimension.

11 Examples

11.1 Any rule at high noise

Let $f \colon \mathcal{A}^{[-r,r]} \to \mathcal{A}$ be a deterministic rule and perturb it by uniform noise: with probability $1 - \varepsilon$ the update applies f, with probability ε it outputs a uniformly random symbol. The

kernel is $p_\varepsilon(a \mid u) = (1-\varepsilon)\mathbf{1}\{a = f(u)\} + \varepsilon/|\mathcal{A}|$, strictly positive for $\varepsilon > 0$. Represent each update by f with probability $1-\varepsilon$ and by a fresh constant map with probability ε. If the depth-t composite map at the origin is nonconstant, tracing an essential input backwards yields a path of t deterministic updates; there are at most $(2r+1)^t$ such paths, and the updates along a fixed path are independent because their times differ. Hence

$$q_t(\varepsilon) \;\le\; \big[(2r+1)(1-\varepsilon)\big]^t, \tag{43}$$

which decays exponentially for $\varepsilon > 1 - 1/(2r+1)$. By Proposition 4.2 and Theorem 3.1, every noisy rule at that noise has a unique space–time DLR state with exponential weak mixing. At $\varepsilon = 1$ every factor is constant and the unique DLR state is i.i.d. uniform. The path count behind (43) is the branching comparison that gives $p_c(N) \ge 1/|N|$ for directed site percolation; Marcovici, Sablik and Taati prove uniform ergodicity under the weaker condition $\varepsilon > 1 - p_c(N)$ [19, Corollary 3.7].

11.2 The noisy shift

Take $f(u) = u_{-1}$, the shift, with uniform noise $\varepsilon > 0$. The composite map at the origin depends only on the diagonal chain of t updates, and it is nonconstant exactly when all of them are deterministic: $q_t = \theta_t = (1-\varepsilon)^t$. The certificate $(2L+1)(1-\varepsilon)^L < 1$ holds for every $\varepsilon > 0$ at a depth $L(\varepsilon)$ that grows as $\varepsilon \downarrow 0$. Thus the noisy shift has a unique space–time Gibbs state with exponential weak mixing at every positive noise, with constants that depend on the noise through the certificate depth. The example shows that the depth of a certificate, and the mixing constants, may depend on the table in an unbounded way even on a one-parameter family.

11.3 The noisy majority rule: exact certificates

Let $\mathcal{A} = \{0,1\}$, $r = 1$, and $f(u) = \mathrm{maj}(u_{-1}, u_0, u_1)$, perturbed by uniform noise ε, so that $p_\varepsilon(a \mid u) = (1-\varepsilon)\mathbf{1}\{a = f(u)\} + \varepsilon/2$. The bound (43) gives uniqueness for $\varepsilon > 2/3$. A sharper analytic bound is classical. In spin notation the rule has magnetization $(1-\varepsilon)\,\mathrm{sgn}(\sigma_{-1}+\sigma_0+\sigma_1) = \frac{1-\varepsilon}{2}(\sigma_{-1}+\sigma_0+\sigma_1-\sigma_{-1}\sigma_0\sigma_1)$, so the quantity δ_1 of the high-noise Condition 1 of Lebowitz, Maes and Speer equals $2(1-\varepsilon)$, and for $\varepsilon \ge 1/2$ their Lemma 1 gives $c_1(P^t\sigma_0) \le \delta_1^t$ [15, (4.1), (4.10), (4.13)], where $c_1(F)$ is the sum of the absolute values of the coefficients of the non-constant monomials in the expansion of F in products of spins. Since $q_t = \frac{1}{2}\big(P^t\sigma_0(\mathbf{1}) - P^t\sigma_0(\mathbf{0})\big)$ by Proposition 4.4, which is the sum of the coefficients of the monomials of odd degree in $P^t\sigma_0$, we get $q_t \le c_1(P^t\sigma_0) \le (2(1-\varepsilon))^t$, and a certificate exists at some depth for every $\varepsilon > 1/2$; for ergodicity this range is already noted in [15, Example 1.1], whose noise parameter is half of ours. Maes and Shlosman applied their criterion to this rule at depth two and obtained a bound equivalent to $\varepsilon > 0.634$ [17, Section 5]. The exact computations below matter at and below $\varepsilon = 1/2$. Majority and the constants are monotone, so by Proposition 4.4 $\theta_L = q_L = \mathbb{P}(F_L(\mathbf{1})(0) = 1) - \mathbb{P}(F_L(\mathbf{0})(0) = 1)$, and the depth-$L$ coefficient is a polynomial in ε of degree L^2 with rational coefficients. The reproducibility package computes it in two independent ways: by enumerating the 3^{L^2} innovation configurations of the cone for $L \le 3$, which gives the polynomials

$$\begin{aligned}
q_1 &= 1-\varepsilon,\\
q_2 &= (1-\varepsilon)^2\left(1+\varepsilon-\tfrac{1}{2}\varepsilon^2\right),\\
q_3 &= \tfrac{1}{8}(1-\varepsilon)^3\left(8+16\varepsilon+4\varepsilon^2-20\varepsilon^3+15\varepsilon^4-6\varepsilon^5+\varepsilon^6\right),
\end{aligned}$$

depth L	certified noise ε_L	$(2L+1)q_L(\varepsilon_L)$	$(2L+1)q_L(3/5)$	$(2L+1)q_L(1/2)$
1	667/1000	0.9990	1.2000	1.5000
2	627/1000	0.9951	1.1360	1.7188
3	587/1000	0.9942	0.9097	1.6697
4	277/500	0.9950	0.6693	1.4910
5	527/1000	0.9944	0.4682	1.2660
6	63/125	0.9974	0.3167	1.0395
7	97/200	0.9930	0.2091	0.8333

Table 2: Exact certificates for the noisy majority rule. At each depth, ε_L is the first grid point $k/1000$ at which the common-map certificate $(2L+1)q_L < 1$ holds, verified in rational arithmetic; the elementary bound (43) certifies $\varepsilon > 2/3$, and the analytic bound from [15] recalled above certifies every $\varepsilon > 1/2$. The two right columns show the certificate at fixed noise as the depth grows. Values are rounded; the exact rationals are in the recorded output.

and by an exact row-by-row transfer over the cone for $L \le 7$, which agrees with the enumeration at $\varepsilon = 2/3$ for $L \le 3$ exactly, as rational numbers. Table 2 lists, for each depth, the first noise level on the grid $\varepsilon = k/1000$ at which $(2L+1)q_L < 1$ holds, verified at that point in exact rational arithmetic, together with the value of $(2L+1)q_L$ at $\varepsilon = 3/5$ and $\varepsilon = 1/2$. At depth seven the certificate holds at $\varepsilon = 1/2$, with $(2L+1)q_7(1/2) = 0.8333$: the noisy majority rule at noise one half has a unique space–time Gibbs state.

Corollary 11.1 (Noisy majority)**.** *The space–time specification of the noisy majority rule has a unique DLR state with exponential weak mixing and exponential clustering at every $\varepsilon > 1/2$, and at $\varepsilon = 1/2$ and $\varepsilon = 97/200 = 0.485$ by the depth-seven certificates of Table 2.*

Each listed point is a separate certificate; monotonicity of q_L in ε is observed on the grid but is not used, and no statement is made about the noise levels between the listed points other than those covered by the analytic bound above and by Proposition 4.5. Ergodicity of this rule at every positive noise is attributed to Gray: Lebowitz, Maes and Speer cite a private communication for a unique invariant measure at every noise [15, Example 1.1], Maes and Shlosman state that Gray proved ergodicity at every noise [17, Section 5], and Marcovici, Sablik and Taati describe Gray's published argument as an outline for small symmetric noise and list the general small-noise question as open [19, Section 6, Problem 7]. We have not found a complete published proof. This computation does not touch that question; the table records how far finite certificates reach, not where ergodicity stops.

11.4 Two further sources of the hypothesis

Two families of positive PCA satisfy (2) for structural reasons, and Theorem 3.1 applies to each of them in one dimension.

Noisy nilpotent rules. Suppose the deterministic rule f is nilpotent: for some T, f^T maps every row to one constant row. Perturb it by uniform noise ε as in Section 11.1. The depth-T composite map is constant whenever no update of its cone is noisy, so with $M_T = T + rT(T-1)$ updates in the cone,

$$q_T \;\le\; 1 - (1-\varepsilon)^{M_T} \;\le\; \varepsilon\, M_T,$$

and Proposition 4.2 applies as soon as $\varepsilon\,(2rT+1)\,M_T < 1$. The threshold deteriorates with T; Marcovici, Sablik and Taati prove small-noise ergodicity of noisy nilpotent automata in

general, with an automaton-dependent threshold [19, Theorem 3.9]. Failure of the displayed inequality is not evidence of non-ergodicity.

Product stationary laws. The following gives an explicit exponential envelope in every dimension; the qualitative statement that positive-rate PCA with a stationary Bernoulli law are rapidly forgetful is Marcovici and Taati's [21].

Proposition 11.2 (Product stationary law)**.** *Let a finite-range PCA on $\mathbb{Z}^d$ have minimum rate $c = \min_{a,u} p(a \mid u) > 0$ and a stationary product law $\lambda = \pi^{\otimes \mathbb{Z}^d}$. Put $\theta = c/2$ and let $N^{(t)}$ be the t-fold sum of the neighbourhood, $N^{(0)} = \{0\}$. For every initial law ν and finite J,*

$$D\big((\nu P^t)_J \,\big\|\, \lambda_J\big) \;\le\; (1-\theta)^t \,\big|J + N^{(t)}\big| \,\log(1/c),$$

where D is relative entropy. In particular (2) *holds, with a polynomial prefactor absorbed into the exponential.*

Proof. Stationarity gives $\pi_a = \sum_u \lambda(\sigma_N = u)\, p(a \mid u) \ge c$. Factor the update as $P = Q\Theta$, where Θ independently resets each site to π with probability θ and $Q(a \mid u) = (p(a \mid u) - \theta\pi_a)/(1-\theta)$, a stochastic nonnegative rule because $p(a \mid u) \ge c > \theta\pi_a$. On every finite marginal Θ is invertible, its one-site eigenvalues being 1 and $1-\theta$, and $\lambda Q\Theta = \lambda = \lambda\Theta$ gives $\lambda Q = \lambda$.

For a law ρ on $\mathcal{A}^J$, $\rho\Theta_J$ is the mixture over random subsets $S \subset J$, each site retained independently with probability $1-\theta$, of $\rho_S \otimes \pi^{\otimes(J\setminus S)}$. Convexity of relative entropy in its first argument gives $D(\rho\Theta_J \| \pi^{\otimes J}) \le \mathbb{E}_S D(\rho_S \| \pi^{\otimes S})$. Writing $D(\rho_S \| \pi^{\otimes S}) = -H(X_S) + \sum_{i\in S} \mathbb{E}[-\log \pi_{X_i}]$ under $X \sim \rho$: the cross-entropy term scales exactly by the retention probability, while the entropy chain rule and the fact that conditioning reduces entropy give $H(X_S) = \sum_{i\in S} H(X_i \mid X_{S\cap\{<i\}}) \ge \sum_{i\in S} H(X_i \mid X_{<i})$ for any ordering, so $\mathbb{E}_S H(X_S) \ge (1-\theta)H(X_J)$. Hence $D(\rho\Theta_J \| \pi^{\otimes J}) \le (1-\theta) D(\rho \| \pi^{\otimes J})$, also when some coordinates of ρ have zero mass. Data processing for the local kernel Q, whose J-marginal output depends on the $(J+N)$-marginal input, gives $D((\nu Q)_J \| \lambda_J) \le D(\nu_{J+N} \| \lambda_{J+N})$. Iterating the two steps t times and using $D(\nu_K \| \pi^{\otimes K}) \le |K| \log(1/c)$ proves the inequality. Pinsker's inequality turns it into a total variation bound, uniform in ν; the factor $|J + N^{(t)}|$ is polynomial in t and is absorbed into the exponential at a smaller rate, and the dependence on $|J|$ is at most linear. □

11.5 A fixed reentrant family: the Marsan–Sablik–Törmä automaton

Marsan, Sablik and Törmä [22] construct one fixed radius-one cellular automaton T on a finite alphabet $\mathcal{A}$ whose uniform-noise perturbations

$$p_\varepsilon(a \mid u) = (1-\varepsilon)\mathbf{1}\{a = T(u)\} + \frac{\varepsilon}{|\mathcal{A}|}, \qquad 0 < \varepsilon \le 1, \tag{44}$$

are ergodic at sufficiently small positive noise, non-ergodic at an intermediate noise, and ergodic at high noise [22, Theorem 3.1]. Their low-noise proof is a finite-state exploration argument, and it supplies a quantitative corollary once combined with the following elementary estimate.

Lemma 11.3 (Exponential lower tails for a finite-state additive chain)**.** *Let (S_n, J_n) be a Markov-additive chain with $S_0 = 0$, finite irreducible phase chain J_n, bounded additive increments and positive stationary drift $v = \sum_i \pi_i \sum_{j,k} k\, p_{ij}(k)$, where $p_{ij}(k) = \mathbb{P}(J_{n+1} = j, S_{n+1} - S_n = k \mid J_n = i)$ and π is the stationary phase law. There are $z > 0$, $\lambda \in (0,1)$ and $D < \infty$ such that for every real A and every initial phase, $\mathbb{P}_i(S_n \le A) \le e^{zA} D \lambda^n$. The phase chain need not be aperiodic.*

Proof. Let $K(z)_{ij} = \sum_k p_{ij}(k)e^{-zk}$. Bounded increments make its entries analytic in z, and for real z its irreducibility pattern is that of the phase chain. At $z = 0$ its Perron eigenvalue is 1 and is algebraically simple, even for a periodic chain, with left eigenvector π and right eigenvector $\mathbf{1}$; so $\lambda'(0) = \pi K'(0)\mathbf{1} = -v < 0$ for the Perron eigenvalue $\lambda(z)$ near zero. Choose $z > 0$ small with $\lambda = \lambda(z) < 1$ and a positive right eigenvector h. Then $\mathbb{E}_i e^{-zS_n} = (K(z)^n\mathbf{1})_i \le \lambda^n \max h / \min h$, and Markov's inequality gives the claim with $D = \max h / \min h$. □

We now state the structural inputs that we read from the low-noise proof of [22, Section 5]; the numbering refers to that paper.

(M1) For each fixed sufficiently small $\varepsilon > 0$, the basic innovation field has two *full-state dependence borders* l_t, r_t and two inward Markov-additive walks S_n^+, S_n^- with finite irreducible phase spaces, bounded spatial increments and positive stationary drifts (Lemmas 5.4–5.5, Propositions 5.8 and 5.11).

(M2) There are deterministic constants $b, h < \infty$ and increasing integer clocks $\tau_n^\pm$, $\tau_0^\pm = 0$, with $1 \le \tau_{n+1}^\pm - \tau_n^\pm \le h$ (Corollaries 5.10 and 5.12).

(M3) $l_{\tau_n^+ + b} \ge S_n^+ - b$ and $r_{\tau_n^- + b} \le -S_n^- + b$, and the borders obey the finite-speed interpolation $l_{t+s} \ge l_t - s$, $r_{t+s} \le r_t + s$.

The bound h is for fixed positive noise and need not be uniform as $\varepsilon \downarrow 0$. The borders are defined as follows: l_t is the least starting-row coordinate that is an *essential input* for some output of the nonnegative half-row at depth t, and r_t the greatest essential input for some output of the nonpositive half-row, with $\pm\infty$ when none exists; an input is essential for an output if changing it alone changes that output for at least one assignment of the other inputs. If the origin output is nonconstant, telescoping changes of its finitely many inputs exhibit an essential one, which belongs to both half-row dependence sets, so

$$\{\text{origin map nonconstant at depth } t\} \;\subset\; \{l_t \le r_t\} \;\subset\; \{l_t \le 0\} \cup \{r_t \ge 0\}. \tag{45}$$

This concerns every coordinate of the full alphabet and every starting row, not a selected observed coordinate.

Proposition 11.4 (Bounded exploration clocks preserve exponential decay)**.** *Under (M1)–(M3), $q_t \le C_\varepsilon e^{-c_\varepsilon t}$ for some $C_\varepsilon, c_\varepsilon > 0$. Neither the two walks nor their clocks need be independent of each other.*

Proof. For $t \ge b$ let $N^\pm = N^\pm(t)$ be defined by $\tau_{N^\pm}^\pm + b \le t < \tau_{N^\pm+1}^\pm + b$. Since clock increments are at most h, $N^\pm \ge n_*(t) := \max\{0, \lfloor (t-b)/h \rfloor - 1\}$, and the gap from t to the last clock time is less than h. Finite-speed interpolation in (M3) gives $l_t \ge S_{N^+}^+ - b - h$ and $r_t \le -S_{N^-}^- + b + h$. The random indices may depend on the whole innovation field, so instead of substituting them into a fixed-time estimate, take the union over all possible indices: by (45),

$$q_t \;\le\; \sum_{n \ge n_*(t)} \mathbb{P}\big(S_n^+ \le b + h\big) + \sum_{n \ge n_*(t)} \mathbb{P}\big(S_n^- \le b + h\big) \;\le\; \frac{D_+ \lambda_+^{n_*(t)}}{1 - \lambda_+} + \frac{D_- \lambda_-^{n_*(t)}}{1 - \lambda_-},$$

by Lemma 11.3 at the fixed threshold $b + h$. Since $n_*(t)$ grows linearly in t this is the claim, after enlarging the constant for $t < b$. No optional stopping is used and no fixed-time estimate is conditioned on an exploration clock. □

Corollary 11.5 (Low-noise all-DLR mixing for the fixed automaton)**.** *Under (M1)–(M3) there is $\varepsilon_0 > 0$ such that for each fixed $0 < \varepsilon < \varepsilon_0$ the PCA (44) satisfies (2), and its normalized space–time interaction has one DLR state, exponential weak spatial mixing and exponential clustering. Its alphabet and its interaction support $\{(-1,0),(0,0),(1,0),(0,1)\}$ are fixed throughout, and all weights are positive and rational whenever ε is positive and rational.*

Proof. Proposition 11.4 gives a finite L with $(2L+1)q_L < 1$, and Proposition 4.2 and Theorem 3.1 give the conclusions. Alternatively, couple every deterministic starting row and a stationary starting row with the same innovations; on the event that all $|I|$ terminal maps are constant their output words agree, so $\sup_\eta d_{\mathrm{TV}}((P_\varepsilon^t \eta)_I, \mu_{\varepsilon,I}) \le |I| C_\varepsilon e^{-c_\varepsilon t}$ directly. □

Corollary 11.6 (A reentrant all-DLR family)**.** *Under (M1)–(M3), the family (44) has all-DLR uniqueness and exponential weak mixing for every sufficiently small positive noise and for every $2/3 < \varepsilon \le 1$; at some intermediate real parameter $\varepsilon_{\mathrm{mid}}$ it has more than one DLR state. The family has one fixed finite alphabet and one fixed finite interaction support.*

Proof. The small-noise statement is Corollary 11.5; the high-noise statement is (43) with $r = 1$. The intermediate PCA is non-ergodic by [22, Theorem 3.1], so Corollary 9.3 gives all-DLR non-uniqueness. Shrink ε_0 if necessary so that $\varepsilon_0 < \varepsilon_{\mathrm{mid}}$. □

The intermediate parameter is not asserted to be rational, nor is its additional Gibbs state asserted to be translation invariant. "Reentrant" describes a one-parameter family of positive interaction weights; the noise parameter has not been identified with an inverse temperature, and the endpoint $\varepsilon = 0$ is excluded from the positive-potential result. The rate upgrade in Proposition 11.4 is a corollary of the published finite-state, strict-drift, bounded-clock and pathwise-comparison statements, and not a fresh verification of every exploration formula of [22]; indexing conventions in that source should be resolved before its exploration rules are treated as an executable specification. None of this qualification concerns the independent cap argument of Section 6, which applies to any PCA already known to satisfy (2).

12 Consequences in equilibrium language

Theorem 3.1 places the space–time models of exponentially ergodic PCA inside the classical theory of uniqueness. We record what that theory then gives, with the attributions.

Let γ be a translation-invariant finite-state finite-range specification on $\mathbb{Z}^2$ with positive local weights, and for a finite block B let $J \subset \mathsf{B}^c$ contain all exterior coordinates on which γ_{B} depends. Define

$$c_j = \max_{\eta = \eta' \text{ off } j} W_{\mathrm{Ham}}\big(\gamma_{\mathsf{B}}^{\eta}, \gamma_{\mathsf{B}}^{\eta'}\big), \qquad C_{\mathsf{B}} = \sum_{j \in J} c_j, \tag{46}$$

where W_{Ham} is optimal transport with the number of disagreements inside B as cost. The condition $C_{\mathsf{B}} < |\mathsf{B}|$ is the Dobrushin–Shlosman constructive uniqueness criterion [8], in the block-coupling presentation of Weitz [24]: it implies a unique Gibbs measure among all DLR states, exponential convergence of finite-volume laws in the distance to the boundary uniformly in the boundary, and exponential clustering; the strict condition is open in the positive local weight tables; and for rational tables it has a finite exact rational certificate, since each c_j is the optimum of a finite rational linear program whose feasible points are dense.

Theorem 12.1 (Equilibrium consequences)**.** *Let p be a uniformly exponentially ergodic positive PCA on $\mathbb{Z}$. Then its normalized space–time specification satisfies the Dobrushin–Shlosman block*

condition $C_{\mathsf{B}} < |\mathsf{B}|$ for some finite block B, has strong mixing on squares, and lies in an open region of all-DLR uniqueness among arbitrary sufficiently small perturbations of its positive weights on the fixed supports, including perturbations that are not PCA-normalized.

Proof. By Theorem 3.1 the specification has arbitrary-volume exponential weak mixing. In two dimensions, Martinelli, Olivieri and Schonmann [23, Theorems 1.1 and 3.2] prove that uniform weak mixing on arbitrary volumes is equivalent to strong mixing on squares and to the existence of a strict Dobrushin–Shlosman block certificate, under their finite-range spin-system hypotheses, which the normalized interaction satisfies. Openness of the strict certificate in the positive weights gives the last assertion. □

The last assertion must not be obtained by normalization. Replacing arbitrary positive factors $w(a,u)$ by $p(a \mid u) = w(a,u)/\sum_b w(b,u)$ inserts a parent-dependent factor at every update, and those factors change the space–time interaction; the full positive-weight conclusion comes through the block criterion. No large numerical block certificate was computed for the examples of Section 11; existence of a certificate from a theorem is different from producing a useful-sized explicit one.

Proposition 12.2 (A past-only certificate in one dimension)**.** *Let p be a finite-alphabet, finite-radius, strictly positive PCA on $\mathbb{Z}$ satisfying (2). For every $\bar{\gamma} > 0$ there are N, a fundamental cell R, an admissible covering of the strip of height N with fundamental cell R, and couplings and coefficients satisfying [17, eq. (3-3)] for the discrete metric on $\mathcal{A}$, whose dependence matrix has $\gamma < \bar{\gamma}$, in the notation of [17, Section 3]. In particular, $C_{N,R}$ holds for some N, R.*

Proof. We first take radius one and use the discrete metric on the alphabet. By [2, Lemma 8], applied with $M = 0$, for all large N there are numbers $\varepsilon_N \le \exp(-N^{1+o(1)})$ such that an upward triangle of height N (Bérard's upward trapezoid with top length $M = 0$, an expanding gap in the language of Figure 1), with a single initial site, admits a coupling of its complete histories for all values of that site. For every fixed lateral boundary the probability that its terminal row is not constant over the initial values is at most ε_N, uniformly over the lateral boundary. Thus $(2N+1)\varepsilon_N \to 0$.

Fix two deterministic initial rows that differ only at y. Let

$$D_y = \{(i,t) : 0 \le t \le N,\ |i-y| \le t\}$$

be its forward cone. No update outside D_y has a parent in D_y. Generate all spins outside D_y once, using the common initial values there and the usual product updates. Their law is the same for both initial rows and does not depend on the value at y. Conditional on this exterior history, the dynamics in D_y have precisely the upward triangle law with its now fixed lateral boundary: in a finite restriction this follows by factoring the product of transition probabilities into the exterior factors and the cone factors. The former contain no cone spins. Only finitely many exterior spins enter the cone factors, so the same statement holds on the infinite strip.

Use the coupling from Lemma 8 in the cone, conditional on this common exterior history. Couplings of whole histories are part of the definition in [2, Section 3.2.2]; equivalently, a terminal coupling is extended inside the cone by the corresponding conditional history laws. For every site (i,t) with $1 \le t \le N$, the resulting pair of histories satisfies

$$\mathbb{P}(X_t(i) \ne X'_t(i)) \le h_{t,i;y}, \qquad h_{t,i;y} = \begin{cases} 0, & |i-y| > t, \\ 1, & 1 \le t < N,\ |i-y| \le t, \\ \varepsilon_N, & t = N,\ |i-y| \le N. \end{cases}$$

Indeed the terminal disagreement event at any one cone site is contained in the failure of terminal coalescence for the sampled boundary, and the bound for that failure is uniform over the boundary. The histories outside the cone are equal by construction.

Choose an integer $\ell > 2N$, put $R = \{0, \dots, \ell - 1\}$, and let the large trapezoid have time-t section

$$\Lambda_t = \{t, \dots, \ell - 1 - t\}, \qquad 0 \le t \le N.$$

Its translates by $\ell\mathbb{Z}$ have disjoint interiors. Add the unit trapezoid at every positive-time site not in one of these interiors. This is an admissible covering in the sense of [17, Definitions 1–2].

For base configurations differing only at $y \in R$, extend them identically outside R, apply the preceding cone construction on the full strip, and restrict to Λ. Each marginal is the required trapezoid law: all parents of every positive-time site of Λ belong to Λ, and its law depends only on its base. This also handles cones meeting an edge of Λ; the coupling is constructed on the full cone before restriction, rather than on a truncated triangle.

For general base configurations η, ζ, change their differing sites one at a time. Glue the consecutive whole-history couplings along their common intermediate marginals. Explicitly, on the finite history space, disintegrate each consecutive pair coupling over its first coordinate and sample a chain of histories with those transition kernels. Its consecutive pair laws are the prescribed couplings, and its endpoint marginals are the laws with bases η and ζ. The endpoint disagreement indicator at any coordinate is at most the sum of the consecutive disagreement indicators. Consequently the single endpoint coupling satisfies, simultaneously for all $x = (i, t) \in I(\Lambda)$,

$$\mathbb{P}(X_x \neq X'_x) \le \sum_{y \in R} k^{\Lambda}_{x,y} \mathbf{1}\{\eta_y \neq \zeta_y\}, \qquad k^{\Lambda}_{(i,t),y} = h_{t,i;y}.$$

This is [17, eq. (3-3)]. On each unit trapezoid use an identical draw when the two parent configurations agree, and any coupling otherwise. Its coefficients may be taken as one at each of its three parents, and zero elsewhere.

It remains to bound the path matrix of this covering. For a site at time $t < N$ in a large trapezoid, its outgoing arrows go directly to time zero and have total weight at most $2t + 1 \le 3^t$. For a unit trapezoid the outgoing arrows go to at most three sites at time $t - 1$. Induction therefore bounds by 3^t the total weight of all paths from any time-t site to time zero. At a terminal site in a large trapezoid the sharper bound is $(2N + 1)\varepsilon_N$, because all its arrows go directly to its base. There are $\ell - 2N$ such sites per cell and $2N$ other terminal sites. Hence

$$\gamma \le \frac{\ell - 2N}{\ell}(2N + 1)\varepsilon_N + \frac{2N}{\ell}3^N \le (2N + 1)\varepsilon_N + \frac{2N3^N}{\ell}.$$

First choose N so that the first term is less than $\bar{\gamma}/2$, and then ℓ so that the second term is less than $\bar{\gamma}/2$.

For radius $r \ge 1$, group r consecutive sites into a block. The blocked PCA has radius at most one, a finite alphabet, strictly positive rates, and (2) with changed constants. Apply the cone construction to this blocked PCA, obtaining the same bounds $h_{t,i;j}$ in block coordinates. We construct the covering and coefficients directly in the original spin coordinates as follows; no invariance of $C_{N,R}$ under blocking is needed.

Take a cell of ℓ blocks, hence $r\ell$ original sites, and let the large trapezoid at time t consist of the complete blocks numbered $t, \dots, \ell - 1 - t$. Fill its complement by original single-site unit trapezoids. The parent-closure condition holds since an update reaches at most one neighbouring block. For an original interior site x in block i and an original base site y in

block j, set $k^{\Lambda}_{x,y} = h_{t,i;j}$. A single original-site change is a change in just one initial block, so the blocked cone coupling and the same gluing argument prove [17, eq. (3-3)] for these coefficients. For original unit trapezoids use coefficient one on each of the at most $2r+1$ parents.

A nonterminal site in a large trapezoid now has outgoing weight at most $r(2t+1) \le (3r)^t$ for $t \ge 1$. Induction at the unit trapezoids, using $2r+1 \le 3r$, bounds all time-t path row sums by $(3r)^t$. A terminal site of a large trapezoid has row sum at most $r(2N+1)\varepsilon_N$. The fraction of terminal sites outside the large trapezoids is still $2N/\ell$. Thus, in the original coordinates,

$$\gamma \le r(2N+1)\varepsilon_N + \frac{2N(3r)^N}{\ell}.$$

Choosing N and then ℓ as before proves the claim. □

Corollary 12.3 (Dynamical and equilibrium criteria agree in one dimension)**.** *For a positive PCA on $\mathbb{Z}$, the Dobrushin–Shlosman condition $C_{\mathsf{B}} < |\mathsf{B}|$ of (46) holds for the normalized space–time specification for some finite block $\mathsf{B} \subset \mathbb{Z}^2$ if and only if the Maes–Shlosman condition $C_{N,R}$ of [17, Section 3] holds for some N and R.*

Proof. If $C_{N,R}$ holds, [17, Main Theorem] together with $\sum_x \delta_x(P^b f) \le (2r+1)^b \sum_x \delta_x f$ gives (2), hence weak mixing (Theorem 3.1) and a block with $C_{\mathsf{B}} < |\mathsf{B}|$ (Theorem 12.1). Conversely, such a block gives weak mixing in every finite volume [8] (in this form, [23, Theorem 3.1]), hence (2) (Proposition 9.1). Proposition 12.2 then gives $C_{N,R}$ for some N and R. □

Thus in one dimension the past-only conditions $C_{N,R}$ and the strict equilibrium block conditions $C_{\mathsf{B}} < |\mathsf{B}|$ in Problem 4 of [15, Section 6.6] hold in the same region of positive PCA tables. Their common region is the region of uniform exponential ergodicity, equivalently of all-DLR uniqueness with exponential weak mixing; bare uniqueness without a rate remains the question of Open problem 13.2.

13 Discussion and open problems

The assumptions of Theorem 3.1 are each used. Positivity makes the cap likelihood ratios (18) finite. Finite alphabet and finite radius give compactness, finite tables and deterministic finite propagation. Uniform exponential convergence is a full-state statement over every initial row; a rate for one logical coordinate, or for one chosen stationary law, does not supply it. The converse from all-DLR uniqueness yields no rate.

Open problem 13.1 (Higher dimensions)**.** Does uniform exponential ergodicity of a positive finite-range PCA on $\mathbb{Z}^d$, $d \ge 2$, imply exponential weak spatial mixing of its space–time specification on $\mathbb{Z}^{d+1}$, and hence all-DLR uniqueness? The converse holds (Proposition 9.1). The forward proof above uses one dimension in the imported triangular coupling and in the ancestor sum (26), which in d dimensions is bounded by a constant multiple of $\sum_\ell (\ell+1)^{-d}$ and cannot absorb an arbitrary fixed cap cost. This is a limitation of the proof, not a counterexample. Corollary 10.12 narrows the question: for an ergodic PCA in any dimension, a second DLR state must fail uniform recurrence, so the problem is exactly whether an exponentially ergodic PCA on $\mathbb{Z}^d$, $d \ge 2$, can have a boundary-selected space–time Gibbs state of that kind (Remark 10.13). Such an object would be striking; an equivalence in all dimensions would be a clean theorem in the tradition of [8, 23].

Open problem 13.2 (Rates)**.** Does uniqueness of the DLR state of the space–time specification imply *exponential* ergodicity? Theorem 9.2 gives uniform local convergence without a rate. For attractive binary PCA on $\mathbb{Z}^d$, an $o(n^{-d})$ difference between the single-site probabilities started from the all-plus and all-minus rows already forces exponential convergence [18, Corollary 2, eq. (4.10)]. Whether an intermediate rate can occur at all for positive finite-range PCA, and whether all-DLR uniqueness excludes it, is open.

Open problem 13.3 (Certificate depth)**.** For a uniformly exponentially ergodic positive table, Proposition 4.1 is sufficient, but it does not bound the depth in terms of the constants in (2), or establish that a certificate appears for a specified basic representation (Remark 2.2). Finite criteria allowing arbitrary couplings are available: the conditions $C_{N,R}$ of [17] are necessary in one dimension by Proposition 12.2, and Bérard's finite-size criterion [2, eq. (5)] is another characterization. The proof of Proposition 12.2 constructs a certificate from Bérard's triangular couplings, but supplies no numerical depth bound in the constants of (2) alone. Can such a bound be obtained? The variational converse stated in [17, Theorem 2] has the coupling qualification described in Section 1.4; its printed construction therefore does not by itself supply a general explicit bound.

Open problem 13.4 (Uniformity in the noise)**.** For the family (44), Corollary 11.5 is pointwise in the noise: the constants $C_\varepsilon, c_\varepsilon$ and the threshold ε_0 are not computed, and h in (M2) need not be uniform as $\varepsilon \downarrow 0$. A computed ε_0, or an explicit coalescence certificate for one small rational noise, would make the reentrant family of Corollary 11.6 fully effective.

Two remarks on scope. First, the result is about the normalized interaction of the dynamics, with the future as a genuine boundary; it says nothing about the spatial Gibbs measure that a reversible dynamics samples, to which Maes and Shlosman apply their criterion [17, Section 4, Example 3]. Second, the forward proof never uses that a basic representation coalesces, and this is not a technicality: for the ergodic PCA of [1] the single-site maps do not coalesce, the coefficient q_t of that representation does not decay, and yet, if that PCA is exponentially ergodic, Theorem 3.1 applies to it through the whole-history tiles.

Credit

The translation-invariant correspondence between stationary PCA laws and space–time Gibbs measures is Goldstein, Kuik, Lebowitz and Maes' [9], and its statistical-mechanical development is Lebowitz, Maes and Speer's [15]. The constructive ergodicity criteria and the contraction argument behind Proposition 4.1 are Maes and Shlosman's [17, 18]; Proposition 4.1 is an elementary finite-alphabet instance of their method. They also state a variational converse [17, Theorem 2]. The past-only conditions that, through Theorem 3.1, characterize in one dimension the region of all-DLR uniqueness with exponential mixing (Problem 4 of [15]) are theirs; Proposition 12.2 establishes their necessity for positive one-dimensional PCA using Bérard's triangular coupling [2, Lemma 8]. The observation that the Dobrushin–Shlosman condition for the space–time interaction gives a unique invariant measure is also Maes and Shlosman's [17, p. 235]. The local coupling from the past, the couplings of complete histories on triangular tiles, and the exponential coalescence of their composition are Bérard's [2]; the example showing that ergodicity does not imply a successful basic coupling is also his [1]. The block criterion and its coupling proof are Dobrushin and Shlosman's [8] and Weitz's [24]; the upgrade from weak to strong mixing in two dimensions is Martinelli, Olivieri and Schonmann's [23]. The reentrant automaton and its low-noise exploration are Marsan, Sablik

and Törmä's [22]. The finite-volume adapter, the cap comparison, the identification of all DLR states, the whole-tile chain-rule coupling, the converses of Section 9, the monotone identity, and the computations are proved here. In Section 10, the proof that ergodic positive PCA are uniformly ergodic is Halawi's [11]; the surface defect bound and its consequences are proved here. The comparison of the specification with the forward kernel by relative entropy is elementary and may well appear elsewhere; we have not found it, and we have not checked the full texts of [13, 25] for it. Its key estimate, that the finite-volume partition function of the normalized interaction lies between $p_{-}^{|\partial_{+}\Lambda|}$ and 1 for every boundary condition, is the observation of Lebowitz, Maes and Speer [15, Section 1.3] that the free energy is of the order of the boundary for any boundary condition. The present results have been compared with the full texts of [17, 18, 15].

How this work was done

The authors chose the question, directed the work and judged every result. AI models from more than one developer explored the problem, derived and drafted the arguments, and wrote and ran the computations, building on and challenging each other's results. The cap comparison went through several versions; the fixed-cluster resampling of Section 6.4 replaced an earlier argument that resampled a random cluster and asserted its independence. Model reviewers checked the arguments outside Section 10, refutation-first and then step by step, and the findings were fixed and re-checked. The exact computations of Section 11.3 were implemented by two different algorithms, enumeration of the innovation field and a row transfer, which agree at every tested point, and a separate program, sharing no code, re-derives every entry of Table 2. Section 10 was derived separately and checked refutation-first by a separate model reviewer; its surface bound was also tested numerically on every subset of small space–time windows. Models are not listed as authors; the authors take responsibility for every claim.

Verification status

- **Proved in the text.** Lemma 2.4, Theorem 3.1 with Lemmas 5.2, 6.1, 6.2 and 8.1, Propositions 8.2 and 8.4, Propositions 4.1 to 4.5, Proposition 9.1, Theorem 9.2, Corollary 9.3, Theorem 10.3 with Lemmas 10.2 and 10.5, Corollary 10.4, Theorem 10.6, Proposition 10.7, Lemma 10.8, Theorem 10.10, Corollaries 10.11 and 10.12, Lemma 11.3, Proposition 11.4, the bounds of Sections 11.1 and 11.4 with Proposition 11.2, and Theorem 12.1, Proposition 12.2 and Corollary 12.3 given their cited inputs. Theorem 10.10(iii) uses the uniqueness of the extremal decomposition of DLR states [10, Theorem 7.26].
- **Imported with stated scope.** Bérard's construction, as Proposition 5.1, and his boundary-uniform upward-triangle coupling [2, Lemma 8] used in Proposition 12.2; the structural inputs (M1)–(M3) of Marsan, Sablik and Törmä's low-noise proof, which are not re-derived here, so that Corollaries 11.5 and 11.6 and Theorem E are conditional on them; the Dobrushin–Shlosman criterion, the Martinelli–Olivieri–Schonmann theorem, and the Main Theorem of [17], used in Corollary 12.3.
- **Checked by machine.** The polynomials and all numbers of Table 2, in exact rational arithmetic, by two different algorithms; recorded outputs and their checksum are in the reproducibility package. No floating-point value enters a stated certificate.

- **Not formalized.** Nothing is verified in Lean.
- **Internal only.** Every audit is internal; none of it is journal peer review.

A Reproducibility

The `verification/` directory of the accompanying repository contains:

- `code/majority_certificates.py`: computes q_L and θ_L for the noisy majority rule. Part 1 enumerates the 3^{L^2} innovation configurations of the depth-L cone for $L \leq 3$, records integer counts grouped by the number of noisy updates, forms the exact polynomials, evaluates them at rational points, and finds the maximizing input pair for θ_L by exhaustive comparison. Part 2 computes $\mathbb{P}(F_L(\mathbf{1})(0) = 1) - \mathbb{P}(F_L(\mathbf{0})(0) = 1)$ by an exact row transfer for $L \leq 7$, cross-checks against Part 1 at $\varepsilon = 2/3$, locates the first passing grid point in floating point, and certifies it in rational arithmetic.
- `code/majority_certificates_output.txt` and `.json`: the recorded outputs. The script prints the SHA-256 digest of the text output; `code/SHA256SUMS` records the digests of the script and of both outputs.
- `independent/table2_check.py`: a separate program, sharing no code with the first, that re-derives every entry of Table 2 by an exact integer transfer over the backward light cone, checks that each ε_L is the first passing grid point, and compares the results with the recorded outputs; it runs in seconds.

Running `python3 majority_certificates.py 7` reproduces every number in Table 2; depth six takes about half a minute and depth seven about twenty minutes on a laptop, almost all of it in the exact rational evaluation. The paper builds from `paper/main.tex` with `pdflatex`, by `python -B build.py` in `paper/`; no computation is used in any proof.

References

[1] J. Bérard. A probabilistic cellular automaton that admits no successful basic i.i.d. coupling. Preprint, arXiv:2207.14569 (2022), Theorem 1.

[2] J. Bérard. Coupling from the past for exponentially ergodic one-dimensional probabilistic cellular automata. *Electron. J. Probab.* 28 (2023), 17 pp. doi:10.1214/23-EJP1013; arXiv:2106.07219v2. Theorem 1, Section 3.2, Lemmas 8 and 9, and the proof of Theorem 1.

[3] C. Borgs. Translation symmetry breaking in four dimensional lattice gauge theories. *Comm. Math. Phys.* 96 (1984), 251–284.

[4] P. Chassaing and J. Mairesse. A non-ergodic probabilistic cellular automaton with a unique invariant measure. *Stochastic Process. Appl.* 121 (2011), 2474–2487.

[5] J.-R. Chazottes, F. Redig and E. Ugalde. Gaussian concentration bounds for probabilistic cellular automata. *J. Stat. Phys.* 192 (2025), 169. doi:10.1007/s10955-025-03552-4.

[6] E. Domany and W. Kinzel. Equivalence of cellular automata to Ising models and directed percolation. *Phys. Rev. Lett.* 53 (1984), 311–314.

[7] P. Dai Pra, P.-Y. Louis and S. Rœlly. Stationary measures and phase transition for a class of probabilistic cellular automata. *ESAIM Probab. Stat.* 6 (2002), 89–104.

[8] R. L. Dobrushin and S. B. Shlosman. Constructive criterion for the uniqueness of Gibbs field. In *Statistical Physics and Dynamical Systems*, Progress in Physics 10, Birkhäuser (1985), 347–370.

[9] S. Goldstein, R. Kuik, J. L. Lebowitz and C. Maes. From PCA's to equilibrium systems and back. *Comm. Math. Phys.* 125 (1989), 71–79.

[10] H.-O. Georgii. *Gibbs Measures and Phase Transitions*, 2nd ed. De Gruyter Studies in Mathematics 9, de Gruyter, Berlin (2011). Theorem 7.26.

[11] A. Halawi. *Asymptotic behaviour of probabilistic cellular automata*. MS thesis, American University of Beirut (2022). https://scholarworks.aub.edu.lb/handle/10938/23549. Theorem 4.2.3 and Proposition 4.2.4.

[12] H. Künsch. Time reversal and stationary Gibbs measures. *Stochastic Process. Appl.* 17 (1984), 159–166.

[13] O. Kozlov and N. Vasilyev. Reversible Markov chains with local interaction. In *Multicomponent Random Systems*, R. L. Dobrushin and Ya. G. Sinai (eds.), Adv. Probab. Related Topics 6, Dekker, New York (1980), 451–469.

[14] J. L. Lebowitz and A. E. Mazel. On the uniqueness of Gibbs states in the Pirogov–Sinai theory. *Comm. Math. Phys.* 189 (1997), 311–321.

[15] J. L. Lebowitz, C. Maes and E. R. Speer. Statistical mechanics of probabilistic cellular automata. *J. Stat. Phys.* 59 (1990), 117–170.

[16] P.-Y. Louis. Ergodicity of PCA: equivalence between spatial and temporal mixing conditions. *Electron. Comm. Probab.* 9 (2004), 119–131.

[17] C. Maes and S. B. Shlosman. Ergodicity of probabilistic cellular automata: a constructive criterion. *Comm. Math. Phys.* 135 (1991), 233–251.

[18] C. Maes and S. B. Shlosman. When is an interacting particle system ergodic? *Comm. Math. Phys.* 151 (1993), 447–466. Section 4, Theorem C; Corollary 2, eq. (4.10).

[19] I. Marcovici, M. Sablik and S. Taati. Ergodicity of some classes of cellular automata subject to noise. *Electron. J. Probab.* 24 (2019), paper no. 41, 1–44.

[20] J. Miękisz and S. Taati. Quasicrystalline Gibbs states in 4-dimensional lattice-gas models with finite-range interactions. Preprint, arXiv:2512.24436v2 (2026).

[21] I. Marcovici and S. Taati. Positive-rate PCA and IPS with stationary Bernoulli measures are rapidly forgetful. Preprint, arXiv:2605.16904 (2026).

[22] H. Marsan, M. Sablik and I. Törmä. A perturbed cellular automaton with two phase transitions for the ergodicity. *J. Stat. Phys.* 193 (2026), 6. doi:10.1007/s10955-025-03564-0; arXiv:2507.03485. Theorem 3.1 and Section 5.

[23] F. Martinelli, E. Olivieri and R. H. Schonmann. For 2-D lattice spin systems weak mixing implies strong mixing. *Comm. Math. Phys.* 165 (1994), 33–47.

[24] D. Weitz. Combinatorial criteria for uniqueness of Gibbs measures. *Random Structures Algorithms* 27 (2005), 445–475.

[25] A. L. Toom, N. B. Vasilyev, O. N. Stavskaya, L. G. Mityushin, G. L. Kurdyumov and S. A. Pirogov. Discrete local Markov systems. In *Stochastic Cellular Systems: Ergodicity, Memory, Morphogenesis*, R. L. Dobrushin, V. I. Kryukov and A. L. Toom (eds.), Manchester University Press (1990), 1–182.