

Technical supplement: the validated Bellman/domain-wall certificate

Seth Douglas

apsiape@gmail.com ORCID: 0009-0007-4708-3252

3 August 2026

Abstract

This supplement specifies the computer-assisted input to the exact I_{3322} wall theorem. It records the characteristic map, hyperbolic plateau, analytic unstable-manifold enclosure, Miranda shooting box, exact contact covariance, global graph covers, boundary and exterior exclusions, and the boundary between certified theorem and numerical conjecture. All inequalities that carry the theorem are interval-certified; symbolic identities are exact.

1 Arithmetic model and theorem boundary

Rigorous inequalities use Arb real balls through `python-flint`; exact identities use SymPy. A displayed decimal is descriptive only. A numerical gate passes only when an interval excludes zero with the registered sign.

The computer-assisted proposition supplies a positive concave $F : [-1, 1] \rightarrow (0, \infty)$, a strictly increasing predecessor map P , and

$$q_* \in [0.250875384513976535514, 0.250875384513976536486]$$

such that, with $b(x) = \sqrt{1 - x^2}/2$, $p(x) = b(x)^2/F(x)$, and $d(x, u) = xu + (x - u)/2 - 1$,

$$p(x) + F(u) \leq q_* - d(x, u), \tag{S1}$$

with equality at the unique point $x = P(u)$. It also produces finite tensor-product strategies approaching q_* . The operator certificate and finite-dimensional nonattainment then proceed analytically in the main paper.

2 Characteristic map

For a state (x, y, r) define $s_x = \sqrt{1 - x^2}$, $s_y = \sqrt{1 - y^2}$ and

$$D = xy + \frac{x - y}{2} - 1, \quad v = \frac{2(q - D - s_x/(2r))}{s_y},$$
$$z = \frac{1}{2} \left(\frac{(1 - 2x) + 2yv/s_y}{v^2} - 1 \right).$$

The characteristic recurrence is

$$\mathcal{M}_q(x, y, r) = (y, z, v). \tag{S2}$$

It is obtained from equality and stationarity in (S1). Direct symbolic substitution verifies the recurrence and its exact reversal symmetry. The selected orbit leaves a positive fixed plateau, hits a two-equation reflection section, and returns along the reflected branch.

3 Plateau and analytic tail

The fixed point (C, C, R) belongs to the exact algebraic branch

$$R = \frac{\sqrt{1 - C^2}(2C - 1)}{(1 - C)(2C + 1)}, \quad q = \frac{4C^4 - 5C^2 + 2}{4C^2 - 1}, \quad \frac{\sqrt{3}}{2} < C < 1. \quad (\text{S3})$$

Exact elimination isolates the branch and proves the relevant root count. At the selected point DM_q has two stable eigenvalues and one unstable eigenvalue. Their reference midpoints are

$$5.8379840653\dots, \quad 0.8603760495\dots, \quad 0.1473755392\dots$$

The interval certificate, not these rounded values, separates them from the unit circle.

The local unstable curve $\Pi(t)$ is normalized by

$$\mathcal{M}_q(\Pi(t)) = \Pi(\mu t). \quad (\text{S4})$$

A high-order polynomial approximation is corrected by a graph transform on a complex ball. The certified data are summarized in Table 1.

quantity	certified bound
complex boundary tiles	400
real radicand margin	> 0.22545
denominator margin	> 1.0726
graph-transform contraction	< 0.180945557
adapted-coordinate correction	$< 1.859 \times 10^{-26}$
original-coordinate correction	$< 2.213 \times 10^{-25}$

Table 1: Analytic unstable-manifold enclosure.

4 Validated shooting zero

The unknowns are the plateau coordinate C and unstable parameter t . After four and five iterates, the two residuals are

$$h_1 = \text{crossing}[0] + \text{crossing}[1], \quad h_2 = \text{after}[2] - 1/\text{crossing}[2]. \quad (\text{S5})$$

They encode reflection of the position and amplitude. A preconditioned Miranda argument validates a zero in

$$C = 0.8782729451808124521 \pm 4.86 \times 10^{-20},$$

$$t = 0.0037582873342893243 \pm 4.36 \times 10^{-20}.$$

Every opposite face has the required strict sign after local-manifold and propagation errors are included. The two tail allowances are below 1.9×10^{-23} and 9.95×10^{-25} . Propagating the box through (S3) gives the enclosure of q_* . Miranda supplies existence; branch isolation and the strict graph below supply the uniqueness used in the Bellman construction.

5 Exact contact covariance

Let the one-form on the Bellman equality surface be

$$\beta = dF(x) - (1/2 - y) dx.$$

An exact symbolic calculation yields

$$\mathcal{M}_q^* \beta = \frac{1}{v^2} \beta. \tag{S6}$$

All coefficient residuals reduce identically to zero. Since the local unstable tangent is in $\ker \beta$, (S6) transports the envelope relation along the certified orbit. This identity is the exact bridge from the shooting recurrence to the scalar Bellman function; it is not inferred from a plot or interpolant.

6 Global graph and endpoints

The central orbit is covered at 300-bit precision by 8,192 local plateau tiles and four central pieces of 32,768 tiles each. Every projected derivative box has the registered strict sign, every pivot is positive, and there are no failed tiles. Reflection supplies the opposite half. Thus the characteristic curve is single-valued and the predecessor P is strictly increasing.

An earlier six-forward-piece parametrization failed. The accepted cover is plateau-to-section plus reflection. Both facts remain in the machine receipt to prevent the corrected theorem from silently inheriting the rejected coordinate convention.

The active right wing begins at the unique root in a width- 2×10^{-14} bracket around

$$t = -0.003719358976358651 \dots$$

The root residual has derivative below -34.725 . Two pieces of 32,768 boxes certify the graph to the endpoint, whose predecessor is near 0.898116482394039. Reflection supplies the left wing.

The inactive exterior uses 32,768 boxes. Successor monotonicity and the stationarity-target derivative exclude every additional contact; the reconstructed F has lower bound 0.2104. A first attempted guard based on a strict line derivative failed at the calibrated endpoint. The theorem uses the repaired successor-monotonicity guard, and the receipt retains the failed test.

7 Assembly of the Bellman proposition

The exact covariance, strict graph signs, endpoint wings, and inactive exclusion show that $F > 0$, that its contact slope is monotone, and hence that F is concave. The contact tangent gives (S1) globally, with exactly one contact for each u .

Hyperbolicity supplies square-summable plateau tails. Truncating the two-sided orbit and normalizing gives finite tensor-product strategies whose boundary errors tend to zero. Their Bell values converge to q_* . The analytic positive-operator decomposition proves the matching commuting upper bound.

8 Equality audit and independent reconstruction

Equality in the three positive remainders forces two decreasing bijections of the same finite ordered support. They must be the same reversal. Eliminating the resulting amplitude ratio gives

$$q_* \leq -s + \sqrt{s} \leq \frac{1}{4},$$

contradicting the lower endpoint of the certified interval. This argument has an independently written exact implementation based on rank tables and a separate symbolic elimination.

The release verifier freezes hashes, checks semantic gates, and can rerun all engines. The production path is deterministic replay. A second path uses `mpmath.iv`, a locally implemented rectangular complex-interval layer, and no imports from the Arb/FLINT production modules. It passes 18,000 rational-enclosure checks and independently reconstructs the plateau, degree-12 invariant series, analytic graph transform, Miranda zero, central graph, endpoint wing, and inactive exterior. All eight registered gates pass. Its directed shooting interval is

$$[0.2508753845139765355147934110068914495\dots, \\ 0.2508753845139765357198798108850040972\dots],$$

which overlaps the production Arb enclosure; both endpoints round to the displayed 18-place value. Its analytic-tail allowance is $2.2592334301979876 \times 10^{-25}$, only 1.020883... times the production allowance. This is method-independent internal reconstruction, not a claim about physical enactment or a substitute for the analytic proof.

9 Dimension-rate boundary

For the aligned centered finite carriers, the gaps at dimensions 31, 63, 127, 191, 255 are respectively approximately

$$3.8267 \times 10^{-4}, 2.4605 \times 10^{-5}, 1.8872 \times 10^{-7}, 1.5326 \times 10^{-9}, 1.2458 \times 10^{-11}.$$

Their successive exponential rates converge to 0.07519366..., while the plateau gives $\log R = 0.07519286\dots$. This is strong evidence for $q_* - Q_n = \Theta(R^{-n})$ in that family. It is not a dimension theorem: the data are lower bounds, and the exact finite-support reversal is not stable as written. No device-independent quantitative dimension bound is claimed.

10 Reproduction

From the repository root, quick custody verification and full deterministic replay are respectively

```
python certificate/release/verify_release.py
python certificate/release/verify_release.py --full
```

The manifest and exact file-to-claim map are distributed beside this supplement.