

DFO-HS: Department of Fisheries and Oceans Canada Hecate Strait survey data processing summary

fishglob, Aurore A. Maureaud, Julianio Palacios Abrantes, Zoë Kitchel, Dan Forrest, & Michelle Stuart

December, 2022

Contents

General info	1
Data cleaning in R	1
1. Overview of the survey data table	8
2. Summary of sampling intensity	9
3. Summary of sampling variables from the survey	10
4. Summary of biological variables	11
5. Extreme values	12
6. Summary of variables against swept area	13
7. Abundance or Weight trends of the six most abundant species	14
8. Distribution mapping	15
9. Taxonomic flagging	15
10. Spatio-temporal standardization	16
a. Standardization method 1	16
b. Standardization method 2	20
c. Standardization summary	20

General info

This document presents the cleaning code and summary of the Hecate Strait Survey (Department of Fisheries Oceans Canada) bottom trawl survey provided by Shelee Hamilton, and Maria Cornthwaite. It contains data from 2005 and up to 2019.

Data cleaning in R

```
#####  
#### R code to clean trawl survey for the DFO Hecate Strait Survey  
#### Public data Ocean Adapt  
#### Contacts: Shelee Hamilton Shelee.Hamilton@dfo-mpo.gc.ca Head,  
#### Fishery & Assessment Data Section, Science Branch, DFO Canada  
#### Maria Cornthwaite Maria.Cornthwaite@dfo-mpo.gc.ca Program Head,  
#### Groundfish Data Unit, Science Branch, DFO Canada  
#### Coding: Dan Forrest, Zoë Kitchel November 2021  
#####  
#-----#  
#### LOAD LIBRARIES AND FUNCTIONS ####  
#-----#  
  
library(tidyverse)
```

```

library(lubridate)
library(googledrive)
library(taxize) # for getting correct species names
library(magrittr) # for names wrangling
library(readr)
library(dplyr)
library(PBSmapping)

source("functions/clean_taxa.R")
source("functions/write_clean_data.R")

#Data for the Hecate Strait Survey can be best accessed using the Pinsky
#Lab Ocean Adapt Git Hub Repository.
#Contact malin.pinsky@rutgers.edu for questions or help accessing

#-----#
#### PULL IN AND EDIT RAW DATA FILES ####
#-----#

HS_catch <- read_csv(
  "https://github.com/pinskylab/OceanAdapt/raw/master/data_raw/HS_catch.csv",
  col_types = cols(
    Survey.Year = col_integer(),
    Trip.identifier = col_integer(),
    Set.number = col_integer(),
    ITIS.TSN = col_integer(),
    Species.code = col_character(),
    Scientific.name = col_character(),
    English.common.name = col_character(),
    French.common.name = col_character(),
    LSID = col_character(),
    Catch.weight..kg. = col_double(),
    Catch.count..pieces. = col_integer()
  ))

HS_effort <- read_csv(
  "https://github.com/pinskylab/OceanAdapt/raw/master/data_raw/HS_effort.csv",
  col_types =
    cols(
      Survey.Year = col_integer(),
      Trip.identifier = col_integer(),
      Vessel.name = col_character(),
      Trip.start.date = col_character(),
      Trip.end.date = col_character(),
      GMA = col_character(),
      PFMA = col_character(),
      Set.number = col_integer(),
      Set.date = col_character(),
      Start.latitude = col_double(),
      Start.longitude = col_double(),
      End.latitude = col_double(),
      End.longitude = col_double(),

```

```

        Bottom.depth..m. = col_double(),
        Tow.duration..min. = col_integer(),
        Distance.towed..m. = col_double(),
        Vessel.speed..m.min. = col_double(),
        Trawl.door.spread..m. = col_double(),
        Trawl.mouth.opening.height..m. = col_double()
    )) %>%
select(Trip.identifier, Set.number, Survey.Year, Set.date, Trip.start.date, Trip.end.date,
       GMA, PFMA, Set.date, Start.latitude, Start.longitude, End.latitude, End.longitude,
       Bottom.depth..m., Tow.duration..min., Distance.towed..m., Trawl.door.spread..m.,
       Trawl.mouth.opening.height..m. )

#-----#
#### REFORMAT AND MERGE DATA FILES ####
#-----#

HS <- left_join(HS_catch, HS_effort, by = c(
    "Trip.identifier", "Set.number", "Survey.Year"))

HS <- HS %>%
  # Create a unique haul_id
  mutate(
    haul_id = paste(formatC(Trip.identifier, width=3, flag=0),
                    formatC(Set.number, width=3, flag=0), sep= "-"),
    # Add "strata" (define by lat, lon and depth bands) where needed # degree bins
    # 100 m bins # no need to use lon grids on west coast (so narrow)
    stratum = paste(floor(Start.latitude), floor(Start.longitude),
                    floor(Bottom.depth..m./100)*100, sep= "-"),
    # catch weight (kg.) per tow/
    # (distance towed in m * trawl door spread m) * 1km^2/1000000m^2
    wgt_cpue = Catch.weight..kg./(Distance.towed..m.*Trawl.door.spread..m.) /1000000,
    # catch weight (kg.) per tow/
    # time of tow in minutes*60 minutes/hour
    wgt_h = Catch.weight..kg./Tow.duration..min.*60,
    # catch abundance per tow/
    # (distance towed in m * trawl door spread m) * 1km^2/1000000m^2
    num_cpue = Catch.count..pieces./(Distance.towed..m.*Trawl.door.spread..m.) /1000000,
    # catch weight (kg.) per tow/
    # time of tow in minutes*60 minutes/hour
    num_h = Catch.count..pieces./Tow.duration..min.*60,
    area_swept = (Distance.towed..m.*Trawl.door.spread..m.)/1000000
  )

HS <- HS %>%
  rename(
    latitude = Start.latitude,
    longitude = Start.longitude,
    depth = Bottom.depth..m.,
    verbatim_name = Scientific.name,
    year = Survey.Year,
    num = Catch.count..pieces.,

```

```

    wgt = Catch.weight..kg.
  ) %>%
  mutate(
    date = as.Date(Set.date),
    haul_dur = Tow.duration..min./60 #convert minutes to hours
  ) %>%
  filter(
    verbatim_name != "" &
    !grepl("egg", verbatim_name)
  ) %>%
  # adjust verbatim_name names
  mutate(verbatim_name = ifelse(grepl("Lepidopsetta", verbatim_name),
                                "Lepidopsetta sp.", verbatim_name),
         verbatim_name = ifelse(grepl("Bathyraja", verbatim_name),
                                'Bathyraja sp.', verbatim_name),
         verbatim_name = ifelse(grepl("Squalus", verbatim_name),
                                'Squalus suckleyi', verbatim_name))

# Does the spp column contain any eggs or non-organism notes?
#As of fall 2021, nothing stuck out as needing to be removed
test <- HS %>%
  select(verbatim_name) %>%
  filter(!is.na(verbatim_name)) %>%
  distinct() %>%
  mutate(verbatim_name = as.factor(verbatim_name)) %>%
  filter(grepl("egg", verbatim_name) & grepl("", verbatim_name))
stopifnot(nrow(test)==0)

# combine the wtcpue for each species by haul which is necessary
#because sometimes there are multiple observations for a single genus or family
#this removes duplicates
#i.e.
#HEXACTINELLIDA, GLASS SPONGES; WILLEMOES'S WHITE SEA PEN; CRANGONS
HS <- HS %>%
  group_by(haul_id, year, latitude, longitude, depth, verbatim_name, area_swept,
           date, haul_dur) %>%
  summarise(wgt_cpue = sum(wgt_cpue, na.rm = T), wgt_h = sum(wgt_h, na.rm = T),
            num_h = sum(num_h, na.rm = T), num_cpue = sum(num_cpue, na.rm = T),
            wgt = sum(wgt, na.rm = T), num = sum(num, na.rm = T)) %>%
  ungroup()

HS <- HS %>%
# add survey column
  mutate(survey = "DFO-HS",
         country = "Canada",
         continent = "n_america",
         stat_rec = NA,
         verbatim_aphia_id = NA,
         aphia_id = NA,
         sub_area = NA,
         station = NA,

```

```

    stratum = NA,
    month = lubridate::month(date),
    day = lubridate::day(date),
    season = NA,
    quarter = NA,
    gear = NA,
    sbt = NA,
    sst = NA
  ) %>%
  select(survey, haul_id, country, sub_area, continent, stat_rec, station, stratum,
         year, month, day, quarter, season, latitude, longitude, haul_dur, area_swept,
         gear, depth, sbt, sst, verbatim_name, num, num_h, num_cpue,
         wgt, wgt_h, wgt_cpue, verbatim_name, verbatim_aphia_id)

#check for duplicates, should not be any with more than 1 obs
#check for duplicates
count_HS <- HS %>%
  group_by(haul_id, verbatim_name) %>%
  mutate(count = n())

#none!

#which ones are duplicated?
unique_name_match <- count_HS %>%
  group_by(verbatim_name) %>%
  filter(count>1) %>%
  distinct(verbatim_name)

unique_name_match
#empty

#-----#
#### INTEGRATE CLEAN TAXA FROM TAXA ANALYSIS ####
#-----#

# Get WoRM's id for sourcing
worm <- gnr_datasources() %>%
  filter(title == "World Register of Marine Species") %>%
  pull(id)

### Automatic cleaning
# Set Survey code
hs_survey_code <- "DFO-HS"

HS <- HS %>%
  mutate(
    taxa2 = str_squish(verbatim_name),
    taxa2 = str_remove_all(taxa2, " spp.| sp.| spp| sp|NO "),
    taxa2 = str_to_sentence(str_to_lower(taxa2))
  )

# Get clean taxa
clean_auto <- clean_taxa(unique(HS$taxa2), input_survey = hs_survey_code,

```

```

save = F, output=NA)

#This leaves out the following species, which are all inverts
#Cancer branneri
#Cancer gracilis
#Cheiraster dawsoni
#Pandalopsis

#-----#
#### INTEGRATE CLEAN TAXA in DFO-HS survey data ####
#-----#

correct_taxa <- clean_auto %>%
  select(-survey)

clean_hs <- left_join(HS, correct_taxa, by=c("taxa2"="query")) %>%
  filter(!is.na(taxa)) %>% # query does not indicate taxa entry that were
  #removed in the cleaning procedure
  # so all NA taxa have to be removed from the surveys because: non-existing,
  #non marine or non fish
  rename(accepted_name = taxa,
         aphia_id = worms_id) %>%
  mutate(verbatim_aphia_id = NA,
         source = "DFO",
         timestamp = "09/2020") %>%
  select(survey, source, timestamp, haul_id, country, sub_area, continent, stat_rec, station, stratum,
         year, month, day, quarter, season, latitude, longitude,
         haul_dur, area_swept, gear, depth, sbt, sst, num, num_h, num_cpue, wgt,
         wgt_h, wgt_cpue,
         verbatim_name, verbatim_aphia_id, accepted_name, aphia_id, SpecCode,
         kingdom, phylum, class, order, family, genus, rank)

#check for duplicates
count_clean_hs <- clean_hs %>%
  group_by(haul_id, verbatim_name, accepted_name) %>%
  mutate(count = n())

#none!

#which ones are duplicated?
unique_name_match <- count_clean_hs %>%
  group_by(verbatim_name, accepted_name) %>%
  filter(count>1) %>%
  distinct(verbatim_name, accepted_name)

unique_name_match
#empty

# -----#
#### SAVE DATABASE IN GOOGLE DRIVE ####
# -----#

```

```
# Just run this routine should be good for all  
write_clean_data(data = clean_hs, survey = "HS", overwrite = T)
```

1. Overview of the survey data table

survey	source	timestamp	haul_id	country	sub_area	continent
DFO-HS	DFO	09/2020	59140-001	Canada	NA	n_america
DFO-HS	DFO	09/2020	59140-001	Canada	NA	n_america
DFO-HS	DFO	09/2020	59140-001	Canada	NA	n_america
DFO-HS	DFO	09/2020	59140-001	Canada	NA	n_america
DFO-HS	DFO	09/2020	59140-001	Canada	NA	n_america

stat_rec	station	stratum	year	month	day	quarter	season
NA	NA	NA	2005	5	27	NA	NA
NA	NA	NA	2005	5	27	NA	NA
NA	NA	NA	2005	5	27	NA	NA
NA	NA	NA	2005	5	27	NA	NA
NA	NA	NA	2005	5	27	NA	NA

latitude	longitude	haul_dur	area_swept	gear	depth
52.68587	-130.6156	0.3666667	0.133224	NA	137
52.68587	-130.6156	0.3666667	0.133224	NA	137
52.68587	-130.6156	0.3666667	0.133224	NA	137
52.68587	-130.6156	0.3666667	0.133224	NA	137
52.68587	-130.6156	0.3666667	0.133224	NA	137


sbt	sst	num	num_h	num_cpue	wgt
NA	NA	0	0.00000	0	2.40
NA	NA	4	10.90909	0	0.50
NA	NA	0	0.00000	0	0.44
NA	NA	0	0.00000	0	2.30
NA	NA	0	0.00000	0	1.80

wgt_h	wgt_cpue	verbatim_name	verbatim_aphia_id	accepted_name
6.545454	0	ATHERESTHES STOMIAS	NA	Atheresthes stomias
1.363636	0	CLUPEA PALLASII	NA	Clupea pallasii
1.200000	0	EOPSETTA JORDANI	NA	Eopsetta jordani
6.272727	0	GADUS CHALCOGRAMMUS	NA	Gadus chalcogrammus
4.909091	0	GADUS MACROCEPHALUS	NA	Gadus macrocephalus

aphia_id	SpecCode	kingdom	phylum	class	order	family
279792	517	Animalia	Chordata	Actinopteri	Pleuronectiformes	Pleuronectidae
151159	1520	Animalia	Chordata	Actinopteri	Clupeiformes	Clupeidae
280690	4237	Animalia	Chordata	Actinopteri	Pleuronectiformes	Pleuronectidae
300735	318	Animalia	Chordata	Actinopteri	Gadiformes	Gadidae
254538	308	Animalia	Chordata	Actinopteri	Gadiformes	Gadidae

2. Summary of sampling intensity


Number of hauls per year performed during the survey after data processing.



3. Summary of sampling variables from the survey

Here we show the yearly total and average of the following variables reported in the survey data:


- *area_swept*, swept area by the bottom trawl gear km^2
- *depth*, sampling depth in *m*
- *haul_dur*, haul sampling duration *hour*
- *number of marine fish taxa*, taxa were cleaned following the last version of taxonomy from the World Register of Marine Species (<https://www.marinespecies.org/>, October 2021)



4. Summary of biological variables

Here we display the yearly total and average across hauls of the following variables recorded in the data:


- *num_cpue*, number of individuals (abundance) in $\frac{\text{individuals}}{\text{km}^2}$
- *num_h*, number of individuals (abundance) in $\frac{\text{individuals}}{h}$
- *num*, number of individuals (abundance)
- *wgt_cpue*, weight in $\frac{kg}{\text{km}^2}$
- *wgt_h*, weight in $\frac{kg}{h}$
- *wgt*, weight in *kg*



5. Extreme values

Here we show a yearly total distribution of the biomass data to visualize outliers:


- *wgt*, total weight in *kg* per haul and year per haul and year, if available in the survey data
- *num*, total number of individuals, if available in the survey data




6. Summary of variables against swept area

Here we show the total abundance and number of taxa relationships with the area swept:

- *nbr_taxa*, number of marine fish taxa after taxonomic data cleaning
- *num*, number of individuals, if available in the survey data
- *wgt*, weight in *kg*, if available in the survey data




7. Abundance or Weight trends of the six most abundant species



8. Distribution mapping


Map of the sampling distribution in space. Note that we only show one year per coordinate.



9. Taxonomic flagging

This species flagging method was adapted from <https://github.com/pinskyLab/OceanAdapt/blob/master/R/add-spp-to-taxonomy.Rmd#L33>

Visualization of flagged taxa



Statistics related to the taxonomic flagging outputs


Total number of species	165.0
Percentage of species flagged	13.3


10. Spatio-temporal standardization

a. Standardization method 1

This standardization method was adapted from https://github.com/zoekitchel/rawl_spatial_turnover/blob/master/data_prep_code/species/explore_NorthSea_trimming.Rmd
It was run for hex resolution 7 and 8.


Plot of number of cells x years with overlaid flagging options








Map of hauls retained and removed per flagging method and threshold

Survey: DFO-HS - trimming 0% Hex res 7






Survey: DFO-HS - trimming 2% Hex res 7





Map of numbers of years removed per grid cell and flagging method/threshold






b. Standardization method 2

This standardization method was adapted from BioTIME code from https://github.com/Wubing-Xu/Range_size_winners_losers

Map of hauls retained and removed



c. Standardization summary

Statistics of hauls removed for each standardization method

summary	grid cell 7, 0% threshold	grid cell 7, 2% threshold	grid cell 8, 0% threshold	grid cell 8, 2% threshold	method 2 (biotime)
number of hauls removed	0	0	15.0	15.0	617.0
percentage of hauls removed	0	0	1.2	1.2	3.1