

# EBS: Eastern Bering Sea US survey data processing summary

fishglob, Aurore A. Maureaud, Juliano Palacios Abrantes, Zoë Kitchel, Dan Forrest, & Michelle Stuart

December, 2022

## Contents

General info . . . . .	1
Data cleaning in R . . . . .	1
1. Overview of the survey data table . . . . .	6
2. Summary of sampling intensity . . . . .	7
3. Summary of sampling variables from the survey . . . . .	8
4. Summary of biological variables . . . . .	9
5. Extreme values . . . . .	10
6. Summary of variables against swept area . . . . .	11
7. Abundance or Weight trends of the six most abundant species . . . . .	12
8. Distribution mapping . . . . .	13
9. Taxonomic flagging . . . . .	13
10. Spatio-temporal standardization . . . . .	14
a. Standardization method 1 . . . . .	14
b. Standardization method 2 . . . . .	18
c. Standardization summary . . . . .	18

## General info

This document presents the summary of the Eastern Bering Sea bottom trawl survey provided by Stan Kotwicki and Jim Thorson. It contains annual data from 1982-2019.

## Data cleaning in R

```
#####
#### R code to clean trawl survey Eastern Bering Sea
#### Public data Ocean Adapt
#### Contacts: Stan Kotwicki stan.kotwicki@noaa.gov Program Manager,
####           Groundfish Assessment Program, NOAA AFSC
####           Jim Thorson james.thorson@noaa.gov Program Leader,
####           Habitat and Ecological Processes Research, NOAA AFSC
#### Coding: Michelle Stuart, Dan Forrest, Zoë Kitchel November 2021
#####
#Alaska Fisheries Science Center - NOAA
#https://www.afsc.noaa.gov/RACE/groundfish/survey\_data/
#metadata_template.php?fname=RACEweb.xml
#This NOAA center provides data for the Aleutian Islands,
#Eastern Bering Sea, and Gulf of Alaska.
#Files provided by the Alaska Fisheries Science Center
#
#-----#
```

```

##### LOAD LIBRARIES AND FUNCTIONS #####
#-----#
library(tidyverse)
library(lubridate)
library(googledrive)
library(taxize) # for getting correct species names
library(magrittr) # for names wrangling
library(readxl)

source("functions/clean_taxa.R")
source("functions/write_clean_data.R")
fishglob_data_columns <- read_excel("standard_formats/fishglob_data_columns.xlsx")

#Data for the Gulf of Alaska can be accessed using the public
#Pinsky Lab OceanAdapt Git Hub Repository.
#Contact malin.pinsky@rutgers.edu for questions or help accessing

#-----#
##### PULL IN AND EDIT RAW DATA FILES #####
#-----#

#make list of csv files from OceanAdapt GitHub
files <- list(
  "https://raw.githubusercontent.com/pinskylab/OceanAdapt/master/data_raw/ebs1982_1984.csv",
  "https://raw.githubusercontent.com/pinskylab/OceanAdapt/master/data_raw/ebs1985_1989.csv",
  "https://raw.githubusercontent.com/pinskylab/OceanAdapt/master/data_raw/ebs1990_1994.csv",
  "https://raw.githubusercontent.com/pinskylab/OceanAdapt/master/data_raw/ebs1995_1999.csv",
  "https://raw.githubusercontent.com/pinskylab/OceanAdapt/master/data_raw/ebs2000_2004.csv",
  "https://raw.githubusercontent.com/pinskylab/OceanAdapt/master/data_raw/ebs2005_2008.csv",
  "https://raw.githubusercontent.com/pinskylab/OceanAdapt/master/data_raw/ebs2009_2012.csv",
  "https://raw.githubusercontent.com/pinskylab/OceanAdapt/master/data_raw/ebs2013_2016.csv",
  "https://raw.githubusercontent.com/pinskylab/OceanAdapt/master/data_raw/ebs2017_2019.csv")

# combine all of the data files into one table
ebs_data <- files %>%
  # read in all of the csv's in the files list
  map_dfr(read_csv) %>%
  # remove any data rows that have headers as data rows
  filter(LATITUDE != "LATITUDE", !is.na(LATITUDE)) %>%
  mutate(stratum = as.integer(STRATUM)) %>%
  # remove any extra white space from around spp and common names
  mutate(COMMON = str_trim(COMMON),
        SCIENTIFIC = str_trim(SCIENTIFIC))

# import the strata data
ebsstrat <-
  "https://raw.githubusercontent.com/pinskylab/OceanAdapt/master/data_raw/ebs_strata.csv"
ebs_strata <- read_csv(ebsstrat, col_types = cols(
  SubareaDescription = col_character(),
  StratumCode = col_integer(),
  Areakm2 = col_integer())

```

```

)) %>%
  rename(stratum = StratumCode)

#-----#
##### REFORMAT AND MERGE DATA FILES #####
#-----#

ebs <- left_join(ebs_data, ebs_strata, by = "stratum")

# are there any strata in the data that are not in the strata file?
stopifnot(nrow(filter(ebs, is.na(Areakm2))) == 0)

ebs <- ebs %>%
  mutate(
    # Create a unique haul_id
    haul_id = paste(formatC(VESSEL, width=3, flag=0), CRUISE,
                    formatC(HAUL, width=3, flag=0), LONGITUDE, LATITUDE, sep=''),
    #get rid of any use of -9999 as a no data marker
    numcpue = ifelse(NUMCPUE < -9000, NA, NUMCPUE),
    sbt = ifelse(BOT_TEMP < -9000, NA, BOT_TEMP),
    sst = ifelse(SURF_TEMP < -9000, NA, SURF_TEMP)) %>%
  rename(year = YEAR,
         latitude = LATITUDE,
         longitude = LONGITUDE,
         depth = BOT_DEPTH,
         spp = SCIENTIFIC,
         station = STATION,
         num_cpue.raw = numcpue, #units = number/hectare
         wgt_cpue.raw = WTCPUE #units = kg/hectare (1 hectare = 0.01 km^2)
) %>%
  mutate(
    #convert date to month and day columns
    datetime = mdy_hm(DATETIME),
    month = month(datetime),
    day = day(datetime),
    quarter = case_when(month %in% c(1,2,3) ~ 1,
                         month %in% c(4,5,6) ~ 2,
                         month %in% c(7,8,9) ~ 3,
                         month %in% c(10,11,12) ~ 4),
    season = 'NA',
    #convert cpue which is currently per hectare to per km^2 by multiplying by 100
    wgt_cpue = 100*wgt_cpue.raw,
    num_cpue = 100*num_cpue.raw
) %>%
  # remove non-fish
  filter(
    spp != '' &
      !grepl("egg", spp)) %>%
  # adjust spp names
  mutate(
    #Manual taxa cleaning (happens later in other get.x.R scripts)
    spp = ifelse(grepl("Lepidopsetta", spp), "Lepidopsetta sp.", spp),
    spp = ifelse(grepl("Myoxocephalus", spp) & !grepl("scorpius", spp),

```

```

        "Myoxocephalus sp.", spp),
spp = ifelse(grepl("Bathyraja", spp) & !grepl("panthera", spp),
             'Bathyraja sp.', spp)
) %>%
#finalize columns
mutate(survey = "EBS",
       country = "United States",
       sub_area = NA,
       continent = "n_america",
       stat_rec = NA,
       verbatim_name = spp,
       haul_dur = NA,
       gear = NA,
       num = NA,
       num_h = NA,
       wgt = NA,
       wgt_h = NA,
       area_swept = NA
) %>%
select(survey, haul_id, country, sub_area, continent, stat_rec, station,
       stratum, year, month, day, quarter, season, latitude, longitude,
       haul_dur, area_swept, gear, depth, sbt, sst,
       num, num_h, num_cpue, wgt, wgt_h, wgt_cpue, verbatim_name)

#check to make sure all looks right
#str(ebs)

#-----#
#### INTEGRATE CLEAN TAXA FROM TAXA ANALYSIS ####
#-----#


# Get WoRM's id for sourcing
wrn <- gnr_datasources() %>%
  filter(title == "World Register of Marine Species") %>%
  pull(id)

### Automatic cleaning
# Set Survey code
ebs_survey_code <- "EBS"

ebs_taxa <- ebs %>%
  select(verbatim_name) %>%
  mutate(
    taxa = str_squish(verbatim_name),
    taxa = str_remove_all(taxa, " spp.| sp.| spp| sp|NO "),
    taxa = str_to_sentence(str_to_lower(taxa))
  ) %>%
  pull(taxa) %>%
  unique()

# Get clean taxa
clean_auto <- clean_taxa(ebs_taxa, input_survey = ebs_survey_code, save = F,
                           output=NA) # takes 4.1 mins!

```

```

#check those with no match from clean_taxa()
#Beringius beringii                                no match
#Crangon communis                                 no match
#Crangon abyssorum                                no match
#Cheiraster dawsoni                                no match

#####clear all invertebrates

#-----#
##### INTEGRATE CLEAN TAXA in EBS survey data #####
#-----#


clean_taxa <- clean_auto %>%
  select(-survey)

clean_ebs <- left_join(ebs, clean_taxa, by=c("verbatim_name"="query")) %>%
  filter(!is.na(taxa)) %>% # query does not indicate taxa entry that were
  #removed in the cleaning procedure
  # so all NA taxa have to be removed from the surveys because: non-existing,
  #non marine or non fish
  rename(accepted_name = taxa,
         aphia_id = worms_id) %>%
  mutate(verbatim_aphia_id = NA,
        source = "NOAA",
        timestamp = my("03/2021")) %>%
  select(fishglob_data_columns$`Column name fishglob`)

#check for duplicates
count_clean_ebs <- clean_ebs %>% count(haul_id, accepted_name)

#no duplicates

# -----#
##### SAVE DATABASE IN GOOGLE DRIVE #####
# -----#


# Just run this routine should be good for all
write_clean_data(data = clean_ebs, survey = "EBS_v2", type = F, overwrite = T)

```

## 1. Overview of the survey data table

survey	haul_id	country	sub_area	continent	stat_rec	station
EBS	037198401019-162.7143358.353	United States	NA	n_america	NA	K-09
EBS	037198401019-162.7143358.353	United States	NA	n_america	NA	K-09
EBS	037198401019-162.7143358.353	United States	NA	n_america	NA	K-09
EBS	037198401019-162.7143358.353	United States	NA	n_america	NA	K-09
EBS	037198401019-162.7143358.353	United States	NA	n_america	NA	K-09

stratum	year	month	day	quarter	season	latitude	longitude
10	1984	6	14	2	NA	58.353	-162.7143
10	1984	6	14	2	NA	58.353	-162.7143
10	1984	6	14	2	NA	58.353	-162.7143
10	1984	6	14	2	NA	58.353	-162.7143
10	1984	6	14	2	NA	58.353	-162.7143

haul_dur	area_swept	gear	depth	sbt	sst
NA	NA	NA	31	4.2	3.6
NA	NA	NA	31	4.2	3.6
NA	NA	NA	31	4.2	3.6
NA	NA	NA	31	4.2	3.6
NA	NA	NA	31	4.2	3.6


num	num_h	num_cpue	wgt	wgt_h	wgt_cpue
NA	NA	30.73	NA	NA	139.38
NA	NA	399.46	NA	NA	11.15
NA	NA	61.46	NA	NA	111.51
NA	NA	1136.92	NA	NA	3094.20
NA	NA	61.46	NA	NA	5.56

verbatim_name	verbatim_aphia_id	accepted_name	aphia_id	SpecCode
Rajidae	NA	Rajidae	105711	NA
Mallotus villosus	NA	Mallotus villosus	126735	252
Gadus chalcogrammus	NA	Gadus chalcogrammus	300735	318
Gadus macrocephalus	NA	Gadus macrocephalus	254538	308
Podothecus accipenserinus	NA	Podothecus accipenserinus	254501	4153

kingdom	phylum	class	order	family	genus	rank
Animalia	Chordata	Elasmobranchii	Rajiformes	Rajidae	NA	Family
Animalia	Chordata	Actinopteri	Osmeriformes	Osmeridae	Mallotus	Species
Animalia	Chordata	Actinopteri	Gadiformes	Gadidae	Gadus	Species
Animalia	Chordata	Actinopteri	Gadiformes	Gadidae	Gadus	Species
Animalia	Chordata	Actinopteri	Perciformes	Agonidae	Podothecus	Species

## 2. Summary of sampling intensity


Number of hauls per year performed during the survey after data processing.



### 3. Summary of sampling variables from the survey

Here we show the yearly total and average of the following variables reported in the survey data:


- *area\_swept*, swept area by the bottom trawl gear  $km^2$
- *depth*, sampling depth in  $m$
- *haul\_dur*, haul sampling duration *hour*
- *number of marine fish taxa*, taxa were cleaned following the last version of taxonomy from the World Register of Marine Species (<https://www.marinespecies.org/>, October 2021)



#### 4. Summary of biological variables

Here we display the yearly total and average across hauls of the following variables recorded in the data:


- $num\_cpue$ , number of individuals (abundance) in  $\frac{individuals}{km^2}$
- $num\_h$ , number of individuals (abundance) in  $\frac{individuals}{h}$
- $num$ , number of individuals (abundance)
- $wgt\_cpue$ , weight in  $\frac{kg}{km^2}$
- $wgt\_h$ , weight in  $\frac{kg}{h}$
- $wgt$ , weight in  $kg$



## 5. Extreme values

Here we show a yearly total distribution of the biomass data to visualize outliers:

- $num\_cpue$ , number of individuals (abundance) in  $\frac{individuals}{km^2}$
- $wgt\_cpue$ , weight in  $\frac{kg}{km^2}$




## 6. Summary of variables against swept area

Here we show the total abundance and number of taxa relationships with the area swept:


- $nbr\_taxa$ , number of marine fish taxa after taxonomic data cleaning
- $num$ , number of individuals, if available in the survey data
- $wgt$ , weight in  $kg$ , if available in the survey data

## 7. Abundance or Weight trends of the six most abundant species



## 8. Distribution mapping


Map of the sampling distribution in space. Note that we only show one year per coordinate.



## 9. Taxonomic flagging

This species flagging method was adapted from <https://github.com/pinskylab/OceanAdapt/blob/master/R/add-spp-to-taxonomy.Rmd#L33>

Visualization of flagged taxa



Statistics related to the taxonomic flagging outputs


Total number of species	166.0
Percentage of species flagged	0.6


## 10. Spatio-temporal standardization

### a. Standardization method 1


This standardization method was adapted from [https://github.com/zookitchel/trawl\\_spatial\\_turnover/blob/master/data\\_prep\\_code/species/explore\\_NorthSea\\_trimming.Rmd](https://github.com/zookitchel/trawl_spatial_turnover/blob/master/data_prep_code/species/explore_NorthSea_trimming.Rmd)  
It was run for hex resolution 7 and 8.


Plot of number of cells x years with overlaid flagging options







Map of hauls retained and removed per flagging method and threshold





Map of numbers of years removed per grid cell and flagging method/threshold






### b. Standardization method 2

This standardization method was adapted from BioTIME code from [https://github.com/Wubing-Xu/Range\\_size\\_winners\\_losers](https://github.com/Wubing-Xu/Range_size_winners_losers)

Map of hauls retained and removed



### c. Standardization summary

Statistics of hauls removed for each standardization method

summary	grid cell 7, 0% threshold	grid cell 7, 2% threshold	grid cell 8, 0% threshold	grid cell 8, 2% threshold	method 2 (biotime)
number of hauls removed	471.0	0	1260.0	5	429.0
percentage of hauls removed	3.3	0	8.9	0	0.3