






Chimaera monstrosa 10.0 -100 -7.5 nolength numh 5.0 -50 **-**2.5 -0.0 -0 -1960 1980 2000 2020 1960 1980 2000 2020 Year year
















Hexanchus griseus 0.30 -0.3 -0.25 -0.2 nolength 0.20 -0.1 -0.15 -0.10 -1980 2000 1980 2000 2020 1960 2020 1960 Year year




Lepidorhombus whiffiagoni 90 -60 nolength numh 30 -0 -1980 1980 2000 1960 2000 2020 1960 2020 Year year




Leucoraja fullonica 0.3 -1.5 -0.2 -1.0 nolength numh 0.5 -0.1 -0.0 -1980 1980 2000 1960 2000 2020 2020 1960 Year year


Leucoraja naevus 8 -3 **-**6 -2 nolength qwnu 4-1 -2-0 -1960 1980 1980 2000 2020 1960 2000 2020 Year year


Lophius budegassa 1.25 -3 -1.00 -0.75 -2 nolength numh 0.50 -1 -





Lophius piscatorius 3 -6 **-**2 nolength 1 -2 -0 -1960 1980 1980 2000 2020 2000 2020 1960 Year year






























Scophthalmus rhombus 2.5 -1.5 -2.0 -1.5 -1.0 nolength numh 1.0 -0.5 -0.5 -0.0 -0.0 -1980 1980 1960 2000 1960 2000 2020 2020 Year year









Torpedo marmorata 0.6 -0.3 -0.4 nolength unu 0.2 -0.2 -0.1 -0.0 -0.0 -1980 2000 2020 1980 2020 1960 2000 Year year