HIGHER-ORDER CENTER CONDITIONS AND THE LIJUN–YUN CONJECTURE FOR POLYNOMIAL ABEL EQUATIONS

ALEXANDRE M. ALVES, ABÍLIO LEMOS, AND ANDERSON L. A. DE ARAUJO

Abstract. We study the center problem for the Abel equation $x' = f(t)x^3 + g(t)x^2$ on $[-1,1]$, with emphasis on $g(t) = t^{n-1}$, n even. Lijun and Yun conjectured that, for $g(t) = 2t$ and f polynomial, $x = 0$ is a center if and only if f is odd; in a previous paper we proved this for $\deg f \le 5$ by showing that a center forces the first three moments $\int_{-1}^{1} fG^k\,dt$ $(k = 0, 1, 2)$ to vanish. We compute the next three center conditions in closed form: a center forces $m_3 = \frac{1}{2}\int gF^2$, $m_4 = 2\int gGF^2$ and an explicit identity for m_5, where F, G are the primitives of f, g. This explains why m_3 need not vanish at a center. For odd g vanishing only at the origin we prove that the vanishing of *all* moments is equivalent to f being odd, so the conjecture is equivalent to the statement that a center forces all moments to vanish. We give an elementary sign-change criterion that unifies and extends the known cases, a closed Cauchy-type formula for the relevant moment determinants, and a "hyperplane" theorem for $\deg f_{\mathrm{e}} \le 6$ showing that the conjecture can only fail if the odd part of f lies in an explicit affine subspace of codimension at most two. Finally, using exact Gröbner basis computations over $\mathbb{Q}$, we prove the conjecture for $g = t^{n-1}$ when $\deg f \le 7$ and $2 \le n \le 20$ (and, more generally, when $\deg f_{\mathrm{e}} \le 6$ and $\deg f_{\mathrm{o}} \le 9$), and for $g = t$ when $\deg f \le 8$.

1. Introduction

Consider the Abel differential equation

$$x' = f(t)\,x^3 + g(t)\,x^2, \qquad t \in [-1,1], \tag{1.1}$$

where f and g are real analytic on a neighbourhood of $[-1,1]$. We say that (1.1) has a *center* at $x = 0$ (on $[-1,1]$) if every solution with $|x(-1)|$ small enough satisfies $x(1) = x(-1)$. This center problem goes back to the classical Poincaré center–focus problem: in polar coordinates, and after Cherkas' transformation [12], a planar system $\dot{x} = -y + P_n(x,y)$, $\dot{y} = x + Q_n(x,y)$ with homogeneous nonlinearities becomes a trigonometric Abel equation, and its centers correspond to centers of the latter. The polynomial version on an interval was proposed by Briskin, Françoise and Yomdin as a model of the classical problem [6, 7, 8]; see also [23] for the related problem of counting solutions with $x(0) = x(1)$.

Throughout we write

$$F(t) = \int_{-1}^{t} f, \qquad G(t) = \int_{-1}^{t} g, \qquad m_k = \int_{-1}^{1} f(t)G(t)^k\,dt \quad (k \ge 0),$$

and $f = f_{\mathrm{e}} + f_{\mathrm{o}}$ for the decomposition into even and odd parts.

Composition and moments. The basic sufficient condition is the *composition condition* of Alwash and Lloyd [4]: if $F = \tilde{F} \circ W$ and $G = \tilde{G} \circ W$ for some W with $W(-1) = W(1)$, then (1.1) has a center. For small degrees and many special classes it is also necessary [6, 7, 8, 13, 3, 18], but not in general: counterexamples are known for trigonometric [3, 19] and,

2020 *Mathematics Subject Classification.* Primary 34C07; Secondary 34A34, 34C25, 30E05, 13P10.

Key words and phrases. Abel equation, center problem, moment conditions, composition condition, Lijun–Yun conjecture, Gröbner bases.

more recently, for polynomial coefficients [20]. If f is replaced by εf, the first-order condition in ε is the vanishing of all moments m_k [7]; the moments do not force composition [24], the polynomial moment problem was solved by Pakovich and Muzychuk [25], and Pakovich proved that $x' = gx^2 + \varepsilon f x^3$ has a center for all ε if and only if the composition condition holds [26]. Higher-order ("double") moments, iterated integrals and the algebraic geometry of the center set are studied in [11, 10, 14, 15, 9, 17, 1].

The Lijun–Yun conjecture. Lijun and Yun [22, Thm. 5.5] proved that $x' = 2tx^2 + \varepsilon f(t)x^3$ has a center for all small ε if and only if the polynomial f is odd (this also follows from [26]), and asked whether this remains true for a fixed ε:

Conjecture 1.1 (Lijun–Yun [22, Rem. 5.6])**.** *Let f be a polynomial. The equation $x' = 2t\,x^2 + f(t)\,x^3$, $t \in [-1,1]$, has a center at $x = 0$ if and only if f is odd.*

Since $x \mapsto \lambda x$ replaces (f, g) by $(\lambda^2 f, \lambda g)$, the coefficient $2t$ may be replaced by ct, $c \neq 0$. Following [2], we consider more generally $g(t) = t^{n-1}$ with n even. The "if" part is the composition condition with $W(t) = t^2$. For the "only if" part, in [2] we proved that a center of (1.1) forces $m_0 = m_1 = m_2 = 0$, and deduced the conjecture when $\deg f \leq 5$, or more generally when g is odd, vanishes only at 0, and the even part of f changes sign at most twice on $[0,1]$. They also observed, using an example of Giné, Grau and Santallusia [19], that a center need not satisfy $m_3 = 0$.

Main results. Our first result computes the next center conditions in closed form. Put $H(t) = \int_{-1}^{t} fG$.

Theorem A (Higher-order center conditions)**.** *Assume $G(1) = 0$. If (1.1) has a center at $x = 0$, then $m_0 = m_1 = m_2 = 0$ and*

$$m_3 = \frac{1}{2}\int_{-1}^{1} g\,F^2\,dt, \tag{1.2}$$

$$m_4 = 2\int_{-1}^{1} g\,G\,F^2\,dt, \tag{1.3}$$

$$m_5 = \frac{1}{2}\int_{-1}^{1} g\,F^3\,dt + \frac{9}{2}\int_{-1}^{1} g\,G^2F^2\,dt - \int_{-1}^{1} f\,G^2H\,dt. \tag{1.4}$$

For the example of [19] (on $[0, 2\pi]$), both sides of (1.2) equal $\pi/2$, which explains the observation of [2]; see Example 3.2.

From now on we say that g satisfies *Condition I* if g is odd and vanishes only at $t = 0$ [2]; $g(t) = t^{n-1}$, n even, is the model case.

Theorem B (Reformulation of the conjecture)**.** *Let g satisfy Condition I and let f be continuous. Then $m_k = 0$ for all $k \geq 0$ if and only if f is odd. Consequently, for such g and polynomial (or analytic) f, the following are equivalent:*

(i) *every center of (1.1) has f odd;*
(ii) *every center of (1.1) has $m_k = 0$ for all k.*

Theorem B shows that, for $g = t^{n-1}$, Conjecture 1.1 is precisely the statement "center $\Rightarrow$ moment conditions" (an implication which is open in general; cf. the diagram in [2, §1.3]). The proof is elementary and rests on a sign-change lemma (Lemma 5.1). The same lemma, combined with Theorem A, gives the following criterion. For $\varphi, \psi \in L^2(0,1)$ and a weight w put $\langle \varphi, \psi \rangle_w = \int_0^1 w\varphi\psi\,dt$, and set

$$A(t) = \int_0^t f_{\mathrm{e}}, \qquad C(t) = \int_t^1 f_{\mathrm{o}} \qquad (0 \leq t \leq 1).$$

Theorem C (Sign-change criterion)**.** *Let g satisfy Condition I and let f be a polynomial. Write $f_{\mathrm{e}}(t) = p(t^2)$. In each of the following cases, (1.1) has a center at $x = 0$ if and only if f is odd:*

(a) *p changes sign at most twice in $(0,1)$;*
(b) *p changes sign at most three times in $(0,1)$ and $\langle A, C\rangle_g = 0$;*
(c) *p changes sign at most four times in $(0,1)$ and $\langle A, C\rangle_g = \langle A, C\rangle_{gG} = 0$.*

Case (a) is [2, Thm. 1.10], here with a short self-contained proof; it contains [2, Thm. 1.12] and the case $\deg f_{\mathrm{e}} \le 4$ with f_{o} arbitrary. Since A depends on f_{e}, the orthogonality conditions in (b) and (c) are implicit. For $\deg f_{\mathrm{e}} \le 6$ they can be replaced by explicit conditions on f_{o} only:

Theorem D (Hyperplane theorem)**.** *Let g satisfy Condition I with $g > 0$ on $(0,1]$. There is a nonzero even polynomial e of degree at most 6, unique up to a constant factor, with $\int_{-1}^{1} eG^k = 0$ for $k = 0, 1, 2$; moreover $\mu := \int_{-1}^{1} eG^3 \neq 0$. Let $\nu := \int_{-1}^{1} eG^4$ and $A_e(t) = \int_0^t e$. If f is a polynomial with $\deg f_{\mathrm{e}} \le 6$ and (1.1) has a center, then either f is odd, or $f_{\mathrm{e}} = a\,e$ with $a \neq 0$ and*

$$\langle A_e, C\rangle_g = -\frac{\mu}{2}, \qquad \langle A_e, C\rangle_{gG} = -\frac{\nu}{8}. \tag{1.5}$$

In particular, if the odd part of f violates either equality in (1.5), then (1.1) has a center if and only if f is odd.

Since C depends linearly on f_{o}, (1.5) defines an affine subspace of codimension at most two in the space of odd parts. Finally, the higher center conditions can be handled by exact computer algebra:

Theorem E (Computer-assisted)**.** *Let $g(t) = t^{n-1}$.*

(i) *For every even n with $2 \le n \le 20$ and every polynomial f with $\deg f_{\mathrm{e}} \le 6$ and $\deg f_{\mathrm{o}} \le 9$, (1.1) has a center if and only if f is odd; for $n \in \{2, 4, 6\}$ the same holds with $\deg f_{\mathrm{o}} \le 11$. In particular Conjecture 1.1 (and its version with t^{n-1}, $2 \le n \le 20$) holds whenever $\deg f \le 7$.*
(ii) *For $n = 2$, Conjecture 1.1 holds whenever $\deg f \le 8$.*

The proof of Theorem E computes the first K center conditions as exact polynomials in the coefficients of f and shows, by Gröbner bases over $\mathbb{Q}$, that they have no common complex zero with $f_{\mathrm{e}} \neq 0$ (Section 7). Interestingly, the number of conditions needed is sharp: for $\deg f \le 7$ the first 8 conditions admit complex common zeros with $f_{\mathrm{e}} \neq 0$ for every n tested, whereas 9 conditions exclude them.

The paper is organized as follows. Section 2 sets up the expansion of the return map. Theorem A is proved in Section 3. Section 4 contains the parity reduction, Section 5 the moment lemmas and Theorem B, and Section 6 the proofs of Theorems C and D. Section 7 describes the computations behind Theorem E, and Section 8 lists further remarks and open problems.

2. The expansion of the return map

Let $x(t, \rho)$ be the solution of (1.1) with $x(-1, \rho) = \rho$.

Lemma 2.1. *For $|\rho|$ small, the problem*

$$u' = g(t) + \frac{\rho\, f(t)}{1 - \rho u}, \qquad u(-1) = 0, \tag{2.1}$$

has a solution $u(t,\rho)$ on $[-1,1]$, analytic in ρ, and $x(t,\rho)=\rho/(1-\rho u(t,\rho))$. In particular, (1.1) has a center at $x=0$ if and only if $u(1,\rho)\equiv 0$, i.e. if and only if $u_k(1)=0$ for all $k\geq 0$, where $u=\sum_{k\geq 0}u_k\rho^k$.

Proof. For $\rho=0$, (2.1) has the solution $u=G$ on $[-1,1]$; the right-hand side is analytic in (u,ρ) near $\{(G(t),0)\}$, so by analytic dependence on parameters $u(t,\rho)$ exists on $[-1,1]$ and is analytic in ρ for $|\rho|$ small. The function $y=\rho/(1-\rho u)$ satisfies $y(-1)=\rho$ and $y'=\rho^2u'/(1-\rho u)^2=y^2(g+fy)$, so $y=x$ by uniqueness. For $\rho\neq 0$, $x(1,\rho)=\rho$ is equivalent to $u(1,\rho)=0$. □

Writing $S:=(1-\rho u)^{-1}=\sum_{k\geq 0}S_k\rho^k$, the identity $S=1+\rho uS$ and (2.1) give the recursion

$$u_0=G,\qquad u_k(t)=\int_{-1}^{t} f\,S_{k-1}\quad(k\geq 1),\qquad S_0=1,\qquad S_k=\sum_{i=0}^{k-1}u_i\,S_{k-1-i}. \tag{2.2}$$

Explicitly,

$$\begin{aligned}
&S_1=u_0,\qquad S_2=u_1+u_0^2,\qquad S_3=u_2+2u_0u_1+u_0^3,\\
&S_4=u_3+2u_0u_2+u_1^2+3u_0^2u_1+u_0^4,\\
&S_5=u_4+2u_0u_3+2u_1u_2+3u_0^2u_2+3u_0u_1^2+4u_0^3u_1+u_0^5.
\end{aligned}$$

In particular $u_0(1)=G(1)$, so $G(1)=0$ is necessary for a center. When f and g are polynomials, every u_k is a polynomial in t whose coefficients are polynomials with rational coefficients in those of f and g.

Remark 2.2. The expansion in powers of ρ of [22, 2] uses $x=\rho+\sum r_k\rho^k$; the variable u is related by $1/x=1/\rho-u$. Its advantage is that $u_{k+1}(1)$ contains the moment m_k *linearly*, with corrections that are at least quadratic in f (Theorem A).

3. Proof of Theorem A

Let $K_2(t)=\int_{-1}^{t}fG^2$. By (2.2),

$$u_1=F,\qquad u_2=H,\qquad u_3=\tfrac12F^2+K_2,$$

hence $u_1(1)=m_0$, $u_2(1)=m_1$ and $u_3(1)=\frac12m_0^2+m_2$. A center thus forces $m_0=m_1=m_2=0$ (this is [2, Thm. 1.7]). From now on assume $G(1)=m_0=m_1=m_2=0$, so that

$$F(\pm1)=G(\pm1)=H(\pm1)=K_2(\pm1)=0.$$

All integrals below are over $[-1,1]$, and integrations by parts produce no boundary terms.

Order 4. By (2.2), $u_4(1)=\int f(H+2GF+G^3)$. Since $F'=f$, $\int fH=-\int FfG$, and since $(F^2)'=2fF$, $\int fFG=-\frac12\int gF^2$. Hence

$$u_4(1)=m_3+\int fFG=m_3-\frac12\int gF^2. \tag{3.1}$$

Order 5. $u_5(1)=\int fS_4$. We have $\int fF^2=\frac13F^3\big|_{-1}^{1}=0$, $\int fK_2=-\int FfG^2$, $2\int fGH=2\int HH'=0$ and $\int fG^4=m_4$. Therefore

$$u_5(1)=m_4+2\int fFG^2=m_4+\int G^2(F^2)'=m_4-2\int gGF^2. \tag{3.2}$$

Order 6. $u_6(1)=\int fS_5$. Integrating by parts, $\int fu_4=-\int Ff(H+2GF+G^3)$, so

$$u_6(1)=m_5+\int fFH+2\int fGF^2+3\int fG^3F+2\int fGK_2+3\int fG^2H.$$

Now $\int fGK_2 = \int H'K_2 = -\int HfG^2$, $\int fFH = \int F'FH = -\frac{1}{2}\int fGF^2$, $\int fGF^2 = \frac{1}{3}\int G(F^3)' = -\frac{1}{3}\int gF^3$ and $\int fG^3F = \frac{1}{2}\int G^3(F^2)' = -\frac{3}{2}\int gG^2F^2$. Collecting terms,

$$u_6(1) = m_5 - \frac{1}{2}\int gF^3 - \frac{9}{2}\int gG^2F^2 + \int fG^2H. \tag{3.3}$$

By Lemma 2.1, a center gives $u_4(1) = u_5(1) = u_6(1) = 0$, which is (1.2)–(1.4). □

Remark 3.1. The proof of (3.1) only uses $m_0 = m_1 = 0$. The identities (3.1)–(3.3) hold on any interval $[a, b]$ with $G(a) = G(b) = 0$, with the obvious changes. We also verified them symbolically for generic polynomials of degree 7 and $g = t^{n-1}$, $n = 2, 4, 6$, and checked the recursion (2.2) against high-precision numerical integration of (1.1).

Example 3.2. The trigonometric Abel equation

$$\frac{dx}{d\theta} = (\sin\theta - \sin 2\theta + \sin 3\theta)x^3 + (\cos\theta + 2\cos 2\theta)x^2, \qquad \theta \in [0, 2\pi],$$

has a center [19], and $m_0 = m_1 = m_2 = 0$, $m_3 = \pi/2$ [2, Rem. 1.8]. Here $G = \sin\theta + \sin 2\theta$ vanishes at 0 and 2π, and a direct symbolic computation gives $\frac{1}{2}\int_0^{2\pi} gF^2\,d\theta = \pi/2$, in agreement with (1.2). Thus $m_3 \neq 0$ at this center is not an anomaly: the fourth center condition is $m_3 = \frac{1}{2}\int gF^2$, not $m_3 = 0$.

Remark 3.3. The integrals in (1.2)–(1.4) are iterated integrals of $f\,dt$ and $g\,dt$; e.g. $\int fG^2H = \iint_{-1<s<t<1} f(s)G(s)\,f(t)G(t)^2\,ds\,dt$. Theorem A is therefore an explicit instance of Brudnyi's expansion of the first return map in iterated integrals [11], and of the second-order ("double moment") conditions studied in [14, 9], in a normalization in which the moments appear linearly.

4. Parity

Assume that g satisfies Condition I. Replacing x by $-x$ changes (f, g) into $(f, -g)$, so we may assume $g > 0$ on $(0, 1]$. Then G is even, $G(\pm 1) = 0$, $G < 0$ on $(-1, 1)$, and G restricted to $[0, 1]$ is an increasing bijection onto $[G(0), 0]$; let $\tau : [G(0), 0] \to [0, 1]$ be its inverse. Since fG^k has the same parity as f, only f_{e} contributes to the moments:

$$m_k = \int_{-1}^{1} f_{\mathrm{e}}\,G^k\,dt = 2\int_0^1 f_{\mathrm{e}}\,G^k\,dt. \tag{4.1}$$

Lemma 4.1. *Assume $m_0 = 0$. For $t \in [0, 1]$,*

$$F(t) = A(t) - C(t), \qquad F(-t) = -A(t) - C(t).$$

Consequently, for every $j \geq 0$,

$$\int_{-1}^{1} gG^jF^2\,dt = -4\langle A, C\rangle_{gG^j}, \qquad \int_{-1}^{1} gG^jF^3\,dt = 2\int_0^1 gG^j\,(A^3 + 3AC^2)\,dt.$$

Proof. By (4.1), $\int_0^1 f_{\mathrm{e}} = 0$. For $t \in [0, 1]$, $\int_{-1}^t f_{\mathrm{e}} = \int_0^t f_{\mathrm{e}} = A(t)$ and $\int_{-1}^t f_{\mathrm{o}} = -\int_t^1 f_{\mathrm{o}} = -C(t)$, which gives $F(t)$; the formula for $F(-t)$ follows from the parities of $f_{\mathrm{e}}, f_{\mathrm{o}}$. Since gG^j is odd, $\int_{-1}^1 gG^j\Phi(F) = \int_0^1 gG^j[\Phi(F(t)) - \Phi(F(-t))]dt$ for any Φ, and $(A - C)^2 - (A + C)^2 = -4AC$, $(A - C)^3 + (A + C)^3 = 2A^3 + 6AC^2$. □

Similarly, if $m_1 = 0$, then $H(t) = B(t) - D(t)$ and $H(-t) = -B(t) - D(t)$ on $[0, 1]$, where $B(t) = \int_0^t f_{\mathrm{e}}G$ and $D(t) = \int_t^1 f_{\mathrm{o}}G$, and a computation as above gives $\int_{-1}^1 fG^2H = 2\int_0^1 G^2(f_{\mathrm{o}}B - f_{\mathrm{e}}D)$. Combining with Theorem A:

Corollary 4.2. *Let g satisfy Condition I. If (1.1) has a center, then $m_0 = m_1 = m_2 = 0$ and*

$$m_3 = -2\langle A, C\rangle_g, \qquad m_4 = -8\langle A, C\rangle_{gG},$$
$$m_5 = \int_0^1 g(A^3 + 3AC^2) - 18\langle A, C\rangle_{gG^2} - 2\int_0^1 G^2(f_{\mathrm{o}}B - f_{\mathrm{e}}D).$$

Remark 4.3. In the variable $s = G(t)$ one has $g\,dt = ds$, so $\langle A, C\rangle_{gG^j} = \int_{G(0)}^0 (AC)(\tau(s))\, s^j\, ds$: the fourth and fifth center conditions say that the 0-th and 1-st moments of AC in the variable G are determined by m_3 and m_4, the corresponding moments of f_{e} (cf. Proposition 5.2). For $g = t^{n-1}$ the weight t^{n-1} on $(0,1)$ is a shifted Jacobi weight, and $\langle A, C\rangle_g$ can be expanded in the polynomials $P_k^{(0,n-1)}(2t-1)$; in particular, disjoint Jacobi spectral supports of A and C imply $\langle A, C\rangle_g = 0$.

5. Moments and sign changes

We say that $\varphi \in L^1(\alpha, \beta)$ has at most N sign changes if there are $\alpha = s_0 < s_1 < \cdots < s_r < s_{r+1} = \beta$, $r \le N$, such that $(-1)^i\varphi \ge 0$ a.e. on (s_i, s_{i+1}), or $(-1)^i\varphi \le 0$ a.e. on all of them.

Lemma 5.1. *If $\varphi \in L^1(\alpha, \beta)$ has at most N sign changes and $\int_\alpha^\beta \varphi(s)s^k\, ds = 0$ for $k = 0, \dots, N$, then $\varphi = 0$ a.e.*

Proof. With $s_1, \dots, s_r$ as above, $\varphi(s)\prod_{i=1}^r (s - s_i)$ has constant sign and, being φ times a polynomial of degree $r \le N$, has zero integral. Hence it vanishes a.e., and so does φ. (This is the classical Chebyshev-system argument; see [21, 27].) □

Proposition 5.2. *Let g satisfy Condition I with $g > 0$ on $(0,1]$, and let $f_{\mathrm{e}}(t) = p(t^2)$ with p continuous on $[0,1]$. Then*

$$m_k = 2\int_{G(0)}^0 \varphi(s)\, s^k\, ds, \qquad \varphi(s) = \frac{p(\tau(s)^2)}{g(\tau(s))},$$

where $\varphi \in L^1(G(0), 0)$ has the same sign changes as p on $(0,1)$. In particular, if p has at most N sign changes in $(0,1)$ and $m_0 = \cdots = m_N = 0$, then $p \equiv 0$.

Proof. Substitute $s = G(t)$ in (4.1). Integrability holds because $\int |\varphi|\, ds = \int_0^1 |p(t^2)|\, dt$; the singularity of $1/g$ at $t = 0$ is compensated by $ds = g\,dt$. Since $g > 0$ and τ is increasing, φ and $p \circ \tau^2$ have the same sign changes. Apply Lemma 5.1. □

Proof of Theorem B. If f is odd, fG^k is odd and $m_k = 0$. Conversely, if all m_k vanish, then by Proposition 5.2 all moments of $\varphi \in L^1$ on the compact interval $[G(0), 0]$ vanish; by the Weierstrass approximation theorem, $\varphi = 0$ a.e., i.e. $f_{\mathrm{e}} \equiv 0$. The equivalence of (i) and (ii) follows. □

Remark 5.3. For $g = t^{n-1}$ the first part of Theorem B can also be derived from [25]: up to affine changes of variable, the right compositional factors of $G = (t^n - 1)/n$ are t^d with $d \mid n$, and $W(-1) = W(1)$ forces d to be even. Note also that $\{t^{nk}\}_{k\ge 0}$ is complete in $C[0,1]$ by the Müntz theorem [5], since $\sum 1/(nk) = \infty$.

For $g = t^{n-1}$ the moment matrices have a closed-form determinant, which generalizes the 3×3 determinant computed in [2, Thm. 1.12].

Proposition 5.4. *Let $n \ge 2$ be even, $G(t) = (t^n - 1)/n$ and $M^{(N)} = \left(\int_{-1}^1 t^{2j} G(t)^k\, dt\right)_{k,j=0}^N$. Then*

$$\det M^{(N)} = \frac{2^{N+1}}{n^{N(N+1)/2}} \cdot \frac{\prod_{0\le i<k\le N} 2n\,(k-i)^2}{\prod_{i,j=0}^N (ni + 2j + 1)} > 0.$$

Proof. Expanding $(t^n-1)^k$, row k of $M^{(N)}$ equals n^{-k} times row k of $\big(2/(ni+2j+1)\big)_{i,j}$ plus a combination of the previous rows. Hence $\det M^{(N)} = 2^{N+1}n^{-N(N+1)/2}\det\big(1/(x_i+y_j)\big)$ with $x_i = ni+1$, $y_j = 2j$, and the Cauchy determinant equals $\prod_{i<k}(x_k-x_i)(y_k-y_i)/\prod_{i,j}(x_i+y_j)$. □

Remark 5.5. Proposition 5.4 shows that $m_0 = \cdots = m_N = 0$ means $f_{\mathrm{e}} \perp t^{nk}$ in $L^2(-1,1)$ for $k = 0,\dots,N$, an orthogonality problem of Müntz type [5]. For $n = 2$ this is orthogonality to all even polynomials of degree $\le 2N$; e.g. for $\deg f_{\mathrm{e}} \le 6$ and $N = 2$, f_{e} is a multiple of the Legendre polynomial P_6.

6. Proofs of Theorems C and D

Proof of Theorem C. If f is odd, write $f(t) = t\hat{f}(t^2)$, $g(t) = t\hat{g}(t^2)$; then F and G are functions of $W(t) = t^2$, with $W(-1) = W(1)$, and the composition condition [4] gives a center. Conversely, assume a center; we may assume $g > 0$ on $(0,1]$. By Corollary 4.2, $m_0 = m_1 = m_2 = 0$ and, under the orthogonality hypotheses of (b) (resp. (c)), also $m_3 = 0$ (resp. $m_3 = m_4 = 0$). Proposition 5.2 with $N = 2,3,4$ gives $p \equiv 0$. □

In particular, Theorem C(b) contains the statement: if $g = t^{n-1}$, $\deg f_{\mathrm{e}} \le 6$ and A, C are orthogonal in $L^2((0,1), t^{n-1}dt)$, then a center forces f to be odd.

Proof of Theorem D. Let V be the 4-dimensional space of even polynomials of degree ≤ 6. By Proposition 5.2 with $N = 3$ (a polynomial of degree ≤ 3 in $y = t^2$ has at most three sign changes), the linear map $V \to \mathbb{R}^4$, $f_{\mathrm{e}} \mapsto (m_0,\dots,m_3)$, is injective, hence bijective. Therefore the kernel of (m_0, m_1, m_2) on V is spanned by a single $e \neq 0$, and $\mu = m_3(e) \neq 0$. If (1.1) has a center and $\deg f_{\mathrm{e}} \le 6$, then $m_0 = m_1 = m_2 = 0$, so $f_{\mathrm{e}} = ae$ and $A = aA_e$. By Corollary 4.2,

$$a\mu = m_3 = -2a\langle A_e, C\rangle_g, \qquad a\nu = m_4 = -8a\langle A_e, C\rangle_{gG}.$$

If $a \neq 0$, this is (1.5). □

Remark 6.1. The orthogonality hypothesis $\langle A, C\rangle_g = 0$ of Theorem C(b) avoids the first hyperplane in (1.5), since $\mu \neq 0$; Theorem D shows that it suffices to avoid either one. On the exceptional set, the sixth center condition (Corollary 4.2) becomes a quadratic equation in (a, f_{o}) with a term $a^2\int_0^1 gA_e^3$; this is the starting point of the computations of Section 7.

Example 6.2. Let $f_{\mathrm{o}} = b_0t + b_1t^3 + b_2t^5 + b_3t^7$. For $g = t$ $(n = 2)$, $e(t) = 231t^6 - 315t^4 + 105t^2 - 5 = 16P_6(t)$ and, with $f_{\mathrm{e}} = a\,e/231$,

$$u_4(1) = \tfrac{64a}{530675145}\,(255b_1 + 238b_2 + 189b_3 + 765),$$
$$u_5(1) = \tfrac{256a}{10082827755}\,(4845b_0 + 969b_1 + 133b_2 - 63b_3 - 3876).$$

For $g = t^3$ $(n = 4)$, $e(t) = 385t^6 - 585t^4 + 231t^2 - 15$ and, with $f_{\mathrm{e}} = a\,e/385$,

$$u_4(1) = -\tfrac{64a}{5601570975}\,(1615b_0 - 399b_2 - 462b_3 - 693),$$
$$u_5(1) = \tfrac{256a}{128836132425}\,(16169b_0 + 5313b_1 + 1817b_2 + 561b_3 - 2376).$$

In both cases the exceptional set (1.5) is a 2-dimensional affine plane in $\mathbb{R}^4$.

7. Computer-assisted proof of Theorem E

Fix n and degrees $2E$, $2R+1$, and write

$$f(t) = \sum_{j=0}^{E} a_j t^{2j} + \sum_{j=0}^{R} b_j t^{2j+1}, \qquad g(t) = t^{n-1}.$$

By (2.2), each $c_k := u_k(1)$ is a polynomial in $\mathbb{Q}[a_0, \dots, a_E, b_0, \dots, b_R]$, and by Lemma 2.1, a center forces $c_1 = \cdots = c_K = 0$ for every K. Let I_K be the ideal generated by $c_1, \dots, c_K$. If, for each i,

$$1 \in I_K + \langle 1 - z\, a_i \rangle \subset \mathbb{Q}[z, a, b], \tag{7.1}$$

then every common complex zero of $c_1, \dots, c_K$ has $a_i = 0$ (Rabinowitsch's trick), so a center forces $f_{\mathrm{e}} = 0$, and Theorem C's sufficiency part finishes the proof. Condition (7.1) holds if and only if the reduced Gröbner basis of $I_K + \langle 1 - z a_i \rangle$ is $\{1\}$.

Reduction of variables. By Theorem A, c_1, c_2, c_3 vanish if and only if $m_0 = m_1 = m_2 = 0$, a linear system in $a_0, \dots, a_E$ of rank 3 (Proposition 5.4). We therefore parametrize its solution space as $f_{\mathrm{e}} = \sum_{i=1}^{E-2} \gamma_i e_i$, with an explicit basis e_i of integer polynomials, check that c_1, c_2, c_3 vanish identically on this family, and apply (7.1) to $c_4, \dots, c_K$ and the variables γ_i. This is equivalent to the unreduced test, and much faster.

Implementations. The polynomials c_k were computed by two independent implementations of (2.2): one in `SymPy` (exact rational arithmetic) and one written directly in `Singular` [16]. We checked that both give identical values of $c_1, \dots, c_6$ at random rational parameters, and that the truncated series $\rho/(1 - \rho \sum_{k \le 8} c_k \rho^k)$ agrees with the numerical solution of (1.1) up to the expected order $O(\rho^{11})$. Gröbner bases were computed with `SymPy`'s `groebner` and with `Singular`'s `std` (degree reverse lexicographic order, characteristic 0). All rows of Table 1 were obtained with `Singular`; the rows with $\deg f_{\mathrm{e}} \le 6$ were also reproduced with `SymPy` for $n \in \{2, 4, 6\}$ (and $\deg f_{\mathrm{o}} \le 7$), and for $n = 2$, $\deg f_{\mathrm{o}} \le 9$, with identical outcomes. The code is available in the repository cited in the Data availability statement.

n	$\deg f_{\mathrm{e}} \le$	$\deg f_{\mathrm{o}} \le$	K	common zeros of $c_1, \dots, c_K$ with $f_{\mathrm{e}} \neq 0$
$2, 4, \dots, 20$	6	7	8	exist (over $\mathbb{C}$)
$2, 4, \dots, 20$	6	7	9	none
$2, 4, \dots, 20$	6	9	9	exist (over $\mathbb{C}$)
$2, 4, \dots, 20$	6	9	10	none
$2, 4, 6$	6	11	10	exist (over $\mathbb{C}$)
$2, 4, 6$	6	11	11	none
2	8	7	11	none

TABLE 1. Results of the exact Gröbner basis computations.

Proof of Theorem E. For (i), use the rows of Table 1 marked "none" with $\deg f_{\mathrm{e}} \le 6$. For (ii), use the last row: a polynomial of degree ≤ 8 has $\deg f_{\mathrm{e}} \le 8$ and $\deg f_{\mathrm{o}} \le 7$. In both cases a center forces $c_1 = \cdots = c_K = 0$, hence $f_{\mathrm{e}} = 0$, and conversely odd f gives a center. □

Remark 7.1. (1) The computations are exact and decide the absence of *complex* solutions; the conclusion therefore holds a fortiori over $\mathbb{R}$. (2) The rows with "exist" show that, in these degrees, the first $K - 1$ conditions do not suffice. In particular, the sixth to eighth center conditions alone do not exclude the exceptional set of Theorem D. (3) The computations for $\deg f_{\mathrm{e}} \le 6$ take seconds (minutes for $\deg f_{\mathrm{o}} \le 11$); the case $\deg f_{\mathrm{e}} \le 8$ took about one hour of a single core with `std`. A modular computation (`modStd`) suggests that already $K = 10$ suffices in that case, but we only claim what was obtained with `std` over $\mathbb{Q}$.

8. Further remarks and open problems

Problem 8.1. Prove Conjecture 1.1 for $\deg f_{\mathrm{e}} \le 6$ and f_{o} of arbitrary degree. By Theorem D, it suffices to show that the higher center conditions exclude the exceptional set (1.5). In all cases of Table 1 with $\deg f_{\mathrm{e}} \le 6$ and $\deg f_{\mathrm{o}} = 2R+1$, exactly $K = R+6$ conditions are needed ($K = 9, 10, 11$ for $R = 3, 4, 5$). Is this true for all R?

Problem 8.2. Find a closed formula for $u_{k+1}(1)$, modulo the previous conditions, of the form $m_k + \mathcal{Q}_k$, where $\mathcal{Q}_k$ is a combination of iterated integrals of order ≥ 2 in f. Theorem A gives $\mathcal{Q}_3, \mathcal{Q}_4, \mathcal{Q}_5$; Brudnyi's formalism [11] suggests that a general expression exists. Together with Lemma 4.1, it would extend Theorem C to arbitrary numbers of sign changes, under additional orthogonality hypotheses.

Problem 8.3. Extend Theorems C–E to trigonometric Abel equations with a reflection symmetry $\theta \mapsto -\theta$, where Example 3.2 shows that the nonlinear correction in (1.2) is essential.

Data availability

The `SymPy` and `Singular` code used for the identities of Theorem A, the determinant formula of Proposition 5.4, Example 6.2 and Table 1 is available at `https://github.com/Araujo2021/araujo2021.github.io/tree/main/pesquisa/abel-lijun-yun/codigo`.

Acknowledgements

References

[1] A. Álvarez, J. L. Bravo, C. Christopher, P. Mardešić, *Infinitesimal center problem on zero cycles and the composition conjecture*, Funct. Anal. Appl. **55** (2021), no. 4, 257–271.

[2] A. M. Alves, A. Lemos, A. L. A. de Araujo, *A partial positive answer to a Lijun–Yun conjecture*, Boll. Unione Mat. Ital. **13** (2020), 49–59.

[3] M. A. M. Alwash, *The composition conjecture for Abel equation*, Expo. Math. **27** (2009), 241–250.

[4] M. A. M. Alwash, N. G. Lloyd, *Non-autonomous equations related to polynomial two-dimensional systems*, Proc. Roy. Soc. Edinburgh Sect. A **105** (1987), 129–152.

[5] P. Borwein, T. Erdélyi, *Polynomials and Polynomial Inequalities*, Graduate Texts in Mathematics **161**, Springer, New York, 1995.

[6] M. Briskin, J.-P. Françoise, Y. Yomdin, *Center conditions, compositions of polynomials and moments on algebraic curves*, Ergodic Theory Dynam. Systems **19** (1999), 1201–1220.

[7] M. Briskin, J.-P. Françoise, Y. Yomdin, *Center conditions II: parametric and model center problems*, Israel J. Math. **118** (2000), 61–82.

[8] M. Briskin, J.-P. Françoise, Y. Yomdin, *Center conditions III: parametric and model center problems*, Israel J. Math. **118** (2000), 83–108.

[9] M. Briskin, F. Pakovich, Y. Yomdin, *Algebraic geometry of the center-focus problem for Abel differential equations*, Ergodic Theory Dynam. Systems **36** (2016), 714–744.

[10] M. Briskin, N. Roytvarf, Y. Yomdin, *Center conditions at infinity for Abel differential equations*, Ann. of Math. (2) **172** (2010), 437–483.

[11] A. Brudnyi, *On the center problem for ordinary differential equations*, Amer. J. Math. **128** (2006), 419–451.

[12] L. A. Cherkas, *Number of limit cycles of an autonomous second-order system*, Differ. Equ. **5** (1976), 666–668.

[13] C. Christopher, *Abel equations: composition conjectures and the model problem*, Bull. London Math. Soc. **32** (2000), 332–338.

[14] A. Cima, A. Gasull, F. Mañosas, *An explicit bound of the number of vanishing double moments forcing composition*, J. Differential Equations **255** (2013), 339–350.

[15] A. Cima, A. Gasull, F. Mañosas, *A simple solution of some composition conjectures for Abel equations*, J. Math. Anal. Appl. **398** (2013), 477–486.

[16] W. Decker, G.-M. Greuel, G. Pfister, H. Schönemann, Singular 4-3-2 — A computer algebra system for polynomial computations, `https://www.singular.uni-kl.de`, 2023.

[17] L. Gavrilov, *On the center-focus problem for the equation* $\frac{dy}{dx} + \sum_{i=1}^{n} a_i(x) y^i = 0$, $0 \le x \le 1$, *where* a_i *are polynomials*, Ann. Henri Lebesgue **3** (2020), 615–648.

[18] J. Giné, M. Grau, J. Llibre, *Universal centres and composition conditions*, Proc. London Math. Soc. (3) **106** (2013), 481–507.
[19] J. Giné, M. Grau, X. Santallusia, *Universal centers in the cubic trigonometric Abel equation*, Electron. J. Qual. Theory Differ. Equ. **2014**, No. 1, 1–7.
[20] J. Giné, M. Grau, X. Santallusia, *A counterexample to the composition condition conjecture for polynomial Abel differential equations*, Ergodic Theory Dynam. Systems **39** (2019), 3347–3352.
[21] S. Karlin, *Total Positivity*, Vol. I, Stanford University Press, Stanford, 1968.
[22] Y. Lijun, T. Yun, *Some new results on Abel equations*, J. Math. Anal. Appl. **261** (2001), 100–112.
[23] A. Lins Neto, *On the number of solutions of the equation* $dx/dt = \sum_{j=0}^{n} a_j(t)x^j$, $0 \leq t \leq 1$, *for which* $x(0) = x(1)$, Invent. Math. **59** (1980), 67–76.
[24] F. Pakovich, *A counterexample to the "composition conjecture"*, Proc. Amer. Math. Soc. **130** (2002), 3747–3749.
[25] F. Pakovich, M. Muzychuk, *Solution of the polynomial moment problem*, Proc. London Math. Soc. (3) **99** (2009), 633–657.
[26] F. Pakovich, *Solution of the parametric center problem for the Abel differential equation*, J. Eur. Math. Soc. **19** (2017), 2343–2369.
[27] G. Pólya, G. Szegő, *Problems and Theorems in Analysis*, Vol. II, Springer, Berlin, 1976.

Departamento de Matemática, CCE, Universidade Federal de Viçosa, Avenida PH Rolfs, s/n, Viçosa, MG, CEP 36570-900, Brazil
Email address: amalves@ufv.br

Departamento de Matemática, CCE, Universidade Federal de Viçosa, Avenida PH Rolfs, s/n, Viçosa, MG, CEP 36570-900, Brazil
Email address: abiliolemos@ufv.br

Departamento de Matemática, CCE, Universidade Federal de Viçosa, Avenida PH Rolfs, s/n, Viçosa, MG, CEP 36570-900, Brazil
Email address: anderson.araujo@ufv.br