


## Tutorial 8 - Mathematics for CS

1. Consider a random variable measuring the following quantities. In each case, state with reasons whether you think it is more appropriate to define the random variable as discrete or as continuous.
  - (a) A person's height: Discrete. It is possible to measuring a person's height continuously, however, a person's height continuous changes, and hence measurements with units smaller than year quickly becomes meaningless.
  - (b) A student's course grade: Discrete. It is impossible grade a student's ability with precision.
  - (c) The thickness of a metal plate: Continuous. Measuring with continuity gives a better picture for how thickness changes overtime.
  - (d) The purity of a chemical solution: Continuous. Measuring with continuity gives a better picture for how the purity changes overtime.
  - (e) The type of personal computer a person owns: Discrete. There is a finite numbers of computer types.
  - (f) A person's age: Discrete. It is possible to measuring a person's age continuously, however, one's age continuously changes, and hence measurements with units smaller than year quickly becomes meaningless.
2. A random variable  $X$  takes values between 4 and 6 with a probability density function

$$f(x) = \frac{1}{x \ln(1.50)}$$

for  $4 \leq x \leq 6$  and  $f(x) = 0$  elsewhere.

- (a) Make a sketch of the probability density function.



- (b) Check that the total area under the probability density function is equal to 1.

$$\int_{-\infty}^{\infty} f(x) dx = \int_4^6 \frac{1}{x \ln(1.5)} dx = \left. \frac{\ln(x)}{\ln(1.5)} \right|_{x=4}^{x=6} = 1$$



(c) What is  $P(4.5 \leq X \leq 5.5)$ ?


$$\int_{4.5}^{5.5} \frac{1}{x \ln(1.5)} dx = \left. \frac{\ln(x)}{\ln(1.5)} \right|_{x=4.5}^{x=5.5} = \ln(5.5/4.5) / \ln(1.5) \approx 0.4949148310$$

(d) Construct and sketch the cumulative distribution function.

For  $x < 4$ ,  $F(x) = 0$ . For  $x > 6$ ,  $F(x) = 1$ .

For  $4 \leq x \leq 6$ ,

$$F(x) = \int_4^x \frac{1}{t \ln(1.5)} dt = \left. \frac{\ln(t)}{\ln(1.5)} \right|_{t=4}^{t=x} = \frac{\ln(x) - \ln(4)}{\ln(1.5)}$$



(e) What is the expected value of this random variable?

$$E(X) = \int_4^6 \frac{x}{x \ln(1.5)} dx = \left. \frac{x}{\ln(1.5)} \right|_{x=4}^{x=6} = \frac{2}{\ln(1.5)} \approx 4.93260692475$$

(f) What is the median of this random variable?

$$F(x) = 0.5 \Leftrightarrow \frac{\ln(x) - \ln(4)}{\ln(1.5)} = 0.5 \Leftrightarrow \ln(x) = 0.5 \ln(1.5) + \ln(4) \Leftrightarrow x = 4\sqrt{1.5} = 2\sqrt{6}$$

(g) What is the variance and the standard deviation of this random variable?

$$\begin{aligned} \text{Var}(X) &= \int_4^6 \frac{x^2}{x \ln(1.5)} dx - E(X)^2 = \int_4^6 \frac{x}{\ln(1.5)} dx - E(X)^2 \\ &= \left. \frac{x^2}{2 \ln(1.5)} \right|_{x=4}^{x=6} - E(X)^2 = \frac{10}{\ln(1.5)} - E(X)^2 \approx 0.332423549644 \\ \sigma &= \sqrt{\text{Var}(X)} \approx 0.576561835057 \end{aligned}$$

3. A random variable  $X$  takes values between 0 and  $\infty$  with a cumulative distribution function

$$F(x) = A + Be^{-x}$$

for  $0 \leq x < \infty$ .



- (a) Find the values of  $A$  and  $B$  and sketch the cumulative distribution function.


Since  $1 = \lim_{x \rightarrow \infty} F(x), 1 = \lim_{x \rightarrow \infty} A + Be^{-x} = A$ .

Hence  $A = 1$ .

Since  $X$  takes values between 0 and  $\infty$ ,  $F(0) = 0$  ( $X$  does not takes 0).

Hence  $F(0) = A + Be^{-x} = A + B = 0 \Rightarrow B = -1$ .

Therefore,  $F(x) = 1 - e^{-x}$




- (b) What is  $P(2 \leq X \leq 3)$ ?

$$P(2 \leq X \leq 3) = F(3) - F(2) = e^{-2} - e^{-3} \approx 0.0855482149$$

- (c) Construct and sketch the probability density function.

For  $0 < x < \infty$ ,

$$f(x) = \frac{d}{dx} (1 - e^{-x}) = e^{-x}$$



Done by Huynh Le An Khanh on November 6, 2025.

