Urban flood prediction and climate resilience

"I'm working on urban flood prediction. I would like to know the suitable application of rainfall nowcasting and drainage network models in climate resilience for Indian cities. COULD YOU PRIOVIDE THE EXTRACT DIRECTION for me on how to do i come up with better service or solution addressing real world problem" *Generated by LeapSpace*

Urban Flood Prediction for Indian Cities at a Glance

Key Findings Table

Topic	What the evidence supports	Most relevant application in Indian cities	Key citations
Rainfall nowcasting	Short-lead rainfall nowcasting is most useful at 0-3 h, with strongest skill typically in the first 0-2 h; performance drops quickly for localized convective storms [1] [2]	Event-day warning, pump/drain operations, hotspot alerting, traffic and emergency response [3] [4]	[1] [2] [3]
Drainage-network models	SWMM, PCSWMM, MIKE, HEC-RAS, and coupled 1D-2D models are suitable for representing flooded nodes, surcharge, overland flow, and inundation extent [5] [6] [7] [8] [9] [10]	Hotspot diagnosis, ward-scale inundation forecasting, drainage redesign, resilience planning [10] [11] [12]	[5] [6] [7] [8] [9] [10]
Best service architecture	The strongest overall pattern is a coupled system: forecast rainfall forces a drainage/hydraulic model to generate location-specific flood outputs [13] [14] [15] [16]	Real-time hotspot forecasting and climate-resilience decision support [17] [18]	[13] [14] [15] [16]
Main cause of repeated flood hotspots	Flooding is often driven as much by drainage failure as by rainfall intensity: undersized conduits, blockages, encroachment, sedimentation, and low-lying terrain repeatedly explain recurrence [10] [19] [20] [21] [22]	Combine weather forecasting with drainage-asset intelligence and maintenance targeting [10] [23]	[10] [19] [20] [21] [22] [23]

Topic	What the evidence supports	Most relevant application in Indian cities	Key citations
Data-scarce deployment	Remote sensing, IMERG, Sentinel SAR, DEM reconstruction, and document-derived sewer inventories are common workarounds where gauges and sewer maps are incomplete [24] [25] [26]	Start with a hybrid minimum viable service, then add local sensors and drainage audits [3] [27]	[3] [24] [25] [26]
Resilience interventions	SuDS, retention ponds, drainage upgrades, lake/wetland protection, and hybrid mitigation can materially reduce runoff, flood volume, and flooded area [28] [29] [30] [31]	Use the same model stack for both warning and adaptation testing [29] [32] [33]	[28] [29] [30] [31] [32] [33]

Direct Answer

The most suitable application split is:

- **Use rainfall nowcasting for 0-3 hour operational warning**, especially 0-2 hour response windows, because this is where radar-based and blended systems show the most reliable short-lead skill [1] [2].
- **Use drainage-network and hydraulic models for flood-process translation**, meaning hotspot diagnosis, flooded-node prediction, surcharge analysis, inundation depth/extent, and infrastructure planning [10] [11] [12] [34].
- **For real-world climate resilience, do not deploy them separately**: the best service concept is a **coupled nowcast-to-drainage system** that converts forecast rainfall into ward- or hotspot-specific flood alerts and planning outputs [13] [14] [15] [16].

A practical direction for building a better service is:

1. **Pick one city and 5-10 recurring hotspots first**, rather than trying to solve the whole city at once, because recurrent flooding is strongly linked to identifiable drainage bottlenecks and low-lying terrain [10] [21] [22].
2. **Build a minimum viable drainage model** in SWMM or PCSWMM using available maps, DEM, land use, and reconstructed network data if full sewer data are missing [6] [10] [24] [25].
3. **Feed that model with short-lead rainfall nowcasts** from radar, gauge-radar blends, or satellite-supported inputs to generate hotspot forecasts every 15-30 minutes [1] [3] [13].

4. **Issue threshold-based municipal actions**, such as pump deployment, drain cleaning dispatch, traffic diversion, and local warnings, based on flooded-node counts, expected depth, or exceedance probability [23] [35] [36].

5. **Reuse the same platform for resilience planning**, testing SuDS, retention ponds, conduit upgrades, and lake/wetland restoration under present and future extreme rainfall scenarios [28] [29] [30] [32] [33].

Taken together, these findings suggest that a strong urban-flood service for Indian cities should be framed less as a "forecasting product" and more as a **city flood operations and resilience platform** combining rainfall intelligence, drainage diagnostics, and mitigation planning [1] [3] [19] [36].

Confidence level: High - The application split and service direction are strongly supported across multiple themes in the evidence, even though end-to-end Indian operational validations remain limited [1] [3] [10] [16].

Study Scope

- **Time window represented**: mainly studies from roughly 2015-2025, with some foundational method papers included where they underpin current practice [37] [38].
- **Disciplines covered**: radar nowcasting, deep learning rainfall prediction, convection-permitting NWP, urban drainage hydraulics, hydrodynamic inundation modeling, GIS/remote sensing hotspot mapping, and operational flood-warning design [1] [5] [13] [39].
- **Geographic emphasis**: Indian cities and Indian monsoon/coastal settings, with some non-Indian methodological evidence used only where the retrieved set framed general system architectures or probabilistic methods [10] [40] [41].

Assumptions & Limitations

- The evidence strongly supports the **architecture** of coupling nowcasts with drainage models, but provides **few fully documented Indian city systems** reporting lead time, trigger thresholds, false alarms, and missed-event rates together [3] [16] [42].
- India-specific validation is stronger for **radar-based extrapolation and physics-based high-resolution forecasting** than for deep-learning rainfall nowcasting at 15 min-3 h leads [1] [43] [44] [45].
- Many drainage studies rely on **reconstructed or incomplete sewer data**, so model realism depends heavily on local verification and updating [24] [25].

Suggested Further Research

- Benchmark **radar nowcasting vs deep learning vs convection-permitting NWP** for Indian cities at 0-3 h lead time using the same events and metrics [1] [43] [46].
- Build Indian demonstrations that report **end-to-end service value**: hotspot hit rate, false alarms, missed events, actionable lead time, and avoided impacts [3] [42].

- Compare **SWMM, PCSWMM, MIKE+, HEC-RAS, and reduced-complexity 1D-2D** under Indian data constraints [10] [12].
- Develop practical **probabilistic warning thresholds** for municipalities using exceedance probabilities and percentile inundation outputs [35] [36] [47].
- Link **hotspot maps, blockage detection, and maintenance records** to dynamic forecasting so services support both emergency response and asset management [21] [23] [48].

Urban Flood Prediction and Climate-Resilient Service Design for Indian Cities

Foundational Context

Urban flood prediction in Indian cities requires combining two different but complementary modeling logics.

First, **rainfall nowcasting** estimates where and how much rain will fall in the next minutes to hours. Its core value is temporal: it provides short warning lead time during fast-onset urban flooding, especially in monsoon and convective conditions [1] [2]. Second, **drainage-network or hydraulic modeling** translates rainfall into physically meaningful flood consequences such as pipe surcharge, manhole overflow, street ponding, and inundation depth [5] [6] [10]. Without this second step, rainfall intensity alone cannot identify which road, ward, junction, or neighborhood will actually flood [10] [21].

This distinction matters because the evidence repeatedly shows that Indian urban flooding is produced by the interaction of **extreme rainfall, rapid urbanization, impervious surfaces, low-lying terrain, and stressed drainage systems**, rather than weather alone [7] [23] [49] [50] [51]. That is why a climate-resilient flood service must solve three problems at once:

- detect short-term rainfall danger [1];
- convert rainfall into local flood impact [10];
- identify structural weaknesses for long-term adaptation [29] [52].

Confidence level: High - This framing is directly supported across weather, hydraulic, and governance-oriented evidence streams [5] [13] [49] [52].

Why Urban Flood Prediction Matters for Indian Cities

The reviewed evidence indicates that Indian cities face rising flood pressure from both climate and land-surface change. Multiple studies project increasing extreme precipitation or future flood hazard under changing rainfall variability and monsoon behavior [9] [32] [49] [50] [53] [54]. At the same time, urbanization increases imperviousness and alters drainage pathways, which can raise flood duration, flooded nodes, and hazard area even where rainfall change alone is not sufficient to explain the impact [7] [9] [55] [56].

City case evidence reinforces this. Hotspot mapping studies found repeated flood concentrations in Mumbai, Kolkata, Hyderabad, Guwahati, and Bengaluru-linked settings, with validation against historical flood points, event footprints, or waterlogged locations [39] [57] [58] [59] [60]. In Mumbai, susceptibility models showed strong alignment with observed hotspots such as Kurla, Chembur, and Sion, and one study found 85.46% of 234 waterlogged locations fell in severe-to-very-high vulnerability zones [57] [61]. In Kolkata, GIS/AHP mapping reached AUC 0.89 against historical flood locations [39]. In Hyderabad, low-lying vulnerable areas matched observed flooding during the September 2016 rains [58].

Taken together, these findings suggest that urban flood prediction in India is best treated as a **localized infrastructure and service-delivery problem under climate stress**, not only as a meteorological forecasting problem [10] [62].

Rainfall Nowcasting for Urban Flood Prediction

Best-performing short-lead approaches

The strongest nowcasting evidence in the set favors **radar-based extrapolation and optical-flow frameworks**, especially for 0-2 hour lead times. A pySTEPS-based system reported strong 30-minute performance with FSS 0.87, HSS 0.84, POD 0.90, CSI 0.81, and FAR 0.19, and maintained high FSS across reflectivity bins for convective storms over India [1]. Another radar ensemble nowcasting study showed that skill remained useful for longer-lived widespread systems but degraded sharply for convective rainfall by 2 h lead time [2]. Broader evidence also shows radar nowcasting reaching FSS up to 0.78 at 2 h for tropical storms, with weaker performance for convective and stratiform cases [2].

Verification practice in these studies relies mainly on:

- **event detection metrics**: POD, FAR, CSI, HSS, ETS, FSS [1] [2] [63] [64];
- **extreme detection metrics**: EDS, EDI, SEDI [64];
- **intensity/error metrics**: correlation, RMSE, MAE, bias [65] [66].

The operational implication is straightforward: for Indian urban flood services, nowcasting is most valuable when aligned to immediate actions such as pumping, traffic control, field crew mobilization, and local alerts, not as a substitute for longer-horizon planning [1] [2] [3].

Deep learning: promise, but thinner India-specific validation

The evidence includes several promising deep-learning nowcasting results, such as ConvLSTM, U-Net, and 3D Conv-LSTM architectures with strong intensity/structure metrics, including Pearson correlation 0.96 for ConvLSTM and RMSE 0.8 mm/hr with PCC 0.78 for U-Net-like models [66] [67] [68]. Transfer learning also improved rare heavy-rain downscaling in Mumbai, outperforming raw GFS and the operational forecast model at multiple lead times [69].

However, the retrieved evidence explicitly notes that it does **not** contain Indian-region or monsoon nowcasting papers with exact 15-minute to 3-hour benchmark metrics for ConvLSTM, 3D Conv-LSTM, or U-Net comparable to the radar-based studies [43] [44] [45] [70]. So while AI methods are promising, local operational trust currently rests more strongly on radar extrapolation and event-specific high-resolution physics setups [1] [46].

Why high-resolution NWP still matters

For Indian monsoon extremes, convection-permitting NWP remains important because forecast usefulness depends strongly on physical scheme selection. For the 2015 Chennai event, 1 km WRF with explicit convection and MYNN PBL better captured rainfall intensity, distribution, moisture convergence, and circulation than YSU or MYJ [40]. Cloud microphysics also mattered strongly: Thompson performed best, followed by Morrison, in reproducing rainfall location, timing, and hydrometeor evolution [46]. Another Chennai study found Goddard microphysics + YSU PBL + Noah LSM best captured rainfall magnitude and reflectivity [71]. Over west-coast atmospheric-river events, Lin or Thompson microphysics combined with KF or Grell3D cumulus schemes performed best [72].

This means NWP can support flood services, but only after **city- and event-specific benchmarking** against gauges, radar, and satellite products [73] [74].

Confidence level: High - Radar-nowcasting utility and physics sensitivity are well supported; confidence is lower for Indian deep-learning operational superiority because validation is thinner [1] [43] [46].

Urban Drainage and Sewer Network Modeling

Core model choices

Urban drainage studies commonly use **SWMM, MIKE URBAN/MIKE+, PCSWMM, and HEC-RAS** to simulate drainage capacity, flooded nodes, and inundation extent [5] [6] [7] [8] [9]. SWMM-based Indian studies often use subcatchment-to-pipe structures with dynamic-wave routing to identify overflowing nodes and assess drainage adequacy in cities such as Vijayawada, Surat, and Delhi [7] [8] [34] [75]. For inundation mapping, coupled setups are more informative: SWMM + PCSWMM was used for Hyderabad flood-risk mapping, while MIKE 11 + MIKE 21 produced depth-based inundation maps for Mumbai [10] [11].

Why 1D-2D and rain-on-grid matter

Research consistently indicates that pure 1D sewer routing is often insufficient for realistic urban flooding because it misses surcharge-to-surface exchange, overland routing, and ponding behavior [76] [77] [78]. Coupled 1D-2D frameworks connect sewer networks to surface flow via manholes, inlets, or exchange nodes, allowing water to move in both directions during surcharge [76] [79] [80]. These approaches better represent flood depths, velocities, inundation patterns, backflow, and in coastal settings, downstream tidal effects on drainage capacity [81] [82].

Indian examples also show the value of coupled frameworks. A HEC-RAS 1D-2D study for Kolhapur reported fairly accurate flood depth and inundation extent [12]. The Mumbai Mithi catchment study used MIKE FLOOD hazard mapping that combined flood depth and velocity [11]. A lower Tapi 1D/2D model and a Surat floodplain application both reported more than 80% submergence in severe event scenarios, illustrating the hazard-mapping value of coupled setups [83] [84].

Calibration and data reconstruction

Automatic calibration is attractive because manual calibration is slow and can miss optimal parameter combinations [85] [86]. In Indian SWMM-focused evidence, GA outperformed other tested evolutionary algorithms in one setup, and Monte Carlo auto-calibration also produced good performance in GIS-based flood modeling [75] [87]. Yet the retrieved Indian-city content does not provide many direct recent examples of automatic calibration pipelines tied to flooded-node prediction [88] [89].

In data-scarce cities, model realism is often improved through reconstruction: DEM-derived elevations, land-use-based subcatchments, document-derived conduit inventories, and road/building connectivity corrections [24] [25]. Studies report that better subcatchment delineation, corrected node/surface elevations, and more faithful conduit/network representation produce the largest realism gains [24] [25] [90].

Drainage failure mechanisms

The strongest and most actionable drainage finding is that recurrent flood hotspots are usually tied to **asset and terrain failures**: inadequate capacity, blocked bottlenecks, structural non-resilience, sedimentation, encroachment, and low-lying topography [10] [19] [20] [21] [22]. This is crucial for product design: a city flood service that predicts rain but ignores drains will miss the actual causes of repeat inundation [23].

Confidence level: High - Model classes, coupled-structure benefits, and drainage-failure mechanisms are consistently supported, though head-to-head tool comparisons are limited [10] [76].

Coupled Nowcasting-Drainage Forecasting Systems

The evidence supports a clear technical pattern: **forecast rainfall is used as forcing for runoff, drainage, or inundation models** [13] [14] [15]. Example coupling modes include:

- **Direct forcing**: nowcasted precipitation time series fed directly into HEC-RAS or similar hydraulic models [13].
- **Sequential cascade**: ensemble rainfall drives a hydrologic model first, then runoff feeds a hydraulic model for flood-hazard mapping [91].
- **Grid-based forcing to urban drainage**: high-resolution forecast rainfall used as input to PCSWMM or related urban flood frameworks [15].

The practical benefits reported across the broader literature are better real-time forecasting, useful advance warning, inundation estimation, and hazard probability mapping [4] [91] [92] [93]. Probabilistic methods extend this by turning rainfall uncertainty into exceedance maps, percentile inundation extents, and probability-threshold warnings [35] [36] [41] [47] [94].

However, there is an important evidence gap: the retrieved set does **not** provide many Indian city studies that fully document operational nowcast-driven pipelines with reported lead time, thresholds, inundation-accuracy gains, false alarms, and missed events [3] [16] [17] [42]. This is not a reason to avoid the architecture; it means implementation should proceed with phased validation.

Confidence level: High for architecture, Medium for Indian operational maturity - The coupling pattern is well supported, but deployment evidence in Indian cities is under-documented [13] [15] [16].

Indian City Case Studies and Deployment Constraints

Hotspot mapping and validation

Indian city studies show that hotspot mapping can already provide municipal value before full real-time hydraulic maturity. Mumbai studies found strong overlap between mapped vulnerability and observed waterlogging, though performance varied by event and method [57] [61] [95]. Kolkata susceptibility mapping validated strongly against historical flood locations [39] [96]. Guwahati GIS+AHP mapping achieved AUC 0.909 on prediction data and identified critical wards [59]. These are useful foundations for where to deploy sensors, calibrate drainage models, and pilot warnings [52].

Real-time sensing and warning deployments

Some near-operational systems are emerging. Bengaluru's integrated setup combines automatic rain gauges, weather data, lake-network inputs, and SWMM/HEC-RAS modeling, with validated forecasts reported at 12-72 h lead time and dissemination through the Megha Sandhesa app and Varunamitra portal [3]. Guwahati's edge-integrated system combined sensing nodes, rainfall prediction, and blockage detection, reporting 92.4% rainfall-prediction performance and 91% blockage-segmentation accuracy [48]. Nationally, Google's flood warning system covered 376 target gauges in 2021, with inundation models for 233 gauges [97].

Yet across these examples, explicit false-alarm and missed-event rates are generally not reported in the retrieved summaries [3] [48] [97].

Data scarcity and workarounds

A consistent deployment constraint is missing high-resolution rainfall data, incomplete drainage maps, and rapidly changing urban surfaces [24] [98] [99] [100] [101]. Common workarounds include IMERG rainfall, ALOS PALSAR DEM, Sentinel-2 land cover, Sentinel-1 SAR flood mapping, statistical rainfall disaggregation, and sewer reconstruction from documents and GIS [24] [25] [26] [102] [103].

This supports an incremental strategy: start with remotely sensed and reconstructed inputs, then improve with local gauges, water-level sensors, and drainage audits [3] [24].

Practical Direction: How to Build a Better Real-World Service

Recommended product architecture

A strong solution for Indian cities is a **three-layer service**:

1. **Preparedness layer**: hotspot susceptibility map and drainage-risk atlas for seasonal planning [39] [59].
2. **Operations layer**: 0-2 h rainfall nowcasting linked to hotspot-level drainage/inundation forecasting [1] [13].
3. **Adaptation layer**: scenario testing for SuDS, retention ponds, upgrades, wetland/lake restoration, and future rainfall stress [28] [29] [32].

Minimum viable pipeline

- **Step 1: Select a city and narrow geography** to top recurring waterlogging zones [10] [57].
- **Step 2: Build a base flood inventory** from past complaints, ward records, SAR flood footprints, and road-closure history [57] [104].
- **Step 3: Construct a simplified drainage model** in SWMM/PCSWMM using available sewer maps; if incomplete, reconstruct with DEM, road network, and planning documents [10] [25].
- **Step 4: Add rainfall nowcasting input** using radar-first where available, with gauge or satellite supplementation [1] [26].
- **Step 5: Validate on known flood events** against hotspot hits, flooded nodes, observed water depths, or inundation footprints [10] [58].
- **Step 6: Create action thresholds** for municipal response, such as “expected surcharge at node cluster,” “depth > threshold,” or “probability of exceedance > threshold” [35] [36].
- **Step 7: Add sensing where it improves decision value most**, especially blockage-prone nodes and low-lying hotspots [23] [48].

What to optimize for

A differentiated service should optimize for **actionable hotspot lead time**, not only rainfall RMSE [4] [66]. Useful KPIs include:

- hotspot hit rate [95];
- flooded-node prediction accuracy [75];
- inundation depth/extent error [12];

- warning lead time [3];
- false alarms and missed events, where possible [42].

Resilience planning use

The same platform can support adaptation testing. Evidence shows measurable benefits from interventions: in Nagpur, rooftop disconnection reduced runoff by about 57%, infiltration trench by about 15%, and the combined measure by nearly 67% [28]. In Gurgaon/Gurugram, proposed retention ponds/SuDS reduced flood volume from 240 to 180 CMS for yearly rainfall and from 500 to 350 CMS for an extreme 7-hour rainfall scenario [29]. A hybrid mitigation strategy reported reductions up to 37.3% in peak flow, 71.5% in flood volume, and 93.6% in flooded area [30].

Confidence level: Medium-High - The service design logic is strongly supported, but some operational KPI recommendations are inferential because Indian end-to-end reporting remains incomplete [3] [42].

Conclusion

The evidence indicates that the most suitable application for Indian cities is a **coupled urban flood service** in which **short-lead rainfall nowcasting provides immediate warning input** and **drainage-network/hydraulic models translate that input into hotspot-specific flood consequences** [1] [10] [13]. Rainfall nowcasting alone is insufficient because recurring urban flooding is repeatedly driven by drainage failures, encroachment, sedimentation, and low-lying terrain [10] [21] [23]. Drainage models alone are insufficient for operations because they need real-time rainfall forcing to produce actionable short-lead forecasts [13] [14].

For a better real-world solution, the most evidence-based path is:

- start small and hotspot-focused [22];
- use hybrid data sources in data-scarce environments [24] [26];
- couple radar/satellite-informed rainfall nowcasts to a practical drainage model [1] [10];
- deploy threshold-based actions for municipal response [36];
- expand the same platform into resilience planning for SuDS and drainage upgrades [29] [30].

Overall confidence level: High - The core recommendation to couple short-lead rainfall nowcasting with drainage/hydraulic modeling for Indian urban resilience is strongly grounded across the retrieved evidence, though documentation of fully operational Indian end-to-end systems remains limited [1] [3] [10] [16].

References

1. Mishra, A.K., Chhari, A., Kumar, G., (...), Shukla, K.K. (2025) Next-gen storm nowcasting: harnessing satellite-based radar reflectivity over India. *Theoretical and Applied Climatology*. https://doi.org/10.1007/s00704-025-05650-5
2. Bibraj, B., Sahoo, S., Puviarasan, N., Chandrasekar, V. (2024) Operational Assessment of High Resolution Weather Radar Based Precipitation Nowcasting System. *Atmosphere*. https://doi.org/10.3390/atmos15020154
3. https://www.sciencedirect.com/science/article/pii/S2363504522000765
4. Liu, J., Cho, H.-S., Osman, S., (...), Lee, K. (2022) Review of the status of urban flood monitoring and forecasting in TC region. *Tropical Cyclone Research and Review*. https://doi.org/10.1016/j.tcrr.2022.07.001
5. Manoj, N.K., Robert, S., Menon, R., George, S. (2026) Preliminary Assessment of Stormwater Flooding in an Urban Area Using MIKE URBAN+: A Case Study of Kalamassery Municipality, Kerala, India. *Lecture Notes in Civil Engineering*. https://doi.org/10.1007/978-981-95-1726-8_24
6. Kumar, S., Kaushal, D.R., Gosain, A.K. (2017) Urbanization Challenges in Emerging Economies. *Urbanization Challenges in Emerging Economies: Energy and Water Infrastructure; Transportation Infrastructure; and Planning and Financing - Selected Papers from the ASCE India Conference 2017*. https://www.scopus.com/pages/publications/85091336447
7. Agarwal, S., Kumar, S. (2023) Assessment of Urban Flood in Vijayawada under Historical and Future Projected Climate Scenarios. *Indian Journal of Environmental Protection*. https://www.scopus.com/pages/publications/85160032087
8. Baudhanwala, D., Kantharia, V., Patel, D., (...), Waikhom, S. (2023) APPLICABILITY OF SWMM FOR URBAN FLOOD FORECASTING A CASE STUDY OF THE WESTERN ZONE OF SURAT CITY. *Larhyss Journal*. https://www.scopus.com/pages/publications/85163025165
9. Kumar, S., Agarwal, A., Ganapathy, A., (...), Sivakumar, B. (2022) Impact of climate change on stormwater drainage in urban areas. *Stochastic Environmental Research and Risk Assessment*. https://doi.org/10.1007/s00477-021-02105-x
10. Rangari, V.A., Gonugunta, R., Umamahesh, N.V., (...), Bhatt, C.M. (2018) 1D-2D modeling of urban floods and risk map generation for the part of Hyderabad city. *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences - ISPRS Archives*. https://www.scopus.com/pages/publications/85057584792
11. Ghosh, M., Karmakar, S., Ghosh, S. (2022) Flood Modelling for an Urban Indian Catchment: Challenges and Way Forward. *Lecture Notes in Civil Engineering*. https://doi.org/10.1007/978-981-16-5501-2_5
12. Kulkarni, A.D., Kale, G.D. (2024) The Development of Coupled 1D–2D Hydrodynamic Flood Model by Using HEC-RAS: a Case Study of the Panchganga River Basin, Kolhapur District, Maharashtra, India. *Water Resources*. https://doi.org/10.1134/S0097807823601322

13. Aaritha, V.V., Kuiry, S.N., Chandrasekar, V. (2026) ConvLSTM-Based Precipitation Nowcasting for Urban Flood Forecasting. *World Environmental and Water Resources Congress 2026: Krewe of Innovation - Proceedings of World Environmental and Water Resources Congress 2026*. https://doi.org/10.1061/9780784486931.022

14. Ashok, S.P., Pekkat, S. (2026) Application of Deep Learning and Rain-On-Grid Hydrodynamic Modeling for Catchment-Scale Flood Forecasting in Guwahati, India. *Journal of Flood Risk Management*. https://doi.org/10.1111/jfr3.70186

15. Davis, S., Pentakota, L., Saptarishy, N., Mujumdar, P.P. (2022) A Flood Forecasting Framework Coupling a High Resolution WRF Ensemble With an Urban Hydrologic Model. *Frontiers in Earth Science*. https://doi.org/10.3389/feart.2022.883842

16. Kannapiran, U.M., Bhaskar, A.S. (2024) “Flood inundation mapping of upstream region in the Adyar River basin: Integrating hydrologic engineering centre's river analysis system (HEC-RAS) approach with groundwater considerations”. *Groundwater for Sustainable Development*. https://doi.org/10.1016/j.gsd.2024.101085

17. Nithila Devi, N., Sridharan, B., Kuiry, S.N. (2019) Impact of urban sprawl on future flooding in Chennai city, India. *Journal of Hydrology*. https://doi.org/10.1016/j.jhydrol.2019.04.041

18. Zope, P.E., Eldho, T.I., Jothiprakash, V. (2015) Impacts of urbanization on flooding of a coastal urban catchment: a case study of Mumbai City, India. *Natural Hazards*. https://doi.org/10.1007/s11069-014-1356-4

19. Mishra, S.P., Sethi, K.C., Siddique, M. (2020) Emerging threats during anthropocene as urban flooding of bhubaneswar city, India. *Water and Energy International*. https://www.scopus.com/pages/publications/85088515130

20. Sadashiv, S., Sar, S. (2025) Urban Flooding in Tier-2 Indian Cities: A Qualitative Exploration of Causes, Impacts and Mitigation Strategies. *Disaster Advances*. https://doi.org/10.25303/1810da090096

21. Tiwari, M., Shukla, S., Rawat, K.S., (...), Kumar Singh, S.K. (2025) High-Resolution Waterlogging Mapping Along Ghazipur Drain in Delhi: A UAV-Based Bathymetric Analysis Approach. *Nature Environment and Pollution Technology*. https://doi.org/10.46488/NEPT.2025.v24i04.B4316

22. Prakash, P., Das, A.K., Sreenivasulu, K., Sreenivasa Reddy, G. (2019) Geospatial technologies as an aid to decision support system for assessment of urban floods in GHMC areas of Hyderabad. *Springer Series in Geomechanics and Geoengineering*. https://doi.org/10.1007/978-3-319-77276-9_34

23. K, K., John, C.M., Shrivasthava, N.K. (2015) Urban flood vulnerability zoning of Cochin City, southwest coast of India, using remote sensing and GIS. *Natural Hazards*. https://doi.org/10.1007/s11069-014-1372-4

24. Guptha, G.C., Swain, S., Al-Ansari, N., (...), Dayal, D. (2021) Evaluation of an urban drainage system and its resilience using remote sensing and GIS. *Remote Sensing Applications: Society and Environment*. https://doi.org/10.1016/j.rsase.2021.100601

25. Li, Ruidong, Liu, Jiapei, Sun, Ting, (...), Ni, Guangheng (2025) Enhancing urban pluvial flood modeling through graph reconstruction of incomplete sewer networks. *Hydrology and Earth System Sciences*. https://doi.org/10.5194/hess-29-5677-2025

26. Kumar Taloor, A.K., Goswami, A., Bahuguna, I.M., (...), Kothyari, G.C. (2022) Remote sensing and GIS applications in water cryosphere and climate change. *Remote Sensing Applications: Society and Environment*. https://doi.org/10.1016/j.rsase.2022.100866

27. Costabile, P., Lombardo, M., Chiaravalloti, F., (...), Carmelina, C. (2026) Predicting pluvial flood impacts in data-scarce urban environments: Uncertainty and interplay between rainfall inputs and conceptual drainage loss models. *Urban Climate*. https://doi.org/10.1016/j.uclim.2025.102724

28. Gaurkhede, N., Adane, V.S. (2025) Siting Selected Sustainable Urban Drainage Systems (SuDS) Through SWMM Simulations in the Sub-Tropical Climate of Nagpur, India. *National Academy Science Letters*. https://doi.org/10.1007/s40009-025-01677-4

29. Mondal, A., Garg, R.D. (2025) Assessing the role of sustainable water bodies in urban drainage systems to mitigate urban flooding: A case study of Gurgaon, Haryana, India. *Physics and Chemistry of the Earth*. https://doi.org/10.1016/j.pce.2024.103803

30. Moon, H.-T., Kim, J.-S., Chen, J., (...), Moon, Y.-I. (2024) Mitigating urban flood Hazards: Hybrid strategy of structural measures. *International Journal of Disaster Risk Reduction*. https://doi.org/10.1016/j.ijdrr.2024.104542

31. Guptha, G.C., Swain, S., Al-Ansari, N., (...), Dayal, D. (2022) Assessing the role of SuDS in resilience enhancement of urban drainage system: A case study of Gurugram City, India. *Urban Climate*. https://doi.org/10.1016/j.uclim.2021.101075

32. Mahato, R.B., Sahu, A., Thomas, D.D. (2026) Integrating rainfall analysis and nature-based solutions for urban flood resilience in Panaji, Goa. *Theoretical and Applied Climatology*. https://doi.org/10.1007/s00704-026-06396-4

33. Guptha, G.C., Swain, S., Al-Ansari, N., (...), Dayal, D. (2022) Assessing the role of SuDS in resilience enhancement of urban drainage system: A case study of Gurugram City, India. *Urban Climate*. https://doi.org/10.1016/j.uclim.2021.101075

34. Ahamed, S.M.F., Agarwal, S. (2019) Urban flood modeling and management using SWMM for new R.R. pet region, Vijayawada, India. *International Journal of Recent Technology and Engineering*. https://www.scopus.com/pages/publications/85067887723

35. Saleh, F., Ramaswamy, V., Wang, Y., (...), Pullen, J. (2017) A multi-scale ensemble-based framework for forecasting compound coastal-riverine flooding: The Hackensack-Passaic watershed and Newark Bay. *Advances in Water Resources*. https://doi.org/10.1016/j.advwatres.2017.10.026

36. Bouttier, François, Marchal, Hugo (2024) Probabilistic short-range forecasts of high-precipitation events: optimal decision thresholds and predictability limits. *Natural Hazards and Earth System Sciences*. https://doi.org/10.5194/nhess-24-2793-2024

37. Bowler, N.E., Pierce, C.E., Seed, A.W. (2006) STEPS: A probabilistic precipitation forecasting scheme which merges an extrapolation nowcast with downscaled NWP. *Quarterly Journal of the Royal Meteorological Society*. https://doi.org/10.1256/qj.04.100

38. Golding, B.W. (1998) Nimrod: A system for generating automated very short range forecasts. *Meteorological Applications*. https://doi.org/10.1017/S1350482798000577

39. Goutham, V., Reddy, M., Padma Priya, K.T., (...), Reddy, S.A.T. (2025) Flood Susceptibility Mapping of Kolkata Municipal Corporation: Using Multi-Criteria Decision-Making Model of GIS and AHP Techniques. *Journal of Engineering Science and Technology Review*. https://doi.org/10.25103/jestr.186.04

40. Srinivas, C.V., Viswanadhapalli, V., Hari Prasad, D., (...), Venkatraman, B. (2018) Simulation of an extreme heavy rainfall event over Chennai, India using WRF: Sensitivity to grid resolution and boundary layer physics. *Atmospheric Research*. https://doi.org/10.1016/j.atmosres.2018.04.014

41. Hardy, J., Gourley, J.J., Kirstetter, P.-E., (...), Flamig, Z.L. (2016) A method for probabilistic flash flood forecasting. *Journal of Hydrology*. https://doi.org/10.1016/j.jhydrol.2016.04.007

42. Acosta-Coll, Melisa, Ballester-Merelo, Francisco, Martinez-Peiró, Marcos, De la Hoz-Franco, Emiro (2018) Real-Time Early Warning System Design for Pluvial Flash Floods—A Review. *Sensors (Basel, Switzerland)*. https://doi.org/10.3390/s18072255

43. Ko, J., Lee, K., Hwang, H., (...), Shin, K. (2022) Effective training strategies for deep-learning-based precipitation nowcasting and estimation. *Computers and Geosciences*. https://doi.org/10.1016/j.cageo.2022.105072

44. Tan, Jinkai, Huang, Qiqiao, Chen, Sheng (2024) Deep learning model based on multi-scale feature fusion for precipitation nowcasting. *Geoscientific Model Development*. https://doi.org/10.5194/gmd-17-53-2024

45. Geng, Huantong, Ge, Xiaoyan, Xie, Boyang, (...), Zhuang, Xiaoran (2023) LSTMAtU-Net: A Precipitation Nowcasting Model Based on ECSA Module. *Sensors (Basel, Switzerland)*. https://doi.org/10.3390/s23135785

46. Mohan, P., Srinivas, C.V., Viswanadhapalli, V., (...), Venkatraman, B. (2018) Simulation of a heavy rainfall event over Chennai in Southeast India using WRF: Sensitivity to microphysics parameterization. *Atmospheric Research*. https://doi.org/10.1016/j.atmosres.2018.04.005

47. Islam, Md Feroz, Bhattacharya, Biswa, Popescu, Ioana (2019) Flood risk assessment due to cyclone-induced dike breaching in coastal areas of Bangladesh. *Natural Hazards and Earth System Sciences*. https://doi.org/10.5194/nhess-19-353-2019

48. https://www.sciencedirect.com/science/article/pii/S2661890726003716

49. Sharma, S., Kumar, S., Kumar, M., Sharma, S. (2026) Machine Learning-Based Prediction of Daily Rainfall in a Monsoonal Urban Environment: A Case Study of Chandigarh, India. *International Conference on Civil, Structural and Transportation Engineering*. https://doi.org/10.11159/iccste26.242

50. Jain, V., Singh, S., Goyal, M.K. (2025) Urban precipitation dynamics risk assessment across Indian smart cities. *Environmental Research Letters*. https://doi.org/10.1088/1748-9326/ade3f0
51. Gupta, A.K., Nair, S.S. (2011) Urban floods in Bangalore and Chennai: Risk management challenges and lessons for sustainable urban ecology. *Current Science*. https://www.scopus.com/pages/publications/79959771587
52. Ranger, N., Hallegatte, S., Bhattacharya, S., (...), Corfee-Morlot, J. (2011) An assessment of the potential impact of climate change on flood risk in Mumbai. *Climatic Change*. https://doi.org/10.1007/s10584-010-9979-2
53. Patel, G., Verma, S., Verma, M.K. (2026) Temporal Disaggregation and Short-Term Forecasting of Hourly Rainfall Data: A Case Study of Raipur City Chhattisgarh, India. *Iranian Journal of Science and Technology - Transactions of Civil Engineering*. https://doi.org/10.1007/s40996-025-01901-7
54. Kumari, A., Kumar, P., Murukesh, M. (2026) Unravelling Changes in Precipitation Extremes over India using CMIP6 Simulations under the High-Emission SSP5-8.5 Scenario. *Pure and Applied Geophysics*. https://doi.org/10.1007/s00024-026-04097-7
55. Manchikatla, M.S., Umamahesh, N.V. (2024) Integrated assessment of future climate and land use changes on urban floods: A Markov chain and PCSWMM-based approach for Hyderabad case study. *Water Science and Technology*. https://doi.org/10.2166/wst.2024.034
56. Guptha, G.C., Swain, S., Al-Ansari, N., (...), Dayal, D. (2021) Evaluation of an urban drainage system and its resilience using remote sensing and GIS. *Remote Sensing Applications: Society and Environment*. https://doi.org/10.1016/j.rsase.2021.100601
57. Manna, H., Das, M., Pramanik, M., (...), Zhran, M. (2025) Ensemble intelligence for urban resilience: flood susceptibility modeling in Mumbai using advanced machine learning. *Geomatics, Natural Hazards and Risk*. https://doi.org/10.1080/19475705.2025.2588718
58. Bhatt, C.M., Srinivasa Rao, G. (2018) HAND (height above nearest drainage) tool and satellite-based geospatial analysis of Hyderabad (India) urban floods, September 2016. *Arabian Journal of Geosciences*. https://doi.org/10.1007/s12517-018-3952-1
59. Roy, A., Roy, P. (2026) Urban flood susceptibility modelling of Guwahati in India using GIS-based AHP-MCDM for sustainable disaster risk management. *Progress in Disaster Science*. https://doi.org/10.1016/j.pdisas.2026.100649
60. Vanama, V.S.K., Shitole, S., Rao, Y.S. (2020) Urban flood mapping with C-band RISAT-1 SAR Images: 2016 Flood Event of Bangalore City, India. *2020 International Conference on Convergence to Digital World - Quo Vadis, ICCDW 2020*. https://doi.org/10.1109/ICCDW45521.2020.9318710
61. Mann, R., Gupta, A. (2023) Mapping flood vulnerability using an analytical hierarchy process (AHP) in the Metropolis of Mumbai. *Environmental Monitoring and Assessment*. https://doi.org/10.1007/s10661-023-12141-5
62. Revi, A. (2008) Climate change risk: An adaptation and mitigation agenda for Indian cities. *Environment and Urbanization*. https://doi.org/10.1177/0956247808089157

63. Singh, N., Paul Shukla, B., Kaushik, N., (...), Bhan, S.C. (2023) Validation of INSAT-3D/3DR based nowcasting rain occurrences for heavy rainfall using Hydro-Estimator product. *Advances in Space Research*. https://doi.org/10.1016/j.asr.2023.05.030

64. Ashrit, R., Mohan, M.S., Dube, A., (...), Mitra, A.K. (2022) Evaluation of five high-resolution global model rainfall forecasts over India during monsoon 2020. *Journal of Earth System Science*. https://doi.org/10.1007/s12040-022-01990-2

65. Trivedi, D., Sharma, O., Pattnaik, S. (2024) Minimization of Forecast Error Using Deep Learning for Real-Time Heavy Rainfall Events over Assam. *IEEE Geoscience and Remote Sensing Letters*. https://doi.org/10.1109/LGRS.2024.3378517

66. Katyal, B., Upadhyaya, S. (2024) EXPLORING DEEP LEARNING TECHNIQUES TO NOWCAST SATELLITE PRECIPITATION RETRIEVALS. *2024 IEEE India Geoscience and Remote Sensing Symposium, InGARSS 2024*. https://doi.org/10.1109/InGARSS61818.2024.10984138

67. Shukla, S., Mishra, S., Mishra, D. (2025) Enhancing BoB cyclone precipitation nowcasting with deep learning: A convolutional LSTM approach using INSAT 3D data. *Journal of Hydrology*. https://doi.org/10.1016/j.jhydrol.2025.133911

68. Upadhyay, A.B., Shah, S.R., Thakker, R.A. (2025) Advanced rainfall nowcasting using 3D convolutional LSTM networks on satellite data. *Journal of Computational Mathematics and Data Science*. https://doi.org/10.1016/j.jcmds.2025.100125

69. Tripathy, P., Murtugudde, R., Chandel, V.S., (...), Ghosh, S. (2026) Hyperlocal urban extreme rainfall forecast system: A convolutional neural network-transfer learning-based downscaling approach. *Urban Climate*. https://doi.org/10.1016/j.uclim.2026.102995

70. Kim, W., Jeong, C.-H., Kim, S. (2024) Improvements in deep learning-based precipitation nowcasting using major atmospheric factors with radar rain rate. *Computers and Geosciences*. https://doi.org/10.1016/j.cageo.2024.105529

71. Singh, K.S., Bonthu, S., Purvaja, R., (...), Ramesh, R. (2018) Prediction of heavy rainfall over Chennai Metropolitan City, Tamil Nadu, India: Impact of microphysical parameterization schemes. *Atmospheric Research*. https://doi.org/10.1016/j.atmosres.2017.11.028

72. Douluri, D.L., Chakraborty, A. (2021) Assessment of WRF-ARW model parameterization schemes for extreme heavy precipitation events associated with atmospheric rivers over West Coast of India. *Atmospheric Research*. https://doi.org/10.1016/j.atmosres.2020.105330

73. de Campos, B., Carvalho, V.S.B., Mattos, E.V. (2023) Assessment of cloud microphysics and cumulus convection schemes to model extreme rainfall events over the Paraiba do Sul River Basin. *Urban Climate*. https://doi.org/10.1016/j.uclim.2023.101618

74. Samanta, S., Prabhakaran, T.V., Murugavel, P., Suneetha, P. (2023) Morphological and microphysical characteristics associated with the lifecycle of a stationary cloud cluster during the Indian Summer Monsoon: A comparative study with numerical simulations and radar observation. *Atmospheric Research*. https://doi.org/10.1016/j.atmosres.2022.106464

75. Kumar, S., Kaushal, D.R., Gosain, A.K. (2019) Evaluation of evolutionary algorithms for the optimization of storm water drainage network for an urbanized area. *Acta Geophysica*. https://doi.org/10.1007/s11600-018-00240-8

76. Zhou, Qianqian, Panduro, Toke Emil, Thorsen, Bo Jellesmark, Arnbjerg-Nielsen, Karsten (2013) Adaption to Extreme Rainfall with Open Urban Drainage System: An Integrated Hydrological Cost-Benefit Analysis. *Environmental Management*. https://doi.org/10.1007/s00267-012-0010-8

77. Rosenberger, L., Leandro, J., Pauleit, S., Erlwein, S. (2021) Sustainable stormwater management under the impact of climate change and urban densification. *Journal of Hydrology*. https://doi.org/10.1016/j.jhydrol.2021.126137

78. Kc, S., Shrestha, S., Ninsawat, S., Chonwattana, S. (2021) Predicting flood events in Kathmandu Metropolitan City under climate change and urbanisation. *Journal of Environmental Management*. https://doi.org/10.1016/j.jenvman.2020.111894

79. Xing, Y., Shao, D., Liang, Q., (...), Ullah, I. (2022) Investigation of the drainage loss effects with a street view based drainage calculation method in hydrodynamic modelling of pluvial floods in urbanized area. *Journal of Hydrology*. https://doi.org/10.1016/j.jhydrol.2021.127365

80. Tran, V.N., Kim, J. (2024) UIDS: A Matlab-based urban flood model considering rainfall-induced and surcharge-induced inundations. *Environmental Modelling and Software*. https://doi.org/10.1016/j.envsoft.2024.106132

81. Barbosa, S.A., Wang, Y., Goodall, J.L. (2025) Exploring infiltration effects on coastal urban flooding: Insights from nuisance to extreme events using 2D/1D hydrodynamic modeling and crowdsourced flood reports. *Science of the Total Environment*. https://doi.org/10.1016/j.scitotenv.2025.178908

82. Tang, B., Nederhoff, K., Gallien, T.W. (2025) Quantifying compound coastal flooding effects in urban regions using a tightly coupled 1D–2D model explicitly resolving flood defense infrastructure. *Coastal Engineering*. https://doi.org/10.1016/j.coastaleng.2025.104728

83. Jibhakate, S.M., Timbadiya, P.V., Patel, P.L. (2023) Flood hazard assessment for the coastal urban floodplain using 1D/2D coupled hydrodynamic model. *Natural Hazards*. https://doi.org/10.1007/s11069-022-05728-7

84. Timbadiya, P.V., Patel, P.L., Porey, P.D. (2015) A 1D-2D coupled hydrodynamic model for river flood prediction in a coastal urban floodplain. *Journal of Hydrologic Engineering*. https://doi.org/10.1061/(ASCE)HE.1943-5584.0001029

85. Tanim, A.H., Smith-Lewis, C., Downey, A.R.J., (...), Goharian, E. (2024) Bayes_Opt-SWMM: A Gaussian process-based Bayesian optimization tool for real-time flood modeling with SWMM. *Environmental Modelling and Software*. https://doi.org/10.1016/j.envsoft.2024.106122

86. Mignot, E., Dewals, B. (2022) Hydraulic modelling of inland urban flooding: Recent advances. *Journal of Hydrology*. https://doi.org/10.1016/j.jhydrol.2022.127763

87. Rai, P.K., Chahar, B.R., Dhanya, C.T. (2017) GIS-based SWMM model for simulating the catchment response to flood events. *Hydrology Research*. https://doi.org/10.2166/nh.2016.260

88. https://www.sciencedirect.com/science/article/pii/S1573295920003524

89. Kim, H., Hong, S., Limos, A.G., (...), Yoon, J. (2023) Improving water quality modelling for green roof runoff using Storm Water Management Model. *Urban Climate*. https://doi.org/10.1016/j.uclim.2023.101717

90. Sun, D., Wang, H., Huang, J., (...), Huang, W. (2024) Urban flood hazard insights from multiple perspectives based on internet of things sensor data. *International Journal of Disaster Risk Reduction*. https://doi.org/10.1016/j.ijdrr.2024.104919

91. Nithila Devi, N., Kuiry, S.N. (2024) Evaluation of Predictive Skill of Short-Term Raw Ensemble Forecasts for Mapping of Extreme Floods. *Lecture Notes in Mechanical Engineering*. https://doi.org/10.1007/978-981-97-6009-1_17

92. Nikam, V., Gupta, K. (2014) SVM-based model for short-term rainfall forecasts at a local scale in the Mumbai Urban Area, India. *Journal of Hydrologic Engineering*. https://doi.org/10.1061/(ASCE)HE.1943-5584.0000875

93. Solanki, H., Vegad, U., Kushwaha, A., Mishra, V. (2025) Improving Streamflow Prediction Using Multiple Hydrological Models and Machine Learning Methods. *Water Resources Research*. https://doi.org/10.1029/2024WR038192

94. Villarini, G., Krajewski, W.F., Ntelekos, A.A., (...), Smith, J.A. (2010) Towards probabilistic forecasting of flash floods: The combined effects of uncertainty in radar-rainfall and flash flood guidance. *Journal of Hydrology*. https://doi.org/10.1016/j.jhydrol.2010.02.014

95. Khatri, S., Kokane, P., Kumar, V., Pawar, S. (2023) Prediction of waterlogged zones under heavy rainfall conditions using machine learning and GIS tools: a case study of Mumbai. *GeoJournal*. https://doi.org/10.1007/s10708-022-10731-3

96. Ghosh, S., Pal, M., Maity, D.K., (...), Ghosal, R. (2025) Mapping stormwater inundation potential in data-scarce urban areas: A machine learning approach for Kolkata, India. *Urban Climate*. https://doi.org/10.1016/j.uclim.2025.102451

97. Nevo, Sella, Morin, Efrat, Gerzi Rosenthal, Adi, (...), Reich, Ofir (2022) Flood forecasting with machine learning models in an operational framework. *Hydrology and Earth System Sciences*. https://doi.org/10.5194/hess-26-4013-2022

98. Zwirglmaier, V., Garschagen, M. (2024) Linking urban structure types and Bayesian network modelling for an integrated flood risk assessment in data-scarce mega-cities. *Urban Climate*. https://doi.org/10.1016/j.uclim.2024.102034

99. Zwirglmaier, Veronika, Reimuth, Andrea, Garschagen, Matthias (2024) How suitable are current approaches to simulate flood risk under future urbanization trends?. *Environmental Research Letters*. https://doi.org/10.1088/1748-9326/ad536f

100. Mondal, A., Garg, R.D. (2025) Assessing the role of sustainable water bodies in urban drainage systems to mitigate urban flooding: A case study of Gurgaon, Haryana, India. *Physics and Chemistry of the Earth*. https://doi.org/10.1016/j.pce.2024.103803

101. Salim, M.Z., Choudhari, N., Kafy, A.-A., (...), Al-Ramadan, B. (2024) A comprehensive review of navigating urbanization induced climate change complexities for sustainable groundwater resources management in the Indian subcontinent. *Groundwater for Sustainable Development*. https://doi.org/10.1016/j.gsd.2024.101115

102. Kundu, A., Roy, S., Chauhan, A. (2026) Spatiotemporal flood hazard mapping in Kolkata based on Sentinel-1 SAR observations. *Advances in Space Research*. https://doi.org/10.1016/j.asr.2025.12.082

103. Vorobevskii, Ivan, Park, Jeongha, Kim, Dongkyun, (...), Kronenberg, Rico (2024) Simulating sub-hourly rainfall data for current and future periods using two statistical disaggregation models: case studies from Germany and South Korea. *Hydrology and Earth System Sciences*. https://doi.org/10.5194/hess-28-391-2024

104. Pandey, P., Palaniswamy, S. (2025) Multi-Modal Flood Detection and Validation: Integrating Sentinel-1 SAR, Deep Learning, and Named Entity Recognition for Enhanced Situational Awareness. *2025 International Conference on Intelligent Computing and Knowledge Extraction, ICICKE 2025*. https://doi.org/10.1109/ICICKE65317.2025.11136558