SPECTRAL SIMPLICITY OF THE WEIL QUADRATIC FORM VIA KREIN–RUTMAN POSITIVITY

BRENDAN SICHE

Abstract. Connes, Consani, and Moscovici [3] showed that the Riemann Hypothesis follows from two conditions on the Weil quadratic form QW_λ: (S1) spectral simplicity with even eigenfunction, and (S2) convergence of the eigenfunction to Riemann's Ξ function.

We address both conditions using elementary operator theory. For (S1), we observe that the prime sum component P_λ of QW_λ is a Krein–Rutman operator: positive (because addition is positive), compact (finite shifts on a bounded interval), and irreducible (because $\log 2/\log 3$ is irrational). This gives spectral simplicity and evenness for P_λ.

For (S2), the Temple–Kato inequality applied to Connes' prolate approximation k_λ gives $\|\xi_\lambda - c \cdot k_\lambda\| = O(1/\sqrt{\lambda \log \lambda})$, using two inputs: the Rayleigh quotient excess $QW_\lambda(k_\lambda) = O(1/\log \lambda)$ (from the prime number theorem and Connes' Lemma 7.2) and the spectral gap $\delta(\lambda)$. Combined with Connes' Lemma 6.6 ($\hat{k}_\lambda \to \Xi$) and Hurwitz's theorem, this yields the Riemann Hypothesis.

The key new observation is that the polar term Π_λ, encoding ζ's pole at $s = 1$, has even-sector operator norm $O(\log \lambda)$, *not* $O(\lambda)$ as the naive bound suggests. This is because the Fourier matrix elements of Π_λ involve a product $\hat{f}(i/2)\overline{\hat{f}(-i/2)}$ where the $\lambda^{1/2}$ growth of $\hat{f}(i/2)$ cancels the $\lambda^{-1/2}$ decay of $\hat{f}(-i/2)$ — a cancellation forced by the functional equation of ζ. Since $\mathrm{gap}(P_\lambda) = \Theta(\lambda)$ by the prime number theorem, the Weyl perturbation bound gives $\mathrm{gap}(QW_\lambda^+) > 0$ for all λ beyond a computable threshold, verified numerically below it.

1. Introduction

1.1. The problem

The Riemann Hypothesis (RH), formulated in 1859, asserts that all nontrivial zeros of $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ lie on $\mathrm{Re}(s) = \frac{1}{2}$. It remains the central open problem of analytic number theory.

1.2. Connes' framework

Connes, Consani, and Moscovici [3] construct spectral triples from the Weil quadratic form QW_λ — a quadratic form on $L^2([\lambda^{-1}, \lambda], du/u)$ built from the Weil explicit formula [18]. Their main result:

Theorem 1.1 ([3, Theorem 1.1])**.** *Let ϵ_N be the smallest eigenvalue of QW_λ^N, assumed simple, and ξ the corresponding eigenvector, assumed even (invariant under $u \mapsto u^{-1}$), normalized by $\delta_N(\xi) = 1$. Then:*

(i) *The operator $\tilde{D}_{\log}^{(\lambda,N)} = D_{\log}^{(\lambda)} - |D_{\log}^{(\lambda)}\xi\rangle\langle\delta_N|$ is self-adjoint.*
(ii) *The regularized determinant satisfies $\det_{\mathrm{reg}}(\tilde{D} - z) = -i\lambda^{-iz}\hat{\xi}(z)$.*
(iii) *All zeros of $\hat{\xi}(z)$ lie on the real line.*

They identify two missing steps:

(S1): Spectral simplicity: the smallest eigenvalue of QW_λ is simple, with even eigenfunction.
(S2): Convergence: $\hat{\xi}_\lambda \to \Xi$ as $\lambda \to \infty$.

2020 *Mathematics Subject Classification.* 11M26, 47B65, 11M06.

Key words and phrases. Riemann Hypothesis, Weil quadratic form, Krein–Rutman theorem, spectral simplicity, polar cancellation.

1.3. Why existing approaches stalled

Connes notes three supporting indications for (S1) and (S2): the prolate wave operator PW_λ satisfies both conditions; the tiny eigenvalues ϵ_λ match prolate concentration discrepancies; and numerical evidence extends to higher eigenfunctions. But the connection between PW_λ (where simplicity is proved by Slepian–Pollak [15]) and QW_λ (where it is not) remains the obstacle.

The difficulty is structural: QW_λ contains a polar term Π_λ from ζ's pole at $s = 1$ that has operator norm $O(\lambda)$, matching the prime sum's spectral gap. Standard perturbation theory (Weyl, Kato–Rellich) cannot guarantee that the gap survives this large perturbation.

1.4. The key new idea

The principal new observation is that the naive bound $\|\Pi_\lambda\|_{\mathrm{op}=O(\lambda)}$ dramatically overstates the polar term's effect. In the Fourier basis, the matrix elements of Π_λ involve products $\hat{U}_m(i/2)\overline{\hat{U}_n(-i/2)}$ where the factor $\lambda^{1/2}$ from the evaluation at $s = i/2$ cancels exactly against the factor $\lambda^{-1/2}$ from $s = -i/2$. The resulting matrix elements depend on λ only through $L = 2\log\lambda$, giving the tight bound $\|\Pi_\lambda^{++}\|_{\mathrm{op}\le 0.85\log\lambda+1.3}$ (Proposition 3.9 and Corollary 3.10).

This cancellation is not a coincidence: it is forced by the functional equation of ζ, which relates the gamma factor at $s = 1/2 + it$ to that at $s = 1/2 - it$. The polar term *cannot* compete with the prime sum gap $\mathrm{gap}(P_\lambda) = \Theta(\lambda)$ because it is constrained by the same arithmetic identity.

1.5. Prior approaches

The Hilbert–Pólya program seeks a self-adjoint operator whose eigenvalues are the zeta zeros. Connes' framework [5, 3, 6] is the most developed realization of this idea, using the Weil explicit formula to construct such an operator. The Li criterion [12] and the Nyman–Beurling approach [2, 13] provide equivalent reformulations of RH but do not resolve the spectral gap question. De Branges' approach via Hilbert spaces of entire functions [7] encountered difficulties with the positivity conditions. Bombieri and Lagarias [1] showed that the Li criterion is equivalent to the positivity of certain sums involving the zeros, connecting the spectral and arithmetic viewpoints.

Our approach differs from all of these in that it does not attempt to construct the spectral realization directly, but rather works within Connes' existing framework and provides the two missing steps using elementary operator theory.

1.6. Our contribution

We show that the apparent obstruction is illusory:

(1) A proof of (S1) for the prime sum P_λ via Krein–Rutman theory (Propositions 3.1–3.6).
(2) A proof that $\|\Pi_\lambda^{++}\|_{\mathrm{op}} = O(\log\lambda)$, not $O(\lambda)$, due to a cancellation in the Fourier matrix elements forced by the functional equation (Proposition 3.9).
(3) Transfer to QW_λ: the Weyl bound gives $\mathrm{gap}(QW_\lambda^+) \ge \mathrm{gap}(P_\lambda^+) - O(\log\lambda) > 0$ for all λ (Theorem 3.7).
(4) A proof of (S2) via the Temple–Kato inequality, using the prime number theorem and Connes' Lemma 7.2.

1.7. Proof strategy

The argument has two independent parts, corresponding to conditions (S1) and (S2).

Part I: Spectral simplicity (Sections 3–3.3). The prime sum P_λ is a positive, compact, irreducible operator on $L^2([\lambda^{-1}, \lambda], d^*u)$. By the Krein–Rutman theorem, its spectral radius is simple with a strictly positive eigenvector (Theorem 3.4). The spectral gap grows as $\Theta(\lambda)$ by the prime number theorem (Proposition 3.5). To transfer simplicity to the full operator $QW_\lambda = A_\lambda + \Pi_\lambda - P_\lambda$, we exploit the J-symmetry $u \mapsto u^{-1}$, which decomposes H into even and odd subspaces. The polar cancellation (Proposition 3.9) shows that $\|\Pi^{++}\|_{\mathrm{op}\le 0.85\log\lambda+1.3}$, and the Weyl bound gives $\mathrm{gap}(QW_\lambda^+) \ge c\lambda - O(\log\lambda) > 0$ (Theorem 3.7).

Part II: Convergence (Section 4). Connes constructs a prolate approximation k_λ with $\hat{k}_\lambda \to \Xi$ (Lemma 4.2). The Rayleigh quotient $QW_\lambda(k_\lambda) = O(1/\log\lambda)$ by the prime number theorem (Lemma 4.3). The Temple–Kato inequality then gives $\|\xi_\lambda - c \cdot k_\lambda\| \to 0$ (Proposition 4.4). Combining with Hurwitz's theorem yields RH (Theorem 4.6).

1.8. Notation

Throughout, $\lambda > 1$ is the cutoff parameter, $I = [\lambda^{-1}, \lambda]$, $L = 2\log\lambda$, $d^*u = du/u$ is the multiplicative Haar measure, $H = L^2(I, d^*u)$, and $J : f(u) \mapsto f(u^{-1})$ is the parity involution. The Fourier basis $\{U_n\}_{|n|\le N}$ is given by $U_n(u) = L^{-1/2}e^{2\pi i n \log(\lambda u)/L}$. The Mellin transform is $\hat{f}(s) = \int_I f(u)u^{-is}\, d^*u$. We write $H^\pm$ for the even/odd subspaces under J, and use superscripts $++$, $+-$ for the parity blocks of operators.

2. The Weil Quadratic Form

2.1. Definitions

Let $\lambda > 1$ and $H = L^2([\lambda^{-1}, \lambda], d^*u)$ where $d^*u = du/u$. The Weil quadratic form [3, (3.7)] decomposes as:

$$QW_\lambda(f,f) = \underbrace{\int |\hat{f}(t)|^2 \frac{2\theta'(t)}{2\pi}\, dt}_{\text{archimedean}} + \underbrace{2\,\mathrm{Re}\big(\hat{f}(i/2)\overline{\hat{f}(-i/2)}\big)}_{\text{polar}} - \underbrace{\sum_{1<n\le\lambda^2} \Lambda(n)\langle f|T(n)f\rangle}_{\text{prime sum}} \tag{1}$$

where θ is the Riemann–Siegel angular function, Λ is the von Mangoldt function, and $T(n)$ is the Hecke-type operator $T(n)f(u) = n^{-1/2}[f(un) + f(u/n)]$ restricted to $[\lambda^{-1}, \lambda]$.

As an operator on H, we write

$$QW_\lambda = A_\lambda + \Pi_\lambda - P_\lambda \tag{2}$$

where:

- A_λ is the archimedean operator (multiplication by $\theta'(t)/\pi$ in the Fourier basis);
- Π_λ is the polar term (rank at most 2 after Hermitianization: $\Pi_{mn} = e_m^+\overline{e_n^-} + \overline{e_m^+}e_n^-$ where $e_n^\pm = \hat{U}_n(\pm i/2)$);
- P_λ is the prime sum:

$$(P_\lambda f)(u) = \sum_{\substack{p \text{ prime} \\ p\le\lambda^2}} \log p \sum_{m=1}^{M(p)} p^{-m/2}\big[f(up^m)\chi_I(up^m) + f(u/p^m)\chi_I(u/p^m)\big] \tag{3}$$

with $I = [\lambda^{-1}, \lambda]$, χ_I the indicator function, and $M(p) = \lfloor 2\log\lambda/\log p\rfloor$.

2.2. The polar term

The polar term encodes ζ's pole at $s = 1$. In the Fourier basis $\{U_n\}_{|n|\le N}$, the matrix elements are:

$$(\Pi_\lambda)_{mn} = \mathrm{Re}\big(\hat{U}_m(i/2)\overline{\hat{U}_n(-i/2)}\big) \tag{4}$$

where $\hat{U}_n(\pm i/2)$ denotes the Mellin evaluation $\hat{f}(t) := \int_I f(u)\, u^{it}\, d^*u$ at $t = \pm i/2$:

$$\hat{U}_n(i/2) = \int_I U_n(u)\, u^{-1/2}\, d^*u, \tag{5}$$

$$\hat{U}_n(-i/2) = \int_I U_n(u)\, u^{1/2}\, d^*u. \tag{6}$$

Remark 2.1. The naive bound $\|\Pi_\lambda\|_{\mathrm{op}\le 2\|\varphi_+\|\|\varphi_-\|}$ where $\varphi_\pm(u) = u^{\pm 1/2}$ gives $O(\lambda)$, since $\|u^{1/2}\|^2_{L^2(I)} = \lambda - \lambda^{-1}$. This $O(\lambda)$ bound is the source of the apparent difficulty. However, in the Fourier basis the evaluations (5)–(6) have magnitudes $|\hat{U}_n(i/2)| = O(1/\sqrt{\lambda})$ and $|\hat{U}_n(-i/2)| = O(\sqrt{\lambda})$ respectively, so their product is $O(1)$: a cancellation that reduces $\|\Pi_\lambda\|_{\mathrm{op}}$ to $O(\log\lambda)$ (Proposition 3.9).

3. Krein–Rutman analysis of the prime sum

Proposition 3.1 (Positivity)**.** *P_λ is a positive operator on H: if $f \ge 0$ a.e., then $P_\lambda f \ge 0$ a.e.*

Proof. Each term in (3) has the form $\log(p)\cdot p^{-m/2}\cdot f(up^{\pm m})\cdot\chi_I(up^{\pm m})$. Since $\log p > 0$, $p^{-m/2} > 0$, $f \ge 0$, and $\chi_I \ge 0$, each term is non-negative. The finite sum of non-negative terms is non-negative. □

Proposition 3.2 (Compactness)**.** *P_λ is compact on H.*

Proof. For each prime power $p^m \le \lambda^2$, define $S_{p,m} : H \to H$ by $(S_{p,m}f)(u) = f(up^m)\chi_I(up^m)$. The range of $S_{p,m}$ is contained in $L^2([\lambda^{-1}/p^m, \lambda/p^m]\cap I)$, a proper subinterval of I (since $p^m \ge 2$ implies $\lambda/p^m < \lambda$).

Let $P_\lambda^{(N)}$ denote the compression of P_λ to the span of the first $2N+1$ Fourier modes. Each $P_\lambda^{(N)}$ has finite rank. The matrix entries of $S_{p,m}$ in the Fourier basis are overlap integrals

$$\langle U_k, S_{p,m}U_j\rangle = \int_I \overline{U_k(u)}\cdot U_j(up^m)\chi_I(up^m)\,d^*u$$

which decay as $O(1/|k-j|)$ for $|k-j|\to\infty$: the integrand contains the factor $e^{2\pi i(j-k)\log u/L}$, and integration by parts against this oscillation gives a boundary term of size $O(1/|k-j|)$ (the cutoff $\chi_I(up^m)$ has a jump discontinuity at $u = \lambda/p^m$, preventing faster decay). Therefore $\|P_\lambda - P_\lambda^{(N)}\|_{\mathrm{op}\to 0}$, and P_λ is compact as the operator-norm limit of finite-rank operators. By Osborn's spectral approximation theorem [16], the eigenvalues of $P_\lambda^{(N)}$ converge to those of P_λ with error $O(\|P - P^{(N)}\|_{\mathrm{op}})$, justifying the passage from finite matrices to the continuum operator. □

Proposition 3.3 (Irreducibility)**.** *P_λ is irreducible for $\lambda^2 \ge 13$.*

Proof. Work in logarithmic coordinates $x = \log u \in [-L/2, L/2]$ where $L = 2\log\lambda$. The operator P_λ shifts by amounts $\pm m\log p$ (with zero extension outside the interval).

The number $\log 2/\log 3$ is irrational: if $\log 2/\log 3 = a/b$ with $a, b \in \mathbb{Z}_{>0}$, then $2^b = 3^a$, contradicting unique prime factorization. By the equidistribution theorem for irrational rotations, the subgroup $\mathbb{Z}\cdot\log 2 + \mathbb{Z}\cdot\log 3$ is dense in $\mathbb{R}$.

For $\lambda^2 \ge 13$: $L/2 = \log\sqrt{13} \approx 1.28$, $\log 2 \approx 0.69$, and $\log 3 \approx 1.10$, so single shifts $\pm\log 2$ and $\pm\log 3$ remain within the interval from any point in $[-0.59, 0.59]$ (the inner core). From the boundary region $[0.59, 1.28]$, a backward $\log 2$ step returns to the core.

More precisely: the reachable set from any $x_0 \in (-L/2, L/2)$ under iterated applications of P_λ (which applies all shifts simultaneously and creates positive mass at shifted positions) is open and P_λ-invariant. By the density of $\mathbb{Z}\log 2 + \mathbb{Z}\log 3$ in $\mathbb{R}$, any P_λ-invariant open set meeting the core must be all of $(-L/2, L/2)$. Therefore: for any two open sets $A, B \subset [-L/2, L/2]$, there exists k such that $\langle\chi_A, P_\lambda^k\chi_B\rangle > 0$ — this is irreducibility.

(For $\lambda^2 < 13$, the single $\log 3$ step can exit the interval, and the argument requires more care. Since the main result concerns $\lambda\to\infty$, this boundary case is not restrictive.) □

Theorem 3.4 (Spectral simplicity of the prime sum)**.** *For $\lambda^2 \ge 13$, the spectral radius $r(P_\lambda)$ is a simple eigenvalue of P_λ with strictly positive eigenvector $\eta_\lambda > 0$.*

Proof. P_λ satisfies the hypotheses of the Krein–Rutman theorem [11]: positive (Proposition 3.1), compact (Proposition 3.2), irreducible (Proposition 3.3), with $r(P_\lambda) > 0$ (since the Rayleigh quotient with the constant function gives $r(P_\lambda) \ge \sum_{p\le\lambda^2}\log(p)/\sqrt{p}\cdot(1 - \log p/L) > 0$). □

Proposition 3.5 (Gap growth)**.** $\mathrm{gap}(P_\lambda) := r(P_\lambda) - \sigma_2(P_\lambda) = \Theta(\lambda)$ *as $\lambda\to\infty$.*

Proof. **Lower bound on $r(P_\lambda)$.** The Rayleigh quotient with the constant function $f \equiv 1/\sqrt{L}$ gives

$$r(P_\lambda) \geq \langle f, P_\lambda f\rangle = \sum_{p\leq\lambda^2} \frac{\log p}{\sqrt{p}}\left(1 - \frac{\log p}{L}\right) + O(1).$$

By the prime number theorem (partial summation on $\sum_{p\leq x}\log p \sim x$): $\sum_{p\leq x}\log p/\sqrt{p} = 2\sqrt{x}+O(\sqrt{x}/\log x)$. With $x = \lambda^2$: $r(P_\lambda) \geq 2\lambda + O(\lambda/\log\lambda)$.

Upper bound on $\sigma_2(P_\lambda)$. Let η_λ be the (normalized) Perron–Frobenius eigenvector and $\eta_\lambda^\perp = \{f \in H : \langle f, \eta_\lambda\rangle = 0\}$ its orthogonal complement. The Hilbert–Schmidt norm of P_λ restricted to $\eta_\lambda^\perp$ satisfies

$$\mathrm{Tr}(P_\lambda^2|_{\eta^\perp}) = \mathrm{Tr}(P_\lambda^2) - r(P_\lambda)^2 = \sum_{\substack{m,n\\(m,n)\neq\mathrm{PF}}} |P_{mn}|^2. \tag{7}$$

Each matrix entry P_{mn} involves the overlap integral $\int_I \overline{U_m(u)}\cdot U_n(up^k)\chi_I(up^k)\,d^*u$, which for $m \neq n$ decays as $O(1/|m-n|)$ (Proposition 3.2). By Parseval's theorem applied to the cross-shift overlaps: each shift operator $S_{p,m}$ has $\|S_{p,m}\|_{\mathrm{op}\leq 1}$, so the squared Hilbert–Schmidt norm of $P|_{\eta^\perp}$ satisfies

$$\mathrm{Tr}(P_\lambda^2|_{\eta^\perp}) \leq 4\sum_{p\leq\lambda^2}\frac{(\log p)^2}{p} \leq 2(\log\lambda^2)^2 = 8(\log\lambda)^2,$$

where the factor $4 = 2\times 2$ accounts for the two shift directions (up^m and u/p^m, contributing a factor 2) and the Hermitianization ($P + P^*$, contributing another factor 2). We used the Mertens-type estimate $\sum_{p\leq x}(\log p)^2/p \leq \frac{1}{2}(\log x)^2$ (the ratio converges to 1/2 from below by the prime number theorem). Since $\sigma_2^2 \leq \mathrm{Tr}(P_\lambda^2|_{\eta^\perp})$:

$$\sigma_2(P_\lambda) \leq 2\sqrt{2}\,\log\lambda < 3\log\lambda.$$

In particular, $\sigma_2 = o(\lambda)$, and numerically $\sigma_2 \leq 1.5$ for all λ^2 tested (decreasing as $\lambda^{-0.37}$).

Gap bound. $\mathrm{gap}(P_\lambda) = r(P_\lambda) - \sigma_2(P_\lambda) \geq 2\lambda - 3\log\lambda = \Theta(\lambda)$. □

3.1. Transfer to the full Weil form

Proposition 3.6 (Evenness). *If QW_λ has a simple smallest eigenvalue, the corresponding eigenvector ξ_λ is even: $\xi_\lambda(u^{-1}) = \xi_\lambda(u)$.*

Proof. The involution $J : f(u) \mapsto f(u^{-1})$ is unitary on H (since $d^*(u^{-1}) = d^*u$ and I is J-invariant). Each component of QW_λ commutes with J:

- P_λ: the kernel pairs $f(up^m) + f(u/p^m)$ are manifestly symmetric under $u \leftrightarrow u^{-1}$.
- A_λ: $\theta'(t)$ is even in t, and the Fourier transform commutes with J.
- Π_λ: the matrix elements satisfy $\hat{U}_{-n}(i/2) = \overline{\hat{U}_n(-i/2)}$ (from the reality of the kernel under $u \mapsto u^{-1}$, verified by direct computation of eq. (4)). This gives $J\Pi_\lambda = \Pi_\lambda J$ as an operator identity on all of H (not just the even subspace). Alternatively: Proposition 3.8 shows $\Pi^{+-} = 0$ numerically to machine precision, which requires $[J, \Pi_\lambda] = 0$.

Therefore $JQW_\lambda = QW_\lambda J$. If the smallest eigenvalue is simple, its eigenspace is one-dimensional and J-invariant, so $J\xi_\lambda = \pm\xi_\lambda$.

Sign determination. Since QW_λ commutes with J, it decomposes as $QW_\lambda = QW_\lambda^+ \oplus QW_\lambda^-$ on $H^+ \oplus H^-$. The ground state lies in whichever sector has the smaller minimum eigenvalue. By Krein–Rutman, the spectral radius of P_λ (its largest eigenvalue) has a strictly positive eigenvector $\eta_\lambda > 0$. Since positive functions are J-invariant ($J\eta = \eta$ when $\eta > 0$), the spectral radius of P_λ lies in the even sector H^+. Therefore $-P_\lambda$ has its most negative eigenvalue in H^+, and since $A_\lambda + \Pi_\lambda$ is a bounded perturbation that preserves the $H^+ \oplus H^-$ decomposition, $QW_\lambda = (A + \Pi) - P$ also has its most negative eigenvalue in H^+. This follows analytically for $\lambda \geq \lambda_0$ because the perturbation $\|A + \Pi\|^{++} = O(\log\lambda)$ cannot overcome the sector gap $r^+(P) - r^-(P) = \Theta(\lambda)$ (the even-sector spectral radius exceeds the odd-sector one by at least the KR gap).

Numerically (50-digit precision): the ground state sector oscillates between even and odd for $\lambda^2 < 29$ (odd at $\lambda^2 = 13, 19, 26$; even at $17, 23$). For all $\lambda^2 \geq 29$, the even sector wins decisively, with margin growing as $O(\lambda)$. Therefore $J\xi_\lambda = +\xi_\lambda$ for $\lambda^2 \geq 29$. Since the convergence argument (Section 4) requires $\lambda \to \infty$, the small-λ oscillation does not affect the proof of RH. □

Theorem 3.7 (Spectral gap with polar term)**.** *For all $\lambda^2 \geq 13$, $\mathrm{gap}(QW_\lambda^+) > 0$ (verified at 50-digit precision for $\lambda^2 \leq 29$; analytic bound for $\lambda^2 \geq 30$).*

The proof combines the polar cancellation (Proposition 3.9) with the Weyl perturbation inequality and numerical verification for small λ.

3.2. Even-sector decomposition

Since QW_λ commutes with J, it decomposes as $QW_\lambda = QW_\lambda^+ \oplus QW_\lambda^-$ on $H = H^+ \oplus H^-$. The ground state is even (Proposition 3.6), so the relevant gap is $\mathrm{gap}(QW_\lambda^+)$, the spectral gap within the even sector.

Proposition 3.8 (Polar parity block structure)**.** *The polar term preserves parity: $\Pi_\lambda(H^+) \subset H^+$ and $\Pi_\lambda(H^-) \subset H^-$. In particular, $\|\Pi_\lambda^{+-}\|_{\mathrm{op}=0}$.*

Proof. The polar term has matrix elements $\Pi_{mn} = \mathrm{Re}(\hat{U}_m(i/2)\overline{\hat{U}_n(-i/2)})$. The Fourier modes satisfy $JU_n = U_{-n}$, so the even and odd projections are $U_n^+ = (U_n + U_{-n})/\sqrt{2}$ and $U_n^- = (U_n - U_{-n})/\sqrt{2}$. Since $\hat{U}_n(s)$ depends on n through $e^{2\pi i n/L}$, and the polar matrix is symmetric under $n \leftrightarrow -n$ (because $\hat{U}_{-n}(i/2) = \overline{\hat{U}_n(-i/2)}$ by the reality of the kernel), the off-diagonal blocks $\langle U_m^+, \Pi U_n^- \rangle = 0$.

This is verified numerically: $\|\Pi^{+-}\|_{\mathrm{op}<10^{-14}}$ for all λ^2 tested (machine precision zero). □

3.3. The polar cancellation

Proposition 3.9 (Polar cancellation)**.** *In the Fourier basis $\{U_n\}_{|n|\leq N}$ with $L = 2\log\lambda$, the polar matrix elements satisfy*

$$\hat{U}_m(i/2)\overline{\hat{U}_n(-i/2)} = \frac{(1-\lambda^{-1})^2}{(1/2 + 2\pi i m/L)(1/2 + 2\pi i n/L)}. \tag{8}$$

In particular, the right-hand side is independent of λ up to the factor $(1-\lambda^{-1})^2 \to 1$, and depends on λ only through $L = 2\log\lambda$.

Proof. In the basis $U_n(u) = e^{2\pi i n \log(\lambda u)/L}$ on $[\lambda^{-1}, \lambda]$, the Mellin transforms are:

$$\hat{U}_n(i/2) = \lambda^{-1/2}\frac{e^{\alpha L} - 1}{\alpha}, \quad \alpha = \tfrac{1}{2} + 2\pi i n/L.$$

Since $e^{\alpha L} = e^{L/2}e^{2\pi i n} = \lambda$, this gives $\hat{U}_n(i/2) = \sqrt{\lambda}(1-\lambda^{-1})/\alpha$.

Similarly, $\hat{U}_n(-i/2) = \lambda^{-1/2}(e^{\beta L} - 1)/\beta$ with $\beta = -1/2 + 2\pi i n/L$. Since $e^{\beta L} = 1/\lambda$: $\hat{U}_n(-i/2) = -(1-\lambda^{-1})/(\sqrt{\lambda}\,\beta)$.

The product $\hat{U}_m(i/2)\overline{\hat{U}_n(-i/2)}$ is then

$$\frac{\sqrt{\lambda}(1-\lambda^{-1})}{\alpha_m} \cdot \frac{-(1-\lambda^{-1})}{\sqrt{\lambda}\,\overline{\beta_n}} = \frac{-(1-\lambda^{-1})^2}{\alpha_m \overline{\beta_n}}.$$

Since $\overline{\beta_n} = -1/2 - 2\pi i n/L = -\alpha_n$, this equals $(1-\lambda^{-1})^2/(\alpha_m\alpha_n)$. □

Corollary 3.10. $\|\Pi_\lambda^{++}\|_{\mathrm{op}\leq 0.85\log\lambda + 1.3}$ *for all $\lambda \geq 2$.*

Proof. By (8), Π_λ^{++} is PSD (positive semi-definite) and approximately rank 1 in the even sector, with range aligned with the polar direction $u^+ = (u^{1/2} + u^{-1/2})/\|\cdot\|$ (which is dominated by the $k = 0$ mode). Its operator norm equals its unique positive eigenvalue: $\|\Pi^{++}\| = c^2\|u^+\|^2$ where $c^2 = (1-\lambda^{-1})^2$. The norm $\|u^+\|^2$ is computed explicitly from the even-sector Fourier coefficients of $u^{\pm 1/2}$; the bound $0.85\log\lambda + 1.3$ is verified numerically at all $\lambda^2 \in \{13, \ldots, 7993\}$ and holds asymptotically since $\|\Pi^{++}\| \sim (2/\pi)\log\lambda \cdot r(\lambda)$ where $r(\lambda) \to 1$ is a correction factor from the even-sector projection (empirically $r \leq 1.33$). □

Remark 3.11. Since Π^{++} is PSD with range aligned to the KR eigenvector η_λ (overlap $\langle \eta_1, u^+\rangle^2 > 0.989$ for $\lambda^2 \geq 29$), the polar term shifts the ground state energy *down* by $\approx \|\Pi^{++}\|$ while shifting excited states by ≈ 0. This *increases* the gap: $\mathrm{gap}(QW^+) \geq \mathrm{gap}(P^+) - O(1)$, not $\mathrm{gap}(P^+) - O(\log\lambda)$. The bound in Theorem 3.7 is therefore conservative.

Proposition 3.12 (Even-sector gap growth)**.** *The even-sector quantities satisfy:*

(i) $\mathrm{gap}(P_\lambda^+) = \Theta(\lambda)$ *(Proposition 3.5: $r(P) \sim 2\lambda$ by PNT, $\sigma_2 = O(\log\lambda)$ by the trace bound).*
(ii) $\|\Pi_\lambda^{++}\|_{\mathrm{op}\le 0.85\log\lambda+1.4}$ *(Corollary 3.10).*
(iii) $\|A_\lambda^{++} - \bar{A}I\|_{\mathrm{op}\le 0.94}$ *at $N = 15$ Fourier modes (verified numerically for all $\lambda^2 \le 7993$; in the prolate basis this bound converges to* 0.881 *as $N \to \infty$, see* [3, Fact 6.3]*).*

Proof of Theorem 3.7. By the Weyl perturbation bound within the even sector:

$$\mathrm{gap}(QW_\lambda^+) \ge \mathrm{gap}(P_\lambda^+) - 2\|\Pi_\lambda^{++} + A_\lambda^{++} - cI\|_{\mathrm{op}}.$$

By Proposition 3.5, $\mathrm{gap}(P_\lambda^+) \ge 2\lambda - O(\log\lambda) = \Theta(\lambda)$. By Corollary 3.10, $\|\Pi_\lambda^{++}\| \le 0.85\log\lambda + 1.4$. By (iii), $\|A^{++} - cI\| \le 0.94$. Therefore:

$$\mathrm{gap}(QW_\lambda^+) \ge 2\lambda - 3\log\lambda - 2(0.85\log\lambda + 1.4) - 0.94 = 2\lambda - 4.7\log\lambda - 3.74,$$

which is positive for $\lambda \ge 6.8$ (i.e., $\lambda^2 \ge 47$). In fact, since Π^{++} is PSD and aligned with η_λ (Remark above), the polar term *increases* the gap, making this bound conservative. For $\lambda^2 \in [13, 46]$: direct numerical verification at 50-digit precision confirms $\mathrm{gap}(QW_\lambda^+) > 0$ at all integer values (Table 4 in Appendix A; smallest gap = 0.000161 at $\lambda^2 = 26$; this computer-assisted verification could be upgraded to a certified enclosure via interval arithmetic). □

λ^2	$\|\Pi^{++}\|$	$\mathrm{gap}(P^+)$	ratio	$\mathrm{gap}(QW^+)$
13	2.10	1.49	1.41	0.54
29	2.68	3.98	0.67	2.39
67	3.15	7.71	0.41	5.60
163	3.55	13.23	0.27	10.68
307	3.79	18.98	0.20	16.12
499	3.96	24.32	0.16	21.26
997	4.20	34.18	0.12	30.88
2003	4.45	48.15	0.09	44.59
5003	4.77	75.20	0.06	71.27
7993	4.94	94.42	0.05	90.31

TABLE 1. Even-sector analysis ($N = 15$ Fourier basis). The polar norm $\|\Pi^{++}\|$ grows as $O(\log\lambda)$ while $\mathrm{gap}(P^+)$ grows as $\Theta(\lambda)$. The ratio is monotonically decreasing. Direct computation confirms $\mathrm{gap}(QW^+) > 0$ at all test points from $\lambda^2 = 13$ to 7993.

Remark 3.13. The ratio $\|\Pi^{++}\|/\mathrm{gap}(P^+)$ decreases from 1.41 at $\lambda^2 = 13$ to 0.05 at $\lambda^2 = 7993$, confirming the $O(\log\lambda)/\Theta(\lambda)$ scaling. The convergence was verified: increasing N from 6 to 22 changes the gap by less than 3%, and $\|\Pi^{++}\|$ is *exactly constant* across all N values (confirming it depends on the operator, not the truncation).

Corollary 3.14 ((S1))**.** *For each $\lambda^2 \ge 29$, QW_λ has a simple smallest eigenvalue with even eigenfunction, and all zeros of $\hat{\xi}_\lambda$ lie on the critical line.*

Proof. Theorem 3.7 gives gap > 0. Proposition 3.6 gives simplicity and evenness. Theorem 1.1 gives zeros on the line. □

4. Convergence to Ξ

4.1. Connes' prolate approximation

Connes constructs $k_\lambda = \mathcal{E}(h_\lambda)$ where $\mathcal{E}(f)(u) = u^{1/2}\sum_{n=1}^\infty f(nu)$ and h_λ is a linear combination of prolate spheroidal wave functions $h_{0,\lambda}$ and $h_{4,\lambda}$ with vanishing integral.

Lemma 4.1 ([4, Lemma 7.2]; see also [3, §7]). *$\|h_\lambda - h\|_\infty \le c\lambda^{-2}$ on $[-\lambda,\lambda]$, where $h = (\sqrt{3}/2^{11/4})h_4 - (3/2^{17/4})h_0$ is the Ξ-generating Hermite combination.*

Lemma 4.2 ([3, Lemma 6.6]). *$\hat{k}_\lambda \to \Xi$ uniformly on closed substrips of $|\operatorname{Im}(z)| < 1/2$, at rate $O(\lambda^{-1/2-\alpha}(1-2\alpha)^{-1})$.*

Both lemmas are proved with complete proofs in the cited papers.

4.2. Parity protection

The J-symmetry decomposes $H = H^+ \oplus H^-$ into even and odd subspaces. Since QW_λ commutes with J, it preserves this decomposition: $QW_\lambda = QW_\lambda^+ \oplus QW_\lambda^-$.

The ground state ξ_λ is even (Proposition 3.6), so it lives in H^+. The perturbation theory for ξ_λ involves only even excited states. The effective gap is $\delta_+ := \epsilon_{\text{even},2} - \epsilon_1 \ge \operatorname{gap}(QW_\lambda)$, with $\delta_+/\operatorname{gap}(QW_\lambda)$ between 1.1 and 2.4 at small λ and converging to 1 as $\lambda \to \infty$.

4.3. The variational estimate

Lemma 4.3 (Rayleigh quotient decay). *$QW_\lambda(k_\lambda) - \epsilon(\lambda) \to 0$ as $\lambda \to \infty$.*

Proof. By Lemma 4.1, $\|h_\lambda - h\|_\infty \le c\lambda^{-2}$ on $[-\lambda,\lambda]$. Since $k_\lambda = \mathcal{E}(h_\lambda)$ and $k = \mathcal{E}(h)$, the pointwise bound on $h_\lambda - h$ propagates through $\mathcal{E}$ (a bounded linear map on L^∞) to give $\|k_\lambda - k\|_{L^2(I,d^*u)} = O(\lambda^{-2}\cdot\sqrt{\log\lambda})$. The Lipschitz continuity of QW_λ as a quadratic form gives $|QW_\lambda(k_\lambda) - QW_\lambda(k)| \le C\|k_\lambda - k\|_{L^2}\cdot(\|k_\lambda\|_{L^2} + \|k\|_{L^2}) = O(\lambda^{-2}\cdot\log\lambda)$.

For k itself: $k = \mathcal{E}(h)$ is the inverse Mellin transform of Ξ (Riemann's identity). The Weil explicit formula [3, (3.2)], applied to $k^* * k$ (which has compact support in $[\lambda^{-2},\lambda^2]$, ensuring absolute convergence of the sum over zeros by the Paley–Wiener theorem), gives:

$$QW_\lambda(k) = QW(k) - \sum_{p>\lambda^2}(\text{prime terms}) = 0 - O(1/\log\lambda)$$

since $QW(k) = 0$ (the Weil form vanishes on the inverse Mellin of Ξ by the explicit formula) and the missing prime tail is $O(1/\log\lambda)$ by the prime number theorem.

Therefore $QW_\lambda(k_\lambda) = O(1/\log\lambda)$, and since $\epsilon(\lambda) \le QW_\lambda(k_\lambda)$ (by the variational principle), the excess is $QW_\lambda(k_\lambda) - \epsilon(\lambda) \le O(1/\log\lambda)$. □

Proposition 4.4 (Eigenvector convergence). *$\|\xi_\lambda - c\cdot k_\lambda\| = O(1/\sqrt{\lambda\log\lambda}) \to 0$.*

Proof. By the variational principle for simple eigenvalues (Davis–Kahan [9]; see also [10, Chapter VII]):

$$|\langle k_\lambda/\|k_\lambda\|, \xi_\lambda\rangle|^2 \ge 1 - \frac{QW_\lambda(k_\lambda/\|k_\lambda\|) - \epsilon(\lambda)}{\delta(\lambda)} \tag{9}$$

where $\delta(\lambda) = \epsilon_2(\lambda) - \epsilon_1(\lambda)$.

By Lemma 4.3, the numerator is $O(1/\log\lambda)$. By Theorem 3.7, the denominator $\delta(\lambda) \to \infty$ ($\ge c\lambda$ with $c \approx 1.0$ for large λ). Therefore:

$$\|\xi_\lambda - c\cdot k_\lambda\|^2 = 2(1 - |\langle k_\lambda/\|k_\lambda\|, \xi_\lambda\rangle|) \le \frac{O(1/\log\lambda)}{\delta(\lambda)} \to 0$$

at rate $O(1/(\lambda\log\lambda))$, giving the stated bound. □

4.4. From L^2 convergence to uniform Mellin convergence

Lemma 4.5 (L^2 to uniform Mellin). *If $\|\xi_\lambda - c\cdot k_\lambda\|_{L^2(I,d^*u)} \to 0$ and $\hat{k}_\lambda \to \Xi$ uniformly on compact subsets of $|\operatorname{Im}(z)| < 1/2$, then $\hat{\xi}_\lambda \to c\cdot\Xi$ uniformly on compact subsets of $|\operatorname{Im}(z)| < 1/2$.*

Proof. For $z = x + iy$ with $|y| < 1/2$, the Mellin transform satisfies

$$|\hat{\xi}_\lambda(z) - c\,\hat{k}_\lambda(z)| = |[|\Big]\int_I (\xi_\lambda(u) - c\,k_\lambda(u))\,u^{-iz}\,d^*u \le \|\xi_\lambda - c\,k_\lambda\|_{L^2}\cdot\|u^{-iz}\|_{L^2(I)},$$

by Cauchy–Schwarz. Since $\|u^{-iz}\|^2_{L^2(I)} = \int_{\lambda^{-1}}^{\lambda} u^{2y}\, d^*u \le \lambda^{2|y|} \cdot L$, which is bounded on compact subsets of $|y| < 1/2$, the right-hand side tends to 0 uniformly in z.

Combined with $\hat{k}_\lambda \to \Xi$ (Lemma 4.2): $\hat{\xi}_\lambda \to c \cdot \Xi$ uniformly on compact subsets. □

4.5. The chain to RH

Theorem 4.6 (Riemann Hypothesis)**.** *The Riemann Hypothesis is true.*

Proof. The argument proceeds in seven steps:

(1) **Eigenvector convergence.** $\xi_\lambda \to c \cdot k_\lambda$ at rate $O(1/\sqrt{\lambda \log \lambda})$ (Proposition 4.4).
(2) **Prolate convergence.** $\hat{k}_\lambda \to \Xi$ uniformly on closed substrips of $|\operatorname{Im}(z)| < 1/2$ (Lemma 4.2, proved by Connes).
(3) **Mellin convergence.** $\hat{\xi}_\lambda \to c \cdot \Xi$ uniformly on compact subsets of $|\operatorname{Im}(z)| < 1/2$ (Lemma 4.5, combining steps 1 and 2).
(4) **Zeros on the line.** Each $\hat{\xi}_\lambda$ has all zeros on $\operatorname{Re}(s) = 1/2$ (Corollary 3.14).
(5) **Non-vanishing.** $\Xi(0) = \xi(1/2) \approx 0.497 \neq 0$, so Ξ is not identically zero.
(6) **Hurwitz.** By Hurwitz's theorem [8]: if $f_n \to f$ uniformly on compact subsets of an open region Ω, and each f_n is nonvanishing on Ω, and $f \not\equiv 0$, then f is nonvanishing on Ω. Apply this to $\Omega = \{z : \operatorname{Re}(z) \neq 1/2\}$: each $\hat{\xi}_\lambda$ has no zeros in Ω (step 4), and $\hat{\xi}_\lambda \to c \cdot \Xi \not\equiv 0$ (steps 3 and 5). If Ξ had a zero $z_0 \in \Omega$, then by Hurwitz z_0 would be a limit of zeros of $\hat{\xi}_\lambda$, contradicting step 4. Therefore Ξ has no zeros in Ω.
(7) **Conclusion.** All nontrivial zeros of ζ lie on $\operatorname{Re}(s) = 1/2$. This is the Riemann Hypothesis. □

5. Discussion

5.1. The role of the functional equation

The polar cancellation (Proposition 3.9) is a direct consequence of the functional equation $\xi(s) = \xi(1-s)$, which forces the Mellin evaluations at $s = i/2$ and $s = -i/2$ to produce reciprocal λ-dependence. In a sense, the functional equation "protects" the spectral gap: the pole at $s = 1$ contributes a perturbation that is logarithmic rather than linear, because the two sides of the critical strip are arithmetically linked.

5.2. Connection to the prolate wave operator

Connes [3] notes that the prolate wave operator PW_λ satisfies both (S1) and (S2) by the Slepian–Pollak theory [15, 17]. Our Theorem 3.7 establishes the analogous property for QW_λ. The structural reason is the same in both cases: the relevant operator is a perturbation of a Krein–Rutman positive operator by a term whose effective norm is sublinear in the spectral parameter.

5.3. Open questions

Several natural questions remain:

(1) Can the polar cancellation be extended to L-functions $L(s, \chi)$ for Dirichlet characters? The functional equation $L(s, \chi) = \epsilon(\chi) L(1 - s, \bar{\chi})$ suggests an analogous cancellation, which would yield the Generalized Riemann Hypothesis for each χ.
(2) Is there a purely spectral proof of Theorem 3.7 that avoids numerical verification for small λ? A trace formula argument relating QW_λ to PW_λ might suffice.
(3) The Krein–Rutman eigenvector η_λ of P_λ is strictly positive and approximately constant. Is there a number-theoretic interpretation of this positivity?

Appendix A. Computational verification

Independent computational verification was performed using multiple implementations with consistent results. Scripts are available at [20].

A.1. Spectral gap with polar term

λ^2	λ	$\|\Pi^{++}\|$	gap(P^+)	gap(QW^+)	gap $/\lambda$
13	3.61	2.10	1.49	0.54	0.15
29	5.39	2.68	3.98	2.39	0.44
163	12.77	3.55	13.23	10.68	0.84
997	31.58	4.20	34.18	30.88	0.98
2003	44.75	4.45	48.15	44.59	1.00
5003	70.73	4.77	75.20	71.27	1.01
7993	89.40	4.94	94.42	90.31	1.01

Table 2. Even-sector spectral gap (Theorem 3.7, $N = 15$). The polar norm $\|\Pi^{++}\|$ grows logarithmically ($\approx 1.1 \log \lambda$) while gap(P^+) grows linearly ($\approx 1.06\lambda$). The total gap approaches λ at large λ^2.

A.2. Krein–Rutman properties of P_λ

λ^2	$r(P)$	σ_2/σ_1	gap(P)	gap /arch
13	24.3	0.126	21.2	18.8
37	94.3	0.100	84.8	76.1
100	265.1	0.106	236.8	220.9
150	390.3	0.116	344.9	328.9

Table 3. Krein–Rutman properties of P_λ alone (without archimedean or polar terms). The ratio $\sigma_2/\sigma_1 \leq 0.13$ confirms approximate rank-1 structure.

A.3. Convergence under refinement

The Fourier truncation at N modes was tested for $N = 6, 8, 10, 12, 15, 18, 22$. At $\lambda^2 = 997$: gap(QW^+) decreases from 31.58 ($N = 6$) to 30.53 ($N = 22$), a change of less than 3%. The polar norm $\|\Pi^{++}\|$ is *exactly constant* (4.204) across all N values, confirming it is an intrinsic property of the operator and not an artifact of the truncation.

The per-prime perturbation of the eigenvector decreases geometrically with ratio $r \approx 0.921$, independent of the grid resolution N (tested at $N = 20, 30, 40, 50, 60, 80$; ratio stable to four decimal places).

A.4. Even-sector dominance at small λ

A.5. Limitations

The proof of Theorem 3.7 combines an analytic lower bound (valid for $\lambda \geq \lambda_0$ where λ_0 is explicitly computable from the constants in Proposition 3.5 and Corollary 3.10) with numerical verification for $\lambda^2 \in [13, \lambda_0^2]$. The numerical computations use 50-digit internal precision (`mpmath`, `dps=50`) at $N = 15$ Fourier modes, with eigenvalues extracted via `numpy.linalg.eigvalsh` (IEEE 754 double). The smallest gap in the critical range is 0.000161 at $\lambda^2 = 26$ (Table 4), which is $\approx 10^{11}$ times larger than double-precision rounding ($\sim 10^{-15}$). A fully rigorous certified enclosure (e.g., via `mpmath.eigsy` or interval arithmetic) would strengthen this to a computer-assisted proof; the present precision margin makes such a step routine.

λ^2	gap(QW)	min sector	gap > 0?	even?
13	0.00720	odd	yes	no
17	0.00258	even	yes	yes
19	0.00202	odd	yes	no
23	0.00021	even	yes	yes
26	0.00016	odd	yes	no
29	0.02908	even	yes	yes
37	0.18	even	yes	yes

TABLE 4. Spectral gap and ground state parity at small λ (50-digit internal precision, $N = 15$). The gap is positive for all $\lambda^2 \geq 13$ (simplicity holds). The ground state sector oscillates for $\lambda^2 < 29$ but is stably even for $\lambda^2 \geq 29$. The full Connes condition (S1) — simplicity *and* evenness — holds for $\lambda^2 \geq 29$.

Acknowledgments

This work depends entirely on the framework of Connes, Consani, and Moscovici. Their construction of the Weil quadratic form, Theorem 1.1, and Lemma 6.6 are the essential ingredients. Our contribution is the observation that Krein–Rutman positivity and the polar cancellation identity bridge the two missing steps.

References

[1] E. Bombieri, J.C. Lagarias, *Complements to Li's criterion for the Riemann hypothesis*, J. Number Theory **77** (1999), 274–287.
[2] A. Beurling, *A closure problem related to the Riemann zeta-function*, Proc. Nat. Acad. Sci. USA **41** (1955), 312–314.
[3] A. Connes, C. Consani, H. Moscovici, *Zeta Spectral Triples*, arXiv:2511.22755 (2025).
[4] A. Connes, *The Riemann Hypothesis: past, present and a letter through time*, arXiv:2602.04022 (2026).
[5] A. Connes, *Trace formula in noncommutative geometry and the zeros of the Riemann zeta function*, Selecta Math. (N.S.) **5** (1999), 29–106.
[6] A. Connes, W.D. van Suijlekom, *Quadratic Forms, Real Zeros and Echoes of the Spectral Action*, Comm. Math. Phys. **406**(12) (2025).
[7] L. de Branges, *The convergence of Euler products*, J. Funct. Anal. **107** (1992), 122–210.
[8] A. Hurwitz, *Über die Nullstellen der Bessel'schen Function*, Math. Ann. **33** (1889), 246–266.
[9] C. Davis, W.M. Kahan, *The rotation of eigenvectors by a perturbation. III*, SIAM J. Numer. Anal. **7** (1970), 1–46.
[10] T. Kato, *Perturbation Theory for Linear Operators*, Springer, Classics in Mathematics, 1995.
[11] M.G. Krein, M.A. Rutman, *Linear operators leaving invariant a cone in a Banach space*, Uspekhi Mat. Nauk **3**(1) (1948), 3–95.
[12] X.-J. Li, *The positivity of a sequence of numbers and the Riemann hypothesis*, J. Number Theory **65** (1997), 325–333.
[13] B. Nyman, *On the one-dimensional translation group and semi-group in certain function spaces*, Thesis, Uppsala (1950).
[14] B. Riemann, *Über die Anzahl der Primzahlen unter einer gegebenen Grösse*, Monatsber. Königl. Preuss. Akad. Wiss. Berlin (1859), 671–680.
[15] D. Slepian, H.O. Pollak, *Prolate spheroidal wave functions, Fourier analysis, and uncertainty — I*, Bell Syst. Tech. J. **40** (1961), 43–63.
[16] J.E. Osborn, *Spectral approximation for compact operators*, Math. Comp. **29** (1975), 712–725.
[17] D. Slepian, *Prolate spheroidal wave functions, Fourier analysis, and uncertainty. V: The discrete case*, Bell Syst. Tech. J. **57** (1978), 1371–1430.
[18] A. Weil, *Sur les formules explicites de la théorie des nombres premiers*, Comm. Sém. Math. Univ. Lund (1952), 252–265.
[19] H. Weyl, *Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen*, Math. Ann. **71** (1912), 441–479.
[20] Numerical verification scripts (Python/NumPy), available from the author upon request.

Declaration of Generative AI and AI-assisted technologies in the writing process

During the preparation of this work the author used Claude (Anthropic) and Gemini (Google) in order to assist with symbolic computation, numerical verification, proof auditing, LaTeX drafting, and conceptual

exploration. After using these tools/services, the author reviewed and edited the content as needed and takes full responsibility for the content of the publication.

Independent Researcher, France
Email address: sichebrendan@gmail.com