

Coordinate Geometry Revisited

A Matrix approach

Raghav Girgaonkar Sagar Jain

Indian Institute of Technology Hyderabad

February 2019

Problem

The straight line $2x - 3y = 1$ divides the circular region $x^2 + y^2 \leq 6$ into two parts. If

$$S = \left\{ \left(2, \frac{3}{4} \right), \left(\frac{5}{2}, \frac{3}{4} \right), \left(\frac{1}{4}, -\frac{1}{4} \right), \left(\frac{1}{8}, \frac{1}{4} \right) \right\}$$

then find the number of points in S lying inside the smaller part of the circle.

Approach

The steps we took to solve this problem were:

- Find the centre of the given circle(in this case it is $(0,0)$)

Approach

The steps we took to solve this problem were:

- Find the centre of the given circle(in this case it is $(0,0)$)
- Find position of centre with respect to the line given

Approach

The steps we took to solve this problem were:

- Find the centre of the given circle(in this case it is $(0,0)$)
- Find position of centre with respect to the line given
- Find position of each point in **S** with respect to the line

Approach

The steps we took to solve this problem were:

- Find the centre of the given circle(in this case it is $(0,0)$)
- Find position of centre with respect to the line given
- Find position of each point in **S** with respect to the line
- Find the points which satisfy our condition using the constraints found in above steps

Matrix Transformation

- Consider a matrix \mathbf{S}_o with all the points in the set S stacked as follows:

$$\mathbf{S}_o = \begin{bmatrix} 2 & \frac{5}{2} & \frac{1}{4} & \frac{1}{8} \\ \frac{3}{4} & \frac{3}{4} & -\frac{1}{4} & \frac{1}{4} \end{bmatrix}$$

Matrix Transformation

- Consider a matrix \mathbf{S}_o with all the points in the set S stacked as follows:

$$\mathbf{S}_o = \begin{bmatrix} 2 & \frac{5}{2} & \frac{1}{4} & \frac{1}{8} \\ \frac{3}{4} & \frac{3}{4} & -\frac{1}{4} & \frac{1}{4} \end{bmatrix}$$

- Now the normal vector to the given line is as follows:

$$\mathbf{N} = [2 \quad -3]$$

Matrix Transformation

- Consider a matrix \mathbf{S}_o with all the points in the set S stacked as follows:

$$\mathbf{S}_o = \begin{bmatrix} 2 & \frac{5}{2} & \frac{1}{4} & \frac{1}{8} \\ \frac{3}{4} & \frac{3}{4} & -\frac{1}{4} & \frac{1}{4} \end{bmatrix}$$

- Now the normal vector to the given line is as follows:

$$\mathbf{N} = [2 \quad -3]$$

- The matrix

$$\mathbf{P} = [1 \quad 1 \quad 1 \quad 1]$$

Matrix Transformation

- Consider a matrix \mathbf{S}_o with all the points in the set S stacked as follows:

$$\mathbf{S}_o = \begin{bmatrix} 2 & \frac{5}{2} & \frac{1}{4} & \frac{1}{8} \\ \frac{3}{4} & \frac{3}{4} & -\frac{1}{4} & \frac{1}{4} \end{bmatrix}$$

- Now the normal vector to the given line is as follows:

$$\mathbf{N} = [2 \quad -3]$$

- The matrix

$$\mathbf{P} = [1 \quad 1 \quad 1 \quad 1]$$

- and the matrix

$$\mathbf{O} = [0 \quad 0]$$

Matrix Transformation

- Consider a matrix S_o with all the points in the set S stacked as follows:

$$S_o = \begin{bmatrix} 2 & \frac{5}{2} & \frac{1}{4} & \frac{1}{8} \\ \frac{3}{4} & \frac{3}{4} & -\frac{1}{4} & \frac{1}{4} \end{bmatrix}$$

- Now the normal vector to the given line is as follows:

$$N = [2 \quad -3]$$

- The matrix

$$P = [1 \quad 1 \quad 1 \quad 1]$$

- and the matrix

$$O = [0 \quad 0]$$

- The equation of a line is $N * x = p$ where x is a 2 dimensional column vector and p is a constant.

Matrix Transformation Continued

Our original problem now transforms as follows:

- The equation of our line is $\mathbf{N} * \mathbf{x} = 1$

Matrix Transformation Continued

Our original problem now transforms as follows:

- The equation of our line is $\mathbf{N} * \mathbf{x} = 1$
- We can see that $\mathbf{N} * \mathbf{O}^T - 1 = -1$ this is the relative position of the centre with respect to the line.

Matrix Transformation Continued

Our original problem now transforms as follows:

- The equation of our line is $\mathbf{N} * \mathbf{x} = 1$
- We can see that $\mathbf{N} * \mathbf{O}^T - 1 = -1$ this is the relative position of the centre with respect to the line.
- Let $\mathbf{M} = \mathbf{N} * \mathbf{S}_o$

Matrix Transformation Continued

Our original problem now transforms as follows:

- The equation of our line is $\mathbf{N} * \mathbf{x} = 1$
- We can see that $\mathbf{N} * \mathbf{O}^T - 1 = -1$ this is the relative position of the centre with respect to the line.
- Let $\mathbf{M} = \mathbf{N} * \mathbf{S}_o$
- Find the number of elements in $\mathbf{M} - \mathbf{P} > 0$ and $(\mathbf{S}_o^T * \mathbf{S}_o)_{ii} \leq 6$

Solution

- $\mathbf{S}_o^T * \mathbf{S}_o$ will give us a $4*4$ matrix with the diagonal elements being the values of $x^2 + y^2$ for the 4 points, hence we will consider only the diagonal entries.

Solution

- $\mathbf{S}_o^T * \mathbf{S}_o$ will give us a 4×4 matrix with the diagonal elements being the values of $x^2 + y^2$ for the 4 points, hence we will consider only the diagonal entries.
- Calculating $\mathbf{M} - \mathbf{P}$ we get:

$$\mathbf{M} - \mathbf{P} = \begin{bmatrix} \frac{3}{4} & \frac{7}{4} & \frac{1}{4} & -\frac{3}{2} \end{bmatrix}$$

Solution

- $\mathbf{S}_o^T * \mathbf{S}_o$ will give us a 4×4 matrix with the diagonal elements being the values of $x^2 + y^2$ for the 4 points, hence we will consider only the diagonal entries.
- Calculating $\mathbf{M} - \mathbf{P}$ we get:

$$\mathbf{M} - \mathbf{P} = \begin{bmatrix} \frac{3}{4} & \frac{7}{4} & \frac{1}{4} & -\frac{3}{2} \end{bmatrix}$$

- The diagonal elements of $\mathbf{S}_o^T * \mathbf{S}_o$ are:

$$(\mathbf{S}_o^T * \mathbf{S}_o)_{ii} = \begin{bmatrix} \frac{73}{16} & \frac{109}{16} & \frac{1}{8} & \frac{1}{32} \end{bmatrix}$$

Solution

- $\mathbf{S}_o^T * \mathbf{S}_o$ will give us a 4×4 matrix with the diagonal elements being the values of $x^2 + y^2$ for the 4 points, hence we will consider only the diagonal entries.
- Calculating $\mathbf{M} - \mathbf{P}$ we get:

$$\mathbf{M} - \mathbf{P} = \begin{bmatrix} \frac{3}{4} & \frac{7}{4} & \frac{1}{4} & -\frac{3}{2} \end{bmatrix}$$

- The diagonal elements of $\mathbf{S}_o^T * \mathbf{S}_o$ are:

$$(\mathbf{S}_o^T * \mathbf{S}_o)_{ii} = \begin{bmatrix} \frac{73}{16} & \frac{109}{16} & \frac{1}{8} & \frac{1}{32} \end{bmatrix}$$

- Now the only points which satisfy the conditions $\mathbf{M} - \mathbf{P} > 0$ and $(\mathbf{S}_o^T * \mathbf{S}_o)_{ii} \leq 6$ are $\left(2, \frac{3}{4}\right)$ and $\left(\frac{1}{4}, -\frac{1}{4}\right)$

Solution

- $\mathbf{S}_o^T * \mathbf{S}_o$ will give us a 4×4 matrix with the diagonal elements being the values of $x^2 + y^2$ for the 4 points, hence we will consider only the diagonal entries.
- Calculating $\mathbf{M} - \mathbf{P}$ we get:

$$\mathbf{M} - \mathbf{P} = \begin{bmatrix} \frac{3}{4} & \frac{7}{4} & \frac{1}{4} & -\frac{3}{2} \end{bmatrix}$$

- The diagonal elements of $\mathbf{S}_o^T * \mathbf{S}_o$ are:

$$(\mathbf{S}_o^T * \mathbf{S}_o)_{ii} = \begin{bmatrix} \frac{73}{16} & \frac{109}{16} & \frac{1}{8} & \frac{1}{32} \end{bmatrix}$$

- Now the only points which satisfy the conditions $\mathbf{M} - \mathbf{P} > 0$ and $(\mathbf{S}_o^T * \mathbf{S}_o)_{ii} \leq 6$ are $\left(2, \frac{3}{4}\right)$ and $\left(\frac{1}{4}, -\frac{1}{4}\right)$
- Hence the number of points lying in the smaller region of the circle is 2.

Figure

