# Coordinate Geometry Revisited A Matrix approach

Raghav Girgaonkar Sagar Jain

Indian Institute of Technology Hyderabad

February 2019

#### **Problem**

The straight line 2x - 3y = 1 divides the circular region  $x^2 + y^2 \le 6$  into two parts. If

$$S = \left\{ \left(2, \frac{3}{4}\right), \left(\frac{5}{2}, \frac{3}{4}\right), \left(\frac{1}{4}, -\frac{1}{4}\right), \left(\frac{1}{8}, \frac{1}{4}\right) \right\}$$

then find the number of points in S lying inside the smaller part of the circle.

The steps we took to solve this problem were:

► Find the centre of the given circle(in this case it is (0,0))

The steps we took to solve this problem were:

- ► Find the centre of the given circle(in this case it is (0,0))
- Find position of centre with respect to the line given

The steps we took to solve this problem were:

- ► Find the centre of the given circle(in this case it is (0,0))
- Find position of centre with respect to the line given
- Find position of each point in *S* with respect to the line

The steps we took to solve this problem were:

- ► Find the centre of the given circle(in this case it is (0,0))
- ► Find position of centre with respect to the line given
- ▶ Find position of each point in *S* with respect to the line
- ► Find the points which satisfy our condition using the constraints found in above steps

#### Matrix Transformation

Consider a matrix  $S_o$  with all the points in the set S stacked as follows:

$$S_o = \begin{bmatrix} 2 & \frac{5}{2} & \frac{1}{4} & \frac{1}{8} \\ & & & \\ \frac{3}{4} & \frac{3}{4} & -\frac{1}{4} & \frac{1}{4} \end{bmatrix}$$

Now the normal vector to the given line is as follows:

$$N = \begin{bmatrix} 2 & -3 \end{bmatrix}$$

The matrix

$$P = \begin{bmatrix} 1 & 1 & 1 & 1 \end{bmatrix}$$

and the matrix

$$O = \begin{bmatrix} 0 & 0 \end{bmatrix}$$

#### Matrix Transformation Contd

Our original problem now transforms as follows:

 $\blacktriangleright \text{ Let } M = N * S_o$ 

#### Matrix Transformation Contd

Our original problem now transforms as follows:

- ightharpoonup Let  $M = N * S_o$
- ▶ We can see that  $N * O^T 1 = -1$  this is the relative position of the centre with respect to the line.

#### Matrix Transformation Contd

Our original problem now transforms as follows:

- ightharpoonup Let  $M = N * S_o$
- ▶ We can see that  $N * O^T 1 = -1$  this is the relative position of the centre with respect to the line.
- ► Find the number of elements in M P > 0 and  $(S_o^T * S_o)_{ii} \leq 6$

▶  $S_o^T * S_o$  will give us a 4\*4 matrix with the diagonal elements being the values of  $x^2 + y^2$  for the 4 points, hence we will consider only the diagonal entries.

- ▶  $S_o^T * S_o$  will give us a 4\*4 matrix with the diagonal elements being the values of  $x^2 + y^2$  for the 4 points, hence we will consider only the diagonal entries.
- ▶ Calculating M P we get:

$$M - P = \begin{bmatrix} \frac{3}{4} & \frac{7}{4} & \frac{1}{4} & -\frac{3}{2} \end{bmatrix}$$

- ▶  $S_o^T * S_o$  will give us a 4\*4 matrix with the diagonal elements being the values of  $x^2 + y^2$  for the 4 points, hence we will consider only the diagonal entries.
- ► Calculating M P we get:

$$M - P = \begin{bmatrix} \frac{3}{4} & \frac{7}{4} & \frac{1}{4} & -\frac{3}{2} \end{bmatrix}$$

▶ The diagonal elements of  $S_o^T * S_o$  are:

$$(S_o^T * S_o)_{ii} = \begin{bmatrix} \frac{73}{16} & \frac{109}{16} & \frac{1}{8} & \frac{1}{32} \end{bmatrix}$$



- ▶  $S_o^T * S_o$  will give us a 4\*4 matrix with the diagonal elements being the values of  $x^2 + y^2$  for the 4 points, hence we will consider only the diagonal entries.
- ► Calculating M P we get:

$$M-P = \begin{bmatrix} \frac{3}{4} & \frac{7}{4} & \frac{1}{4} & -\frac{3}{2} \end{bmatrix}$$

► The diagonal elements of  $S_o^T * S_o$  are:

$$(S_o^T * S_o)_{ii} = \begin{bmatrix} \frac{73}{16} & \frac{109}{16} & \frac{1}{8} & \frac{1}{32} \end{bmatrix}$$

Now the only points which satisfy the conditions M-P>0 and  $(S_o^T*S_o)_{ii}\leqslant 6$  are  $\left(2,\frac{3}{4}\right)$  and  $\left(\frac{1}{4},-\frac{1}{4}\right)$ 



- ▶  $S_o^T * S_o$  will give us a 4\*4 matrix with the diagonal elements being the values of  $x^2 + y^2$  for the 4 points, hence we will consider only the diagonal entries.
- ► Calculating M P we get:

$$M - P = \begin{bmatrix} \frac{3}{4} & \frac{7}{4} & \frac{1}{4} & -\frac{3}{2} \end{bmatrix}$$

► The diagonal elements of  $S_o^T * S_o$  are:

$$(S_o^T * S_o)_{ii} = \begin{bmatrix} \frac{73}{16} & \frac{109}{16} & \frac{1}{8} & \frac{1}{32} \end{bmatrix}$$

- Now the only points which satisfy the conditions M-P>0 and  $(S_o^T*S_o)_{ii}\leqslant 6$  are  $\left(2,\frac{3}{4}\right)$  and  $\left(\frac{1}{4},-\frac{1}{4}\right)$
- ▶ Hence the number of points lying in the smaller region of the circle is 2.



## Figure




Figure: Shaded part denotes the smaller side of the circle.