

sampbias, a method for quantifying geographic sampling biases in species distribution data

Alexander Zizka^{1,2}, Alexandre Antonelli^{3,4,5}, Daniele Silvestro^{3,4,6}

1. German Centre for Integrative Biodiversity Research Halle-Jena-Leipzig (iDiv), University of Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany
2. Naturalis Biodiversity Center, Leiden University, Leiden, Darwinweg 2, 2333 CR Leiden The Netherlands
3. Gothenburg Global Biodiversity Centre, University of Gothenburg, Box 461, 405 30 Gothenburg, Sweden
4. Department for Biological and Environmental Sciences, University of Gothenburg, Box 461, 405 30 Gothenburg, Sweden
5. Royal Botanic Gardens Kew, TW9 3AE, Richmond, Surrey, United Kingdom
6. Department of Biology, University of Fribourg, Ch. du Musée 10, 1700 Fribourg, Switzerland

Abstract

Geo-referenced species occurrences from public databases have become essential to biodiversity research and conservation. However, geographical biases are widely recognized as a factor limiting the usefulness of such data for understanding species diversity and distribution. In particular, differences in sampling intensity across a landscape due to differences in human accessibility are ubiquitous but may differ in strength among taxonomic groups and datasets. Although several factors have been described to influence human access (such as presence of roads, rivers, airports and cities), quantifying their specific and combined effects on recorded occurrence data remains challenging. Here we present *sambias*, an algorithm and software for quantifying the effect of accessibility biases in species occurrence datasets. *Sambias* uses a Bayesian approach to estimate how sampling rates vary as a function of proximity to one or multiple bias factors. The results are comparable among bias factors and datasets. We demonstrate the use of *sambias* on a dataset of mammal occurrences from the island of Borneo, showing a high biasing effect of cities and a moderate effect of roads and airports. *Sambias* is implemented as a well-documented, open-access and user-friendly R package that we hope will become a standard tool for anyone working with species occurrences in ecology, evolution, conservation and related fields.

Keywords

Collection effort, Global Biodiversity Information Facility (GBIF), Presence only data, Roadside bias, Sampling intensity

Background

Publicly available datasets of geo-referenced species occurrences, such as provided by the Global Biodiversity Information Facility (www.gbif.org) have become a fundamental resource in biological sciences, especially in biogeography, conservation, and macroecology. However, these datasets are typically not collected systematically and rarely include information on collection effort. Instead, they are often compiled from a variety of sources (e.g. scientific expeditions, census counts, genetic barcoding studies, and citizen-science observations). Species occurrences are therefore often subject to multiple sampling biases (Meyer et al. 2016).

Sampling biases that may affect the recording of species occurrences (presence, absence and abundance, Isaac and Pocock 2015, Boakes et al. 2010) include the under-sampling of specific taxa (“taxonomic bias”, e.g., birds *vs.* nematodes), specific geographic regions (“geographic bias”, i.e. easily accessible *vs.* remote areas), and specific temporal periods (“temporal bias”, i.e. wet season *vs.* dry season). In particular geographic sampling bias—the fact that sampling effort is spatially biased, rather than equally distributed over the study area—is likely to be widespread in all non-systematically collected datasets of species distributions.

Many aspects can lead to sampling biases, including socio-economic factors (i.e. national research spending, history of scientific research; Zizka et al. 2020, Meyer et al. 2015, Daru et al. 2018), political factors (armed conflict, democratic rights; Rydén et al. 2020), and physical accessibility (i.e. distance to a road or river, terrain conditions, slope; Yang et al. 2014, Botts et al. 2011). Especially physical accessibility by people is omnipresent as a bias factor (e.g. Lin et al. 2015, Kadmon et al. 2004, Engemann et al. 2015), across spatial scales,

as the commonly used term “roadside bias” testifies. In practice, this means that most species observations are made in or near cities, along roads, paths, and rivers, and near human settlements. Relatively fewer observations are expected to be available from inaccessible areas in e.g. a tropical rainforest or a mountain top. Since the recording of different taxonomic groups poses different challenges, geographic sampling bias and the effect of accessibility may differ among taxonomic groups (Vale and Jenkins 2012).

The implications of not considering geographic sampling biases in biodiversity research are likely substantial (Barbosa et al. 2013, Yang et al. 2013, Meyer et al. 2016). The presence of geographic sampling biases is broadly recognized (e.g. Kadmon et al. 2004), and approaches exist to account for it in some analyses—such as species-richness estimates (Engemann et al. 2015), occupancy models (Kery and Royle 2016), and abundance estimates (Shimadzu and Darnell 2015). In the case of species distribution modelling—the statistical estimation of species geographic distributions based on known occurrences and environmental conditions—geographically biased sampling is problematic because it often causes environmentally biased sampling which decreases model performance (Kadmon et al. 2004, Lobo and Tognelli 2011, Bystriakova et al. 2012, Kramer-Schadt et al. 2013, Varela et al. 2014). Many approaches exist to remedy the effect of biased sampling on species distribution models (Fourcade et al. 2014), including rarefaction to reduce clumped sampling in geographic (Beck et al. 2014, Boria et al. 2014, Aiello-Lammens et al. 2015) or environmental space (Varela et al. 2014), collecting background points for presence-only models to reflect the same sampling bias as the presence records (Phillips et al. 2009), and explicitly modelling sampling bias (Fithian et al. 2015, Stolar and Nielsen 2015, Komori et al. 2020). In contrast, few attempts have been

made to compare the geographic sampling bias among datasets (Fernández and Nakamura 2015, Ruete 2015, Monsarrat et al. 2019) and to our knowledge, no tools exist to quantify the effect size of specific bias factors and compare it among them. We define as *bias factors* any anthropogenic or natural features that facilitate human access and sampling, such as roads, rivers, airports, and cities.

It is unrealistic to expect that accessibility bias in biodiversity data will ever disappear even after more automated observation technologies are developed. It is therefore crucial that researchers realise the intrinsic biases associated with the data they deal with, especially in cross-taxonomic studies, since occurrence datasets from different taxa are likely differently affected by sampling biases due to differences in specimen collection and transportation. This is the first step towards estimating to which extent these biases may affect their analyses, results, and conclusions. Any study dealing with species occurrence data should arguably assess the strength of accessibility biases in the underlying data. Such a quantification can also help researchers to target further sampling efforts.

Here, we present *sambias* 1, a probabilistic method to quantify accessibility bias in datasets of species occurrences. *Sambias* is implemented as a user-friendly R-package and uses a Bayesian approach to address three questions:

- 1) How strong is the accessibility bias in a given dataset?
- 2) How strong is the effect of different bias factors in causing the overall accessibility bias?
- 3) How is accessibility bias distributed in space, i.e. which areas are a priority for targeted sampling?

Sambias is implemented in R (R Core Team 2019), based on commonly used packages for data handling (*ggplot*, Wickham 2009, *forcats*, 2019, *tidyr*, Wickham and Henry 2019, *dplyr*, Wickham et al. 2019, *magrittr*, Bache and Wickham 2014, *viridis*, Garnier 2018), handling geographic information and geo-computation (*raster*, Hijmans 2019, *sp*, Pebesma and Bivand 2005, Bivand et al. 2013) and statistical modelling (*stats*, R Core Team 2019). *Sambias* offers an easy and largely automated means for biodiversity scientists and non-specialists alike to explore bias in species occurrence data, in a way that is comparable across datasets. The results may be used to identify priorities for further collection or digitalization efforts, assess the reliability of scientific results based on publicly available species distribution data.

Methods and Features

General concept

Under the assumption that organisms exist across the entire area of interest, we can expect the number of sampled occurrences in a restricted area, such as a single biome, to be distributed uniformly in space (even though, of course, the density of individuals and the species diversity may be heterogeneous). With *sambias* we assess to which extent variation in sampling rates can be explained by distance from bias factors.

Sambias works at a user-defined spatial scale, and any dataset of multi-species occurrence records can be tested against any geographic gazetteer. Reliability increases with increasing dataset size. Default global gazetteers for airports, cities, rivers and roads are provided

with *sambias*, and user-defined gazetteers can be added easily. Species occurrence data as downloaded from the data portal of GBIF can be directly used as input data for *sambias*. The output of the package includes measures of the sampling rates across space, which are comparable between different gazetteers (e.g. comparing the biasing effect of roads and rivers), different taxa (e.g. birds *vs.* flowering plants) and different data sets (e.g. specimens *vs.* human observations).

Distance calculation

Sambias uses gazetteers of the geographic location of bias factors (hereafter indicated with B) to generate a regular grid across the study area. By default the study area is defined by the geographic extent of the study dataset, but it can also be customized via user-defined polygons, for instance to limit the analyses to an environmentally homogeneous region (e.g. rainforest) or an area of special interest (e.g. a national park). In this case all occurrence records outside the user-defined area will be disregarded for the analysis.. For each grid cell i , we then compute a vector $X_i(j)$ of minimum distances (straight aerial distance, “as the crow flies”) to each bias factor $j \in B$. The resolution of the grid defines the precision of the distance estimates, for instance a 1x1 degree raster will yield approximately a 110 km precision at the equator. Due to the assumption of homogeneous sampling and a computational trade-off between the resolution of the regular grid and the extent of the study area (for instance, a 1 second resolution for a global dataset would become computationally prohibitive in most practical cases), *sambias* is best suited for local or regional datasets at high resolution (c. 100 – 10,000 m). Since the differences in grid cell size are negligible on the local and regional scale,

sambias uses a latitude/longitude grid by default, but a custom grid in any projection and coordinate reference system—for instance an equal area grid—may be provided by the user.

Quantifying accessibility bias using a Bayesian framework

We describe the observed number of sampled occurrences S_i within each cell i as the result of a Poisson sampling process with rate λ_i . We model the rate λ_i as a function of a parameter q , which represents the expected number of occurrences per cell in the absence of biases, i.e. when $\sum_{j=1}^B X_i(j) = 0$. Additionally, we model λ_i to decrease exponentially as a function of distance from bias factors, such that increasing distances will result in a lower sampling rate. For a single bias factor the rates of cell i with distance X_i from a bias is:

$$\lambda_i = q \times \exp(-wX_i)$$

where $w \in \mathbb{R}^+$ defines the steepness of the Poisson rate decline, such that $w \approx 0$ results in a null model of uniform sampling rate q across cells. In the presence of multiple bias factors (e.g. roads and rivers), the sampling rate decrease is a function of the cumulative effects of each bias and its distance from the cell:

$$\lambda_i = q \times \exp\left(-\sum_{j=1}^B w_j X_i(j)\right) \quad (1)$$

where a vector $\mathbf{w} = [w_1, \dots, w_B]$ describes the amount of bias attributed to each specific factor.

To quantify the amount of bias associated with each factor, we jointly estimate the parameters q and \mathbf{w} in a Bayesian framework. We use Markov Chain Monte Carlo (MCMC) to sample

these parameters from their posterior distribution:

$$P(q, \mathbf{w} | \mathbf{S}) \propto \prod_{i=1}^N Poi(S_i | \lambda_i) \times P(q) P(\mathbf{w}) \quad (2)$$

where the likelihood of sampled occurrences S_i within each cell $Poi(S_i | \lambda_i)$ is the probability mass function of a Poisson distribution with rate per cell defined as in Eqn. (1). The likelihood is then multiplied across the N cells considered. We used exponential priors on the parameters q and \mathbf{w} , $P(q) \sim \Gamma(1, 0.01)$ and $P(\mathbf{w}) \sim \Gamma(1, 1)$, respectively. We chose exponential priors because they represent the standard choice for rate parameters such as q and the weights \mathbf{w} , all of which must be positive and have support $[0, +\infty]$. We designed the priors to be informative (i.e. not allowing negative values) and yet vague enough to encompass a much wider range of parameter space than the range of values observed in empirical tests. For instance, the posterior weights estimated in our empirical study ranged from 0 to 0.013, while the prior distribution applied on these parameters, $\text{Exp}(1)$ has a 95% of its density in range 0 – 3. Custom priors, within the flexible family of gamma distributions, which include the exponential priors used here, are possible via the `prior_q` and `prior_w` arguments of the `calculate_bias` function.

We summarize the parameters by computing the mean of the posterior samples and their standard deviation. We interpret the magnitude of the elements in \mathbf{w} as a function of the importance of the individual biases. We note that although this test is not explicitly intended to assess the significance of each bias factor, for which a Bayesian variable selection method could be used, it can quantify the expected amount of bias in the data predicted by single or multiple predictors in order to identify under-sampled and unexplored areas. Because several

bias factors might be correlated (e.g. cities, and airports), cumulating their bias estimates from independent analyses of each factor would result in an overestimation of the bias. For this reason, it is important to jointly estimate the effects of correlated factors, as this is based on the likelihood of the data given the combined effects of all biasing factors.

We summarize the results by mapping the estimated ‘ λ_i ’ (the relative deviation of sampling rate from the maximum sampling rate (or on user choice the estimated sampling rates), λ_i) across space. These rates represent the expected number of sampled occurrences for each grid cell and provide a graphical representation of the spatial variation of sampling rates. Provided that the cells are of equal size, the estimated rates will be comparable across data sets, regions, and taxonomic groups. Analysing different regions, biomes, or taxa in separate analyses allows to account for differences in sampling rates, which are not linked with bias factors. For instance, the unbiased sampling rate q is expected to differ between a highly sampled clade like birds and under-sampled groups of invertebrates, but their sampling biases (\mathbf{w}) might be similar across the two groups.

Example and Empirical validation

A default *sambias* analysis can be run with few lines of code in R. The main function `calculate_bias` creates an object of the class "sambias", for which the package provides a plotting and summary method. Based on a `data.frame` including species identity and geographic coordinates. Additional options exist to provide custom gazetteers, study area, spatial grid and grain size of the analysis, as well as some operators for the calculation of the bias distances, including priors for q and \mathbf{w} . A tutorial on how to use *sambias* is available

with the package and in the electronic supplement of this publication (Appendix S1).

To exemplify the use of *sambias*, we downloaded the occurrence records of all mammals available from the island of Borneo ($n = 6,262$, GBIF.org 2016), and ran *sambias* using the default gazetteers as shown in the example code below, to test the biasing effect of the main airports, cities and roads in the dataset. The example dataset is provided with *sambias*.

We found a strong effect of cities on sampling intensity, a moderate effect of roads and airports and a negligible effect of rivers (Fig. 1). All models predict a low number of collection records in the centre of Borneo (Fig. 2), which reflects the original data, and where accessibility means are low (Figure S1 in Appendix S2). The empirical example illustrates the use of *sambias*, for detailed analyses or a smaller geographic scale, higher resolution gazetteers, including smaller roads and rivers and a higher spatial resolution would be desirable. Results might change with increasing resolution, since roads and rivers might have a stronger effect on higher resolutions (facilitating most the access to their immediate vicinity), whereas cities and airports might have a stronger effect on the larger scale (facilitating access to a larger area).

```
library(sambias)

# a data table with species identity, longitude, and latitude
example.in <- read.csv(system.file("extdata",
                                   "mammals_borneo.csv",
                                   package="sambias"),
                      sep = "\t")
```

```
# running sambias

## 'res' defines the resolution of the spatial grid

## for distance calculation in degrees latitude and longitude

## 'buffer' defines the buffer around the study area to account for biasing
## structure adjacent to the study area, in degrees latitude and longitude
## All other options at default, see ?calculate_bias for a description

example.out <- calculate_bias(x = example.in,

                             res = 0.05,

                             buffer = 0.5)

# summary

summary(example.out)

plot(example.out)

# projecting the bias effect in space

proj <- project_bias(example.out)

map_bias(proj)
```

213 *Sambias* is designed to work with sparsely sampled datasets, and to estimate bias effects
214 from datasets with low coverage. To test the performance of *sambias* on small datasets we
215 ran two simulation experiments: 1) we ran the analyses as presented above on the distribution
216 of mammals in Borneo, sub-sampling of the initial dataset (to 6,262, 3,131, 1,566, 626, 62

records for all of Borneo respectively, 5 replicates each). The results showed that parameter estimates and the projection of the bias effect in space were robust to decreases in data, although uncertainty increased (Figs. S2nd S3, in Appendix S2). 2) in a second simulation we randomly sampled occurrence records across Borneo as an example of an unbiased dataset. We sampled 100,000, 50,000, 25,000, and 10,000 records and ran a *sambias* analysis with the same settings as for the empirical dataset. The results show that *sambias* reliably rejected a biasing effect even for small datasets (Figs. S4 and S5, in Appendix S2). In general, the more records, the more precise the parameter estimates will be.

Data accessibility

Sambias is available under a GNU General Public license v3 from <https://github.com/azizka/sambias>, and includes the example dataset as well as a tutorial (Appendix S1) and a summary of possible warnings produced by the package (Appendix S3).


229 **Figures**

Figure 1: Results of the empirical validation analysis, estimating the accessibility bias in mammal occurrences from Borneo). A) bias weights (w) defining the effects of each bias factor, B) sampling rate as function of distance to the closest instance of each bias factor (i.e. expected number of occurrences) given the inferred *sambias* model. At the study scale of 0.05 degrees (c. 5km) *sambias* finds the strongest biasing effect for the proximity of cities and roads.




Figure 2: Spatial projection of the sampling bias in an empirical example dataset of mammal occurrences on the Indonesian island of Borneo (downloaded from www.gbif.org. GBIF.org, 2016). The colours show the projection of the [the relative deviation of sampling rate from the maximum sampling rate given the inferred *sambias* model](#). For instance, a value of -25 indicates a drop of 25% compared to the highest sampling rate. The highest undersampling is in the centre of the island.

230 **Supplementary material**

231 Appendix S1 - Tutorial running *sambias* in R

232 Appendix S2 - Supplementary Figures

233 Appendix S3 - Possible warnings and their solutions

References

- Aiello-Lammens, M. E. et al. 2015. spThin: An R package for spatial thinning of species occurrence records for use in ecological niche models. - *Ecography* 38: 541–545.
- Bache, S. M. and Wickham, H. 2014. magrittr: A Forward-Pipe Operator for R.
- Barbosa, A. M. et al. 2013. Species-people correlations and the need to account for survey effort in biodiversity analyses. - *Diversity and Distributions* 19: 1188–1197.
- Beck, J. et al. 2014. Spatial bias in the GBIF database and its effect on modeling species' geographic distributions. - *Ecological Informatics* 19: 10–15.
- Bivand, R. S. et al. 2013. *Applied spatial data analysis with R*, Second edition. - Springer.
- Boakes, E. H. et al. 2010. Distorted views of biodiversity: Spatial and temporal bias in species occurrence data. - *PLoS Biology* 8: e1000385.
- Boria, R. A. et al. 2014. Spatial filtering to reduce sampling bias can improve the performance of ecological niche models. - *Ecological Modelling* 275: 73–77.
- Botts, E. A. et al. 2011. Geographic sampling bias in the South African Frog Atlas Project: Implications for conservation planning. - *Biodiversity and Conservation* 20: 119–139.
- Bystriakova, N. et al. 2012. Sampling bias in geographic and environmental space and its effect on the predictive power of species distribution models. - *Systematics and Biodiversity* 10: 305–315.
- Daru, B. H. et al. 2018. Widespread sampling biases in herbaria revealed from large-scale digitization. - *New Phytologist* 217: 939–955.

- Engemann, K. et al. 2015. Limited sampling hampers “big data” estimation of species richness in a tropical biodiversity hotspot. - *Ecology and Evolution* 5: 807–820.
- Fernández, D. and Nakamura, M. 2015. Estimation of spatial sampling effort based on presence-only data and accessibility. - *Ecological Modelling* 299: 147–155.
- Fithian, W. et al. 2015. Bias correction in species distribution models: pooling survey and collection data for multiple species. - *Methods in Ecology and Evolution* 6: 424–438.
- Fourcade, Y. et al. 2014. Mapping species distributions with MAXENT using a geographically biased sample of presence data: A performance assessment of methods for correcting sampling bias. - *PLoS ONE* 9: e97122.
- Garnier, S. 2018. viridis: Default color maps from 'matplotlib'.
- GBIF.org 2016. (08 September 2016) GBIF occurrence download, doi.org/10.15468/dl.7fg4zx.
- Hijmans, R. J. 2019. geosphere: Spherical Trigonometry.
- Isaac, N. J. B. and Pocock, M. J. O. 2015. Bias and information in biological records. - *Biological Journal of the Linnean Society* 115: 522–531.
- Kadmon, R. et al. 2004. Effect of roadside bias on the accuracy of predictive maps produced by bioclimatic models. - *Ecological Applications* 14: 401–413.
- Kery, M. and Royle, J. A. 2016. Applied hierarchical modeling in ecology - Analysis of distribution, abundance and species richness in R and BUGS: Volume 1: Prelude and Static Models. - Academic Press, Elsevier.
- Komori, O. et al. 2020. Sampling bias correction in species distribution models by quasi-linear

- 274 Poisson point process. - *Ecological Informatics* 55: 101015.
- 275 Kramer-Schadt, S. et al. 2013. The importance of correcting for sampling bias in MaxEnt
276 species distribution models. - *Diversity and Distributions* 19: 1366–1379.
- 277 Lin, Y.-p. et al. 2015. Uncertainty analysis of crowd-sourced and professionally collected
278 field data used in species distribution models of Taiwanese moths. - *Biological Conservation*
279 181: 102–110.
- 280 Lobo, J. M. and Tognelli, M. F. 2011. Exploring the effects of quantity and location of
281 pseudo-absences and sampling biases on the performance of distribution models with limited
282 point occurrence data. - *Journal for Nature Conservation* 19: 1–7.
- 283 Meyer, C. et al. 2015. Global priorities for an effective information basis of biodiversity
284 distributions. - *Nature Communications* 6: 8221.
- 285 Meyer, C. et al. 2016. Multidimensional biases, gaps and uncertainties in global plant
286 occurrence information. - *Ecology Letters* 19: 992–1006.
- 287 Monsarrat, S. et al. 2019. Accessibility maps as a tool to predict sampling bias in historical
288 biodiversity occurrence records. - *Ecography* 42: 125–136.
- 289 Pebesma, E. J. and Bivand, R. S. 2005. Classes and methods for spatial Data: the sp Package.
290 - *R News* 5: 21–41.
- 291 Phillips, S. J. et al. 2009. Sample Selection Bias and Presence-Only Distribution Models :
292 Implications for Background and Pseudo-Absence. - *Ecological Applications* 19: 181–197.
- 293 R Core Team 2019. R: A language and environment for statistical computing.

- Ruete, A. 2015. Displaying bias in sampling effort of data accessed from biodiversity databases using ignorance maps. - *Biodiversity Data Journal* 3: e5361.
- Rydén, O. et al. 2020. Linking democracy and biodiversity conservation: Empirical evidence and research gaps. - *Ambio* 49: 419–433.
- Shimadzu, H. and Darnell, R. 2015. Attenuation of species abundance distributions by sampling. - *Royal Society Open Science* 2: 140219.
- Stolar, J. and Nielsen, S. E. 2015. Accounting for spatially biased sampling effort in presence-only species distribution modelling. - *Diversity and Distributions* 21: 595–608.
- Vale, M. M. and Jenkins, C. N. 2012. Across-taxa incongruence in patterns of collecting bias. - *Journal of Biogeography* 39: 1744–1744.
- Varela, S. et al. 2014. Environmental filters reduce the effects of sampling bias and improve predictions of ecological niche models. - *Ecography*: 1084–1091.
- Wickham, H. 2009. *ggplot2 - Elegant graphics for data analysis*. - Springer.
- Wickham, H. 2019. *forcats: Tools for working with categorical variables (Factors)*.
- Wickham, H. and Henry, L. 2019. *tidyr: Tidy messy data*.
- Wickham, H. et al. 2019. *dplyr: A grammar of data manipulation*.
- Yang, W. et al. 2013. Geographical sampling bias in a large distributional database and its effects on species richness-environment models. - *Journal of Biogeography* 40: 1415–1426.
- Yang, W. et al. 2014. Environmental and socio-economic factors shaping the geography of floristic collections in China. - *Global Ecology and Biogeography* 23: 1284–1292.

- 314 Zizka, A. et al. 2020. Exploring the Impact of Political Regimes on Biodiversity. - VDem
315 working papers 98: 1–13.