

**\*Problem 13.42:** To determine if a graph  $G$  with 50 vertices is 3-colorable, you test all possible 3-colorings. Your computer checks a million 3-colorings per second. Estimate how long it is going to take, in the worst case.

Test all 50 vertices with 3 colors:

By sum and product law, we know that the number of subsequent choices aren't affected by the previous, we can conclude that each vertices will have 3 possible choices. All possible 3-colorings are calculated by  $3^{50}$

Given that the computer checks a million 3-colorings per second, we can calculate:

$$T = \frac{3^{50}}{10^7} \\ \approx 7.179 \times 10^{16}$$

This means that all possible 3 colorings can be calculated in about  $\boxed{7.179 \times 10^{16}}$  seconds.

**\*Problem 13.50. How many 7-digit phone-numbers are non-decreasing (each digit is not less than the previous one.)**

For this problems, we can think of it as choosing 7 random number out of a bucket, each number is placed 7 times, so there is a possible of a single number being chosen 7 times. We can think of it as this 10 containers and we have 7 items to place in 10 different containers. Since we have 10 different containers for the 7 items, we can select  $k = 7$  numbers from  $r = 10$  containers. With this, we can

derive the k-subset formula with repetition equal to  $\binom{10+7-1}{10-1} = \boxed{\binom{16}{9}}$

**\*Problem 14.15(b-c). Consider the binary strings consisting of 10 bits.**

(a) How many contain (i) 5 or more consecutive 1's (ii) 5 or more consecutive 0's?

(i) 5 or more consecutive 1's

For a string of 5 bits, we have 6 possible start locations to begin placing 1's to make sure that we reach 5 consecutive 1's. For the remaining 5 numbers, we have 2 options, represented by:

$\{ \{1, 1, 1, 1, 1\}, x, x, x, x, x \}$   
 $x = \text{dont care}$

There are  $6 \times 2^5$  choices for this subset. A double counting issue such as this arises:

$\{ x \{1, 1, 1, 1, 1\}, x, x, x, x \}$

where this subset can produce the same string as the one above.

Given that there are 5 different "dont cares", we can restrict each of them, leaving the other 4 free to have all the repeated, forming the equation  $2^4 \times 5$

Removing them from our original subset, we get  $6 \times 2^5 - 5 \times 2^4 = \boxed{112}$

(ii) 5 or more consecutive 0's

following the same logic, we can deduct this is the same, at  $\boxed{112}$  possible binary strings.

(b) **How many contain 5 or more consecutive 0's or 5 or more consecutive 1's?** For every binary string that have 5 or more consecutive 1's, just replace those with 0's to achieve  $2 \times 112$  possibilities. There are two possibilities that are double counted: 1111100000 and 0000011111 =  $\boxed{2 \times 112 - 2}$

**\*Problem 14.34. Consider all permutations of {1, 2, 3, 4, 5, 6}. A permutation is good if any of the sub-sequences 12, 23, or 56 appear. How many good permutations are there?**

Let us create subsets like such:

1. {1, 2}, 3, 4, 5, 6} subsequence 1,2 appear
2. {1, {2, 3}, 4, 5, 6} subsequence 2,3 appear
3. {1, 2, 3, 4, {5, 6}} subsequence 5,6 appear
4. {{1, 2, 3}, 4, 5, 6} subsequence 1,2 and 2,3 appear
5. {{1, 2}, 3, 4, {5, 6}} subsequence 1,2 and 5,6 appear
6. {1, {2, 3}, 4, {5, 6}} subsequence 2,3 and 5,6 appear
7. {{1, 2, 3}, 4, {5, 6}} all subsequence appear

1-3: Total number of subsequences that appears, with duplicated =  $3 \times 5!$

4-6: Number that 2 sequences appear =  $4!$ , total of  $3 \times 4!$

7: Number that all sequences appear =  $3!$

We can get rid of subset 4-6 by subtracting, but for subset 7, since all sequences appear, we removed it before so we have to add it back.

$$3 \times 5! - 3 \times 4! + 3! = 294$$

There are  $\boxed{294}$  possible permutations that contains 12, 23, or 56.

**\*Problem 14.63(g). Here are some counting problems on graphs to challenge you.**

- (g) **How many Hamiltonian cycles are in  $K_{n,n}$ ? [Hint: a Hamiltonian cycle is a cycle on graph  $G = (V, E)$  that starts and ends at vertex  $v_0 \in V$ , visiting each vertex in set  $V - \{v_0\}$  (i.e., all other vertices) exactly once.]**

There are  $n$  vertices on each side. If I start from the left side, I can choose  $n$  vertices on the right side. After choosing from the right side, I have  $n-1$  vertices to choose from the left side, and so on. If I have  $n = 4$ , the possibilities sequence will go from  $\{4, 3, 3, 2, 2, 1, 1, 1\}$ , starting from any vertex. Each vertex has  $n \times ((n-1)!)^2$  possibility of hamiltonian cycles. For a bipartite graph  $K_{n,n}$ , there are  $2n$  total vertices. There is a total of  $2n(n \times ((n-1)!)^2)$  possibilities.

**\*Problem 15.12. Roll a 6-sided die 5 times. What is the probability: (a) some number repeats (b) you get no sixes?**

- (a) some number repeats

Total number of combinations when rolling a die 5 times =  $6^5$

For a number not to repeat, for each roll, the number of possibilities after is one less. To achieve 5 rolls without any repeated the numbers, the product would be  $6 \times 5 \times 4 \times 3 \times 2$ , or  $6!$ .

To solve this, we can remove the total number of rows without a repeated number from the total combinations, leaving us with  $6^5 - 6!$ . To get the probability, we can divide it by the total:

$$\frac{6^5 - 6!}{6^5} = 0.907$$

Probabilities of rolling a 6 sided die 5 times and some number repeats =  $90.7\%$

- (b) no sixes


Total number of combinations stay the same at  $6^5$

6 is eliminated, only 5 possible numbers are left to choose from =  $5^5$


Probabilities of rolling a 6 sided die 5 times and having no 6's =  $\frac{5^5}{6^5}$  or  $40.2\%$

**\*Problem 24.11(f,h,w). Give DFAs for the following languages, a.k.a., computing problems.**


- (f)  $\mathcal{L} = \{1^{2n}01^{2k+1} \mid n, k \geq 0\}$ .



(h) Strings which begin with 10 and end with 01.




(w) Strings whose length is divisible by 3.




**\*Problem 25.7 Give a DFA and a CFG for each problem.**


(a)  $\mathcal{L} = \{01^n \mid n \geq 0\}$



(b)  $\mathcal{L} = \{0^{\bullet n}1^{\bullet n} \mid 0 \leq n \leq 5\}$


$$S \rightarrow \quad \epsilon \mid 01 \mid 0011 \mid 000111 \mid 00001111 \mid 0000011111$$

(c)  $\mathcal{L} = \{ \text{strings which end in a 1} \}$


$$S \rightarrow 1 \mid T_0S \mid T_1S$$
$$T_0 \rightarrow 0$$
$$T_1 \rightarrow 1$$