

***Problem 13.42:** To determine if a graph G with 50 vertices is 3-colorable, you test all possible 3-colorings. Your computer checks a million 3-colorings per second. Estimate how long it is going to take, in the worst case.

Test all 50 vertices with 3 colors:

By sum and product law, we know that the number of subsequent choices aren't affected by the previous, we can conclude that each vertices will have 3 possible choices. All possible 3-colorings are calculated by 3^{50}

Given that the computer checks a million 3-colorings per second, we can calculate:

$$T = \frac{3^{50}}{10^7} \\ \approx 7.179 \times 10^{16}$$

This means that all possible 3 colorings can be calculated in about $\boxed{7.179 \times 10^{16}}$ seconds.

***Problem 13.50.** How many 7-digit phone-numbers are non-decreasing (each digit is not less than the previous one.)

For 10 total numbers,

***Problem 14.15(b-c).** Consider the binary strings consisting of 10 bits.

(a) How many contain (i) 5 or more consecutive 1's (ii) 5 or more consecutive 0's?

(i) 5 or more consecutive 1's

For a string of 5 bits, we have 6 possible start locations to begin placing 1's to make sure that we reach 5 consecutive 1's. For all 6, you can only choose 1 number = 1. For the remaining 4 options, we can choose 2 numbers.

There are 6×2^4 possible binary strings consisting of 10 bits that contain 5 or more consecutive 1's.

(ii) 5 or more consecutive 0's

following the same logic, we can deduct this is the same, at 6×2^4 possible binary strings.

(b) How many contain 5 or more consecutive 0's or 5 or more consecutive 1's?

For every binary string that have 5 or more consecutive 1's, just replace those with 0's to

achieve $\boxed{2 \times 6 \times 2^4}$ possibilities.

***Problem 14.34.** Consider all permutations of $\{1, 2, 3, 4, 5, 6\}$. A permutation is good if any of the sub-sequences 12, 23, or 56 appear. How many good permutations are

there?

Let us create subsets like such:

1. $\{\{1, 2\}, 3, 4, 5, 6\}$ subsequence 1,2 appear
2. $\{1, \{2, 3\}, 4, 5, 6\}$ subsequence 2,3 appear
3. $\{1, 2, 3, 4, \{5, 6\}\}$ subsequence 5,6 appear
4. $\{\{1, 2, 3\}, 4, 5, 6\}$ subsequence 1,2 and 2,3 appear
5. $\{\{1, 2\}, 3, 4, \{5, 6\}\}$ subsequence 1,2 and 5,6 appear
6. $\{1, \{2, 3\}, 4, \{5, 6\}\}$ subsequence 2,3 and 5,6 appear
7. $\{\{1, 2, 3\}, 4, \{5, 6\}\}$ all subsequence appear

1-3: Total number of subsequences that appears, with duplicated = $3 \times 5!$

4-6: Number that 2 sequences appear = $4!$, total of $3 \times 4!$

7: Number that all sequences appear = $3!$

We can get rid of subset 4-6 by subtracting, but for subset 7, since all sequences appear, we have to remove it from 2 subsets:

$$3 \times 5! - 3 \times 4! - 2 \times 3! = 276$$

There are **276** possible permutations that contains 12, 23, or 56.

***Problem 14.63(g). Here are some counting problems on graphs to challenge you.**

- (g) **How many Hamiltonian cycles are in $K_{n,n}$? [Hint: a Hamiltonian cycle is a cycle on graph $G = (V, E)$ that starts and ends at vertex $v_0 \in V$, visiting each vertex in set $V - \{v_0\}$ (i.e., all other vertices) exactly once.]**

***Problem 15.12. Roll a 6-sided die 5 times. What is the probability: (a) some number repeats (b) you get no sixes?**

- (a) some number repeats

Total number of combinations when rolling a die 5 times = 6^5

A die is rolled 5 times, for a number to repeat, you need two rolls to be the same number, we can choose any of the 6 numbers to be that number that repeats = $(1 \times 1) \times 6$

For the rest of the 3 numbers, we have 6 possibilities of what they can be = 6^3

Multiplied together, we get $(1 \times 1) \times 6 \times 6^3 = 6^4$

Probabilities of rolling a 6 sided die 5 times and ending with a repeated number =

$\frac{6^4}{6^5}$

(b) no sixes

Total number of combinations stay the same at 6^5

6 is eliminated, only 5 possible numbers are left to choose from = 5^4

Probabilities of rolling a 6 sided die 5 times and ending with a repeated number and having

$$\text{no 6's} = \frac{5^4}{6^5}$$

***Problem 24.11(f,h,w).** Give DFAs for the following languages, a.k.a., computing problems.

(f) $\mathcal{L} = \{1^{2n}01^{2k+1} \mid n, k \geq 0\}$.

(h) Strings which begin with 10 and end with 01.

(w) Strings whose length is divisible by 3.

***Problem 25.7** Give a DFA and a CFG for each problem.

(a) $\mathcal{L} = \{01^n \mid n \geq 0\}$

(b) $\mathcal{L} = \{0^n1^n \mid 0 \leq n \leq 5\}$

(c) $\mathcal{L} = \{ \text{strings which end in a 1} \}$