

Habib University
CS-113 Discrete Mathematics
Spring 2018
HW 4 Solutions

Released: 19th March, 2018

1. (a) Prove that $1 = -1 \rightarrow 2 = 1$

Solution:

i. Since the antecedent is false, the implication is true. □

ii. Alternately, suppose

$$\begin{aligned}1 &= -1 \\1 + 3 &= -1 + 3 \\4 \times 0.5 &= 2 \times 0.5 \\2 &= 1 \quad \square\end{aligned}$$

- (b) Let $a, b \in \mathbb{N}$. What's wrong with the following proof:

$$a = b \tag{1}$$

$$a^2 = ab \tag{2}$$

$$a^2 - b^2 = ab - b^2 \tag{3}$$

$$(a - b)(a + b) = (a - b)b \tag{4}$$

$$a + b = b \tag{5}$$

$$a = 0 \tag{6}$$

Solution: Eq. 4 is being divided by $(a - b)$, which is 0 when $a = b$.

- (c) What's wrong with the following proof that shows $2^n = 1, n = \{0, 1, 2, \dots\}$, using strong induction on n :

Base step: When $n = 0, 2^0 = 1$, so the result holds.

Inductive step: Suppose the result holds for $0 \leq n \leq k$. We will show that it holds for $n = k + 1$, i.e. that $2^{k+1} = 1$.

$$\begin{aligned}
 2^{k+1} &= \frac{2^{2k}}{2^{k-1}} \\
 &= \frac{2^k \cdot 2^k}{2^{k-1}} \\
 &= \frac{1 \cdot 1}{1} \\
 &= 1
 \end{aligned}$$

Solution: The Inductive step breaks for $n = 1$:

$$\begin{aligned}
 2^1 &= \frac{2^{2 \cdot 0}}{2^{-1}} \\
 2^1 &= \frac{2^0 \cdot 2^0}{2^{-1}} \\
 2^1 &= \frac{1 \cdot 1}{1}
 \end{aligned}$$

However $2^{-1} \neq 1$ because the inductive case is undefined for $n < 0$. □

2. Buckminster Fuller once said: “None of the worlds problems will have a solution until the worlds individuals become thoroughly self-educated.”

The world is full of problems, but which ones should you focus on? According to 80000.org, the most urgent problems are “not only **big**, they’re also **neglected** and **solvable** - the fewer people working on a problem, the easier it is to make a big contribution. An issue can be big but comparatively well-known and crowded, like climate change, or it can be small but neglected, like land use zoning reform, and therefore also worth considering.”

We present a list of the biggest, most solvable and most neglected global issues that would benefit the most from your contribution. You are encouraged to read more about how these problems measure up at 8000hours.org.

Biosecurity	Climate change	Promoting effective altruism
Institutional decision-making	Risk from AI	Nuclear Security
Developing world health	Land use reform	Factory farming

You are given a set of 9 global challenges - to which you may add one more of your choice - and three order relations on them: **NB**: “not bigger than”, **NS**: “not more solvable than” and **NN**: “not more neglected than”. Using all 10 challenges and your intuition, give unique answers for each of the following. (Well-formatted, syntactically sound answers are encouraged)

- (a) A poset of width 5
 - (b) A decomposition of size 4
 - (c) A lattice
 - (d) A totally-ordered set
 - (e) A well-ordered set
- (You may refer to the **Appendix** for a list of definitions)

Solution: Suppose each problem is represented with an English letter. Call this set $S = \{a, b, c, d, e, f, g, h, i, j\}$.

(a) $\mathcal{P} = (S, NB)$

$$NB = \{(a, b), (c, d), (e, f), (g, h), (i, j)\} \cup \{(x, x) | x \in S\}$$

(b) $\mathcal{P} = (S, NS)$

$$NS = \{(a, b), (a, c), (a, d), (a, e), (a, f), (a, g)\} \cup \{(x, x) | x \in S\}$$

(c) $\mathcal{P} = (S, NN)$

$$NN = \{(a, b), (a, c), (a, d), (a, e), (a, f), (a, g), (a, h), (a, i)\} \\ \cup \{(a, j), (b, j), (c, j), (d, j), (e, j), (f, j), (g, j), (h, j), (i, j)\} \\ \cup \{(x, x) | x \in S\}$$

(d) $\mathcal{P} = (S, NB)$

The adjacency matrix of NB contains 1's at the leading diagonal, and to the right of it, and 0 otherwise.

(e) $\mathcal{P} = (S, NS)$

The adjacency matrix of NS contains 1's at the leading diagonal, and to the right of it, and 0 otherwise.

3. A poset (R, \prec) is **well-founded** if there is no infinite decreasing sequence of elements in the poset, i.e. elements x_1, x_2, \dots, x_n such that $\dots \prec x_n \prec \dots \prec x_2 \prec x_1$. A poset (R, \prec) is **dense** if for all $x \in S$ and $y \in S$ with $x \prec y$, there is an element $z \in R$ such that $x \prec z \prec y$.

Show that the set of strings of lowercase English letters with lexicographic order is neither well-founded nor dense.

Solution:

Theorem: Let ENG_L be the set of strings of lowercase English letters along with the lexicographic order relation \prec . The poset $\mathcal{P} = (ENG_L, \prec)$ is **neither well-founded, nor dense**.

Proof: \mathcal{P} has a minimal element, namely the string a . Therefore, all decreasing sequences of \mathcal{P} are finite.

Further, consider the two strings $a, b \in ENG_L$. There is no string s such that $a \prec s \prec b$. Therefore, \mathcal{P} is not dense. \square

4. Let S and T be two partial orders on a set A . Define a new relation R on A by $(x, y) \in R$ iff both $(x, y) \in S$ and $(x, y) \in T$. Prove that R is a partial order on A .

Solution: R is a partial order iff R is:

1) Reflexive:

$$\begin{aligned} \forall a \in A ((a, a) \in S \wedge (a, a) \in T) & \qquad (S, T \text{ are reflexive}) \\ \rightarrow \forall a \in A ((a, a) \in R) & \qquad (R = S \cap T) \end{aligned}$$

2) Anti-symmetric:

$$\begin{aligned} & \forall a, b ((a, b) \in R \wedge (b, a) \in R) \\ & \rightarrow ((a, b) \in S \wedge (b, a) \in S) & (R \subseteq S) \\ & \rightarrow (a = b) & (S \text{ is anti-symmetric}) \end{aligned}$$

3) Transitive:

$$\begin{aligned} & \forall a, b, c ((a, b) \in R \wedge (b, c) \in R) \\ & \rightarrow ((a, b) \in S \cap T \wedge (b, c) \in S \cap T) & (R = S \cap T) \\ & \rightarrow ((a, b) \in S \wedge (a, b) \in T \wedge (b, c) \in S \wedge (b, c) \in T) \\ & \rightarrow ((a, c) \in S \wedge (a, c) \in T) & (S, T \text{ are transitive}) \\ & \rightarrow ((a, c) \in R) \end{aligned}$$

5. Your overly-attached girlfriend / boyfriend has concocted a cruel game to keep you around forever. The game involves piles of excuses - each pile twice as high as the last - of all the times you didn't hang out with them. Stacked in increasing order, the piles look truly endless!




Figure 1: That escalated quickly.

The game of **Gazillion Excuses** is a turn based 2-player game such that at each turn, a player removes a non-zero number of excuses from one of the piles. The player who removes the last excuse wins. Your partner thinks that a sufficiently large number of piles (think gazillion) would keep you playing forever. Little do they suspect that a Discrete Mathematician like yourself has formidable proof skills! Given that you are allowed to go first, **prove using induction** that you will always win the game of Gazillion Excuses.

Solution:

Theorem: Player 1 has a winning strategy with k piles of lengths $2^0, 2^1, \dots, 2^{k-1}$.

Proof: To prove our theorem, we prove the more generalized version of our setup first.

Lemma 1: Player 1 has a winning strategy with N excuses in k piles of lengths n_1, n_2, \dots, n_k , when $n_1 \oplus n_2 \oplus \dots \oplus n_k \neq 0$.

Proof: We perform induction on the number of excuses N . Our proof relies on two important results which we present first.

Lemma 2:

- a) There exists a move that removes x from n_i when $n_1 \oplus n_2 \oplus \dots \oplus n_i \oplus \dots \oplus n_k \neq 0$, resulting in $n_1 \oplus n_2 \oplus \dots \oplus m_i \oplus \dots \oplus n_k = 0$, where $m_i = n_i - x$.
- b) All moves that remove x from n_i when $n_1 \oplus n_2 \oplus \dots \oplus n_i \oplus \dots \oplus n_k = 0$, result in $n_1 \oplus n_2 \oplus \dots \oplus m_i \oplus \dots \oplus n_k \neq 0$, where $m_i = n_i - x$.

Proof: Consider binary representations of n_1, n_2, \dots, n_k written in table form:

	n th bit	$n - 1$ th bit	\dots	0th bit
n_1	b_{1n}	$b_{1(n-1)}$	\dots	b_{10}
n_2	b_{2n}	$b_{2(n-1)}$	\dots	b_{20}
\vdots	\vdots	\vdots	\ddots	\vdots
n_k	b_{kn}	$b_{k(n-1)}$	\dots	b_{k0}
\oplus	$b_{\oplus n}$	$b_{\oplus(n-1)}$	\dots	$b_{\oplus 0}$

The last row is the bitwise XOR of n_1, n_2, \dots, n_k . Note that toggling any one bit b_{ij} in the table toggles $b_{\oplus j}$ as well.

a) Suppose $n_1 \oplus n_2 \oplus \dots \oplus n_i \oplus \dots \oplus n_k \neq 0$. Then, $\exists b \in \{b_{\oplus n}, b_{\oplus(n-1)}, \dots, b_{\oplus 0}\} (b = 1)$. Here's a strategy that ensures $\oplus n_x = 0$: Scan row \oplus from left to right noting column numbers of every 1 encountered. Scan column 1 in this list from top to bottom until 1 is encountered. In this row, toggle all bits that correspond to the listed column numbers. Since toggling any one bit b_{ij} in the table toggles $b_{\oplus j}$ as well, we ensure that all bits in \oplus that were previously 1, are now 0. Thus $n_1 \oplus n_2 \oplus \dots \oplus m_i \oplus \dots \oplus n_k = 0$.

b) Next, suppose that $n_1 \oplus n_2 \oplus \dots \oplus n_i \oplus \dots \oplus n_k = 0$. Then $\forall b \in \{b_{\oplus n}, b_{\oplus(n-1)}, \dots, b_{\oplus 0}\} (b = 0)$. Suppose x excuses are removed from pile n_i . Then at least one bit, say b_{ij} , was toggled. Now $\exists b \in \{b_{\oplus n}, b_{\oplus(n-1)}, \dots, b_{\oplus 0}\} (b = 1)$. Thus, $n_1 \oplus n_2 \oplus \dots \oplus m_i \oplus \dots \oplus n_k \neq 0$.

Now we are ready to prove **Lemma 1** via induction on N :

Base case, $N = 0$:

$\oplus n_x = 0$. Player 1 loses by definition.

Inductive step: Suppose Lemma 1 is true for all values less than N . In the case of N excuses, there are two cases:

- 1) $n_1 \oplus n_2 \oplus \dots \oplus n_k \neq 0$: By Lemma 2a, there exists a move for player 1 whereby x excuses are removed from pile n_i to give $n_1 \oplus n_2 \oplus \dots \oplus m_i \oplus \dots \oplus n_k = 0$. By I.H., Player 1 wins.
- 2) $n_1 \oplus n_2 \oplus \dots \oplus n_k = 0$: By Lemma 2b, all moves by player 1 result in $n_1 \oplus n_2 \oplus \dots \oplus m_i \oplus \dots \oplus n_k \neq 0$. By I.H., Player 2 wins.

Since our **Theorem** gives starting piles $2^0, 2^1, \dots, 2^{k-1}$ where $2^0 \oplus 2^1 \oplus \dots \oplus 2^{k-1} \neq 0$, Lemma 1 states that Player 1 will win. \square

1 Appendix

Let $\mathcal{P} = (P, \preceq)$,

Chain: A chain, $C \subseteq P$, is a subset of mutually comparable elements of P .

Anti-Chain: An anti-chain, $A \subseteq P$, is a subset of mutually incomparable elements of P .

Width: The maximum cardinality of an anti-chain of \mathcal{P} .

Decomposition: A decomposition C of \mathcal{P} into chains, is a family $C = \{C_1, C_2, \dots, C_q\}$ of disjoint chains such that their union is P . The size of a decomposition is the number of chains in it.