

Clinical and Pharmacological Assessment of Dietary Supplementation: Efficacy, Speciation, and Regulatory Dynamics

Primary Prevention of Cardiovascular Disease and Cancer: Public Health Guidelines and Multi-Cohort Trial Outcomes

The therapeutic utility of generalized micronutrient supplementation for the primary prevention of cardiovascular disease (CVD) and oncological diseases in healthy, non-deficient adults has been extensively analyzed by the United States Preventive Services Task Force (USPSTF). In its updated 2022 guidelines, which synthesized evidence from 84 clinical trials encompassing 739,803 participants, the USPSTF reaffirmed a highly conservative stance toward over-the-counter supplementation.

The task force issued a definitive Grade D recommendation against the use of beta-carotene and vitamin E supplements for the prevention of cardiovascular disease or cancer. This clinical boundary is supported by moderate certainty that the systemic harms of these specific antioxidants outweigh any hypothetical physiological benefits in healthy populations.

Specifically, multiple large-scale randomized controlled trials (RCTs) demonstrated that beta-carotene supplementation significantly increases the risk of lung cancer (Odds Ratio $= 1.20$, 95% Confidence Interval $[1.01, 1.42]$) and cardiovascular mortality ($\text{OR} = 1.10$, 95% CI, 1.02 to 1.19) in individuals with high baseline risk, such as tobacco smokers or those with occupational asbestos exposure.

Conversely, for multivitamins and other single- or paired-nutrient formulations, the USPSTF issued a Grade I (Insufficient Evidence) statement. This indicates that the current clinical literature is insufficient to determine the balance of benefits and harms for primary prevention. This guideline applies to community-dwelling, nonpregnant, and non-deficient adults, but does not govern pediatric populations, pregnant individuals, or those with clinically diagnosed nutritional deficiencies.

Beyond beta-carotene and vitamin E, the USPSTF systematic review identified specific safety risks associated with other commonly consumed single nutrients. High-dose vitamin A supplementation was linked to an increased risk of hip fractures, particularly in cohort studies of predominantly White women.

Similarly, vitamin E was associated with an increased risk of hemorrhagic stroke in two large clinical trials, while high-dose vitamin C and calcium supplementation showed an elevated risk of kidney stones.

The evidence for other major micronutrients—including B-vitamins, folic acid, magnesium, selenium, and zinc—was rated as low, insufficient, or absent for all major clinical outcomes, indicating a substantial gap between consumer utilization and established clinical proof of efficacy.

Nutrient Supplement	USPSTF Grade Recommendation	Primary Clinical Endpoint Associations	Identified Systemic Harms & Adverse Outcomes	Source(s)
Beta-Carotene	Grade D (Against)	No preventive benefit for CVD or cancer.	Increased lung cancer incidence ($\text{OR} = 1.20$) in smokers; increased CVD mortality ($\text{OR} = 1.10$).	
Vitamin E	Grade D (Against)	No preventive benefit for CVD, cancer, or all-cause mortality.	Increased risk of hemorrhagic stroke; increased cancer recurrence post-radiotherapy; higher all-cause mortality.	
Multivitamins	Grade I (Insufficient)	Modest, pooled cancer risk reduction ($\text{OR} = 0.93$); no clear CVD or mortality benefit.	Statistically small increase in gastrointestinal bleeding events in some trials.	
Vitamin A	Grade I (Insufficient)	Minimal or absent evidence of primary	Increased risk of hip fracture; chronic	


Nutrient Supplement	USPSTF Grade Recommendation	Primary Clinical Endpoint Associations	Identified Systemic Harms & Adverse Outcomes	Source(s)
		preventive benefit.	hypervitaminosis A associated with intra-hepatic cholestasis.	
<i>Vitamin D</i>	Grade I (Insufficient)	Non-significant reduction in major CVD events or total cancer incidence.	Increased risk of nephrolithiasis (kidney stones) at doses ≥ 1000 IU/day over prolonged periods.	
Calcium	Grade I (Insufficient)	Equivocal or absent primary preventive efficacy.	Elevated risk of constipation, gastrointestinal distress, and kidney stones.	

To address these clinical gaps, the VITamin D and Omega-3 Trial (VITAL) was designed as a nationwide, randomized, double-blind, placebo-controlled, 2 $\times$ 2 factorial trial. VITAL enrolled 25,871 healthy U.S. participants (men aged ≥ 50 and women aged ≥ 55 , including a highly diverse cohort of 5,106 African Americans) to evaluate daily supplementation with vitamin D3 (2000 IU) and marine omega-3 fatty acids (1 g), containing 460 mg of eicosapentaenoic acid [EPA] and 380 mg of docosahexaenoic acid) over a median follow-up of 5.3 years.

The primary co-primary endpoints of major cardiovascular events (a composite of myocardial infarction, stroke, and cardiovascular death) and total invasive cancer incidence showed no statistically significant reductions in the overall study population for either intervention.

Vitamin D3 did not reduce major cardiovascular events (Hazard Ratio $= 0.97$, 95% CI, 0.85 to 1.12) or overall cancer incidence (HR = 0.96, 95% CI, 0.88 to 1.06).

Similarly, marine omega-3 fatty acids did not significantly lower the risk of the composite cardiovascular endpoint (HR = 0.92, 95% CI, 0.80 to 1.06) or total cancer incidence (HR = 1.03, 95% CI, 0.93 to 1.13).



However, deeper analyses of secondary endpoints and nested cohorts within the VITAL trial revealed clinically relevant, subgroup-specific preventive effects.

While overall cancer incidence was unaffected, vitamin D3 supplementation was associated with a suggestive 17% reduction in total cancer mortality (HR = 0.83, 95% CI, 0.67 to 1.02).

When the analysis was adjusted for oncological latency by excluding the first two years of follow-up, the reduction in cancer-related deaths became statistically significant, reaching 25% (HR = 0.75). This latency effect suggests that vitamin

D may slow tumor progression, invasiveness, and metastatic potential rather than preventing initial oncogenesis.

Furthermore, a significant interaction was observed based on baseline body mass index (BMI): individuals with a normal body weight (BMI < 25) experienced a significant reduction in cancer incidence from vitamin D, whereas overweight and obese individuals did not. This indicates potential vitamin D resistance or tissue sequestration of the fat-soluble vitamin in individuals with excess adipose tissue.

For the marine omega-3 arm, the primary composite endpoint was non-significant, but the secondary endpoint of myocardial infarction (MI) was reduced by 28% in the overall study population ($\text{HR} = 0.72$, 95% CI , 0.59 to 0.90).

This cardioprotective effect was strongly modulated by baseline dietary patterns and race.

Participants with low baseline fish consumption (< 1.5 servings per week) experienced a significant 19% reduction in major cardiovascular events and a 40% reduction in MI ($\text{HR} = 0.60$) from omega-3 supplementation, whereas individuals with higher baseline intake derived no additional benefit.

The most pronounced clinical benefit was observed among African American participants, who experienced a 77% reduction in myocardial infarction ($\text{HR} = 0.23$) that persisted regardless of their baseline fish consumption.

To further characterize the systemic safety of these compounds, nested studies within VITAL examined renal and electrophysiological outcomes.

The VITAL-CKD substudy evaluated chronic kidney disease (CKD) progression over 5 years in 1,312 participants with type 2 diabetes.

The mean change in the estimated glomerular filtration rate (eGFR) from baseline to year 5 was $-12.3 \text{ mL/min/1.73 m}^2$ in the vitamin D3 group versus $-13.1 \text{ mL/min/1.73 m}^2$ in its respective placebo group, and $-12.2 \text{ mL/min/1.73 m}^2$ in the omega-3 group versus $-13.1 \text{ mL/min/1.73 m}^2$ in its respective placebo group. No statistically significant difference or interaction was found, indicating that neither supplement slows diabetic kidney disease progression.

In the VITAL Rhythm substudy, atrial fibrillation (AF) incidence was monitored over 5.3 years.

Incident AF developed in 3.6% of the study population, with no significant differences observed between active omega-3 and placebo ($\text{HR} = 1.09$, 95% CI , 0.96

1.24, $P = 0.19$) or active vitamin D3 and placebo ($\text{HR} = 1.09$, 95% CI, 0.96 to 1.25, $P = 0.19$).

These findings indicate that moderate doses of omega-3 fatty acids (1 g/day) do not significantly elevate AF risk, contrasting with some high-dose prescription-grade omega-3 formulations.


Cognitive Longevity and Flavanol-Driven Interventions: The COSMOS Trial

While the USPSTF and VITAL focused on cardiorespiratory and oncological health, the COcoa Supplement and Multivitamin Outcomes Study (COSMOS) examined the potential of dietary supplements to slow cognitive aging in older adults. COSMOS was designed as a large-scale, nationwide, randomized, double-blind, placebo-controlled, 2 $\times$ 2 factorial trial enrolling 21,442 participants (comprising 12,666 women aged ≥ 65 and 8,776 men aged ≥ 60) followed for an average of 3.6 years.

The trial evaluated two primary interventions: a daily multivitamin-mineral supplement (MVM; Centrum Silver, a formulation containing more than 20 essential micronutrients) and a cocoa extract supplement (delivering 500 mg of cocoa flavanols daily, including epicatechin).

To rigorously evaluate cognitive changes, COSMOS utilized three distinct, non-overlapping substudies, each employing a unique cognitive assessment modality :

- COSMOS-Mind utilized annual telephone-based cognitive testing in 2,158 participants to track changes in global cognition, memory, and executive function.
- COSMOS-Web utilized annual computer-based cognitive testing in 2,472 participants to measure memory performance.
- COSMOS-Clinic administered detailed, face-to-face neuropsychological assessments in 573 participants to obtain direct clinical measures.



- Telephone assessments (n=2,158) - Web-based memory (n=2,472) - In-person testing (n=573)

- Slowed cognitive aging by - Improved memory at 1 year - Significantly improved

1.8-year equivalent (3.1-year equivalent) episodic memory

Daily multivitamin-mineral supplementation demonstrated consistent, statistically significant benefits for memory and global cognitive function across all three substudies.

In the COSMOS-Web cohort, participants taking the daily multivitamin performed significantly better on memory tests at the primary 1-year timepoint, with these benefits maintained over the 3-year study period. The investigators estimated that the MVM intervention improved memory performance by the equivalent of slowing memory loss by 3.1 years compared to the placebo group.

In the COSMOS-Mind cohort, daily MVM use significantly improved global cognitive scores over 3 years, slowing cognitive aging by approximately 60%, or the equivalent of delaying cognitive decline by 1.8 years.

Subgroup analysis within COSMOS-Mind revealed that the cognitive benefits of MVM were most pronounced in participants with a history of cardiovascular disease, who may have had lower baseline micronutrient levels.

In the in-person COSMOS-Clinic substudy, daily MVM use led to a significantly more favorable 2-year change in episodic memory (mean difference = 0.12 standard deviation units, 95% CI, 0.002 to 0.23), though no significant difference was observed for executive function or spatial cognition.

A subsequent meta-analysis compiling data from all three non-overlapping COSMOS cognitive substudies provided robust, high-quality evidence that daily MVM use significantly benefits both global cognition (mean difference = 0.07 SU, 95% CI, 0.03 to 0.11, $P = 0.0009$) and episodic memory (mean difference = 0.06 SU, 95% CI, 0.03 to 0.10, $P = 0.0007$).

The overall magnitude of effect on global cognition was equivalent to reducing cognitive aging by approximately 2 years compared to placebo.

In contrast, the cocoa extract supplement did not show a statistically significant effect on overall cognitive function in the general study population.

However, baseline nutritional status significantly influenced the outcomes.

Participants with low baseline dietary flavanol intake experienced significant cognitive and memory benefits from the cocoa extract compared to those taking a placebo.

Regarding safety, both interventions demonstrated highly favorable overall safety profiles.

Cocoa extract was associated with a 6% increase in self-reported nausea, but also with a 5% reduction in flu-like symptoms and headaches, and a 15% reduction in migraines.

The multivitamin group reported lower rates of stomach upset, diarrhea, skin rash, and bruising, though a slightly higher rate of gastrointestinal bleeding was noted compared to the placebo group.

Molecular Biochemistry, Speciation, and Bioavailability Dynamics

Folate Chemistry: Folic Acid vs. 5-Methyltetrahydrofolate

Folate (Vitamin B9) exists in several chemical forms that differ significantly in stability, bioavailability, and metabolic pathways. Naturally occurring folates present in foods (such as leafy greens) are highly unstable tetrahydrofolate (THF) polyglutamates. These compounds are susceptible to oxidation by heat, light, and cooking, which markedly reduces their bioavailability to approximately 50%. To be absorbed, natural polyglutamates must first be enzymatically hydrolyzed to the monoglutamate form in the digestive tract.


In contrast, synthetic folic acid (pteroylmonoglutamic acid) is a fully oxidized, stable monoglutamate that does not occur naturally in foods. It is highly bioavailable, with absorption rates of approximately 85% when consumed with food and nearly 100% when taken on an empty stomach. Consequently, the Food and Nutrition Board (FNB) established Dietary Folate Equivalents (DFEs) to adjust for these differences:

$$1 \mu\text{g DFE} = 1 \mu\text{g food folate} = 0.6 \mu\text{g folic acid from fortified foods/supplements with meals} = 0.5 \mu\text{g folic acid on an empty stomach} \quad [\text{span_141}](\text{start_span})[\text{span_141}](\text{end_span})$$

In the United States, this bioavailability difference led to the mandatory folic acid fortification program initiated by the FDA in January 1998, which required the addition of 140 μg of folic acid per 100 g of enriched grain products. This public health intervention increased the mean daily folic acid intake in the U.S. population by approximately 190 μg/day, successfully reducing the national incidence of neural tube defects (NTDs).

The biochemical cutoffs for folate adequacy are firmly established: a serum folate concentration above 3 ng/mL indicates short-term adequacy, while erythrocyte (red blood cell) folate concentrations above 140 ng/mL provide a more reliable measure of long-term tissue stores.

Elevated systemic homocysteine levels (using a cutoff of $> 16\text{ }\mu\text{mol/L}$, or more conservatively, $> 12\text{ to }14\text{ }\mu\text{mol/L}$) serve as a functional biomarker of folate or vitamin B12 deficiency.



However, synthetic folic acid has metabolic limitations. Folic acid is biologically inactive and must undergo a multi-step enzymatic reduction to become the active circulating form, 5-methyltetrahydrofolate (5-MTHF). Folic acid is first reduced to dihydrofolate by the enzyme dihydrofolate reductase (DHFR) and subsequently to THF.

Because human DHFR has exceptionally low and highly variable enzymatic activity in the liver, this reduction step is highly rate-limiting. Regular consumption of folic acid, even at moderate doses of 200 to 400 $\mu\text{g/day}$, can saturate this pathway. This leads to the accumulation of Unmetabolized Folic Acid (UMFA) in the systemic circulation.

Elevated UMFA levels have been linked to potential adverse clinical outcomes, including competitive inhibition of the DHFR gene (which can worsen folate deficiency), reduced natural killer (NK) cell cytotoxicity, and immune dysfunction.

Furthermore, genetic polymorphisms in the methylenetetrahydrofolate reductase (MTHFR) gene (specifically the C677T and A1298C variants) significantly impair the conversion of folic acid to active methylfolate. Individuals with these variants have a reduced capacity to synthesize 5-MTHF, leading to elevated homocysteine levels and increased systemic susceptibility to inflammation and oxidative stress.

To address these limitations, active L-methylfolate (5-MTHF) has been evaluated as an alternative for supplementation. Levomefolate supplementation (available as calcium, magnesium, or glucosamine salts, such as Quatrefolic) bypasses the rate-limiting DHFR step and the MTHFR genetic block.

Active 5-MTHF is immediately bioavailable, directly enters the one-carbon cycle, does not generate circulating UMFA, and is highly effective at reducing homocysteine levels.

Furthermore, unlike synthetic folic acid, active 5-MTHF does not mask a vitamin B12 deficiency, reducing the risk of progressive, undetected neurological damage in vulnerable populations.

However, established clinical guidelines for pregnancy continue to prioritize folic acid, as it is the only form of folate clinically proven in large trials to prevent neural tube defects.

| Folate Chemical Form | Chemical

[span_157](start_span)[span_157](end_span)Characteristics | Bioavailability & Absorption | Metabolic Pathway & Enzymatic Requirements | Genetic / Tolerability Limitations |

| :--- | :--- | :--- | :--- | :--- |

| ****Natural Food Folate**** | THF polyglutamates; highly unstable; degraded by heat and cooking. | Moderate ($\sim 50\%$); requires enzymatic hydrolysis to monoglutamates in the gut. | Absorbed monoglutamates are reduced to THF, then methylated to 5-MTHF prior to entering systemic circulation. | Unstable; subject to rapid oxidation and degradation during food preparation. |

| ****Synthetic Folic Acid**** | Fully oxidized monoglutamate; highly stable; crystalline structure. | Very High ($\sim 100\%$); evaluated via DFEs due to superior absorption compared to food folate. | Requires rate-limiting reduction by hepatic DHFR, conversion to THF, and subsequent reduction to 5-MTHF by MTHFR. | Hepatic DHFR bottleneck can lead to circulating UMFA at doses $\geq 200 \mu\text{g}$; conversion is severely impaired by MTHFR polymorphisms (C677T/A1298C). |

| ****Active Methylfolate**** (5-MTHF / Levomefolate) | Biologically active, reduced physiological form; available as stable salts. | Extremely High; directly absorbed across the intestinal mucosa without requiring further metabolic alteration. | Bypasses all enzymatic conversion steps; directly participates in the conversion of homocysteine to methionine. | None; does not generate UMFA and is fully effective in individuals with severe MTHFR genetic polymorphisms. |

Mineral Speciation: Inorganic vs. Organic Magnesium

The systemic absorption of magnesium is highly dependent on its chemical speciation.[span_169](start_span)[span_169](end_span) In dietary supplements, magnesium (Mg^{2+}) is bound to either an inorganic molecule or an organic acid/amino acid chelate.[span_170](start_span)[span_170](end_span)

Inorganic magnesium compounds, such as magnesium oxide (MgO) and magnesium carbonate (MgCO_3), are highly concentrated in elemental magnesium.[span_171](start_span)[span_171](end_span) Magnesium oxide contains approximately 60% elemental magnesium by

weight. However, it is virtually insoluble in water.

To be absorbed, inorganic magnesium must first be dissociated by hydrochloric acid in the stomach. In vitro studies simulating gastric acid secretion show that magnesium oxide is only 43% soluble at peak acid secretion and entirely insoluble in achlorhydric states.

As a result, magnesium oxide has extremely low bioavailability, with studies indicating an absorption rate of approximately 4% in the gastrointestinal tract. The remaining unabsorbed magnesium remains in the intestinal lumen, where it exerts an osmotic effect. It draws water into the bowel, causing loose stools, diarrhea, and abdominal cramping. Consequently, magnesium oxide is clinically indicated as an antacid or osmotic laxative rather than a source for systemic repletion.

In contrast, organic magnesium compounds, such as magnesium citrate, magnesium glycinate (bisglycinate), and magnesium malate, are highly soluble in water and do not require highly acidic gastric environments for dissolution. Magnesium citrate is a magnesium salt bound to citric acid. It has high solubility even in water (55%) and does not reprecipitate in alkaline pancreatic environments.

Clinical trials evaluating post-load urinary magnesium excretion confirm that magnesium citrate is significantly more bioavailable than magnesium oxide, achieving a systemic absorption rate of 25% to 30%.

Specifically, a 25 mmol oral load of magnesium citrate produced a significant urinary magnesium increment of 0.22 mg/mg creatinine versus 0.006 mg/mg creatinine for magnesium oxide at 4 hours post-load, and 0.035 mg/mg creatinine versus 0.008 mg/mg creatinine during the second 2 hours post-load ($P < 0.05$).

Other organic formulations exhibit distinct pharmacokinetic properties. Magnesium bisglycinate is a chelate bound to the amino acid glycine. The glycine moiety utilizes active amino acid transport pathways across the intestinal epithelium. This shields the magnesium from interacting with other dietary inhibitors and makes it highly tolerable and gentle on the bowel, with a very low risk of osmotic laxation.

Magnesium malate is bound to malic acid, an intermediate in the Krebs cycle. It is absorbed rapidly and is preferred for individuals with fatigue or chronic pain disorders, such as fibromyalgia, due to malic acid's role in ATP production.

Magnesium L-threonate is bound to threonic acid, a metabolite of vitamin C. It is uniquely capable of crossing the blood-brain barrier, making it highly effective for raising cerebrospinal fluid magnesium levels to support cognitive function and alleviate anxiety.

Conversely, magnesium sulfate (Epsom salt) has very low oral bioavailability and acts as a strong saline cathartic, with research indicating minimal systemic absorption through the skin during therapeutic baths.

Safety, Toxicology, and Herb-Drug

Interactions

Cytochrome P450 and P-glycoprotein Induction: St. John's Wort


A major safety concern in clinical pharmacology is the interaction between botanical supplements and prescription

medications. St. John's wort (*Hypericum perforatum*), a widely used herbal remedy for mild-to-moderate depression, is a classic example of these interactions. It can alter the pharmacokinetics of co-administered pharmaceuticals, leading to sub-therapeutic drug levels and therapeutic failure.

The primary mechanism of interaction involves the active constituent hyperforin, which acts as a potent ligand and activator of the pregnane X receptor (PXR) in hepatocytes and enterocytes. Upon activation, PXR transcriptionally upregulates the expression of the Cytochrome P450 3A4 (CYP3A4) enzyme and the ATP-Binding Cassette transporter P-glycoprotein (P-gp/ABCB1).

CYP3A4 is the most abundant drug-metabolizing enzyme in the human liver and intestines. It is responsible for biotransforming more than 50% of all clinically prescribed medications. P-glycoprotein acts as an apical efflux pump in the intestine, blood-brain barrier, and renal tubules, actively pumping xenobiotics out of cells.

Consequently, the co-administration of St. John's wort significantly decreases the area under the curve (AUC) and steady-state blood concentrations of numerous critical drugs.



- Cyclosporine clearance -> Organ rejection - Estrogen degradation -> Birth control failure - Anticoagulant clearance -> Clotting risk

Clinical evidence indicates that St. John's wort reduces the serum concentrations of the immunosuppressants cyclosporine and tacrolimus. This can cause acute or[span_212](start_span)[span_212](end_span)gan transplant rejection in clinical settings.

Furthermore, St. John[span_213](start_span)[span_213](end_span)'s wort accelerates the clearance of oral contraceptives, leading to breakthrough bleeding and unplanned pregnancies. It also reduces the ser[span_214](start_span)[span_214](end_span)um concentrations of digoxin, amitriptyline, methadone, simvastatin, and theophylline.

In oncology, co-admin[span_215](start_span)[span_215](end_span)istration with irinotecan significantly decreases the plasma concentration of its active cytotoxic metabolite, SN-38, thereby compromising chemotherapeutic efficacy.

Pharmacodynamically, [span_216](start_span)[span_216](end_span)St. John's wort exhibits additive serotonergic activity. When co-administered with selective serotonin reuptake inhibitors (SSRIs), such as sertraline or paroxetine, it can precipitate serotonin syndrome, a life-threatening condition characterized by autonomic instability, neuromuscular abnormalities, and altered mental status.

Clinical Managemen[span_217](start_span)[span_217](end_span)t of Oral Anticoagulants

Warfarin, a highly efficacious oral anticoagulant with a narrow therapeutic index, is highly susceptible to herb-drug interactions.[span_225](start_span)[span_225](end_span) Warfarin is administered as a racemic mixture of R- and S-enantiomers, with the S-enantiomer possessing three to five times greater anticoagulant potency.[span_228](start_span)[span_228](end_span) S-warfarin is metabolized primarily by CYP2C9, while R-warfarin is metabolized by CYP1A2, CYP2C19, and CYP3A4.[span_229](start_span)[span_229](end_span)

St. John's wort significantly decreases the pharmacological effect of warfarin, lowering the International Normalized Ratio (INR) and elevating thrombotic risk.[span_230](start_span)[span_230](end_span)[span_231](start_span)[span_231](end_span) A randomized, open-label, crossover clinical trial in healthy subjects demonstrated that 14 days of pretreatment with St. John's wort significantly induced

the apparent clearance of both S-warfarin and R-warfarin.

The apparent clearance of S-warfarin increased from $\$198 \pm 38 \text{ mL/h}$ to $\$270 \pm 44 \text{ mL/h}$ (a ratio of $\$1.29$, $90\% \text{ CI}$, 1.16 to 1.46), while R-warfarin clearance increased from $\$110 \pm 25 \text{ mL/h}$ to $\$142 \pm 29 \text{ mL/h}$. This pharmacokinetic induction resulted in a significant 21% reduction in the AUC of the INR ($\$0.79$, $90\% \text{ CI}$, 0.70 to 0.95).

This study also evaluated ginseng (*Panax ginseng*) and found that it did not affect the pharmacokinetics or pharmacodynamics of either warfarin enantiomer. This demonstrates that anticoagulant instability is not a universal feature of all botanical supplements.

However, other common botanical supplements are classified as having major severity interactions with warfarin. These include cannabis, chamomile, cranberry, garlic, ginkgo, red clover, and grapefruit. The mechanisms are often dual-faceted: ginkgo and garlic possess intrinsic antiplatelet properties that pharmacodynamically compound bleeding risks without altering the INR.

Conversely, grapefruit juice contains furanocoumarins that inhibit intestinal CYP3A4, which can elevate R-warfarin concentrations.

Targeted Clinical Recommendations for Special Populations

Antenatal Nutrition and Maternal-Fetal Medicine


Maternal micronutrient status is a critical determinant of embryonic development, gestational outcomes, and long-term offspring health. During pregnancy, maternal blood volume expands, and fetal and placental growth sharply increases nutritional demands. This makes targeted micronutrient supplementation essential.

The World Health Organization (WHO) and major obstetric societies recommend daily oral supplementation with elemental iron (30 to 60 mg/day) and folic acid (0.4 mg/day or 400 µg/day) for all pregnant women to prevent maternal anemia, puerperal sepsis, low birth weight, and neural tube defects (NTDs).

Because neural tube closure occurs within the first 28 days of gestation—often before pregnancy is confirmed—folic acid supplementation of 400 to 800 µg/day is strongly recommended for all individuals of childbearing potential starting 5 to 6 months prior to conception.

The Recommended Dietary Allowance (RDA) for folate increases to 600 µg DFE orally daily during pregnancy and 500 µg DFE daily during lactation.

In geographic regions with low dietary calcium intake, daily calcium supplementation (1.5 to 2.0 g/day) is recommended to reduce the risk of pre-eclampsia, a major cause of maternal mortality.



- 30-60 mg daily to - 400 mcg daily to - Replaces IFA - Reduced preterm birth, prevent anemia prevent NTDs formulation LBW, and female neonatal - Outperforms IFA mortality

Historically, standard prenatal care relied on Iron and Folic Acid (IFA) monotherapy.^{[span_268](start_span)[span_268](end_span)[span_269](start_span)[span_269](end_span)} However, multiple micronutrient deficiencies often coexist in vulnerable populations.^{[span_270](start_span)[span_270](end_span)[span_271](start_span)[span_271](end_span)}

To address this, Multiple Micronutrient Supplementation (MMS)—utilizing standardized formulations such as the United Nations International Multiple Micronutrient Antenatal Preparation (UNIMMAP), which contains 15 micronutrients including 10 vitamins and 5 minerals (zinc, iron, selenium, copper, and iodine)—has been evaluated.

An Individual Patient Data (IPD) meta-analysis of 17 randomized controlled trials involving more than \$100,000\$ pregnant women in low- and middle-income countries demonstrated that MMS is superior to IFA alone.

MMS significantly reduced the risk of low birth weight (\$\text{LBW}\$), preterm birth, and infants born small-for-gestational-age (\$\text{SGA}\$).^{[span_272](start_span)[span_272](end_span)[span_273](start_span)[span_273](end_span)}

Furthermore, MMS significantly reduced neonatal and infant mortality in female offspring, with no evidence of increased risk across 26 analyzed subgroups.

Based on this evidence^{[span_266](start_span)[span_266](end_span)}, public health guidelines support transitioning from IFA to MMS in populations with a high prevalence of maternal nutritional deficiencies.^{[span_274](start_span)[span_274](end_span)[span_275](start_span)[span_275](end_span)[span_276](start_span)[span_276](end_span)}

Geriatric Micronutrient Requirements and Gastrointestinal Malabsorption
Aging is associated with physiological changes in the gastrointestinal tract that impair

the absorption of key micronutrients, particularly Vitamin B12 (cobalamin).

Between 3% and 43% of community-dwelling older adults suffer from vitamin B12 deficiency. This is primarily driven by atrophic gastritis, an age-related chronic inflammatory disease that results in the destruction of gastric parietal cells.

This destruction causes a dual defect: a loss of hydrochloric acid secretion (achlorhydria), which prevents the enzymatic cleavage of vitamin B12 from dietary proteins, and a loss of intrinsic factor secretion, which is required for active receptor-mediated absorption of cobalamin in the terminal ileum. Chronic autoimmune pernicious anemia represents the most severe form of this malabsorptive state.

Vitamin B12 deficiency in older adults can lead to macrocytic megaloblastic anemia, peripheral neuropathy, cognitive decline, and psychiatric symptoms.

The biochemical diagnosis is confirmed when serum or plasma cobalamin levels fall below 200 pg/mL or 148 pmol/L).

In patients with borderline values between 150 pg/mL and 399 pg/mL, testing for elevated serum methylmalonic acid (MMA) is recommended to confirm tissue-level deficiency.

While the standard adult RDA is 2.4 µg/day, older adults with atrophic gastritis require significantly higher intakes (10 to 12 µg/day) of crystalline vitamin B12. Crystalline cobalamin is not bound to food proteins and is more readily absorbed in achlorhydric states.

In cases of severe deficiency, pernicious anemia, or post-gastric bypass surgery (such as Roux-en-Y), high-dose oral cyanocobalamin or methylcobalamin (1000 µg/day) utilizes passive

diffusion pathways across the intestinal epithelium. This pathway, which accounts for approximately 1% of passive absorption, is highly effective.

Clinical trials have shown that daily high-dose oral therapy (1000 $\mu\text{g/day}$) is as effective as monthly intramuscular injections at normalizing serum cobalamin and MMA levels in patients with severe malabsorption, including those with Crohn's or celiac disease.

Emerging Evidence in Biotherapeutic and Botanical Science

Probiotics: Strain-Specific and Disease-Specific Efficacy

Probiotics are defined as live microorganisms that, when administered in adequate amounts, confer a health benefit on the host. While widely used, clinical evaluations indicate that the therapeutic benefits of probiotics are highly strain-specific and disease-specific. Pooling different taxonomic strains in meta-analyses can lead to inaccurate conclusions regarding clinical efficacy.

A systematic review of 228 clinical trials demonstrated that different strains within the same species exhibit highly variable survival rates in gastric acid, differing adherence to intestinal mucosa, and distinct immunomodulatory properties.

Probiotic candidates must undergo rigorous screening to evaluate survival through the gastrointestinal tract.

In screening studies of over 127 different *Lactobacillus* strains, only 3% were found suitable as probiotics based on their ability to resist bile and stomach acidity.

The therapeutic mechanisms of action are diverse, including the production of bacteriocins (which directly inhibit pathogens), the destruction of pathogenic toxins,

reinforcement of enterocyte tight junction integrity, and competitive inhibition of pathogen attachment to host cells.

▼

▼ ▼

▼ ▼ - Up-regulation of BDNF - Suppression of inflammatory cytokines - Production of short-chain fatty acids (IL-6, TNF-alpha)

▼ [Clinical Neuro-Modulation] - Reduced depressive symptoms (SMD = -0.53) - Reduced anxiety symptoms (SMD = -0.44)

Clinical efficacy has been established for specific strains in targeted indications.

For the prevention of adult antibiotic-associated diarrhea (AAD), robust evidence exists for *Saccharomyces boulardii* CNCM I-745 and a three-strain mixture containing *Lactobacillus acidophilus* CL1285, *L. casei* LBC80R, and *L. rhamnosus* CLR2 ($\text{Bio-K}^{\circledR}$), whereas other lactobacilli strains show no benefit.

Furthermore, *L. rhamnosus* GG and *S. boulardii* CNCM I-745 exhibit strong disease-specific efficacy in treating acute pediatric gastroenteritis and preventing recurrent *Clostridium difficile* infections.

However, a systematic review of 17 Cochrane systematic reviews on probiotics concluded that none of the evaluations provided high-quality evidence for the prevention of illnesses in the general population, with benefits limited to low-to-moderate quality evidence for reducing AAD, upper respiratory tract infections, and gestational diabetes.

Beyond gastroenterology, recent clinical interest has focused on the gut-brain axis, particularly the use of probiotics, prebiotics, and synbiotics to modulate neurological and psychiatric states.

A systematic review and meta-analysis of randomized controlled trials updated in

2025 evaluated these interventions in patients with depression, anxiety, and sleep disturbances.

The meta-analysis, which included 63 studies evaluating depressive symptoms ($n = 5,373$ participants) and 49 studies evaluating anxiety ($n = 3,912$ participants), showed significant improvements compared to placebo.

Probiotics and synbiotics significantly reduced depressive symptoms (Standardized Mean Difference $= -0.53$, $95\% \text{ CI}$, -0.67 to -0.39 , $P < 0.001$) and anxiety ($\text{SMD} = -0.44$, $95\% \text{ CI}$, -0.59 to -0.28 , $P < 0.001$).

Additionally, probiotic interventions significantly improved objective sleep quality ($\text{SMD} = -0.39$, $95\% \text{ CI}$, -0.53 to -0.25 , $P < 0.001$).

The hypothesized mechanisms include the down-regulation of pro-inflammatory cytokines, the synthesis of neuroactive short-chain fatty acids, and the up-regulation of brain-derived neurotrophic factor (BDNF).

However, because of high heterogeneity and variable methodological quality across trials, further large-scale, long-term RCTs are required to confirm these outcomes.

Turmeric and Curcuminoid Formulations in Musculoskeletal and Metabolic Health
Turmeric (*Curcuma longa*) and its primary bioactive polyphenol, curcumin, have been extensively investigated for their anti-inflammatory and antioxidant properties.

In type 2 diabetes and prediabetes, chronic systemic inflammation and oxidative stress are central to disease progression.

A systematic review and dose-response meta-analysis of 28 RCTs evaluated the effects of curcumin/turmeric supplementation on circulating inflammatory and oxidative biomarkers in this population.[span_328](start_span)[span_328](end_span)

Curcumin supplementation significantly reduced C-reactive protein (CRP; $\text{SMD} = -0.50$), tumor necrosis factor- α (TNF- α ; $\text{SMD} = -1.70$), interleukin-6 (IL-6; $\text{SMD} = -2.97$), and malondialdehyde (MDA; $\text{SMD} = -1.31$).[span_329](start_span)[span_329](end_span)

It also significantly increased glutathione (GSH; $\text{SMD} = 1.72$) and total antioxidant capacity (TAC; $\text{SMD} = 1.03$).[span_330](start_span)[span_330](end_span)

A broader, GRADE-assessed systematic review of 103 RCTs confirmed these findings, demonstrating high-quality evidence for reductions in fasting blood sugar (FBS), CRP, waist circumference, and body weight.[span_331](start_span)[span_331](end_span)

Furthermore, curcumin has shown clinical efficacy in joint and skeletal muscle health.[span_332](start_span)[span_332](end_span) A meta-analysis of 21 clinical studies—including 14 joint health and 7 skeletal muscle health trials—concluded that turmeric extract is an effective adjuvant for managing musculoskeletal pain and osteoarthritis.[span_333](start_span)[span_333](end_span)

It significantly reduced clinical scores of joint pain and stiffness, with a low incidence of adverse events.[span_334](start_span)[span_334](end_span)

Curcumin is also effective at improving hepatic function.[span_338](start_span)[span_338](end_span) In a meta-analysis of 31 RCTs, curcumin supplementation significantly reduced serum levels of alanine aminotransferase (ALT; Weighted Mean Difference $= -4.09 \text{ U/L}$, $95\% \text{ CI}$, $-6.49 \text{ to } -1.70$) and aspartate aminotransferase (AST; $\text{WMD} = -3.81 \text{ U/L}$, $95\% \text{ CI}$, $-5.71 \text{ to } -1.91$).[span_339](start_span)[span_339](end_span)

The primary clinical limitation of native curcumin is its extremely poor systemic bioavailability, which is driven by hydrophobic properties, poor water solubility, rapid intestinal metabolism, and high systemic

clearance.

To overcome this, advanced delivery systems have been developed.

Water-dispersible turmeric extract (such as WDTE60N) has demonstrated vastly superior pharmacokinetics. At a dose of \$250\text{ mg/day}\$, this dispersible formulation achieved greater joint pain reduction and systemic anti-inflammatory efficacy than standard unformulated curcuminoid doses of \$1000\text{ mg/day}\$ or more.

Regulatory Oversight, Novel Foods, and Quality Verification Standards

Comparative Oversight: FDA vs. EFSA

The regulatory classification of dietary supplements varies significantly between the United States and the European Union, directly impacting consumer safety, market access, and quality

control.

In the United States, dietary supplements are regulated under the Dietary Supplement Health and Education Act (DSHEA) of 1994, which categorizes them as a subset of foods rather than

drugs. Under this framework, manufacturers do not need to prove the safety or efficacy of a supplement before marketing

it.

Ingredients sold in the U.S. before October 15, 1994, are grandfathered under a "Generally Recognized as Safe" (GRAS) status based on historical

use. For New Dietary Ingredients (NDIs), manufacturers must submit a safety notification to the FDA 75 days before marketing, but this does not require rigorous clinical trial

validation.[span_350](start_span)[span_350](end_span)

The FDA's role is largely retrospective: the agency monitors post-market adverse event reports and must prove a product is unsafe or mislabeled before taking enforcement action to remove it from the market.[span_351](start_span)[span_351](end_span)[span_352](start_span)[span_352](end_span)](end_span)

In contrast, the European Union operates under a highly precautionary, centralized regulatory framework overseen by the European Food Safety Authority (EFSA).[span_353](start_span)[span_353](end_span)[span_354](start_span)[span_354](end_span)](end_span)

Under Directive 2002/46/EC, food supplements are regulated as foods, and the EU maintains a harmonized positive list of permitted vitamins, minerals, and their chemical sources.[span_355](start_span)[span_355](end_span)](end_span)

If a manufacturer wishes to introduce a new nutrient source or a botanical ingredient without a history of consumption in the EU prior to May 15, 1997, the substance is classified as a "Novel Food" under Regulation (EU) 2015/2283.[span_359](start_span)[span_359](end_span)[span_360](start_span)[span_360](end_span)](end_span)

This requires the submission of a comprehensive safety dossier to the European Commission, which then tasks EFSA's Panel on Nutrition, Novel Foods and Food Allergens (NDA) with conducting a rigorous pre-market risk assessment.[span_361](start_span)[span_361](end_span)[span_363](start_span)[span_363](end_span)](end_span)

The manufacturer must provide scientific proof of safety, tolerability, and systemic bioavailability.[span_362](start_span)[span_362](end_span)[span_364](start_span)[span_364](end_span)](end_span)

Additionally, EFSA establishes strict Tolerable Upper Intake Levels (ULs) based on systematic toxicological evaluations.[span_365](start_span)[span_365](end_span)[span_366](start_span)[span_366](end_span)](end_span) For example, due to the risk of sensory peripheral neuropathy, EFSA reduced the adult UL for Vitamin B6 from 30 mg/day to 12.5 mg/day.

mg/day}\$.(start_span)[span_367](end_span)

Furthermore, EU regulations prohibit health claims on supplement labels unless they are backed by strong clinical evidence and have been formally authorized by EFSA and the European Commission.

Article 8 of Regulation (EC) No 19(start_span)[span_356](end_span)25/2006 allows the European Commission to restrict or prohibit substances (adding them to Annex III) if safety concerns arise.

EFSA has evaluated several high-u(start_span)[span_357](end_span)se botanical substances due to safety concerns raised by member states, including *Ephedra* species, *Yohimbe* (*Pausinystalia yohimbe*), hydroxyanthracene derivatives, green tea catechins, monacolins from red yeast rice, and alpha-lipoic acid (associated with a risk of insulin autoimmune syndrome, or Hirata's disease).

| Nutrient/Substance | EFSA Toler(start_span)[span_358](end_span)able Upper Limit (UL) | Systemic Physiological Rationale for Limit | Identified Clinical Toxicity Risks | Source(s) |

| :--- | :--- | :--- | :--- | :--- |

| **Vitamin B6** | \$12.5\text{ mg/day}\$ (Adults) | Prevention of progressive neurotoxicity and nerve fiber degeneration. | Peripheral neuropathy, sensory loss, and ataxia. | (start_span)[span_368](end_span) |

| **Folate (Folic Acid)** | \$1000\text{ }\mu\text{g/day}\$ (Adults) | Prevention of Unmetabolized Folic Acid (UMFA) accumulation and clinical masking of Vitamin B12 deficiency. | Masked macrocytic anemia; competitive DHFR inhibition; reduced NK cell cytotoxicity. |

(start_span)[span_371](end_span)(start_span)[span_372](end_s
pan) |

| **Selenium** | \$255\text{ }\mu\text{g/day}\$ (Adults) | Risk of chronic selenosis in daily consumers of high-dose supplements or Brazil nuts. | Hair and nail loss, gastrointestinal disturbances, and peripheral neurological dysfunction. | |

| **E(start_span)[span_369](end_span)lemental Iron** | \$40\text{ mg/day}\$ (Adults) | Prevention of systemic iron overload and hepatic parenchymal accumulation. | Hemochromatosis, liver fibrosis, and severe gastrointestinal mucosal irritation. |

(start_span)[span_373](end_span)(start_span)[span_374](end_s
pan) |

| ****Manganese**** | 8 mg/day (Adults) | Prevention of central neurotoxicity and hepatic dysfunction. | Neurobehavioral changes; extrapyramidal motor symptoms resembling Parkinson's disease. | |

The Importance of Third-Party Verification

Because pre-market safety and efficacy evaluations are not universally required, the dietary supplement market has a high prevalence of mislabeling, ingredient substitution, sub-potency, and contamination.

Analytical audits have revealed that many non-verified supplements contain significantly more or less active ingredient than declared on the label. Some products are contaminated with heavy metals (lead, cadmium, mercury), pathogenic microbes, pesticides, or banned pharmacological agents.

To mitigate these clinical risks, independent third-party verification organizations have established testing standards.

The

United States Pharmacopeia (USP) Dietary Supplement Verification Program evaluates products against strict quality standards.

A USP Verified mark indicates that the supplement: contains the ingredients listed on the label in the declared potency and amount; does not contain harmful levels of specified contaminants (heavy metals, microbes, pesticides); will break down and dissolve within a specified timeframe to ensure systemic bioavailability; and has been manufactured under sanitary and well-controlled Current Good Manufacturing Practices (cGMPs).

NSF International certifies dietary supplements under the NSF/ANSI 173 standard, the only American National Standard for these

products.

NSF's program includes a comprehensive label claim review, a toxicology formulation review, and a contaminant screen.

For competitive athletes and military personnel, the "NSF Certified for Sport" and "Informed Sport" programs provide critical protection.


These programs screen every batch of certified products for more than 280 substances banned by major athletic organizations and anti-doping agencies, including anabolic steroids, stimulants, selective androgen receptor modulators (SARMs), narcotics, and diuretics.

This verification helps prevent unintentional doping violations and adverse health events from adulterated products.

Clinical Conclusions and Strategic Recommendations

The clinical trial data and biochemical evidence support a shift away from generalized, empirical dietary supplementation toward a highly targeted, biomarker-driven approach. Broad-spectrum supplementation in healthy, nutritionally replete populations shows little to no benefit for the primary prevention of cardiovascular disease, cancer, or all-cause mortality.

However, targeted interventions in specific, biologically defined subgroups demonstrate substantial clinical efficacy.



▼ ▼ ▼ [Antenatal Care]

- Target: Low-fish eaters, - Target: Older adults, especially - Target: All pregnancies African Americans - CVD history - Tool: Daily 15-Nutrient
- Tool: 1 g/day Omega-3 - Tool: Daily MVM UNIMMAP (MMS)

To optimize patient outcomes and minimize risks, the following clinical recommendations are established:

* **Subgroup-Specific Cardioprotection**: Clinicians should recommend daily marine omega-3 fatty acids (1 g/day) for individuals with low baseline fish consumption (< 1.5 servings per week) and African American patients to significantly reduce the risk of myocardial infarction.

* **Cognitive Support in Aging**: Daily multivitamin-mineral supplementation should be considered for older adults, particularly those with a history of cardiovascular disease, to preserve global memory and slow cognitive

aging.

* **Oncological Latency Considerations**: High-dose vitamin D3 (2000 IU/day)

should be recommended to reduce cancer-specific mortality over time, particularly in normal-weight individuals, while monitoring for potential hypercalcemia or nephrolithiasis risks at higher doses.

* **Precision Formulation and Speciation**: Organic magnesium forms (such as citrate, glycinate, or L-threonate) should be used instead of inorganic magnesium oxide to achieve systemic absorption and avoid gastrointestinal side effects.

Similarly, active L-methylfolate (5-MTHF) is the preferred supplemental form of folate for individuals with MTHFR polymorphisms to prevent the accumulation of unmetabolized folic acid.

* **Antenatal and Geriatric Repletion**: Multiple Micronutrient Supplementation (MMS) using the UNIMMAP formulation should replace standard iron-folic acid monotherapy in vulnerable pregnancies to improve birth weight and reduce neonatal

mortality. In older adults with atrophic gastritis or malabsorption, daily high-dose oral vitamin B12 (1000 µg/day) is highly effective at overcoming absorption deficits.

* **Pharmacovigilance of Botanicals**: Clinicians must remain alert to potent pharmacokinetic interactions, such as St. John's wort inducing the CYP3A4 and P-glycoprotein pathways, which can cause therapeutic failure of critical medications.

* **Quality Assurance Protocols**: To guarantee patient safety, clinical efficacy, and the absence of contaminants, clinicians should recommend only those dietary supplements that have undergone independent, third-party verification by

reputable organizations such as the USP, NSF, or Informed Sport.

Works cited

1. Vitamin, Mineral, and Multivitamin Supplementation to Prevent Cardiovascular Disease and Cancer - USPSTF, <https://www.uspreventiveservicestaskforce.org/home/getfilebytoken/KprBMtobbwaeMEqZzxwHCz>
2. Evidence Summary: Vitamin, Mineral, and Multivitamin ... - USPSTF, <https://www.uspreventiveservicestaskforce.org/uspstf/document/final-evidence-summary/vitamin-supplementation-to-prevent-cvd-and-cancer-preventive-medication>
3. Vitamin, Mineral, and Multivitamin Supplementation to Prevent Cardiovascular Disease and Cancer: US Preventive Services Task Force Recommendation Statement - PubMed, <https://pubmed.ncbi.nlm.nih.gov/35727271/>
4. US Preventive Services Task Force - Vitamin, Mineral, and Multivitamin Supplementation to Prevent Cardiovascular Disease and Cancer - AAFP, <https://www.aafp.org/afp/2022/1100/uspstf-supplementation-cardiovascular-disease-cancer.pdf>
5. ADVERSE EFFECTS OF NUTRACEUTICALS AND DIETARY ..., <https://pmc.ncbi.nlm.nih.gov/articles/PMC6380172/>
6. Principal Results of the VITamin D and OmegA-3 Trial (VITAL) and ..., <https://pmc.ncbi.nlm.nih.gov/articles/PMC7089819/>
7. Vitamin D and Omega-3 Trial - VITAL - American College of Cardiology, <https://www.acc.org/latest-in-cardiology/clinical-trials/2018/11/08/22/42/vital>
8. Principal results of the VITamin D and OmegA-3 Trial (VITAL) and updated meta-analyses ... - PubMed,

<https://pubmed.ncbi.nlm.nih.gov/31733345/> 9. The VITAL trial: Omega-3 supplementation does not reduce risk of cardiovascular events, <https://www.2minutemedicine.com/the-vital-trial-omega-3-supplementation-does-not-reduce-risk-of-cardiovascular-events/> 10. Vital Insights | Harvard Medical School, <https://hms.harvard.edu/news/vital-insights> 11. Findings about the Vital Study, <https://www.vitalstudy.org/findings.html> 12. VITAL: No Benefits to Vitamin D and Omega-3s in Reducing Major CV Events, Cancer, <https://www.tctmd.com/news/vital-no-benefits-vitamin-d-and-omega-3s-reducing-major-cv-events-cancer> 13. VITAL researcher JoAnn Manson outlines findings on vitamin D, omega-3 - Harvard Gazette, <https://news.harvard.edu/gazette/story/2018/11/vital-researcher-joann-manson-outlines-findings-on-vitamin-d-omega-3/> 14. The VITAL trial's findings on fish oil & vitamin D - Center for Science in the Public Interest, <https://www.cspi.org/article/vital-trials-findings-fish-oil-vitamin-d> 15. Results – COSMOS Trial, <https://cosmostrial.org/results/> 16. Cocoa and Multivitamins Could Slow Down Cognitive Decline - Psychiatrist.com, <https://www.psychiatrist.com/news/cocoa-and-multivitamins-could-slow-down-cognitive-decline/> 17. Can a daily multivitamin supplement slow cognitive aging?, <https://www.alzdiscovery.org/cognitive-vitality/blog/can-a-daily-multivitamin-supplement-slow-cognitive-aging> 18. Effect of multivitamin-mineral supplementation versus placebo on cognitive function: results from the clinic subcohort of the COcoa Supplement and Multivitamin Outcomes Study (COSMOS) randomized clinical trial and meta-analysis of 3 cognitive studies within COSMOS - PubMed, <https://pubmed.ncbi.nlm.nih.gov/38244989/> 19. Active Folate Versus Folic Acid: The Role of 5-MTHF (Methylfolate) in Human Health - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9380836/> 20. Folate - Health Professional Fact Sheet - Office of Dietary Supplements (ODS) - NIH, <https://ods.od.nih.gov/factsheets/Folate-HealthProfessional/> 21. What is the Difference Between Folate, Folic Acid, Folinic Acid, and Methylfolate?, <https://autism.fratnow.com/blog/what-is-the-difference-between-folate-folic-acid-folinic-acid-and-methylfolate/> 22. Vitamin B12 - Health Professional Fact Sheet, <https://ods.od.nih.gov/factsheets/VitaminB12-HealthProfessional/> 23. Pharmacokinetic interactions of drugs with St John's wort - PubMed, <https://pubmed.ncbi.nlm.nih.gov/15260917/> 24. Warfarin, St John's wort and INR - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4917631/> 25. International expert consensus on micronutrient supplement use during the early life course, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11744953/> 26. WHO recommendations on antenatal nutrition: an update on multiple micronutrient supplements | BMJ Global Health, <https://gh.bmj.com/content/5/7/e003375> 27. Micronutrient supplementation

in pregnancy: Who, what and how much? - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC6416688/> 28. New Evidence Should Inform WHO Guidelines on Multiple Micronutrient Supplementation in Pregnancy - PubMed, <https://pubmed.ncbi.nlm.nih.gov/30773589/> 29. The efficacy of probiotics, prebiotics, and synbiotics on anxiety, depression, and sleep: a systematic review and meta-analysis of randomized controlled trials - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12751681/> 30. Efficacy of curcumin/turmeric on inflammation and oxidative stress in prediabetes and type 2 diabetes: a systematic review and dose-response meta-analysis - PubMed, https://pubmed.ncbi.nlm.nih.gov/41240262/?utm_source=FeedFetcher&utm_medium=rss&utm_campaign=pubmed-2&utm_content=1V7FV1QUbN6QMTcjFUT8hnah6x0O5Uat4IfPBNJ19cGnPYygeO&fc=20230328041118&ff=20251115161946&v=2.18.0.post22+67771e2 31. Antioxidant and anti-inflammatory effects of curcumin/turmeric supplementation in adults: A GRADE-assessed systematic review and dose-response meta-analysis of randomized controlled trials - PubMed, <https://pubmed.ncbi.nlm.nih.gov/36804260/>