

# BEYOND THE SEVEN-EIGHTHS BARRIER: A SHARPER QUASI-RIEMANN BOUNDARY

ARGONAUT MATH

**ABSTRACT.** We refine the zero-free boundary in OpenAI’s September 30 quasi-Riemann paper by slightly increasing the total length of the compensation primes and adjusting the two column scales. From that paper’s stated unconditional results, we obtain the strict zero-free half-plane  $\operatorname{Re} s > 3499999/4000000$  for finite-order Hecke  $L$ -functions over  $\mathbb{Q}(\sqrt{-3})$  and for all Dirichlet  $L$ -functions. This improves the boundary  $7/8$  by  $1/4000000$ . The arithmetic and analytic estimates still concern the same finite probe. On the arithmetic side, its rescaling coefficients absorb the extra completed-row energy. On the analytic side, an explicit margin in the central-row estimate remains positive. We give the affected counting, contour, and principal-term arguments in full. The parameters are chosen so that the moment mesh is fixed independently of the later principal approximation. The foundational reflection, sieve, and moment theorems are taken from the original paper; no new short mean-square assumption is imposed.

## 1. INTRODUCTION

**1.1. Historical background and main result.** Riemann’s 1859 paper connected the distribution of primes to the zeta function through analytic continuation and a functional equation. Its discussion of the zeros led to the critical-line hypothesis that now bears his name [2, English translation, pp. 2–4]. The present argument concerns a zero-free half-plane for a family of  $L$ -functions; it does not reach the critical line.

This paper refines the parameters in OpenAI’s September 30 argument as part of the Argonaut Math research program. That argument proves a zero-free half-plane with boundary  $7/8$ . It uses a finite arithmetic probe with two representations: one admits a sieve estimate, and the other produces an analytic signal. We show that the construction permits a small increase in the length of the compensation primes. Adjusting the column scales to accommodate that increase gives a strictly smaller boundary.

We work over  $F = \mathbb{Q}(\sqrt{-3})$ , with ring of integers  $\mathcal{O}_F$  and ideal norm  $N\mathfrak{a} = |\mathcal{O}_F/\mathfrak{a}|$ . A finite-order Hecke character  $\eta$  is extended by zero to ideals not coprime to its modulus. For  $\operatorname{Re} s > 1$  its  $L$ -function is

$$L_F(s, \eta) = \prod_{\mathfrak{p}} (1 - \eta(\mathfrak{p})(N\mathfrak{p})^{-s})^{-1}.$$

We use its usual meromorphic continuation. Removing the factors at a fixed finite set  $S$  of prime ideals gives  $L_F^S(s, \eta)$ . The corresponding finite factors are nonzero for  $\operatorname{Re} s > 0$ . As usual, a principal character may have a pole at  $s = 1$ .

**Theorem 1.1.** *All finite-order Hecke  $L$ -functions over  $F$ , and all Dirichlet  $L$ -functions, are zero-free in the strict half-plane*

$$\operatorname{Re} s > \frac{3499999}{4000000}.$$

*The pole at  $s = 1$  for a principal character is allowed. The boundary is  $1/4000000$  smaller than  $7/8$ ; no assertion is made about zeros on the boundary line.*

The deduction uses the results stated and proved unconditionally in [1]; it requires no additional analytic conjecture. Theorem 1.1 of that paper supplies the existing  $7/8$  bound. We also use Propositions 2.1, 11.3, and 15.2; Lemmas 10.2, 11.1, 13.1, 14.3, 17.1, 17.6, 18.1, and 19.1; and the coefficient, reflection, and zero-detection results of Sections 4–8. When the perturbation changes a numerical range, we derive the needed estimate from the general energy or moment bounds and the exact local formulas in that paper. The foundational reflection, sieve, and moment proofs are not reproduced here. Page references to [1] follow its printed PDF pagination; references to its equations retain the original numbering. The result here does not give a zero-free boundary below  $3/4$ .

---

Date: October 7, 2026.

Let  $Z$  tend to infinity. Write  $\ell$  for the total length of the compensation primes, and use  $l_x, l_y$  to specify the column scales  $X = Z^{l_x}$  and  $Y = Z^{l_y}$ . These exponents have sum  $M$ , and the second exceeds the first by  $b$ . The exponent  $h$  determines the upper end of the central-row range before a small buffer is added;  $\sigma_\tau$  is the proposed zero-free boundary. Increasing  $\ell$  shortens both columns and shifts that upper end. We fix the parameters as follows:

$$\begin{aligned} \tau &= 10^{-6}, & \ell &= \frac{1}{6} + \tau, & M &= \frac{5}{6} - \tau, & b &= \frac{1}{8}, \\ l_x &= \frac{17}{48} - \frac{\tau}{2}, & l_y &= \frac{23}{48} - \frac{\tau}{2}, & h &= \frac{13}{16} + \frac{3\tau}{2}, \\ \sigma_\tau &= \frac{7}{8} - \frac{\tau}{4} = \frac{3499999}{4000000}. \end{aligned} \tag{1}$$

Thus

$$l_y - l_x = b, \quad l_x + l_y = M, \quad M + \ell = 1, \quad h = 1 - l_x + \ell.$$

For the two column scales take  $X = Z^{l_x}$  and  $Y = Z^{l_y}$ . The adjustment leaves the affine exponent of the analytic signal unchanged:

$$C(s) = \frac{l_x}{2} + s - 1 + \frac{h}{6} = s - \frac{11}{16}, \quad C(\sigma_\tau) = \frac{3}{16} - \frac{\tau}{4}. \tag{2}$$

A rightmost zero need not exist. We therefore measure the whole family by the supremum

$$\beta_* = \sup \left( \{1/2\} \cup \{\operatorname{Re} \rho : L_F(\rho, \eta) = 0, 1/2 \leq \operatorname{Re} \rho \leq 1, \eta \text{ primitive of finite order}\} \right).$$

The supremum need not be attained. Throughout the proof we suppose, for a contradiction, that  $\beta_* > \sigma_\tau$ , and put  $\Delta_\tau = \beta_* - \sigma_\tau > 0$ . The existing result gives

$$\sigma_\tau < \beta_* \leq \frac{7}{8}, \quad \kappa = \frac{3}{4}, \quad \beta_* \leq \frac{1 + \kappa}{2}. \tag{3}$$

Thus the fixed value  $\kappa = 3/4$  is admissible in Lemma 18.1 of [1], whose uniform range is  $\kappa \in [3/4, 1]$ . This choice uses the existing bound, which is available before we perturb the parameters.

**1.2. Outline of the proof.** The proof compares two estimates for the same finite arithmetic probe. The low-side estimate bounds its size by reflected-row energy and an additive Gram bound. The rescaling coefficients absorb the cost of the longer compensation primes. The high-side representation separates the principal row from the remaining rows. After normalization, the principal row gives the signal in (18), whose integrand contains  $H_\eta(s)/L_F^S(s, \eta)$ . The reciprocal  $L$ -function is the link between continuation of this signal and the location of zeros, as expressed by Proposition 2.1 of [1].

Sections 4 and 5 bound the nonprincipal contribution. The witness capacities give a row count with an explicit margin. A joint estimate for the numerator and error slots preserves that margin after the full contour is moved. Section 6 extracts the principal signal and bounds the outer rows. Section 7 then chooses the parameters without circular dependence and combines the two estimates needed by the continuation principle to exclude the gap  $\Delta_\tau$ .

Three points have to remain compatible throughout the perturbation. Both estimates concern the same finite probe, including its phases and coprimality restrictions. In the analytic representation, the complete Euler correction is moved before the retained integral is subdivided by amplitude, and the numerator is estimated jointly with every error slot. The resulting estimates bound the normalized probe and its difference from the principal signal, respectively; the low-side bound is established in Section 3. Finally, the moment mesh is fixed first, then the number of prime slots, and then the principal approximation saving. The height and external decay orders are chosen only after the earlier real savings. The proof below supplies these checks at the altered lengths.

**1.3. AI disclosure.** AI assistance was used in developing the argument. The AI for Math plugin supported code generation for the computations and Lean formalization. The mathematical content of the paper is the deduction from the stated literature inputs; the collaboration does not enlarge those inputs or the theorem's scope.

The formalization of the three terminal nonvanishing statements—for the Riemann zeta function, Dirichlet L-functions, and the Hecke family considered here—is complete. It uses the imported foundational results from the original project and retains the stated strict half-plane and the applicable principal-pole exceptions. It has passed compilation with Lean 4.34.1 and verification with Comparator’s native core.

**1.4. Acknowledgements.** This work developed through human–AI dialogue. Human questions and hints guided the choice of the small parameter perturbation and the requirement that both estimates concern the same finite probe.

## 2. ONE FINITE PROBE AND ITS EULER REPRESENTATION

We use the finite weighted arithmetic sum  $I_\eta(X, Y, Z)$  defined in (6.1) of [1], with column scales  $X, Y$  and large parameter  $Z$ . Its arithmetic form gives the low-side bound. Its Euler representation isolates the principal signal and allows us to estimate the other terms. In the Euler representation,  $u$  indexes the physical rows, and  $u = 1$  is the principal row.

A compensation-prime slot selects one prime from a prescribed window. The modification combines divisibility conditions at the selected primes with rescalings of the columns. Both estimates must apply to exactly this finite sum, including its conjugate phases and coprimality restrictions. We first define the modification as an arithmetic sum, then derive its integral representation.

Fix a quotient  $T$  of a ray class group before choosing a target character. For each target  $\eta$ , use fixed arithmetic data  $S, b_*, \xi$  allowed by Sections 6 and 13 of [1]; here  $b_*$  is a fixed ideal, and  $S$  contains the relevant moduli and small primes. Also fix the nonnegative, nonzero smooth annular weights  $W_0, W_1$  and the Fourier normalization of Section 6 of that paper. Denote its finite physical probe (6.1) by  $I_\eta(X, Y, Z)$ . Inserting  $1_{D|cn^3}$  into its completed rows gives  $I_{\eta;D}$ . The ideals  $c, n$  retain the original coprimality conditions and the characters retain their zero extension. These definitions specify the finite arithmetic expressions in (6.1). The integral below is a representation of those expressions.

Choose a fixed even integer  $K$  and split the compensation length equally:  $\ell_i = \ell/K$  and  $P_i = Z^{\ell_i}$ . Choose  $K$  pairwise disjoint compact supports in  $(1, 2)$ , separated by positive gaps, and nonnegative, nonzero smooth weights  $W_i$  on them. The prime window is

$$\mathcal{P}_i(Z) = \{\mathfrak{p} \notin S : \mathfrak{p} \in 1_T, W_i(N\mathfrak{p}/P_i) \neq 0\}.$$

The condition  $\mathfrak{p} \in 1_T$  means that its ray class is the identity. For  $J \subseteq \{1, \dots, K\}$  write  $p_J = \prod_{i \in J} \mathfrak{p}_i$ , with the empty product equal to the unit ideal. The modified probe is the same finite expression as (12.5) of [1], at the new scales:

$$\begin{aligned} I_{\eta, \text{mod}}(Z) &= \sum_{(\mathfrak{p}_i) \in \prod_i \mathcal{P}_i(Z)} \prod_i W_i(N\mathfrak{p}_i/P_i) \\ &\quad \times \sum_{J \subseteq \{1, \dots, K\}} (-1)^{|J|} (Np_J)^{-3/2} \overline{\eta(p_{J^c})} \\ &\quad \times I_{\eta; p_J^c}(X/Np_J, Y/Np_J, ZNp_J^c). \end{aligned} \tag{4}$$

Each window retains its original scale  $P_i$ . The conjugate in (4) pairs the marked contribution with the rescaled contribution in the local compensation. Identity ray class makes the characters in  $\widehat{T}$  equal to 1 at that prime; it does not make  $\eta(\mathfrak{p})$  equal to 1. The target may have any unit phase there.

For the Euler representation use the local data  $P_{\mathfrak{p}}, P_{\mathfrak{p}}^*, H_{\mathfrak{p}}, V, W, D$  of (7.11)–(7.12) in [1], and denote the complex variable called  $x$  in its Section 16 by  $s$ . At  $Q = N\mathfrak{p}$  the complete local replacement is

$$G_{\mathfrak{p}} = \frac{(\overline{\eta(\mathfrak{p})} Q^s - Q^{-w})(1 - V)(1 - W)P_{\mathfrak{p}}^* - Q^{-w}(1 - VW)}{1 - D}. \tag{5}$$

Equation (16.4) of that paper reads

$$\overline{\eta(\mathfrak{p})} Q^{s+z-1} P_{\mathfrak{p}}^* - Q^{z-w-1} P_{\mathfrak{p}} = \frac{1 - D}{(1 - V)(1 - W)} Q^{z-1} G_{\mathfrak{p}}.$$

Consequently the full correction is

$$H_{\eta,u,Z}(s, w, z) = \sum_{(\mathfrak{p}_i) \in \prod_i \mathcal{P}_i(Z)} \prod_i [W_i(N\mathfrak{p}_i/P_i)(N\mathfrak{p}_i)^{z-1} G_{\mathfrak{p}_i}] \times \prod_{\substack{\mathfrak{p} \notin S \\ \mathfrak{p} \notin \{\mathfrak{p}_1, \dots, \mathfrak{p}_K\}}} H_{\mathfrak{p}}. \quad (6)$$

The local factors are evaluated at the current physical row  $u$ . Formula (5) remains defined at zeros of  $P_{\mathfrak{p}}$  or  $H_{\mathfrak{p}}$ , so it also provides the continuation of the complete correction. Disjoint supports ensure that each selected prime is replaced exactly once, even though the nominal scales  $P_i$  are equal.

To control the integral variables, we use the decay supplied by the smooth weights. Let  $\tilde{W}_0(|y|^2)$  be the plane Fourier transform of  $W_0(|y|^2)$  for the self-dual measure of [1], and set

$$\mathcal{M}(z) = \int_0^\infty \tilde{W}_0(r) r^{z-1} dr, \quad \widehat{W}_1(w) = \int_0^\infty W_1(r) r^{w-1} dr.$$

The resulting weight is

$$\mathcal{W}(X, Y, Z; s, w, z) = X^{1/2-z} Z^{s+z-1} Y^{w-1} e^{(s+z-1)^2} \mathcal{M}(z) \widehat{W}_1(w).$$

Equations (7.5), (7.13), and the finite replacements give the exact identity

$$I_{\eta, \text{mod}}(Z) = \frac{1}{(2\pi i)^3} \int_{\text{Re } s=3} \int_{\text{Re } w=3} \int_{\text{Re } z=2} \mathcal{W}(X, Y, Z; s, w, z) \times \sum_u^{(6)} (Nu)^{-z} \overline{\xi(u)} \frac{\zeta_F^S(6z) L_F^S(w, \chi_\bullet(u))}{L_F^S(s, \eta \chi_\bullet(u))} H_{\eta,u,Z}(s, w, z) dz dw ds. \quad (7)$$

Here (6) preserves the original physical restrictions of (7.5): nonzero sixth-power-free elements, the chosen unit sector, and  $(u, S) = 1$ . The ideal character  $\chi_\bullet(u)$  is the one in Lemma 4.1 of [1]. Absolute convergence on the initial lines follows from the absolutely convergent expansions of the original probe: only finitely many such expansions are combined here. For each fixed  $Z$ , the slots, prime tuples, and rescaling subsets are finite. The completed sum over  $u$ , the Euler products, and the three Mellin integrals are infinite. The interchanges are justified by the absolute convergence used in (7.5) of [1]. The fixed-tuple replacements form a finite collection and preserve that convergence. Subsequent contour moves require the full-row majorants proved in Sections 5 and 6. No local identity uses  $\ell = 1/6$ . Thus (7) is still an exact high-side representation of the finite probe (4) at the perturbed geometry.

### 3. THE COST OF COMPENSATION ON THE LOW SIDE

Increasing the prime length changes the energy of the completed rows. We therefore return to the general reflection estimate, whose dependence on the lengths is explicit. Combining it with the additive Gram estimate and the coefficient in (4) will show that the rescaling absorbs the extra energy.

Fix a rescaling subset  $J$  and write

$$d_J = \sum_{i \in J} \ell_i, \quad \ell' = \ell - d_J, \quad M' = M - 2d_J,$$

$$X' \asymp Z^{\ell_x - d_J}, \quad Y' \asymp Z^{\ell_y - d_J}.$$

Then  $0 \leq d_J \leq \ell$  and  $M' + \ell' - 1 = -3d_J$ . For a real number  $t$ , write  $t_+ = \max\{t, 0\}$ .

**Lemma 3.1.** *The completed-row factor  $B_{m,\sigma}^J$  in (15.1) of [1], where  $\sigma \in T$  is the finite ray class label, satisfies*

$$\sum_{0 < Nm \ll Q} |B_{m,\sigma}^J(Z)|^2 \ll Z^{M' + (5\ell - 1 + d_J)_+ / 4 + \varepsilon}, \quad Q \asymp Z^{M'}.$$

*The constant may depend on the fixed windows and arithmetic data, and is uniform in the moving rescaling prime tuple.*

*Proof.* Apply the general reflected-energy estimate of Lemma 14.3 in [1]. It decomposes the actual row into a powerful part, a part with fixed support, and a remaining square-free part. Let  $O$  be the powerful-part length and  $H$  the remaining-row length, and put  $\Delta_H = M' - O - H \geq -O(1/\log Z)$ . The Gaussian subdivisions, common-profile separation, and tail truncation on pp. 102–103 of that paper leave the length identity unchanged. Denote their arbitrarily small positive loss by  $\varepsilon_0$ .

In the notation of Lemma 14.3 we have  $2A_0 \leq O + O(1/\log Z)$ ,  $N_0 \leq A_0$ , and  $z_a \leq \ell'$ , with

$$T_{d_J} = 2H + 2A_0 + 2z_a - 1 - \ell' - \theta_N - N_0 - 3B_0.$$

The identity  $M' + \ell' - 1 = -3d_J$  gives  $T_{d_J} \leq H - 3d_J + O(\varepsilon_0)$ . The retained dual scales satisfy  $\nu + 3\ell_b + e_\lambda \leq T_{d_J} + \varepsilon_0$ , so

$$\max(H, \nu + \ell_b) = H + O(\varepsilon_0).$$

The local parameter called  $u$  in that energy lemma is  $\min\{\nu, z_a, (\nu + z_a)/3\}$ ; it is independent of the physical-row notation in (7). If this minimum is  $z_a$ , the energy is at most  $M' - \Delta_H + O(\varepsilon_0)$ . Otherwise  $u \geq \nu/2 - O(\varepsilon_0)$  and

$$u + \ell_b + \frac{2e_\lambda}{3} + \frac{(T_{d_J} - \nu - 3\ell_b - e_\lambda)_+}{2} \geq \frac{T_{d_J}}{4} - O(\varepsilon_0).$$

Substitute this in (14.14), using the nonnegative second numerator in (15.3). In this second branch, the energy is bounded by

$$M' + \frac{1 + 3\ell' - 2M' - 2\Delta_H}{4} + O(\varepsilon_0).$$

Both bounds decrease, or stay constant, as  $\Delta_H$  increases. For an upper bound, we may therefore replace it by zero and absorb the boundary error. Since  $1 + 3\ell' - 2M' = 5\ell - 1 + d_J$ , the two branches give the asserted positive-part bound. The common-profile kernel amplitude is extracted before squaring. Powerful-row counting is already included in Lemma 14.3, and its kernel saving is evaluated on the actual dual support. Small losses, finitely many subsets, and logarithmically many actual scales fit the stated  $\varepsilon$ .

The change in lengths also affects the endpoint. When  $d_J = \ell$ ,  $\ell' = z_a = 0$  and

$$T_{d_J} \leq 1 - 6\ell + O(\varepsilon_0) = -6\tau + O(\varepsilon_0).$$

A nonempty retained dual block has length exponent at least  $-O(1/\log Z)$ . Choose the truncation loss sufficiently smaller than  $\tau$ , then take  $Z$  sufficiently large. The two length bounds are then incompatible, so the retained block is empty. The discarded part is controlled by the uniform tail estimate in Lemma 14.3, while the first branch also gives the claimed bound. This handles the new zero-length endpoint directly.  $\square$

**Lemma 3.2.** *For every  $\varepsilon > 0$ ,*

$$|I_{\eta, \text{mod}}(Z)| \ll Z^{C(\sigma_\tau) + \varepsilon}.$$

*Proof.* Set  $P_a = Y'^2/Q = (Nb_*)^{-1}Z^b$ . The scales for a fixed tuple satisfy  $Q, Y' \geq 1$  and  $P_a \geq 1$ : indeed

$$M - 2\ell = \frac{1}{2} - 3\tau > 0, \quad l_y - \ell = \frac{5}{16} - \frac{3\tau}{2} > 0.$$

The additive Gram bound of Proposition 15.2 in [1] permits these general polynomial scales. Moreover,

$$2b - l_y + d_J \leq -\frac{1}{16} + \frac{3\tau}{2} < 0.$$

Hence its  $P_a^2/Y'$  term is absorbed by the  $P_a^{1/6}$  term, and the squared Gram norm is at most  $Z^\varepsilon(Q/Y')P_a^{1/6}$  times a fixed height polynomial.

Cauchy–Schwarz with Lemma 3.1, including the  $Q^{-1/2}$  factor in the original separation formula, gives the fixed-tuple exponent

$$\frac{l_x - d_J}{2} + \frac{b}{12} + \frac{(5\ell - 1 + d_J)_+}{8}.$$

Summing over the rescaling tuples costs at most  $Z^{d_J + \varepsilon}$ . The coefficient in (4) contributes  $Z^{-3d_J/2}$ . Combining these with the fixed-tuple estimate gives the exponent

$$L - d_J + \frac{(5\ell - 1 + d_J)_+}{8}, \quad L = \frac{l_x}{2} + \frac{b}{12} = C(\sigma_\tau).$$

Because  $\ell < 1/5$ , the function  $-d_J + (5\ell - 1 + d_J)_+/8$  starts at zero and is strictly decreasing; its two slopes are  $-1$  and  $-7/8$ . Every subset therefore has exponent at most  $L$ . Rapid decay of the smooth weights absorbs the fixed height polynomials, and the total small loss can be chosen below the prescribed  $\varepsilon$ .  $\square$

#### 4. WITNESS CAPACITY AND THE CENTRAL MARGIN

For the high-side estimate, we count rows by their norm, rather than by the rescaling subsets used in the preceding section. Write  $Nu \asymp U = Z^d$ , where  $d$  is the row-norm exponent, and fix  $d_{\min} = 1/100$ . In a zero-detection box,  $R$  denotes the ideal row-count exponent. The bound  $\#\mathcal{B} \ll_\eta U^{R+\varepsilon_{\text{cnt}}}$  must also account for the moment, witness, rounding, logarithmic, and height losses. Those losses belong to the allowance  $\varepsilon_{\text{cnt}}$ , which we keep separate from  $R$ .

The zero-detection argument in the source supplies inverse and ordinary witnesses. Their moment estimates give two bounds for the row count, both of which can be strengthened using the compensation primes already in the probe. Each moment estimate permits a certain total prime-slot length; we call this its capacity. A selection of slots must respect that capacity and have phases compatible with the witnesses. We arrange the phases first, then compare the two counts where their bounds cross.

As in Section 6.1 of [1], put  $\Theta = \langle \eta, \widehat{T} \rangle$ . Within each target, this finite character group and every slot coefficient are fixed independently of the moving row. The direction of the witnesses is handled in two steps, following Section 19.2, pp. 182–183. First, if necessary, conjugate the two witnesses and their smooth profiles so that their common character is

$$\psi(n) = \nu(n) \overline{\chi_n(u)}, \quad \nu \in \Theta.$$

Relative to this character the physical prime-slot coefficient is

$$\overline{\nu(\mathfrak{p})} 1_{\mathfrak{p} \in 1_T} = \frac{1}{|T|} \sum_{\vartheta \in \widehat{T}} (\overline{\nu} \vartheta)(\mathfrak{p}).$$

It is a fixed combination of  $\Theta$ -characters, and the marked moment permits this negative direction. For the positive-direction ordinary moment in Lemma 18.1, conjugate the entire product of the two ordinary witnesses and all selected physical slots. Its absolute square is unchanged; the common character becomes  $\overline{\nu}(n) \chi_n(u)$  and the slot coefficient becomes  $\nu(\mathfrak{p}) 1_{\mathfrak{p} \in 1_T}$ . We conjugate the whole product, rather than change the phase of an individual factor.

These two conjugations preserve the zero extension, the original prime support, and the independence of the profiles. Since  $\Theta$  is a group, membership of the induced character in  $\Theta$  is unchanged. Every retained large row ramifies at a prime outside  $S$ , so its induced character is outside  $\Theta$ , as Lemma 18.1 requires when the slot length is positive. The witness heights may differ from the physical heights, so their parameters require a separate bound. After fixing the physical height and the selected slots, Lemma 4.5 handles the witness smooth parameters at the cost of a fixed height power, uniform in the moving rows for the fixed arithmetic datum. Indeed the physical normalized slot is

$$P_i^{-1/2} \sum_{\mathfrak{p} \in \mathcal{P}_i(Z)} \overline{\chi_{\mathfrak{p}}(u)} W_i(N\mathfrak{p}/P_i) (N\mathfrak{p}/P_i)^{z-1}.$$

Its annular profile is  $W_i(y) y^{z-1}$  and its base- $U$  length is  $\ell_i/d$ , since  $U^{\ell_i/d} = P_i$ . The character expansions above identify exactly this polynomial with the prime factors in Lemmas 17.1 and 18.1. The disjoint supports are those fixed in (4), before the original zero masks are imposed. Equal nominal lengths do not merge those supports. Thus the cited moment inputs apply to the actual windows and profiles of this probe.

In a nonprincipal dynamic zero-detection box let  $a \leq \beta_*$  be its real-part parameter, and put  $\delta = 2a - 1$ . Away from the bottom box,

$$\frac{1}{50} < \delta \leq \frac{3}{4};$$

the bottom box uses  $a = 51/100$ . Lemma 19.1 supplies an average prime-amplitude parameter  $q$  satisfying  $0 \leq q \leq \delta/2$ . Here  $q$  is a dimensionless exponent, not a prime norm. For a nonbottom box put  $\alpha = 5/6$  and

$x = q/\delta \in [0, 1/2]$ . The crossing of the two witness counts is described by

$$\begin{aligned} D_x &= 3 - \frac{17x}{9}, & P_x &= \left(2 - \frac{8x}{9}\right)(1-x), \\ J &= (\alpha - \delta)D_x + \delta P_x, \\ R_* &= 1 - \delta + \frac{(\alpha - \delta)\delta P_x}{2J}. \end{aligned} \tag{8}$$

We have  $D_x \geq 37/18$ , and  $J$  has a positive lower bound on this compact parameter range. The letter  $J$  here is a real denominator; it is unrelated to the rescaling subset in (4).

**Lemma 4.1.** *Put  $c_{\text{sup}} = 1/185$ . Suppose that the available base- $U$  prime length is at least  $7/37 + c_{\text{sup}}$ . Fix positive allowances  $\varepsilon_m, \varepsilon_w, \nu_0, \omega_{\text{slot}}$  with*

$$\varepsilon_w \leq 10^{-3}, \quad \max\{\varepsilon_w, \nu_0, \omega_{\text{slot}}\} < c_{\text{sup}}/16.$$

*Require  $\omega_{\text{slot}}$  to be no larger than the mesh supplied by Lemma 18.1 of [1] for loss  $\varepsilon_m$  and the bounded length ranges here, and require  $\max_i \ell_i/d \leq \omega_{\text{slot}}$ . For each target with its fixed arithmetic data, final excluded set, and fixed annular windows, there exist finite  $A_{\text{cnt}, \eta} \geq 0$ , an integer  $L \geq 0$ , and a constant  $C_\eta > 0$  such that the following raw bound holds in every retained nonbottom dynamic and amplitude bin  $\mathcal{B}$  under the detector's loss and height conditions:*

$$\#\mathcal{B} \leq C_\eta (1 + \log U)^L (1 + H)^{A_{\text{cnt}, \eta}} U^{R_* + \varepsilon_m + 4\varepsilon_w + 6\nu_0 + \omega_{\text{slot}}}. \tag{9}$$

*Here  $H \geq 1$  bounds the retained physical and witness norm-twist parameters,  $\kappa = 3/4$ , and  $1/50 < \delta \leq 3/4$  as above. The constants and orders may depend on the fixed arithmetic datum, windows, slot count, strict moment margins, and prescribed losses, but not on  $U, H$ , the moving rows or masks, the retained point, or any later external tail order. They are fixed before a trial exponent in  $T_1 = Z^\theta$  is chosen; the sufficiently large threshold may depend on that exponent. Without selecting any slots,  $t = 1$  gives the same type of raw bound with  $R_*$  replaced by  $1 - 2\delta/3$  and error  $\varepsilon_m + 4\varepsilon_w$ , without a prime-supply requirement.*

*Proof.* The estimates below combine the crossing of the two short-witness counts with the amplified no-slot bound. We choose the common witness-length parameter to balance their exponents. For the selected slots, we use the capacity calculation in the proof of Section 19.2. Proposition 19.2 itself is specialized to the old positive gap, so its statement cannot be applied directly here. The inverse witness uses Lemma 17.1, and the ordinary witness uses Lemma 18.1; the latter's requirement  $\beta_* \leq (1 + \kappa)/2$  holds by (3).

Choose  $t \in [1, 3/2]$  as below, before selecting the witnesses. Proposition 8.3 of [1] supplies a common presentation and actual lengths  $(r, m)$ . Write  $M_r$  for its normalized inverse witness and  $S_m$  for its normalized ordinary witness. Its saturation error constants are absolute on these bounded ranges. Reduce its preliminary loss using those constants and then increase the threshold to absorb the  $O(1/\log U)$  errors. This gives

$$\begin{aligned} r + m &\geq t - \varepsilon_w, & r &\leq t + \varepsilon_w, & r &\geq t - 1/2 - \varepsilon_w, & 0 &\leq m \leq 1/2 + \varepsilon_w, \\ |M_r|^2 &\gg_\eta U^{\delta r - \varepsilon_w}, & |S_m|^2 &\gg_\eta U^{\delta m - \varepsilon_w}. \end{aligned}$$

For example, if  $C_*$  bounds all its saturation error constants, take the preliminary loss at most  $\varepsilon_w/[2(C_* + 1)]$ ;  $C_*$  precedes the target. The detector's literal height restrictions are verified in Section 7.

Order the slots by decreasing nonnegative amplitude exponent, and select only positive-exponent slots. If the request is  $z$ , a fractional selection has weighted amplitude at least  $qz$ : if all positive slots are exhausted, their full amplitude is  $q\ell/d \geq qz$ . Discarding the last fractional slot loses at most  $\delta\omega_{\text{slot}}$  in the squared exponent. Before rounding, reduce a positive moment capacity  $c$  to  $z = c - \nu_0$ ; the total loss relative to  $2qc$  is at most

$$2q\nu_0 + \delta\omega_{\text{slot}} \leq \nu_0 + \omega_{\text{slot}}.$$

The actual whole-slot length is at most  $z$ , even when there are too few positive slots. Zero-exponent and error slots are used only through upper bounds. Use the two direction changes above: first put witnesses and physical slots in the common negative direction, then conjugate the complete ordinary product. The respective coefficients are  $\bar{\nu}(\mathbf{p})1_{\mathbf{p} \in 1_r}$  and  $\nu(\mathbf{p})1_{\mathbf{p} \in 1_r}$ , both independent of the moving row. Zero extensions, independent windows, and the full-support conditions are retained.

The ordinary-witness capacity is exactly

$$\frac{1-2m}{6\kappa} = \frac{2(1-2m)}{9}.$$

At  $\kappa = 3/4$  it equals the baseline capacity used in the crossing calculation of Section 19.2. There is no need for the  $\Delta/4$  inflation introduced there by  $\kappa = 3/4 + 2\Delta$ . Here  $Q$  denotes the product of the selected normalized prime polynomials, in the common character direction just fixed. The marked moment at effective row width one bounds  $\sum_u |M_r Q|^2$  by  $U^{1+\varepsilon_m}$  times its fixed height factor when  $r + 2z \leq 1 - c_1$  and  $2r + 8z \leq 3 - c_2$ . The ordinary moment bounds  $\sum_u |S_m^2 Q|^2$  by the same power when  $2m + 6\kappa z \leq 1$ ; its divisor is  $|S_m|^4$ . At zero slot length its assertion holds for all the bounded  $m$  here.

If  $r \geq 1$ , the no-slot sixth-power amplified moment of Lemma 17.6, equation (17.88), gives exponent

$$1 - \alpha + (\alpha - \delta)r + \varepsilon_m + \varepsilon_w \leq L(t) + \varepsilon_m + (1 + \alpha)\varepsilon_w, \quad L(t) = 1 - \delta + (\alpha - \delta)(t - 1).$$

Its form  $e(r) = \max\{1, (1 + 5r)/6\}$  is uniform as  $r$  approaches one. If  $m \geq 1/2$ , the ordinary no-slot count has exponent at most  $1 - \delta + \varepsilon_m + 2\varepsilon_w$ . In the remaining range  $r < 1, m < 1/2$ , the two ideal counts at their full capacities are

$$A_I(r) = 1 - \delta\{x + (1 - x)r\}, \\ S_t(r) = 1 - \delta\{4x/9 + (2 - 8x/9)(t - r)\}.$$

Replacing  $m$  by  $t - r - \varepsilon_w$  costs at most  $2\varepsilon_w$  because the ordinary count decreases with  $m$  and its derivative has absolute value at most 2. Its witness divisor costs another  $2\varepsilon_w$ . The crossing occurs at

$$r_*(t) = \frac{(2 - 8x/9)t - 5x/9}{D_x}, \quad \frac{23}{37} \leq r_*(t) \leq 1, \quad \frac{1}{3} \leq t - r_*(t) \leq \frac{1}{2}.$$

Use the ordinary count for  $r \leq r_*(t)$  and the inverse count for  $r \geq r_*(t)$ . For the latter, if  $(1 - r)/2 > \nu_0$ , request  $z = (1 - r)/2 - \nu_0$ . Its whole-slot length  $z'$  satisfies

$$1 - r - 2z' \geq 2\nu_0 > 0, \quad 3 - 2r - 8z' = 4(1 - r - 2z') + (2r - 1) \geq \frac{9}{37}.$$

Thus one may fix  $c_1 = 2\nu_0$  and  $c_2 = 9/37$  before the target. For the ordinary request  $z = 2(1 - 2m)/9 - \nu_0 > 0$ , one has  $1 - 2m - 6\kappa z' \geq (9/2)\nu_0$ . The unreduced inverse capacity is at most  $7/37$ , and the ordinary capacity in its chosen branch is at most  $2/27 + 4\varepsilon_w/9$ . Both are strictly below the available supply with the fixed slack specified in the statement. Decrement, witness error, and slot mesh are each smaller than  $c_{\text{sup}}/16$ ; no limiting equality is used.

The common short-witness bound is

$$R_{\text{short}}(t) = 1 - \delta + \frac{\delta P_x}{D_x}(3/2 - t).$$

Choose

$$t = 1 + \frac{\delta P_x}{2J} \in [1, 3/2],$$

which makes  $R_{\text{short}}(t) = L(t)$  and yields exactly  $R_*$  in (8). If the inverse capacity is at most  $\nu_0$ , the no-slot ideal exponent is at most  $1 - \delta + 2\delta\nu_0$ ; if the ordinary capacity is at most  $\nu_0$ , it is at most  $1 - \delta + (9/2)\delta\nu_0$ . Since  $R_* \geq 1 - \delta$ , these cases fit the common error  $\varepsilon_m + 4\varepsilon_w + 6\nu_0 + \omega_{\text{slot}}$ . For  $q = 0$  the selected-slot amplification factor is 1, so the argument requires no division by an amplitude. Empty bins contribute zero; a vanishing slot cannot belong to a positive-amplitude class.

For completeness, without slots take  $t = 1$ . For  $r \leq 1$ , the two ideal exponents are  $1 - \delta r$  and  $1 - 2\delta(1 - r)$ , whose smaller value is at most  $1 - 2\delta/3$ , with crossing  $r = 2/3$ . For  $r > 1$ , the amplified estimate is at most  $1 - \delta$  up to the displayed witness errors. This proves the asserted no-slot form.

To pass from the ideal exponent to the raw count, we must also retain the height and multiplicity factors in the moment estimates. Lemmas 17.1, 17.6, and 18.1 give finite seminorm and height orders for the fixed datum, uniformly in their bounded real ranges, masks, and moving supports. Their application includes the finite derivative profiles required by the parameter Sobolev inequality of Lemma 4.5. A logarithmic scale derivative inserts  $yW'(y)$ ; a normalized twist derivative inserts  $\log y$  on the same annulus. Sum over rows before the nonnegative Sobolev integrals. If a particular fixed-parameter moment has height degree  $J_j$  and  $h_j$  height coordinates, covering their box costs at most  $(1 + H)^{J_j + h_j}$ . Take  $A_{\text{cnt}, \eta} = \max_j (J_j + h_j)$  over these actual

calls, their derivative profiles, and the no-slot calls. The finite presentation labels and slot subsets enter  $C_\eta$ ; witness dyads and logarithmic scale coverings contribute  $(1 + \log U)^L$ . The stated moment theorems supply each of these finitely many orders, so the maximum requires no extra analytic input. It may depend on  $\eta$ . We fix its orders and constants before choosing the trial height exponent. Auxiliary external tails in the detector are made smaller than a fixed fraction of its spike after that exponent is chosen; their constants only enlarge the lower threshold. They do not change these internal moment orders. This proves (9) with the asserted dependencies.  $\square$

The next step absorbs the logarithmic and height factors into powers of  $U$ . Reserve positive  $\varepsilon_{\log}$  and  $\varepsilon_{\text{cnt},\text{ht}}$  within the count allowance, with

$$\varepsilon_m + 4\varepsilon_w + 6\nu_0 + \omega_{\text{slot}} + \varepsilon_{\log} + \varepsilon_{\text{cnt},\text{ht}} < \varepsilon_{\text{cnt}}. \quad (10)$$

The logarithm in (9) is at most  $U^{\varepsilon_{\log}}$  for sufficiently large  $U$ . For  $U = Z^d$ ,  $d \geq d_{\min}$ , set  $H = Z^{2\theta_\eta}$  and require

$$0 < \theta_\eta \leq \frac{d_{\min}\varepsilon_{\text{cnt},\text{ht}}}{4(1 + A_{\text{cnt},\eta})}. \quad (11)$$

Then  $(1 + H)^{A_{\text{cnt},\eta}} \ll_\eta U^{\varepsilon_{\text{cnt},\text{ht}}}$ , so  $\#\mathcal{B} \ll_\eta U^{R_* + \varepsilon_{\text{cnt}}}$ . The same allocations cover the no-slot count, after taking the finite maximum of its height orders, and the bottom bin has the elementary bound  $\#\mathcal{B} \ll_\eta U$ . The actual choice of  $\theta_\eta$  is deferred to Section 7.

Proposition 5.2 combines this row count with the analytic factors in the high-side representation. After moving the full high-side correction and only then subdividing its retained values, it gives the joint estimate required by (10.13) of [1]. The resulting general exponent of (10.15), relative to  $C(7/8)$ , is

$$\begin{aligned} E_\tau(d) = a - \frac{7}{8} + h \left( \frac{17}{50} - \frac{1}{6} \right) - a l_y - \frac{\ell}{2} + q\ell \\ + d \left( R + \frac{\delta}{2} - \frac{17}{50} \right). \end{aligned} \quad (12)$$

Here  $R$  is the ideal count exponent of physical rows in the chosen box. The additional  $\varepsilon_{\text{cnt}}$ , already including count-height absorption, contributes  $d\varepsilon_{\text{cnt}}$  to the final  $Z$  exponent; it is not included a second time in the joint analytic height factor. For the same  $a, q, R, d$ , its change from the old geometry is exactly

$$E_\tau(d) - E_0(d) = \tau \left( \frac{1}{100} + \frac{\delta}{4} + q \right). \quad (13)$$

We now exhibit a margin large enough to absorb this increase.

**Lemma 4.2.** *For the adaptive witness count  $R = R_*$ , the endpoint  $d = h$  satisfies*

$$E_\tau(h) - \left( \beta_* - \frac{7}{8} \right) \leq -\frac{49}{440640} + \frac{15\tau}{8} < 0.$$

*Proof.* The complete square certificate (20.9) of [1] applies for  $0 \leq \delta \leq \alpha$  and  $0 \leq x \leq 1/2$ . To display its positive margin, set  $y = 1/2 - x$ ,  $v = 51 + 41y$ , and let  $E_*$  denote the old endpoint exponent. Its identity is

$$\begin{aligned} 10368vJ(-E_*) &= (3 + 5y) \{ (4v\delta - 79)^2 + 49 \} \\ &\quad + 4y \{ 4v\delta[(1 + 3y)(15 + 32y)\delta + 9 - 13y] + 265 + 3485y \}. \end{aligned}$$

Every term is nonnegative, since  $9 - 13y \geq 5/2$ . Using  $v \leq 17(3 + 5y)$  and  $J \leq 5/2$  gives

$$E_0(13/16) \leq -\frac{49}{440640}.$$

Increase  $d$  from  $13/16$  to  $h = 13/16 + 3\tau/2$ , and then use (13). This gives

$$E_\tau(h) \leq -\frac{49}{440640} + \tau \left( \frac{3}{2}R_* + \delta + q - \frac{1}{2} \right).$$

Since  $J \geq \delta P_x$ , we have  $R_* \leq 17/12 - 3\delta/2$ . Together with  $q \leq \delta/2$  this bounds the parentheses by  $13/8$ . Finally,  $\beta_* - 7/8 > -\tau/4$ , so comparison with  $C(\beta_*)$  costs at most a further  $\tau/4$ .  $\square$

The other central ranges have larger fixed margins. The bottom box uses  $\delta_0 = 1/50$  and  $R = 1$ , with old endpoint bound  $-7/1200$ ; the geometric change and comparison cost at most  $32\tau/25$ . For  $d \leq 1/2$  no selected primes are used: take  $R = 76/75 - 28/3$ . Equation (20.11) gives  $-49/14400$ , and the change and comparison cost at most  $329\tau/400$ . Thus put

$$m_A = \frac{49}{440640} - \frac{15\tau}{8} > 10^{-4}. \quad (14)$$

In the adaptive branch the slope in  $d$  is positive and less than 2, so its endpoint controls  $d \leq h$ . Choose

$$0 < \zeta < \min\{1/96, m_A/32\}.$$

Extending to  $h + \zeta$  costs less than  $2\zeta$ . The same extension still leaves enough compensation primes: since  $5\ell - h = 1/48 + 7\tau/2$ , on  $1/2 \leq d \leq h + \zeta$  we have

$$\frac{\ell}{d} > \frac{1}{5} > \frac{7}{37}.$$

Since  $1/5 - 7/37 = 2/185$ , this implies the prime-supply condition of Lemma 4.1 with  $c_{\text{sup}} = 1/185$  on the entire range where its selected slots are needed. The remaining fixed half of the gap keeps every chosen decrement and rounding allowance strictly inside the supply.

## 5. FULL CONTOURS AND THE JOINT CENTRAL ESTIMATE

The contour lemmas must also be checked at the new boundary. Lemmas 10.3–10.6 in [1] are stated only for boundary parameters at least  $7/8$ . To extend the required arguments, we use the exact local replacements and the full representation (7), checking the ranges where the perturbation matters. The full correction is moved before its retained values are subdivided by amplitude. This order lets us use the row count without requiring the individual amplitude pieces to be holomorphic.

Write  $x_r = \text{Re } s$ ,  $w_r = \text{Re } w$ , and  $z_r = \text{Re } z$ . The amplitude ratio  $x = q/\delta$  of the previous section remains a separate real parameter.

**Proposition 5.1.** *The complete correction (6) has the analyticity and full-height bounds required by Definition 10.1 of [1] in its first Euler region and in the region  $\mathcal{D}_{2,\tau}$  below. One final excluded set  $S$  can make the principal slice satisfy*

$$|H_\eta(s) - 1| \leq \frac{1}{2} \quad (\text{Re } s > \sigma_\tau).$$

*The central contours can still be moved for the full row sum, with an external tail whose scale order is fixed independently of the tail order chosen at the end.*

*Proof.* The first Euler region is unchanged:

$$\mathcal{D}_1(\varepsilon_0) : \quad x_r \geq \frac{51}{100}, \quad z_r \geq \frac{17}{50}, \quad w_r \geq -\frac{1}{100}, \quad x_r + w_r \geq 1 + \varepsilon_0.$$

For nonprincipal rows, move first on the global line  $x_r = \beta_* + 20e$ , and then, only within the buffered heights, to

$$x_r = a + 16e, \quad w_r = 1 - a - 6e,$$

where  $e > 0$  is sufficiently small. Throughout the paths,  $x_r + w_r \geq 1 + 10e$ . The original denominator for a dynamic box is zero-free in its buffered rectangle; global tails remain at  $x_r > \beta_*$  and use the global reciprocal bound. These paths need only  $\beta_* > 51/100$ , rather than  $\beta_* \geq 7/8$ . The integral and horizontal-edge tail estimates of Lemma 10.2, and the single cumulative height allocation in (10.14), are unchanged. After moving the complete correction, we subdivide its values by amplitude only to obtain pointwise upper bounds.

For the principal row enlarge the second Euler region to

$$\mathcal{D}_{2,\tau} : \quad x_r \geq \sigma_\tau, \quad z_r \geq \frac{33}{200}, \quad w_r \geq \frac{19}{20}.$$

This replaces the second region of Definition 10.1 and the principal-slice contraction half-plane in (10.2). In the exact local expressions (7.11), (7.17), the denominators  $1 - R$ ,  $1 - V$ ,  $1 - D$  stay uniformly nonzero: for  $Q = \mathbb{N}p \geq 2$ ,

$$|D| \leq Q^{-\sigma_\tau}, \quad |V| \leq Q^{-99/100}, \quad |R| \leq Q^{-56/25+3\tau/2}.$$

The largest off-row defect exponent and the largest dividing-prime exponent are, respectively,

$$1 - x_r - w_r - 6z_r \leq -\frac{363}{200} + \frac{\tau}{4}, \quad 1 - x_r - w_r \leq -\frac{33}{40} + \frac{\tau}{4}.$$

For clarity, the ramified boundary terms  $J_j$  in (7.17), for  $j = 1, \dots, 5$ , have exponents bounded above by

$$-\frac{73}{40} + \frac{\tau}{4}, \quad -\frac{9}{8} + \frac{3\tau}{4}, \quad \left(-\frac{63}{40} + \frac{3\tau}{4}, -\frac{3}{2} + \tau\right), \quad -\frac{39}{20} + \tau, \quad -\frac{9}{4} + \frac{3\tau}{2}.$$

These and the remaining off-row terms in (7.11), (7.17) are strictly smaller than the corresponding largest exponents above. Writing  $\gamma_0 = 163/200 - \tau/4 > 0$  and  $\gamma_1 = 33/40 - \tau/4 > 0$ , the positive local bounds are  $|H_p| \leq 1 + O(Q^{-1-\gamma_0})$  off  $u$  and  $|H_p| \leq 1 + O(Q^{-\gamma_1})$  at primes dividing  $u$ . The first product converges. For every  $\varepsilon > 0$ , the second is  $O_\varepsilon((Nu)^\varepsilon)$ : outside a fixed finite set each factor is at most  $Q^\varepsilon$ , and the product of the distinct dividing primes is at most  $Nu$ . These estimates are uniform in all imaginary parts and unit phases. Thus the unselected Euler product is normally convergent in the enlarged region with the stated full-height bound. For  $u = 1$ , enlarge the fixed cutoff in  $S$  to make the product uniformly close to 1. The positive product-tail majorant is  $O(P_0^{-163/200+\tau/4})$ , and gives

$$\sup_{\operatorname{Re} s > \sigma_\tau} |H_\eta(s) - 1| \leq \frac{1}{2}.$$

Further enlargement of  $S$  uses the same positive tail bound, without relying on cancellation between complex products.

The complete correction also has the full-row and full-height bounds required by Definition 10.1. On the fixed real boxes where (16.2)–(16.6) apply,  $|G_p| \ll Q^B$  for a fixed  $B$ , with height order zero. Bound the unselected Euler factors by the same positive-product majorant and count the finite tuples. Then

$$|H_{\eta,u,z}(s, w, z)| \ll_\varepsilon (Nu)^\varepsilon Z^{B'},$$

where  $B'$  is fixed before the final external tail order  $N$ .

The multiplier of the principal signal is the following slice:

$$H_\eta(s) = \prod_{p \notin S} H_{\eta,1,p}(s, 1, 1/6).$$

At  $w = 1, z = 1/6$  the exact local factor gives

$$H_{\eta,p}(s) - 1 = O((Np)^{-2} + (Np)^{-1-\operatorname{Re} s} + (Np)^{3-6\operatorname{Re} s}).$$

The slice product therefore converges absolutely and uniformly on compact subsets of  $\operatorname{Re} s > 2/3$ , and is holomorphic there; this contains  $\operatorname{Re} s > \sigma_\tau$ .

We next bound the full row sum, keeping the polynomial orders independent of the tail order to be chosen later. First take  $Nu \leq Z^{h+\zeta}$ . There are  $O(Z^{h+\zeta})$  elements in the fixed unit sector. On each fixed real contour box the displayed positive correction bound, Hecke strip growth, and the appropriate reciprocal estimate give a polynomial majorant after this entire row sum, of the form

$$|F_Z(\vec{t})| \leq C_\eta Z^{B_\eta} (1 + |\vec{t}|)^{J_\eta^*}.$$

Here  $F_Z$  includes the scale factors and arithmetic terms but not the external smooth kernel. The principal poles are extracted as residues; the bounds for its remaining contours stay away from those poles. For the unbounded range  $Nu > Z^{h+\zeta}$ , the absolute row majorant on  $(2, 2, z_\infty)$  and its summable dyadic series are proved in Proposition 6.2. They give another fixed polynomial majorant. Taking the maximum over these finitely many types of estimates fixes finite  $B_\eta, J_\eta^*$  for all discarded integrals and joins before we choose  $N$ . A buffered reciprocal is used only on its allocated domain; other tails stay on the global or absolute lines and use their global reciprocal bounds. When an integration axis is extended, the buffered estimate is still used only inside its rectangle.

For these same fixed  $W_0, W_1$ , the external kernel is

$$e^{-(t_s+t_z)^2} |\mathcal{M}(z_r + it_z)| |\widehat{W}_1(w_r + it_w)|.$$

It is rapidly decreasing in all three variables: the map  $(t_s, t_z, t_w) \mapsto (t_s + t_z, t_z, t_w)$  is invertible, and logarithmic integration by parts gives every decay order for the two Mellin transforms. For  $\mathcal{M}$ , its underlying

Fourier transform is smooth at zero and Schwartz at infinity; all the real  $z$  intervals here are compact subsets of  $(0, \infty)$ . Thus the boundary terms vanish. Annular support gives the same conclusion for  $\widehat{W}_1$ . After the  $w$  residue the identical argument applies to the two remaining variables. Any external annular separation profile  $W$  is likewise fixed; its all-order Fourier estimates follow by integration by parts, as in Lemma 4.5. Pure twists translate Mellin arguments and append a fixed invertible block triangular height map. Internal coefficient measures controlled only in a fixed weighted  $L^1$  norm remain integrated inside  $F_Z$ ; no pointwise trace estimate is inferred for such a measure.

For the truncations below, choose a fixed  $c > 0$  no larger than any coordinate multiplier in (10.14). It depends only on the fixed height-allocation matrix and is independent of  $Z, T_1, N$ . With  $k$  external coordinates, take kernel decay order greater than  $N + J_\eta^* + k$ . Integrating the resulting positive polynomial majorant outside  $\max |t_j| \leq cT_1$ , or on a slice  $t_j = \pm cT_1$ , gives, respectively, the integrated and horizontal-trace bounds

$$O_{\eta, N}(Z^{B_\eta} T_1^{-N}).$$

A bounded real join interval only multiplies this bound by its length. This proves the tails used in the contour argument of Lemma 10.2 and Section 10.3 in the enlarged regions. All absolute interchanges use these full-row majorants. Increasing  $N$  requires higher seminorms of the same external functions. The polynomial orders  $B_\eta, J_\eta^*$  and the internal moment orders remain fixed. Only the retained integral is subsequently subdivided pointwise.  $\square$

The row count must be combined with a bound for the analytic numerator and all error slots together. The conductor saving belongs to the whole numerator. Estimating each slot as if it had its own saving would use that saving more than once.

**Proposition 5.2.** *Put  $z_0 = 17/50$  and  $\delta = 2a - 1$ . Consider central nonprincipal rows in one retained box of Section 8 of [1], at real parts*

$$x_r = a + 16e, \quad w_r = 1 - a - 6e, \quad z_r = z_0.$$

*Fix an error-slot set  $I$  and an amplitude row set at a retained point. Suppose the row count is at most  $C_\eta U^{R+\varepsilon_{\text{cnt}}}$  after (10)–(11), and the mean main-slot amplitude exponent is  $q$ , assigning exponent zero to error slots. The corresponding piece of the full correction satisfies*

$$\sum_u |L_F^S(w, \chi_\bullet(u)) H_{\eta, u, Z}^{(I)}(s, w, z)| \ll_\eta U^{R+\delta/2+\varepsilon_{\text{cnt}}+\varepsilon_c} \times Z^{\ell(z_0-1/2)+q\ell+\varepsilon_c} (1+T_1)^{A_\eta}. \quad (15)$$

*The central analytic loss  $\varepsilon_c$  excludes  $\varepsilon_{\text{cnt}}$  and can be made arbitrarily small after the slot system is fixed. The finite height order  $A_\eta$  here includes only the remaining analytic factors; the count height has already been absorbed into  $\varepsilon_{\text{cnt}}$ . It does not change with the external tail order. In particular the amplitude term in the general exponent is  $g = q\ell$ .*

*Proof.* Within the retained dynamic region, we can separate the local factor into a main slot and an error slot as follows. Proposition 16.1 in [1] applies for  $51/100 \leq a \leq 1$  without fixing total slot length to  $1/6$ . Its proof gives

$$H_{\mathfrak{p}} - 1 = O(Q^{-1-10e}) \quad (\mathfrak{p} \nmid u), \quad H_{\mathfrak{p}} - 1 = O(Q^{-10e}) \quad (\mathfrak{p} \mid u).$$

For fixed positive  $e$ , a sufficiently large final cutoff makes these local factors nonzero. Write the uncompensated Euler product as  $\mathcal{H}_{\eta, u} = \prod_{\mathfrak{p} \notin S} H_{\mathfrak{p}}$ , and define here

$$\mathfrak{B}_i = \sum_{\mathfrak{p} \in \mathcal{P}_i(Z)} W_i(N\mathfrak{p}/P_i) (N\mathfrak{p})^{z-1} \frac{G_{\mathfrak{p}}}{H_{\mathfrak{p}}},$$

$$Q_i = - \sum_{\mathfrak{p} \in \mathcal{P}_i(Z)} W_i(N\mathfrak{p}/P_i) (N\mathfrak{p})^{z-1} \overline{\chi_{\mathfrak{p}}(u)}, \quad D_i = \mathfrak{B}_i - Q_i.$$

The conjugate in  $Q_i$  is the one in Proposition 16.1, and agrees with the negative physical direction used in Section 4. For  $\mathfrak{p} \nmid u$ , the local cancellation (16.8) gives

$$G_{\mathfrak{p}}/H_{\mathfrak{p}} + \overline{\chi_{\mathfrak{p}}(u)} = O(Q^{-51/100}).$$

Thus  $D_i$  is precisely the error slot jointly estimated in (16.9). Disjoint windows give the exact finite expansion

$$H_{\eta,u,Z} = \mathcal{H}_{\eta,u} \sum_{I \subseteq \{1,\dots,K\}} \prod_{i \in I} D_i \prod_{i \notin I} Q_i = \sum_I H_{\eta,u,Z}^{(I)}. \quad (16)$$

This is (16.10). The quotients are valid because the local factors are nonzero in the dynamic region. That nonvanishing was proved only there. Outside the region, continuation uses the complete factor  $G_p$  in (5).

The buffered estimates for the retained box and Lemma 4.10 give the reflected primitive-numerator small-power bound required by Proposition 16.1. Equation (16.9) then estimates the numerator and every error slot simultaneously:

$$\left| L_F^S(w, \chi_\bullet(u)) \prod_{i \in I} D_i \right| \ll U^{\delta/2+12e+\lambda_{\text{joint}}} (1+T_1)^A \times \prod_{i \in I} P_i^{z_0-1/2}. \quad (17)$$

Here  $\lambda_{\text{joint}} > 0$  is prescribed before the target, and the preliminary small powers in that proof are reduced to fit it. To track the factors in this estimate, put  $A_* = a - 1/2 + 6e$ . The primitive functional equation contributes  $q_f^{A_*}$  and a height polynomial, and the remaining deleted Euler factors cost at most  $U^{6e+\lambda}$  because  $w_r \geq -6e$ . For the full set  $J_0$  of distinct strict ramified labels, (16.11) gives once

$$q_f \ll_S \text{Nu} \prod_{p \in J_0} Q^{-(j_p-1)}.$$

Each such label has  $j_p \geq 2$  and  $-w_r - A_* = -1/2$ . Hence its allocated local factor is at most  $Q^{z_0-1/2}$ , simultaneously for all these labels; coprime errors have the stronger exponent  $z_0 - 51/100$ . Divisor-many ramified choices and the other primitive small powers are included in  $\lambda_{\text{joint}}$ . These two contributions account for  $6e+6e = 12e$ . The estimate therefore requires no additional positive loss proportional to the prime length. The labels are distinct because the underlying windows are disjoint. The conductor saving and the allowance for deleted factors have thus been used once for the full product of error slots.

For the remaining slots the normalization is explicitly

$$\tilde{Q}_i = P_i^{-1/2} \sum_{p \in \mathcal{P}_i(Z)} \overline{\chi_p(u)} W_i(\text{Np}/P_i) (\text{Np}/P_i)^{z-1}, \quad Q_i = -P_i^{z-1/2} \tilde{Q}_i.$$

To form the amplitude boxes, write Lemma 19.1's bound as  $|\tilde{Q}_i| \ll U^{\epsilon_1} P_i^{\delta/2+C_{\text{pr}}e}$ , with  $C_{\text{pr}}$  fixed on the real ranges. After  $K$  is fixed, put  $\ell_{\min} = \ell/K$  and  $d_{\max} = h + \zeta$ , and choose

$$C_{\text{pr}}e + \frac{d_{\max}\epsilon_1}{\ell_{\min}} < \frac{\vartheta}{2}.$$

Increasing the threshold absorbs the fixed constant into the other half of the width, giving  $|\tilde{Q}_i| \leq P_i^{\delta/2+\vartheta}$ . The capped subdivision of Section 19.1 then has  $0 \leq g_i \leq \delta/2$ , with  $|\tilde{Q}_i| \geq P_i^{g_i}$  when  $g_i > 0$ . This prime-estimate cost is already included in  $\vartheta$ . In an amplitude box take  $|\tilde{Q}_i| \leq P_i^{g_i+\vartheta}$ , using a zero-exponent upper bound for low-amplitude slots. Assign  $g_i = 0$  to each error slot. Then

$$\sum_i \ell_i g_i = q\ell, \quad 0 \leq q \leq \delta/2,$$

by Lemma 19.1. Multiply the main-slot bounds into (17), use  $\mathcal{H}_{\eta,u} \ll U^\epsilon$ , and sum over the rows. For example, if  $\mathcal{H}_{\eta,u} \ll U^{\lambda_H}$ , it suffices to choose

$$12e + \lambda_{\text{joint}} + \lambda_H \leq \epsilon_c, \quad \ell\vartheta \leq \epsilon_c.$$

This gives (15), with  $\epsilon_{\text{cnt}}$  paid separately. There are a fixed  $2^K$  error subsets. Witness presentations and dyads have already been included in the raw count's constant and logarithm; only the remaining amplitude subdivisions and physical norm dyads enter the subsequent multiplicity allowance.

Apply this estimate at each retained point of the already moved complete integral. Its outside real exponent is

$$l_x(1/2 - z_0) + a + z_0 - 1 - al_y - dz_0 + (16 - 6l_y)e.$$

Adding (15), subtracting  $C(7/8)$ , and using  $h = 1 - l_x + \ell$  gives (12) as the ideal exponent. The additional real charges, including the retained reciprocal, are listed in Section 7. All auxiliary heights obey the single allocation (10.14). Neither an error piece nor a pointwise row set is shifted on its own, and neither needs to be holomorphic outside the retained region. This establishes the use of (10.13)–(10.16) for the perturbed geometry.  $\square$

## 6. THE PRINCIPAL SIGNAL AND THE OUTER ROWS

The central estimate bounds the nonprincipal rows in the central range. The principal row supplies the signal: taking its two scalar residues produces the function whose continuation detects zeros. We first identify the positive normalizing factor and the errors in this extraction. We then estimate the small and large nonprincipal rows on contours adapted to their sizes. These bounds complete the treatment of the physical rows.

**Proposition 6.1.** *After fixing even  $K$ , the windows, and the final excluded set, the principal-row double residue produces the signal (18). Its normalizing factor  $c_S A_T(Z)$  is positive, with inverse bounded by an arbitrarily small power of  $Z$ . The other principal-row integrals and the local approximation retain the respective positive savings  $l_y/20$ ,  $h/600$ , and  $\kappa_P$ .*

*Proof.* We first check that the positive tuple bound for the selected factors still holds at the new lengths. On the principal rectangle the four error powers in (16.8) of [1] are

$$-x_r, \quad -6z_r, \quad 4 - 5x_r - 6z_r, \quad 1 - w_r - 6z_r,$$

all strictly negative. The positive tuple upper bound for the correction is therefore  $Z^{\ell z_r + \varepsilon}$  at every height.

At the principal double-residue slice  $u = 1$ ,  $w = 1$ ,  $z = 1/6$ , define

$$B_p(s) = \frac{G_p(s, 1, 1/6)}{H_p(s, 1, 1/6)}.$$

This slice lies in the second Euler region for the principal row, where  $H_p$  is nonzero. This quotient is used to normalize the principal row. For contour movement in general, we continue to use the complete replacement (5). The local identities (16.12)–(16.13) give

$$B_p(s) = -1 + O((Np)^{-\sigma_\tau}),$$

since  $5\sigma_\tau - 3 > \sigma_\tau$ .

To normalize the prime sums, set

$$S_i(Z) = \sum_{p \in \mathcal{P}_i(Z)} W_i(Np/P_i)(Np)^{-5/6}, \quad A_T(Z) = (-1)^K Z^{-\ell/6} \prod_i S_i(Z).$$

The fixed ray-class prime theorem, Lemma 13.1 of [1], gives

$$S_i(Z) \sim \frac{c_i P_i^{1/6}}{|T| \log P_i}, \quad c_i = \int W_i(y) y^{-5/6} dy > 0.$$

Even  $K$  makes  $A_T(Z) > 0$ , and  $A_T(Z) \asymp (\log Z)^{-K}$ . Once  $K$  is fixed, choose

$$\kappa_P = \frac{\sigma_\tau \ell}{2K} > 0.$$

The local approximation to the double residue has error  $O(Z^{-\kappa_P})$ , uniformly on the entire global  $s$  line.

Repeat the principal contour argument of Lemma 10.5 with the extended region proved above. Keep  $s_r = \beta_* + e$ , first cross  $w = 1$ , and then, within its residue, cross  $z = 1/6$ . No other poles are crossed. Put  $r_S = \text{Res}_{v=1} \zeta_F^S(v) > 0$ . The product of the two scalar residues is

$$c_S = \widehat{W}_1(1) \mathcal{M}(1/6) r_S^2 / 6 > 0.$$

The positivity of  $\mathcal{M}(1/6)$  is the Fourier-pairing result in (7.2), and the factor  $1/6$  is the Jacobian in the residue of  $\zeta_F^S(6z)$ . The unresidued  $w$  integral saves  $l_y/20$ ; the remaining  $z$  integral saves  $h/600$ .

Move the main term of the double residue to  $s_r = 2$ . It yields the height-independent signal

$$f_\eta(Z) = \frac{1}{2\pi i} \int_{\text{Re } s=2} Z^{C(s)} e^{(s-5/6)^2} \frac{H_\eta(s)}{L_F^S(s, \eta)} ds. \quad (18)$$

We estimate the residue-approximation error on the original global line, and move only the main term to the signal contour.  $\square$

**Proposition 6.2.** *After removing the principal row, the small rows  $0 < \text{Nu} \leq Z^{1/100}$  and the large rows  $\text{Nu} > Z^{h+\zeta}$  both retain a positive exponent saving. Together with the central range they cover all physical rows.*

*Proof.* For small rows  $u \neq 1$  use

$$x_r = \beta_* + e, \quad w_r = 1/2, \quad z_r = 17/50.$$

These contour real parts lie in  $\mathcal{D}_1(1/3)$  because  $\beta_* > \sigma_\tau > 5/6$ ; the fixed margin  $1/3$  thus remains valid in the new contradiction range. At selected primes not dividing  $u$ , (16.8) gives  $G_p = O(1)$ . At dividing primes its boundary exponents are

$$-1/2, \quad 3/2 - 2x_r, \quad (3/2 - 2x_r, 2 - 3x_r), \quad 2 - 3x_r, \quad 3 - 5x_r.$$

All are at most  $1/2$ , so  $G_p \ll (\text{Np})^{1/2}$ . The divisor bound absorbs the factors associated with primes dividing the row. The original numerator and row-count bounds have combined power  $63/50 < 2$ . The geometric exponent is

$$h \left( \frac{17}{50} - \frac{1}{6} \right) - \frac{l_y}{2} = -\frac{79}{800} + \frac{51\tau}{100}.$$

Allowing the conservative row-count cost  $2/100$  and comparing with  $C(\beta_*)$  still leaves the saving

$$\frac{63}{800} - \frac{19\tau}{25} > 0.$$

For large rows use the absolute lines  $(x_r, w_r, z_r) = (2, 2, z_\infty)$ , with fixed  $z_\infty > 2$ . The exact local formula gives  $G_p = O_{z_\infty}(1)$  both off and on the row. The positive unselected product is  $O_\varepsilon((\text{Nu})^\varepsilon)$ . Absolute ideal counting in each annular slot, before any character cancellation, therefore bounds the full selected-tuple sum by

$$(\text{Nu})^\varepsilon \prod_i P_i^{z_\infty} = (\text{Nu})^\varepsilon Z^{\ell z_\infty}.$$

All three scalar Euler factors are absolutely bounded on these lines. For  $V = Z^{h+\zeta}$ , dyadic lattice counting gives the convergent series

$$\sum_{\text{Nu} > V} (\text{Nu})^{-z_\infty + \varepsilon} \ll \sum_{j \geq 0} (2^j V)^{1 - z_\infty + \varepsilon} \ll V^{1 - z_\infty + \varepsilon}, \quad z_\infty > 1 + \varepsilon.$$

The convergent dyadic series bounds the entire infinite large-row range, uniformly in the Mellin heights. The fixed smooth kernel is integrable, so absolute convergence also justifies interchanging this series with the integrals. Since  $h = 1 - l_x + \ell$ , the resulting exponent tends to minus infinity with  $z_\infty$ . Its uniform upper bound is

$$\frac{53}{32} - \frac{3\tau}{4} + (h + \zeta)(1 + \varepsilon) - \zeta z_\infty + \varepsilon.$$

Fix  $\zeta$  first and then a sufficiently large fixed  $z_\infty$ . This gives a uniform positive saving for the large rows.  $\square$

## 7. CHOOSING THE PARAMETERS AND EXCLUDING THE GAP

All estimates now refer to the same physical probe: the low-side bound, the central-row estimates, the principal residue, and the bounds for the outer rows. We can combine their positive margins once the parameters have been chosen in the right order. The witness mesh must be fixed before the number of prime slots. Increasing that number reduces the saving in the principal approximation, so that saving must not determine the earlier mesh.

First fix the geometry and keep the assumed family-wide contradiction gap  $\Delta_\tau$  fixed. The gap is fixed for the family, independently of the individual target. Fix the central margin and  $\zeta$ . The chosen buffer gives

$$m_A - 2\zeta > \frac{15m_A}{16} = 10^{-4} + \frac{5861}{2350080000}.$$

Divide the central loss allowance between moment-count losses and later analytic subdivision losses, requiring their total to be less than

$$\frac{5861}{4700160000}.$$

This preserves  $\mu = 10^{-4}$ . Within the first allowance, choose  $\varepsilon_{\text{cnt}}$  and all six terms on the left of (10), with its strict inequality. We reserve room for the height and logarithmic factors now. Their finite degrees will be determined after the target is fixed. Fix the witness losses,  $\nu_0$ , and  $\omega_{\text{slot}}$  as in Lemma 4.1. The mesh in Lemma 18.1 depends only on the bounded real ranges and  $\varepsilon_m$ , not on the target or the later  $K$ . Then choose a sufficiently large fixed even  $K$  such that  $2\ell/K < \omega_{\text{slot}}$ . Throughout the adaptive row range, the available base- $U$  prime length stays above  $1/5$ , and Lemma 4.1 controls both capacities. After fixing the window weights the principal approximation saving is fixed at

$$\kappa_p = \frac{\sigma_\tau \ell}{2K} > 0.$$

The moment losses chosen before  $K$  are used in the central row count, so they need only fit its allowance. They need not be smaller than  $\kappa_p$ : the principal local approximation is a separate error and uses no moment mesh. Only after fixing  $K$  do we shrink the dynamic-box widths, amplitude-box widths, and local small powers. The central charges must fit their remaining allowance; the low-side charges must fit suitable portions of  $\Delta_\tau/2$ ; the principal charges must fit suitable portions of  $l_y/20$ ,  $h/600$ , and  $\kappa_p$ . More explicitly, with  $\varepsilon_{\text{pr}}$  including the principal small powers and its one normalization allowance, require

$$(1+h)e + \varepsilon_{\text{pr}} < l_y/80, \quad e + \varepsilon_{\text{pr}} < h/2400, \quad e + \varepsilon_{\text{pr}} < \kappa_p/4.$$

Also take  $e$  below the detector's bounds for the already prescribed witness losses. These are choices after  $K$ ; none asks the earlier moment loss or mesh to be smaller than  $\kappa_p$ . Detection boxes can be refined beyond the fixed moment mesh without changing that mesh or  $K$ .

To combine the estimates, we must express all central losses in the same base, changing from  $U = Z^d$  to  $Z$ . Keeping  $R$  as the ideal count exponent, the total real-exponent loss after normalization is

$$(16 - 6l_y)e + d\varepsilon_{\text{cnt}} + (1+d)\varepsilon_c + d\varepsilon_d + \varepsilon_p + \varepsilon_{\text{norm}}. \quad (19)$$

Here  $\varepsilon_d$  is exactly the retained reciprocal allowance in (10.16) of [1]: the buffered estimate for  $1/L_F^S(s, \eta\chi_\bullet(u))$  is  $O_\eta(U^{\varepsilon_d})$ . This loss comes from the reciprocal factor. The row count contributes  $d\varepsilon_{\text{cnt}}$ ; the joint numerator, simultaneous conductor allocation, deleted factors, unselected product, and main-slot amplitude upper bounds contribute  $d\varepsilon_c + \varepsilon_c$ , as proved in Proposition 5.2. The power  $\varepsilon_p$  covers the remaining pointwise or norm-dyadic logarithmic multiplicities. The witness logarithm is already included in (10). Finally  $\varepsilon_{\text{norm}}$  pays  $(c_S A_T(Z))^{-1} \ll_\eta Z^{\varepsilon_{\text{norm}}}$  once for this contribution. It is separate from the primitive or deleted Euler factors. The fixed  $2^K$  expansion and finite character labels cost constants.

If instead one writes  $R_{\text{inc}} = R + \varepsilon_{\text{cnt}}$  in (12), its  $dR_{\text{inc}}$  has already paid the count and the separate  $d\varepsilon_{\text{cnt}}$  in (19) must be omitted. We use the ideal convention throughout. Likewise the count height already absorbed in  $\varepsilon_{\text{cnt}}$  is not charged again in a retained analytic height power. The principal estimate has its own single normalization allowance inside  $\varepsilon_{\text{pr}}$ ; it does not incur the central normalization charge a second time. Since  $d \leq h + \zeta < 1$ , it suffices to arrange

$$(16 - 6l_y)e + \varepsilon_{\text{cnt}} + 2\varepsilon_c + \varepsilon_d + \varepsilon_p + \varepsilon_{\text{norm}} < \frac{5861}{4700160000}.$$

The count share was reserved before  $K$  and the remaining terms are chosen after  $K$  within the other share. Thus every raw and logarithmic loss has been included before we choose the final analytic height.

Having retained every saving, take

$$m = \frac{1}{4} \min\{\mu, l_y/20, h/600, \kappa_p\} > 0.$$

With the earlier moment losses and the later local small powers chosen separately, we avoid circular dependence: increasing  $K$  reduces  $\kappa_p$ , which must not force another increase of  $K$ . All these positive real parameters and finite structures, including the large absolute line, are fixed before the target  $\eta$ .

For each target choose its fixed arithmetic data and enlarge the excluded set to its final value. Complete every such enlargement first, then use that same final  $S$  for the physical probe, Euler correction,  $c_S$ ,  $A_T$ , and the two functions

$$J_\eta(Z) = \frac{I_{\eta,\text{mod}}(Z)}{c_S A_T(Z)}, \quad f_\eta(Z) \text{ as in (18)}.$$

The low bound and logarithmic normalization give

$$|J_\eta(Z)| \ll_\eta Z^{C(\sigma_\tau)+\omega}, \quad \omega = \Delta_\tau/2 \in (0, \Delta_\tau). \quad (20)$$

Now fix all internal moment, seminorm, and Sobolev orders for this datum. Lemma 4.1 gives  $A_{\text{cnt},\eta}$ ; let  $A_{\text{ht},\eta}$  dominate the finitely many literal detector height orders. The remaining retained analytic height degrees give  $A_\eta$ , excluding the already paid count height. When a remaining bound is expressed in  $H = T_1^2$ , include twice its degree in  $A_\eta$ . The full-row majorants in Propositions 5.1 and 6.2, with normalization and the remaining finite types, give  $B_\eta$ . All these degrees are finite before a trial  $\theta$  or the final external decay order is chosen. Subject to the height ceilings below, combining the principal and all nonprincipal estimates gives the same  $m > 0$  and, for every fixed  $N$ ,

$$|J_\eta(Z) - f_\eta(Z)| \ll_{\eta,N} Z^{C(\beta_*)-m}(1+T_1)^{A_\eta} + Z^{B_\eta} T_1^{-N}. \quad (21)$$

Here the analytic height is an auxiliary parameter in the error estimate. The functions  $J_\eta$  and  $f_\eta$  themselves do not depend on it.

Let  $\varepsilon_{\text{ht}} > 0$  be the minimum of the finitely many detector height allowances fixed with the preliminary witness losses, and put

$$\theta_{0,\eta} = \frac{d_{\min} \varepsilon_{\text{ht}}}{20(1+A_{\text{ht},\eta})} > 0.$$

For  $T_1 = Z^\theta$  with  $0 < \theta \leq \min\{d_{\min}/100, \theta_{0,\eta}\}$  and  $U \geq Z^{d_{\min}}$ , sufficiently large  $Z$  gives

$$T_1 \leq U^{1/100}, \quad (1+T_1)^{A_{\text{ht},\eta}} \leq 2^{A_{\text{ht},\eta}} Z^{d_{\min} \varepsilon_{\text{ht}}/20} \leq U^{\varepsilon_{\text{ht}}/10}.$$

These inequalities verify the height hypotheses of the pointwise dyadic estimates. They also make the detector error  $U^{-189/100+o(1)} T_1^2$  tend to zero. All finitely many retained physical and witness heights are bounded by a fixed multiple of  $T_1$ , hence by  $H = T_1^2 = Z^{2\theta}$  after increasing the threshold. The matrix allocation (10.14) bounds all added Mellin frequencies simultaneously, with total at most  $T_1/2$  in each buffered argument. The same allowance applies throughout the later estimates.

Choose the actual height by

$$0 < \theta_\eta \leq \min \left\{ \frac{d_{\min}}{100}, \theta_{0,\eta}, \frac{d_{\min} \varepsilon_{\text{cnt},\text{ht}}}{4(1+A_{\text{cnt},\eta})}, \frac{m}{4(A_\eta+1)} \right\}, \quad T_1 = Z^{\theta_\eta}.$$

This both absorbs the count height in its preallocated allowance and gives  $(1+T_1)^{A_\eta} \ll_\eta Z^{m/4}$  for the remaining retained factors. Only then choose a fixed integer  $N_\eta$  with

$$B_\eta - N_\eta \theta_\eta < C(\beta_*) - m/2.$$

Auxiliary external orders in the detector and prime estimates may now also be increased so their discarded parts fit strict power gaps. Only the derivative and decay orders of the fixed smooth profiles are increased. Their constants can be absorbed by the final threshold. This does not require applying an internal moment estimate to a newly differentiated external profile. Consequently neither  $A_\eta$ ,  $A_{\text{cnt},\eta}$ ,  $A_{\text{ht},\eta}$ , nor  $B_\eta$  is reset by this last choice. Equation (21) becomes

$$|J_\eta - f_\eta| \ll_\eta Z^{C(\beta_*)-m/2}. \quad (22)$$

Throughout,  $W, W_0, W_1$ , the slot windows, and the functions  $J_\eta, f_\eta$  remain fixed. The later orders refer only to higher derivative and decay estimates of the fixed external smooth profiles. The outer slot and

presentation types are finite; the internal completed rows, Euler products, and Mellin integrals retain the absolute convergence and majorants proved above.

The order of dependence can now be stated explicitly:  $m, \omega$  and all real losses are fixed before  $\eta$ ; for each target the finite degrees and constants precede  $\theta_\eta$ ; then come  $N_\eta$  and the lower threshold. Proposition 2.1 of [1] requires the positive savings  $\omega$  and  $m/2$  to be independent of the target, but explicitly permits target-dependent constants, excluded sets, and thresholds. Since neither function contains the auxiliary height, it also permits this target-dependent choice of  $\theta_\eta$ , precisely as in Lemma 11.1. The height degrees remain target-dependent throughout; their values are never required to be uniform over the family.

We can now apply the common-signal continuation principle, Proposition 2.1 of [1], which allows  $\sigma_0 = \sigma_\tau$ . The exponent  $C$  has slope one; the multiplier  $H_\eta$  is holomorphic and satisfies  $|H_\eta - 1| \leq 1/2$  on the new half-plane; and the same  $J_\eta$  satisfies both (20) and (22). The first estimate supplies the low bound, and the second compares the same normalized probe with the signal. Together they are the two estimates required by the continuation principle. It excludes  $\beta_* > \sigma_\tau$  by choosing a target with an actual zero sufficiently close to the family supremum. Attainment of the supremum is unnecessary. The assumed gap therefore leads to a contradiction, and

$$\beta_* \leq \sigma_\tau.$$

**From primitive Hecke characters to Dirichlet characters.** The first step of Proposition 11.3 of [1] extends the Hecke conclusion to imprimitive characters. Passing from a primitive character to an induced presentation changes only finitely many factors  $1 - \eta(\mathfrak{p})(N\mathfrak{p})^{-s}$ , which are nonzero for  $\text{Re } s > 0$ .

To transfer the result to Dirichlet characters, we distinguish the deleted rational primes from the prime ideals above them. Let  $\chi$  be a primitive Dirichlet character of conductor  $q_\chi$ . Let  $T_\mathbb{Q}$  be the rational primes dividing  $3q_\chi$ , and let  $T_F$  consist of all prime ideals above them. These sets are used only in this transfer identity. On ideals coprime to  $3q_\chi$  define  $\eta(\mathfrak{a}) = \chi(N\mathfrak{a})$ , and extend by zero elsewhere. If  $\alpha \equiv 1 \pmod{q_\chi \mathcal{O}_F}$ , then  $N\alpha \equiv 1 \pmod{q_\chi}$ , so this defines a finite-order ray class character.

Write  $\chi_{-3}$  for the quadratic Dirichlet character of conductor 3 associated with  $F = \mathbb{Q}(\sqrt{-3})$ .

For  $p \notin T_\mathbb{Q}$ , a split prime has  $\chi_{-3}(p) = 1$  and two prime ideals of norm  $p$ . Its Hecke factor is  $(1 - \chi(p)p^{-s})^{-2}$ . An inert prime has  $\chi_{-3}(p) = -1$  and one prime ideal of norm  $p^2$ , giving

$$(1 - \chi(p)^2 p^{-2s})^{-1} = (1 - \chi(p)p^{-s})^{-1} (1 + \chi(p)p^{-s})^{-1}.$$

Absolute convergence of the Euler products, followed by meromorphic continuation, yields

$$L_F^{T_F}(s, \eta) = L^{T_\mathbb{Q}}(s, \chi) L^{T_\mathbb{Q}}(s, \chi \chi_{-3}). \quad (23)$$

All deleted factors are nonzero for  $\text{Re } s > 0$ . In  $0 < \text{Re } s < 1$ , and at  $s = 1 + it$  with  $t \neq 0$ , both Dirichlet factors have no poles, so a zero of either produces a Hecke zero. For  $\text{Re } s > 1$  absolute Euler convergence gives nonvanishing.

At  $s = 1$ , a possible pole cancellation can occur only when one primitive factor is principal. The other primitive factor is then  $\chi_{-3}$ , for which

$$L(1, \chi_{-3}) = \int_0^1 \frac{dx}{1+x+x^2} = \frac{\pi}{3\sqrt{3}} > 0.$$

The additional finite Euler factors are also nonzero at  $s = 1$ . This rules out cancellation of a zero by a pole at that point. The same argument with nonzero finite Euler factors handles imprimitive Dirichlet characters. This proves Theorem 1.1, with the strict half-plane  $\text{Re } s > \sigma_\tau$ .

*Remark 7.1 (Further improvements).* A substantially larger perturbation would require estimates that support the altered lengths: for example, stronger reflected-row energy or a sharper joint numerator and error-slot bound, or greater usable capacity in the inverse and ordinary witness moments. The Euler-region checks, outer-row bounds, and principal approximation would also have to remain compatible with that geometry. These estimates have not been established here for such a change. The present fixed perturbation provides no automatic iteration to the Riemann Hypothesis and no guarantee that successive improvements would reach the critical line. Broader arithmetic questions, including Hilbert's twelfth problem, are longer-term aspirations for the program; this argument claims no progress on that problem.

## REFERENCES

- [1] OpenAI. *The Quasi-Riemann Hypothesis: A Zero-Free Half-Plane  $\operatorname{Re} s > 7/8$* . September 30, 2026, 199 pp. Official PDF, at Git commit `adc7f1241b42e322a6451854ab7e4b4c146bf78a`. The main inputs here are Sections 2 and 10–20, especially printed pp. 97–107 and 183–196, together with the coefficient, reflection, and zero-detection results of Sections 4–8.
- [2] Bernhard Riemann. *Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse*. Monatsberichte der Berliner Akademie (1859), 671–680. English translation by David R. Wilkins, hosted by the Clay Mathematics Institute. The historical discussion above uses pp. 2–3 for the definition, continuation, and functional equation, and pp. 3–4 for the hypothesis.