

Morphological and physiological response of soybean treated with the microsymbiont *Bradyrhizobium japonicum* pre-incubated with genistein

A. Dolatabadian ^a, S.A.M.M. Sanavy ^{a,*}, F. Ghanati ^b, P.M. Gresshoff ^c

^a Agronomy Department, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran

^b Plant Biology Department, Faculty of Basic Science, Tarbiat Modares University, Tehran, Iran

^c ARC Centre of Excellence for Integrative Legume Research, the University of Queensland, Brisbane, Australia

Received 1 September 2011; received in revised form 26 October 2011; accepted 6 November 2011

Abstract

Genistein, a major root-secreted isoflavone of soybean (*Glycine max* (L.) Merr), is critical for the legume-*Bradyrhizobium* symbiosis as it induces several bacterial *nod*-gene systems. An experiment with soybean grown under salt stress was conducted to evaluate the effect of exogenous genistein addition to the *Bradyrhizobium* culture medium on subsequent nodulation, nitrogen fixation and selected plant physiological attributes. Five day-old plants (in pots) were inoculated with a liquid *B. japonicum* broth culture and irrigated with B&D solution containing either 0, 25, 50 and 100 mM NaCl. Four weeks after inoculation, maximum photochemical efficiency of PSII (Fv/Fm), photosynthetic rate, stomatal conductance, and transpiration rate were measured. Number of nodules per plant and apparent nitrogen fixation (as acetylene reduction activity) were determined. Salt stress decreased nodule number/plant and nitrogenase activity/plant and induced large changes of both photosynthetic parameters and antioxidant enzyme activity, compared to the control, genistein reversed the effect in each level of salinity tested. Moreover, pre-treatment of the microsymbiont with genistein enhanced maximum photochemical efficiency, photosynthetic rate, stomatal conductance and transpiration rate, while the enzymatic activities of catalase, superoxide dismutase and peroxidase in leaves and roots were not affected. It can be concluded that preincubation of the *B. japonicum* inoculant with genistein probably contributed towards growth in soybean via enhancement of nodulation and nitrogen fixation under both normal and salt stress conditions.

© 2011 SAAB. Published by Elsevier B.V. All rights reserved.

Keywords: Antioxidant enzymes; Development; Isoflavone; Photosynthetic parameters; Salinity; Symbiosis

1. Introduction

Soybean (*Glycine max* L. Merr.) is the most important legume crop in the world (Ferguson and Gresshoff, 2009), offering high-quality protein (about 40% of seed) and oil (about 20% of seed), and increasing the input of combined nitrogen as well as carbon into the soil. An integrated interaction between the soil bacterium *Bradyrhizobium japonicum* and its plant host results in the formation of nitrogen fixing root

nodules (Ferguson et al., 2010). The symbiosis benefits both partners as the prokaryotic partner receives carbohydrate in the form of sucrose-derived malate (Udvardi et al., 1988), and the symbiotic bacteria provide the plant with nitrogenous compounds. When in symbiotic association with *B. japonicum*, soybean plants can fix up to 200 kg ha⁻¹ yr⁻¹ of nitrogen (Smith and Hume, 1987), reducing the need for expensive and potentially environmentally damaging nitrogen fertilizer (Zhang et al., 2002; Sutton et al., 2011).

In general, legume plants exude into their rhizosphere complex cocktails of sugars, flavones or isoflavones, which are perceived as *nod*-gene inducers in 'Rhizobium' bacteria. Several *nod*-genes collaborate to synthesize lipo-oligo-saccharides

* Corresponding author. Tel.: +98 21 44196522 23, +98 21 44194911 4; fax: +98 21 44196524.

E-mail address: Modaresa@modares.ac.ir (S.A.M.M. Sanavy).

(Nod Factors), decorated in strain-specific fashion, to induce two parallel developmental pathways, namely root epidermal or root hair infection as well as cortical and pericycle cell division (Mathews et al., 1989a, 1989b). Perception of Nod-factor requires a dimeric receptor protein (the Nod factor receptor) made up of NFR1 and NFR5 in soybean (Indrasumunar et al., 2010, 2011) and a complex downstream signaling cascade controlled by the plant (Ferguson et al., 2010; Reid et al., 2011). Invading rhizobia penetrate the new cells of the nascent nodule, finally entering a closer symbiotic relationship in inverted plant membrane-bound vesicles called 'symbiosomes'. Here carbon and nitrogen compounds are exchanged. The combined processes lead to the formation of lumps or nodules along the root in which nitrogen fixation takes place.

Specifically for soybean, the isoflavones genistein and daidzein released by plant roots, induce the expression of common nodulation genes (*nodYABC*) of the bacterium (Kosslak et al., 1987) and also the bacterial host specific genes (such as *nodZ* and *nodFE*; Horvath et al., 1986). Almost nothing is known about the membrane transport involved in flavonoid secretion from legume roots (Sugiyama et al., 2008). Nonetheless, in response to signals from the plant, the bacteria synthesize lipochito-oligosaccharide (LCO) nod factors that deform root hairs and initiate host cell differentiation (Fisher and Long, 1992). The major lipochito-oligosaccharide for soybean nodulation consists of five β -1, 4-linked N-acetylglucosamines with various modifications (such as methyl-fucose) of the reducing and non-reducing ends, in a species-specific manner that plays a key role in determining the host specificity of the rhizobia (Supanjani et al., 2006). Daidzein and genistein are the major signal components of soybean root extracts (Kosslak et al., 1987, 1990; Sutherland et al., 1990), though daidzein has less *nod* gene-inducing ability than genistein (Mathews et al., 1989b; Sutherland et al., 1990). These substances are operative at very low concentrations (10^{-6} – 10^{-7} M) and stimulate bacterial *nod* gene expression within minutes (Ali et al., 2001). The isoflavone genistein also has antifungal activity (Rivera-Vargas et al., 1993), is the precursor to the phytoalexin kievitone produced by *Phaseolus vulgaris* (Garcia-Arenal et al., 1978) and is involved in the pathway leading to the glyceollin response of soybean cells (Graham and Graham, 2000).

Studies reported by Zhang et al. (1996), Zhang and Smith (1997) and several other studies (Bandyopadhyay et al., 1996; Pan and Smith, 1998) showed that pre-incubation of *B. japonicum* with genistein increased nodule number and nitrogen fixation.

The genistein was reported to inhibit binding of the auxin transport inhibitor naphthyl phthalamic acid (NPA) in zucchini (*Cucurbita pepo*) hypocotyl segments (Jacobs and Rubery, 1988). In legumes, flavonoid compounds were found to accumulate in cells near the site of nodule initiation. It was suggested that this leads to localized auxin accumulation, resulting in nodule organogenesis (Mathesius et al., 2000).

All stages of the soybean nitrogen fixation symbiosis are inhibited by suboptimal conditions such as drought, temperature, acidity and salinity. *Rhizobium* growth is sensitive to high osmotic pressures. So salinity can affect microbial activity via hyperosmotic stress and depress symbiotic performance (Rao et al.,

2002; Rout and Shaw, 2001). On the other hand salt stress limits plant productivity through diminished photosynthetic efficiency, carbon metabolism, leaf-chlorophyll content (Seeman and Critchley, 1985) as well as nitrogen fixation in legumes (Delgado et al., 1994; Ferri et al., 2000; Soussi et al., 1999). Salt stress (Miransari and Smith, 2007) mainly hinders the early events, more related to the physiology of the root hair such as its growth, diameter, structure and curling (Miransari et al., 2006).

Since genistein plays an important role as a signal molecule in the early stages of symbiosis establishment between soybean and *B. japonicum*, we were interested in whether salinity stress disrupt signalling, and whether pre incubation of *B. japonicum* with genistein could increase soybean nodulation and nitrogen fixation. The objective of this study was to evaluate soybean responses to salinity stress and genistein pre-treated *B. japonicum* inocula in terms of nodulation, nitrogen fixation and selected physiological and biochemical parameters.

2. Material and methods

2.1. Chemical materials

Hydrogen peroxide, Ethanol 96%, perlite and vermiculite (grade 2 or 3), 10 cm diameter plastic pots Broughton and Dilworth solutions [Add 500 ml of each stock solution per liter: solution A (2 M CaCl_2), solution B (1 M KH_2PO_4), solution C (20 mM Fe-citrate), solution D (0.5 M MgSO_4 , 0.5 M K_2SO_4 , 2 mM MnSO_4 , 4 mM H_3BO_3 , 1 mM ZnSO_4 , 4 mM CuSO_4 , 0.2 mM MnCoSO_4 , 0.2 mM Na_2MoO_4)], *Bradyrhizobium japonicum* strain CB1809, yeast extract-mannitol broth (YMB), Genistein, NaCl, Acetone, Riboflavin, L- Methionine, EDTA, Na_2CO_3 , Nitro blue tetrazolium (NBT), Sulphosalicylic acid, Guaiacol, Glacial acetic acid, Ninhydrin, Toluene, Thiobarbituric acid, Trichloroacetic acid. In this study the chemicals used were obtained from Sigma Chemical Company (N.Y., USA) or Merck Chemical Company (Deisenhofen, Deutschland).

2.2. Plant growth

Soybean seeds (*Glycine max* L. Merr.) cv. L17 were surface-sterilized in a hydrogen peroxide/ethanol solution for 2 min (10 ml of 30% H_2O_2 and 75 ml of 96% ethanol filled up to 100 ml with sterile distilled water) and rinsed several times with sterile water. Four seeds were sown in 10 cm diameter plastic pots containing autoclaved perlite and vermiculite (1:1 ratio) at depth of 1–2 cm. The pots were placed in a growth cabinet (L/D=16/8 h, T=28/25 °C), and watered with full strength of Broughton and Dilworth solution (Broughton and Dilworth, 1971). Throughout the growth period, each pot received 50 ml of B&D nutrient solution. Each treatment was replicated three times with four plants per pot.

2.3. Inocula preparation and inoculation

The inoculant was produced by culturing *Bradyrhizobium japonicum* strain CB1809 in yeast extract-mannitol broth in

250 ml flasks shaken at 150 rpm at 28 °C. After 24 h genistein was dissolved in methanol and added into yeast extract-mannitol broth to reach a final concentration of 10 μM . The lipochito-oligosaccharide biosynthesis was induced by genistein over the course of 48 h. The culture appeared greyish-white when it was compared with yeast extract-mannitol broth without genistein. The five day old plants were inoculated with yeast extract-mannitol broth. One millilitre of inoculant per plant was applied by pipette onto the rooting medium and watered with B&D solution.

2.4. Salt stress induction

Once the symbiosis was well-established, the plants were subjected to salt stress by adding NaCl to the nutrition solution at 25, 50 and 100 mM. Control plants were maintained in a NaCl-free solution. Twice a week, the plants were watered with distilled water to prevent salt accumulation.

2.5. Photosynthetic parameters

Four weeks after inoculation, maximum photochemical efficiency of PSII (Fv/Fm), photosynthetic rate (P_n : $\mu\text{mol m}^{-2}\text{s}^{-1}$), stomatal conductance (g_s : $\text{mol m}^{-2}\text{s}^{-1}$) and transpiration rate (E : $\text{mmol m}^{-2}\text{s}^{-1}$) were measured. Maximum photochemical efficiency was determined by a portable fluorometer (PAM-2000, H Wals GmbH, Effeltrich, Germany) connected with a leaf-clip holder and with a trifurcated fibre-optic (2010-F, Walz). Before measurement, the leaves were dark-adapted for 15 min. The maximum photochemical efficiency of PSII was determined from the ratio of variable (Fv) to maximum (Fm) fluorescence. A Li-6400 instrument (Li-Cor, Lincoln, USA) was used for measurement of photosynthetic rate, stomatal conductance and transpiration rate.

2.6. Plant measurement

Plants were removed from pots 30 days after inoculation. The roots were gently washed with water to remove all perlite and vermiculite and then nodule number, nodule weight, root volume as well as shoot and root weight were determined. Afterwards the shoot and root were detached and dried at 70 °C for 24 h to calculate dry weight. A sample of decapitated root with nodules of each plant was used for the nitrogenase (ARA) assay. Additionally, fresh samples were collected, quickly frozen in liquid nitrogen and stored in a deep-freezer (−80 °C) for enzyme assays and analytical determinations.

2.7. Nitrogen fixation

Nitrogen fixation of roots was assayed using the nitrogenase acetylene method reduction (Vessey, 1994). Nodulated roots from freshly harvested plants were placed in a 600 ml bottle closed with a rubber cap. Immediately, 60 ml of air were withdrawn from the closed bottle by a syringe and replaced by acetylene gas. Ethylene production was assayed in a gas chromatograph (UNICAM 4600, UK) over a short time period to prevent artefacts by closure of the variable oxygen barrier.

2.8. Chlorophyll

Chlorophyll was extracted in 80% acetone from the leaf samples according to the method of Arnon (1949). Extracts were filtered and total chlorophyll content was determined spectrophotometrically at 645 and 663 nm, respectively. Chlorophyll content was expressed as mg g^{-1} fresh weight according to Eq. (1) (Arnon, 1949).

$$\text{Total chlorophyll} = [20.2(D_{645}) + 8.02(D_{663})] \times V/1000 W \quad (1)$$

where: V=final volume; W sample weight

2.9. Antioxidant enzyme activity

Catalase activity was estimated by the method of Cakmak and Horst (1991). The reaction mixture contained 100 μl crude extract, 500 μl 10 mM H_2O_2 and 1.4 ml 25 mM potassium phosphate buffer. The decrease in absorbance was recorded at 240 nm for 1 min using a spectrophotometer (Cintra GBC, Dandenong, Victoria, Australia). Catalase activity of the extract was expressed as $\text{Abs mg}^{-1} \text{protein min}^{-1}$.

Superoxide dismutase activity was determined by measuring the ability of the enzyme extract to inhibit the photochemical reduction of nitroblue tetrazolium according to the method of Giannopolitis and Ries (1977). The reaction mixture contained 100 μl 1 μM riboflavin, 100 μl 12 mM L-methionine, 100 μl 0.1 mM EDTA (pH 7.8), 100 μl 50 mM Na_2CO_3 (pH 10.2), 100 μl 75 μM NBT in 2.3 ml 25 mM sodium phosphate buffer (pH 6.8) and 200 μl crude enzyme extract, in a final volume of 3 ml. Glass test tubes that contained the reaction mixture were illuminated with a fluorescent lamp (120 W), while identical tubes that were not illuminated served as blanks. After illumination for 45 min, absorbance was measured at 560 nm. One unit of superoxide dismutase activity was defined as the amount of enzyme that caused 50% inhibition of photochemical reduction of nitroblue tetrazolium.

Peroxidase activity was estimated by the method of Ghanati et al. (2002) via oxidation of guaiacol in the presence of H_2O_2 . The increase in absorbance at 470 nm was recorded spectrophotometrically for 1 min. The reaction mixture contained 100 μl crude extract, 500 μl 10 mM H_2O_2 , 500 μl 28 mM guaiacol and 1.9 ml 60 mM potassium phosphate buffer (pH 6.1). Peroxidase activity of the extract was expressed as $\text{Abs mg}^{-1} \text{protein min}^{-1}$. The activity of each enzyme was expressed on a protein basis. Protein concentration of the crude extract was measured by the method of Bradford (1976).

2.10. Proline

Proline content of leaves and roots was determined according to method of Bates et al. (1973). Samples (0.2 g) were homogenized in a mortar and pestle with 3 ml sulphosalicylic acid (3% w/v), and then centrifuged at 12000 rpm for 15 min. Two ml of the supernatant was added to a test tube and then two ml glacial acetic acid and two ml freshly prepared acid

ninhydrin solution were added. The test tubes were incubated in a water bath for one h at 100 °C and then allowed to cool to room temperature. Four ml of toluene was added to the tubes and mixed on a vortex mixer for 20 s. The test tubes were allowed to stand for at least 10 min, to allow separation of the toluene and aqueous phases. The toluene phase was carefully pipetted out into a glass test tube and its absorbance was measured at 520 nm in a spectrophotometer. The content of proline was calculated from a standard curve, and was expressed as mg g⁻¹ fresh weight.

2.11. Malondialdehyde

The level of membrane damage was determined by measuring the amount of malondialdehyde, which is the end product of lipid peroxidation according to De Vos et al. (1991). Samples were homogenized in 10% trichloroacetic acid (w/v) and aliquots of the filtrates were heated in 0.25% thiobarbituric acid 100 °C for 30 min. The amount of malondialdehyde in the samples was determined from the absorbance at 532 nm, followed by correction for non-specific absorbance at 600 nm using a spectrophotometer. The concentration of malondialdehyde was determined by MDA extinction coefficient ($\epsilon=155 \mu\text{M cm}^{-1}$) and the results were expressed in nmol MDA g⁻¹ fresh weight.

2.12. Leaf water potential

The amount of pressure required to force water from the cut end of the petiole equals the leaf's water potential (Smith and Prichard, 2003). A fully expanded leaf was severed and placed in a pressure bomb with its petiole protruding through the seal. Pressure was applied to the tissue until water just appeared at the cut end of the petiole, at which time pressure was recorded.

2.13. Statistical analysis

The experiment was structured following a completely randomized design arranged in a 2×4 factorial with three replications. For all variables, analysis of variance (ANOVA) was performed to test for differences between salinity, genistein treatment and their interactions using the GLM procedure in SAS version 9.1. Main and interaction effects of experimental factors were determined. Where interactions between two factors were significant, we presented the results in the form of a combination of treatments and not separately or individually. The significance of differences among treatment means was compared by LSD at the 5% probability level.

3. Results and discussion

Various morphological and physiological factors in soybean were negatively affected by salinity stress while treatment with *B. japonicum*, pre-incubated in the presence of genistein, tended to circumvent these stress effects (Table 1). The interaction between genistein and NaCl was significant on maximum photochemical efficiency (Fv/Fm), transpiration rate, photosynthesis rate, nodule number, nodule weight, acetylene reduction

activity and chlorophyll content (Table 2). On the other hand stomatal conductance, photosynthetic rate, and transpiration rate were reduced by the addition of NaCl while pre-incubation of *B. japonicum* with genistein increased these parameters when compared with the exposure of plants to NaCl alone (Table 1 and 2). Many studies have reinforced that NaCl causes plant growth inhibition by changes in photosynthetic rate (Gupta et al., 2002), stomatal conductance (Schröppel-Meier and Kaiser, 1988) and transpiration rate (Walters and Horton, 1991). Salt stress causes stomatal closure, which reduces the CO₂/O₂ ratio in leaves and inhibits CO₂ fixation. In this study, we found statistically significant differences among salinity levels for stomatal conductance. Photosynthesis and transpiration rate declined with increasing NaCl concentration due to its adverse impact on stomatal conductance this effect was more pronounced when genistein was not applied into inoculum. The highest and the lowest transpiration and photosynthesis rate were recorded in control plants inoculated with 10 μM genistein pre-incubated *B. japonicum* cells and the highest salt concentration without genistein, respectively (Table 2). Here-observed increases of stomatal conductance and transpiration rate with concomitant increased photosynthesis rate can be attributed to more nitrogen fixation and nitrogen availability due to an enhancing effect of genistein on nodulation this result is in agreement with Miransari and Smith (2008) reports.

We observed a close correlation between the photosynthetic rate and nitrogenase activity ($r^2=0.90^{**}$). Moreover, the increase in nitrogen fixation, satisfying plant demands, would stimulate photosynthesis. On the other hand, large portions of leaf nitrogen are associated with Rubisco, the key enzyme of CO₂ assimilation (Horton, 2000), therefore it would be expected that improvement of nodulation and nitrogen fixation can promote photosynthesis. These results showed that salinity and genistein application had a significant effect on Fv/Fm (Table 2). Fv/Fm ratio decreased with increasing salt concentration. In contrast, genistein treatment alleviated the effects of salt stress and enhanced Fv/Fm (Table 2). Alternatively, salinity induced significant changes in Fv/Fm in sorghum. According to our results, it seems that the effects of salt stress on Fv/Fm in soybean plants depend on the enhancing effects of genistein on nodulation and plant growth.

Significant differences among four different salinity levels and two genistein levels on shoot and root dry weight of soybean were observed (Table 1). Salinity decreased shoot, root and total dry weight while pre-inoculation of *B. japonicum* with genistein led to an increase of dry matter in both portions of the soybean plants (Table 1). The detrimental effects of NaCl on dry weight may be due to the direct effect on photosynthesis (Parida et al., 2003) and plant's ability to seek nutrients in the soil and transport them to growing shoots. Decrease in shoot dry weight accompanied by a decline in root dry weight is a normal growth phenomenon (Hawkins and Lewis, 1993). On the other hand, effect of salinity on plant growth may result from impairment of supply of photosynthetic assimilates (Kraus and Weis, 1991) and cell expansion in leaves can be inhibited by salt stress (Chartzoulakis and Klapaki, 2000). Decreases in root and shoot mass have been reported in chickpea

Table 1

The main effects of salinity and genistein levels on stomatal conductance, root dry weight, shoot dry weight, total dry weight, root volume, catalase, peroxidase and superoxide dismutase activity, malondialdehyde and proline accumulation (Plants harvested at 30 days after inoculation).

Treatments	level	Stomatal conductance (mol m ² s ⁻¹)	Root dry weight (g)	Shoot dry weight (g)	Total dry weight (g)	Root volume (cm ³)	Leaf water potential (–Mpa)	Catalase (Abs mg ⁻¹ protein min ⁻¹)		Peroxidase (Abs mg ⁻¹ protein min ⁻¹)		Superoxide dismutase (% inhibition)		Malondialdehyde (nmol MDA g ⁻¹ fresh weight)		Proline (mg g ⁻¹ fresh weight)	
								Leaf	Root	Leaf	Root	Leaf	Root	Leaf	Root	Leaf	Root
Genistein level	0 µM	0.02b	0.39b	0.73b	1.12b	9.00b	6.71a	143.65a	167.79a	159.25a	183.33a	1.43a	1.38a	1.75a	2.35a	0.03a	0.04a
	10 µM	0.04a	0.55a	1.44a	1.99a	12.41a	5.35b	140.74a	161.59a	157.45a	182.89a	1.37a	1.35a	1.77a	2.32a	0.03a	0.04a
LSD (5%)		0.005	0.14	0.12	0.23	1.24	0.29	5.98	16.27	3.84	2.01	0.15	0.09	0.08	0.07	0.003	0.003
Salinity level	0 mM	0.05a	0.97a	1.84a	2.82a	11.83a	3.73d	104.88c	113.71c	134.33d	152.60d	1.26b	1.04c	1.35c	1.96d	0.01c	0.02d
	25 mM	0.04b	0.52b	1.15b	1.67b	11.83a	4.51c	107.73c	124.15c	147.49c	167.62c	1.34b	1.33b	1.42c	2.15c	0.01c	0.03c
	50 mM	0.03c	0.27c	0.91c	1.19c	10.66a	7.20b	161.88b	192.56b	165.58b	199.91b	1.37b	1.34b	1.81b	2.44b	0.03b	0.05b
	100 mM	0.02d	0.10c	0.43d	0.54d	8.50b	8.70a	194.27a	228.38a	185.99a	212.30a	1.76a	1.63a	2.45a	2.77a	0.06a	0.06a
LSD (5%)		0.007	0.20	0.18	0.33	1.76	0.42	8.45	23.02	5.43	2.85	0.21	0.13	0.12	0.09	0.004	0.005
Sources of variation	Degree of freedom																
Genistein (G)	1	**	*	**	**	**	**	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
Salinity (S)	3	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**
G × S	3	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns
Error	16	0.00	0.02	0.02	0.07	2.08	0.11	47.76	353.74	19.74	5.43	0.03	0.01	0.00	0.00	0.00	0.00
Coefficient of variations (%)		15.06	34.98	13.78	17.57	13.47	5.70	4.86	11.41	2.80	1.27	12.68	8.12	5.62	3.49	11.72	9.13

ns; not significant, * significant at P<0.1; **significant at P<0.05. Values within the each column and followed by the same letter are not different at P<0.05 by an ANOVA protected LSD Test

Table 2
Interaction of salinity and genistein levels on maximum photochemical efficiency, transpiration rate, photosynthetic rate, nodule number, nodule weight, nitrogenase activity, chlorophyll content and leaf water potential (Plants harvested at 30 days after inoculation).

Genistein level	Salinity level (mM)	Fv/Fm	Transpiration rate (mmol m ² s ⁻¹)	Photosynthetic rate (μmol m ² s ⁻¹)	Nodule number	Nodule weight (g)	Nitrogenase activity (μmol C ₂ H ₄ core ⁻¹ h ⁻¹)	Chlorophyll (mg g ⁻¹ fresh weight)
(0 μM)	0	656.33e	1.48c	14.45 d	42.00 d	0.24b	7.47e	2.91c
	25	644.66f	1.35de	12.52e	36.00e	0.23c	6.59f	2.33 d
	50	632.00 g	1.28e	10.85ef	24.66f	0.16 d	6.22 fg	1.54 g
	100	613.00 h	0.99f	10.60f	16.66 g	0.09e	5.76 g	1.39 g
(10 μM)	0	716.66a	1.86a	25.50a	74.66a	0.35a	18.16a	3.65a
	25	687.33b	1.58b	20.13b	68.33b	0.35a	17.29b	3.42b
	50	677.33c	1.53bc	18.33bc	47.00c	0.24b	10.28c	2.15e
	100	664.66 d	1.44 cd	16.78c	43.66 cd	0.24bc	8.10 d	1.76f
LSD (5%)		8.00	0.10	1.81	4.92	0.007	0.62	0.17
Sources of variation	Degree of freedom							
Genistein	1	**	**	**	**	**	**	**
Salinity	3	**	**	**	**	**	**	**
Genistein × Salinity	3	*	*	**	*	**	**	**
Error	16	21.37	0.00	1.09	8.08	0.00	0.12	0.01
Coefficient of variations (%)		0.69	4.10	6.47	6.44	1.89	3.59	4.28

* Significant at P<0.1; **significant at P<0.05, Values within the each column and followed by the same letter are not different at P<0.05 by an ANOVA protected LSD Test.

plants (Singh et al., 2001). We observed that 10 μ M genistein significantly increased shoot and root dry weight relative to the control. In addition, at all levels of salinity, genistein resulted in higher nodule number and nodule weight compared with lack of genistein in stressed plants (Table 2). Indicating that genistein pre-inoculation caused activation and hence more efficiency of nodule bacteria to nitrogen fixation and hence, increased soybean growth. It has been reported that genistein stimulated plant growth (e.g., greater leaf area) through activation of *nod* genes in interacting *B. japonicum* (Zhang and Smith, 1995).

Root volume was reduced in response to salinity (Table 1). Significant difference was observed when NaCl concentration was increased from 50 mM to 100 mM. The inhibition of root growth in term of root volume can be attributed to the inhibition of mitosis, reduced synthesis of cell wall components, damage to the Golgi apparatus and changes in polysaccharide metabolism (Berkelaar and Beverley, 2000). Our findings are consistent with the findings of Siroka et al. (2004) who have reported that salinity decreases root cell development. The effect of genistein on root volume was also significant (Table 1). One possible explanation is that root volume increases with genistein because more photosynthate is transferred to the roots as a result of increased nitrogen fixation and photosynthesis. Moreover, this could be due to the Nod factor production by the inoculum. Pre-incubating *B. japonicum* promotes expression of the *nod* gene (Kosslak et al., 1987) and, thus the production of Nod factor, which can regulate plant morphogenesis at very low concentrations (Spaink, 1996).

Nodule initiation in the legume-*Rhizobium* symbiosis involves a complex interaction between host root, rhizobial strain, and the environment. In this study, we showed that salt treatments significantly reduced nodule number and nodule weight per plant while inoculated soybean plants with genistein pre-incubated *B. japonicum* cells improved soybean nodulation (Table 2). Reduced nodule formation could have been due to the adverse effects of salinity on nodule initiation. In addition, nodule weight decreased under salinity stress it seems that it is because of less photosynthate transport to the roots. The adverse effects of salinity on the legume-*Bradyrhizobium* symbiosis is through affecting rhizobial survival and growth, hindering the infection process, suppressing nodule function, and reducing plant ability to photosynthesis (Elsheikh and Wood, 1995). Our results also show that pre-incubation of *B. japonicum* with 10 μ M genistein increased nodule number and nodule weight per plant. This could be due to an increase either in the number of infections initiated or in the proportion of infections leading to nodule formation (Zhang and Smith, 1995). The significant interaction effects also verify that genistein plays a direct role in overcoming of salinity. Many studies have been reported that genistein addition stimulates the production of lipochito-oligosaccharide under normal and stressful conditions (Lerouge et al., 1990; Zhang and Smith, 1995) which in turn elicits root hair deformation, cortical cell division and nodulin gene expression in soybean roots (Spaink, 1996). Faster formation of nodules in soybean plants and increased

nodule weight following genistein treatment were also found by Pan et al. (1998).

The effects of genistein on the establishment of soybean-*Rhizobium* symbiosis (Eckardt, 2006), increased the nodule numbers and nodule weight hence resulted in the acceleration of C_2H_2 reduction under salt stress conditions. This showed that the effect of salinity on soybean nodulation and nitrogen fixation is less when plants were treated with pre-incubated *B. japonicum* with genistein. Acetylene reduction assays for nitrogenase activity showed that the maximum and the minimum C_2H_4 concentration were determined from the treated plants with pre-incubated *B. japonicum* with 10 μ M genistein under no stress condition and treated plants with 100 mM NaCl without genistein application, respectively (Table 2). It would seem that increase of C_2H_2 reduction can be due to an increase in nodule number as well as nodule weight affected by genistein. Decreased ability of nodules to reduce C_2H_2 under salinity has been well-documented for other legumes (Ferri et al., 2000). The inhibition of acetylene reduction by salt stress may be due to a limitation of oxygen diffusion in nodules or due to toxic effects of Na or Cl accumulation (Serraj et al., 1998). On the other hand, reduction of photosynthetic activity by salt can also reduce nitrogen fixation.

In this experiment there are two possible explanations for the stimulative effects of genistein on nitrogen fixation improvement under conditions of salt stress. First, since genistein has been identified as a major inducer of *nod* genes in *B. japonicum* (Kosslak et al., 1987), genistein could have increased the infection rate by endogenous *B. japonicum*, resulting in increased soybean nodulation and nitrogen fixation. Second, increased C_2H_2 reduction could be partially due to other functions of genistein such as role of genistein in overcoming adverse effect of salinity (Miransari and Smith, 2009). Our results imply that increases of nitrogen fixation would lead to more pronounced nodulation by the addition of genistein. Consistently, increase in nitrogen fixation would correspond to the dilution of *nod* gene inducers from the root exudates and *nod* factors formed by bacteria.

Salinity significantly increased the leaf water potential while genistein decreased. It is well-known that, salt stress reduces root hydraulic conductivity resulting in decreased water flow from roots to shoot. This decrease in water flow due to salt stress may cause a lowering in leaf water content that would result in stomatal closure in order to maintain their water status (Robinson et al., 1997). Reduction in leaf water potential because of genistein application into inocula could be due to this probability that genistein can seriously change the photosynthetic carbon metabolism, leaf chlorophyll content, photosynthetic efficiency as well as root growth, through improvement of symbiosis relationships, nodulation and nitrogen fixation. Furthermore, the decrease in leaf water potential may be due to the accumulation of osmolytes, which are direct products of photosynthesis (Monneveux and Belhassen, 1996).

Saline stress led to the yellowing of leaves, which ultimately resulted in significant damage to the chlorophyll pigments. Similar results have been reported for other legumes (Al-Khanjari et al., 2002). Leaf chlorophyll content was significantly reduced in

soybean plants either with or without genistein in bacterial growth medium. The inhibitory effects of salt on chlorophylls could be due to suppression of specific enzymes responsible for the synthesis of chlorophyll. Alternatively, the decrease in chlorophyll may be attributed to increased chlorophyllase activity (Sudhakar et al., 1997).

Chlorophyll content was increased on account of genistein application under conditions of salt stress (Table 2). Since nitrogen is a critical component of chlorophyll, and without sufficient quantities of this element, chlorophyll cannot be formed (Tucker, 2004); legume–*Rhizobium* symbiosis plays an important role in nitrogen supplementation in these plants. Genistein as a signal molecule stimulates lipochito-oligosaccharide production by the bacterial symbiont (Lerouge et al., 1990) and improve legume–*Rhizobium* interactions during nodule formation and development.

Malondialdehyde content can serve as an indicator of the rate of oxidative processes in cells. The effects of NaCl and genistein on malondialdehyde content are presented in Table 1. Genistein application had no significant effect on malondialdehyde accumulation in roots and leaves. In contrast, salinity stress increased malondialdehyde whether in leaf tissue or root. Malondialdehyde results indicated that, lipid peroxidation started to increase at 50 mM NaCl in leaves. In other words, there was no significant difference between control and 25 mM NaCl treatments. Our results indicated that root cells were more sensitive in response to salt stress because malondialdehyde content in roots was significantly higher than in leaves (Table 1). Accumulation of malondialdehyde was reported in a number of salt-sensitive plants (Gehlot et al., 2003; Luna et al., 2002; Mittova et al., 2002).

In this study, proline content increased in NaCl treated plants while it was not affected by genistein (Table 1). In leaves, proline accumulation was initiated at 50 and 100 mM NaCl stress as compared to control and 25 mM NaCl treatments. Root proline content increased at 25 mM NaCl. Proline accumulates in plant tissues under saline stress and this substance is suspected of contributing to osmotic adjustment (Delauney and Verma, 1993). Proline protects membranes and proteins against the adverse effects of high concentrations of inorganic ions and temperature extremes (Rudolph et al., 1986; Santoro et al., 1992). Proline may also function as a protein-compatible hydrotrope (Srinivas and Balasubramanian, 1995), and as a hydroxyl radical scavenger (Smimoff and Cumbe, 1989).

Activities of antioxidant enzymes in roots and shoots exhibited a similar change. That is to say, genistein application had not significant effect on enzyme activity but obvious differences existed due to salinity stress (Table 1). Specifically, peaks of activities of antioxidant enzymes in roots and shoots appeared at the highest NaCl concentration. Activities of catalase and peroxidase in the two parts of plant showed a similar trend, but the activity of superoxide dismutase in leaves was significantly greater than roots (Table 1). Additionally, the increase in superoxide dismutase activity in roots arose earlier. In other words, superoxide dismutase activity in roots is more sensitive than shoots. As mentioned previously, the peak of antioxidant enzyme activity in roots and shoots appeared at the

highest salinity level. It indicates that reactive oxygen species scavenging system was activated efficiently and removed reactive oxygen species. Hernandez et al. (2001) suggested that salt stress can cause oxidative stress and enhance the antioxidant capacity in the leaf apoplast of pea cultivars. Superoxide dismutase is an antioxidant enzyme with the ability to repair oxidation damage caused by reactive oxygen species. Thus, superoxide dismutase is considered a key enzyme for maintaining normal physiological conditions and coping with oxidative stress in the regulation of intracellular levels of reactive oxygen species (Mittler, 2002). Also peroxidase is widely distributed in higher plants where it is involved in various processes, including lignification, auxin metabolism, salt tolerance and heavy metal stress (Passardi et al., 2005). Catalase, which is involved in the degradation of hydrogen peroxide into water and oxygen, is the most effective antioxidant enzymes in preventing oxidative damage (Mittler, 2002). Even under optimal conditions many metabolic processes produce reactive oxygen species. Plants possess efficient systems for scavenging reactive oxygen species that protect them from destructive oxidative reactions (Foyer et al., 1994). As part of this system, antioxidant enzymes are key elements in defense mechanisms. Many changes have been observed in the activities of antioxidant enzymes in plants under salt stress. The activity of antioxidant enzymes has been reported to increase under saline conditions in the case of salt-tolerant cotton (Meloni et al., 2003), rice (Fadzila et al., 1997) and wheat (Meneguzzo et al., 1999).

4. Conclusion

Overall, the results reported here show that salinity significantly reduced photosynthetic parameters, shoot and root dry weights, root volume, nodule number, nodule weight, nitrogenase activity, and chlorophyll content while antioxidant enzyme activity, proline and malondialdehyde content increased. Additionally, the results indicate that genistein was able to increase photosynthesis, nodulation and nitrogen fixation and subsequently soybean growth under saline and non-saline conditions. However, antioxidant enzymes activity was not affected by genistein application. In conclusion, our results indicate that pre-incubation of *B. japonicum* with genistein improves nodulation and nitrogen fixation of soybean subjected to salt stress and it would be recommended to use under saline soils to attain desirable symbiosis and soybean production.

References

- Ali, M., Belkheir, X.Z., Smith, D.L., 2001. Variability in yield and yield component responses to genistein pre-incubated *Bradyrhizobium japonicum* by soybean [*Glycine max* (L.) Merr] cultivars. *Plant and Soil* 229, 41–46.
- Al-Khanjari, S., Al-Kathiri, A., Esehie, H.A., 2002. Variation in chlorophyll meter readings, nodulation and dry matter yields of alfalfa (*Medicago sativa* L.) cultivars differing in salt tolerance. *Crop Research* 24, 350–356.
- Arnon, D.I., 1949. Copper enzymes in isolated chloroplasts, polyphenoloxidase in *Beta vulgaris*. *Plant Physiology* 24, 1–150.
- Bandyopadhyay, A.K., Jain, V., Nainawate, H.S., 1996. Nitrate Alters the Flavonoid Profile and Nodulation in Pea (*Pisum sativum* L.). *Biology and Fertility of Soils* 21, 189–192.

- Bates, L.S., Waldern, R.P., Teave, I.D., 1973. Rapid determination of free proline for water stress studies. *Plant and Soil* 39, 205–207.
- Berkelaar, E., Beverley, H., 2000. The relationship between morphology and cadmium accumulation in seedlings of two durum wheat cultivars. *Canadian Journal of Botany* 78, 381–387.
- Bradford, M.A., 1976. Rapid and sensitive method for the quantitation of protein utilizing the principle of protein-dye binding. *Annual Review of Biochemistry* 72, 248–254.
- Broughton, W.J., Dilworth, M., 1971. Control of leghaemoglobin synthesis in snake beans. *Biochemistry Journal* 125, 1075–1080.
- Cakmak, I., Horst, W., 1991. Affect of aluminium on lipid peroxidation, superoxide dismutase, catalase and peroxidase activities in root tip of soybean (*Glysin max*). *Plant Physiology* 83, 463–468.
- Chartzoulakis, K., Klapaki, G., 2000. Response of two green house pepper hybrids to NaCl salinity during different growth stages. *Scientia Horticulturae* 86, 247–260.
- De Vos, C., Schat, H.M., De Waal, M.A., Vooijs, R., Ernst, W., 1991. Increased to copper-induced damage of the root plasma membrane in copper tolerant *Silene cucubalus*. *Plant Physiology* 82, 523–528.
- Delauney, A.J., Verma, D.P.S., 1993. Proline biosynthesis and osmoregulation in plants. *The Plant Journal* 4, 215–223.
- Delgado, M.J., Ligerio, F., Lluch, C., 1994. Effects of salt stress on growth and nitrogen fixation by pea, faba-bean, common bean, and soybean plants. *Soil Biology and Biochemistry* 26, 371–376.
- Eckardt, N.A., 2006. The role of flavonoids in root nodule development and auxin transport in *Medicago truncatula*. *The Plant Cell* 18, 1539–1540.
- Elsheikh, E.A.E., Wood, M., 1995. Nodulation and N₂ fixation by soybean inoculated with salt-tolerant rhizobia or salt-sensitive *bradyrhizobia* in saline soil. *Soil Biology and Biochemistry* 27, 657–661.
- Fadzilla, N.M., Finch, R.P., Burdon, R.H., 1997. Salinity, oxidative stress and antioxidant responses in shoot cultures of rice. *Journal of Experimental Botany* 48, 325–331.
- Ferguson, B.J., Gresshoff, P.M., 2009. Soybean as a model legume. *Grain Legumes* 53, 7.
- Ferguson, B.J., Indrasumunar, A., Hayashi, S., 2010. Genetic analysis of legume nodule development and autoregulation. *Journal of Integrated Plant Biology* 52, 61–76.
- Ferri, A., Lluch, C., Ocana, A., 2000. Effect of salt stress on carbon metabolism and bacteroid respiration in root nodules of common bean (*Phaseolus vulgaris* L.). *Plant Biology* 2, 396–402.
- Fisher, R.F., Long, S.R., 1992. *Rhizobium*-plant signal exchange. *Nature* 357, 655–660.
- Foyer, C.H., Lelandais, M., Kunert, K.J., 1994. Photooxidative stress in plants. *Physiologia Plantarum* 92, 696–717.
- Garcia-Arenal, F., Fraile, A., Sagasta, E.M., 1978. The multiple phytoalexin response of bean (*Phaseolus vulgaris*) to infection by *Botrytis cinerea*. *Physiological Plant Pathology* 13, 151–156.
- Gehlot, H.S., Dinesh, R., Choudhary, R., Joshi, S., Sankhla, N., 2003. In vitro studies on chickpea seedlings: Effect of salt on growth and antioxidant activity. *Proceeding Plant Growth Regulation Society of America* 30, 119–121.
- Ghanati, F., Morita, A., Yokota, H., 2002. Induction of suberin and increase of lignin content by excess boron in tobacco cell. *Soil Science and Plant Nutrition* 48, 357–364.
- Giannopolitis, C., Ries, S., 1977. Superoxide dismutase occurrence in higher plant. *Plant Physiology* 59, 309–314.
- Graham, T.L., Graham, M.Y., 2000. Defense potentiation and competency: redox conditioning effects of salicylic acid and genistein. In: Keen, N., Stacey, G. (Eds.), *Plant-Microbe Interactions*, Vol. 5. APS Press, St. Paul.
- Gupta, N.K., Meena, S.K., Gupta, S., Khandelwal, S.K., 2002. Gas exchange, membrane permeability and ion uptake in two species of Indian jujube differing in salt tolerance. *Photosynthetica* 40, 535–539.
- Hawkins, H.J., Lewis, O.A.M., 1993. Combination effect of NaCl salinity, nitrogen form and calcium concentration on the growth, ionic content and gaseous exchange properties of *Triticum aestivum* L.cv. Gamtoos. *New Phytologist* 124, 161–170.
- Hernandez, J.A., Ferrer, M.A., Jimenez, A., Ros-Barcelo, A., Sevilla, F., 2001. Antioxidant systems and O₂-H₂O₂ production in the apoplast of *Pisum sativum* L. leaves: its relation with NaCl induced necrotic lesions in minor veins. *Plant Physiology* 27, 817–831.
- Horton, P., 2000. Prospects for Crop Improvement through the Genetic Manipulation of Photosynthesis: Morphological and Biochemical Aspects of Light Capture. *Journal of Experimental Botany* 51, 475–485.
- Horvath, B., Kondorosi, E., John, M., Schmidt, J., Török, I., Györgypal, Z., Barabas, I., Wieneke, U., Schell, J., Kondorosi, A., 1986. Organization, structure and symbiotic function of *Rhizobium meliloti* nodulation genes determining host specificity for alfalfa. *Cell* 46, 335–343.
- Indrasumunar, A., Kereszt, A., Searle, I., Miyagi, M., Li, D., Nguyen, C., Men, A., Carroll, B., Gresshoff, P., 2010. Inactivation of duplicated nod factor receptor 5 (NFR5) genes in recessive loss-of-function nonnodulation mutants of allotetraploid soybean (*Glycine max* L. Merr.). *Plant & Cell Physiology* 51, 201–214.
- Indrasumunar, A., Searle, I., Lin, M., Kereszt, A., Men, A., Carroll, B.J., Gresshoff, P.M., 2011. Limitation of nodule organ number by nodulation factor receptor kinase 1a transcription in soybean (*Glycine max* L. Merr.). *The Plant Journal* 65, 39–50.
- Jacobs, M., Rubery, P.H., 1988. Naturally occurring auxin transport regulators. *Science* 241, 346–349.
- Kosslak, R.M., Rookland, R., Barkei, J., Paaren, H.E., Appelbaum, E.R., 1987. Induction of *Bradyrhizobium japonicum* common nod genes by isoflavones isolated from *Glycine max*. *Proceedings of the National Academy of Sciences of the United States of America* 34, 7428–7432.
- Kosslak, R.M., Joshi, R.S., Bowen, B.A., Paaren, H.E., Appelbaum, E.R., 1990. Strain-specific inhibition of nod gene induction in *Bradyrhizobium japonicum* by flavonoid compounds. *Applied and Environmental Microbiology* 56, 1333–1341.
- Kraus, G.H., Weis, E., 1991. Chlorophyll fluorescence and photosynthesis: the basis. *Annual Review of Plant Physiology* 136, 472–479.
- Lerouge, P., Roche, P., Faucher, C., Maillet, F., Truchet, G., Prome, J.C., Denarie, J., 1990. Symbiotic host specificity of *Rhizobium meliloti* is determined by a sulphated and acylated glucosamine oligosaccharide signal. *Nature* 344, 781–784.
- Luna, C., Luca, M.D., Taleisnik, E., 2002. Physiological causes for decreased productivity under high salinity in bom tetraploid *Chloris gayana* cultivar. II. Oxidative stress. *Australian Journal of Agricultural Research* 53, 669–674.
- Mathesius, U., Weinman, J.J., Rolfe, B.G., Djordjevic, M.A., 2000. Rhizobia can induce nodules in white clover by ‘hijacking’ mature cortical cells activated during lateral root development. *Molecular Plant-Microbe Interactions* 13, 170–182.
- Mathews, A., Carroll, B.J., Gresshoff, P.M., 1989a. A new recessive gene conditioning non-nodulation in soybean. *Journal of Heredity* 80, 357–360.
- Mathews, A., Kosslak, R.M., Senguptagopalan, C., Appelbaum, E.R., Carroll, B.J., Gresshoff, P.M., 1989b. Biological characterization of root exudates and extracts from non-nodulating and super-nodulating soybean mutants. *Molecular Plant-Microbe Interaction* 2, 283–290.
- Meloni, D.A., Oliva, M.A., Martinez, C.A., Cambraia, J., 2003. Photosynthesis and ability of superoxide dismutase, peroxidase and glutathione reductase in cotton under salt stress. *Environmental and Experimental Botany* 49, 69–76.
- Meneguzzo, S., Navari-Izzo, F., Izzo, R., 1999. Antioxidative responses of shoots and roots of wheat to increasing NaCl concentrations. *Journal of Plant Physiology* 155, 274–280.
- Miransari, M., Smith, D.L., 2007. Overcoming the stressful effects of salinity and acidity on soybean [*Glycine max* (L.) Merr.] nodulation and yields using signal molecule genistein under field conditions. *Journal of Plant Nutrition* 30, 1967–1992.
- Miransari, M., Smith, D.L., 2008. Using signal molecule genistein to alleviate the stress of suboptimal root zone temperature on soybean-*Bradyrhizobium* symbiosis under different soil textures. *Journal of Plant Interactions* 3, 287–295.
- Miransari, M., Smith, D.L., 2009. Alleviating salt stress on soybean (*Glycine max* (L.) Merr.) *Bradyrhizobium japonicum* symbiosis, using signal molecule genistein. *European Journal of Soil Biology* 4, 146–152.
- Miransari, M., Smith, D.L., Mackenzie, A.F., Bahrami, H.A., Malakouti, M.J., Rejali, F., 2006. Overcoming the stressful effect of low pH on soybean root hair curling using lipochitoooligosaccharides. *Communications in Soil Science and Plant Analysis* 37, 1103–1110.
- Mittler, R., 2002. Oxidative stress, antioxidants and stress tolerance. *Trends in Plant Science* 7, 405–410.

- Mittova, V., Tal, M., Volokita, M., Guy, M., 2002. Salt stress induces up-regulation of an efficient chloroplast antioxidant system in the salt-tolerant wild tomato species *Lycopersicon pennellii* but not in the cultivated species. *Physiologia Plantarum* 115, 393–400.
- Monneveux, P., Belhassen, E., 1996. The diversity of drought adaptation in the wide. In: Belhassen, E. (Ed.), *Drought Tolerance in Higher Plants: Genetic, Physiological and Molecular Biological Analysis*. Kluwer Academic Publishers.
- Pan, B.F., Smith, D.L., 1998. Genistein Addition to the Rooting Medium of Soybean at the Onset of Nitrogen Fixation Increases Nodulation. *Journal of Plant Nutrition* 21, 1631–1639.
- Parida, A.K., Das, A.B., Mittra, B., 2003. Effects of NaCl stress on the structure, pigment complex composition, and photosynthetic activity of mangrove *Bruguiera parviflora* chloroplasts. *Photosynthesis* 41, 191–200.
- Passardi, F., Cosio, C., Penel, C., Dunand, C., 2005. Peroxidases have more functions than a Swiss army knife. *Plant Cell Reports* 24, 255–265.
- Rao, D.L.N., Giller, K.E., Yeo, A.R., Flowers, T.J., 2002. The effects of salinity and sodicity upon nodulation and nitrogen fixation in chickpea (*Cicer arietinum*). *Annual Botany* 89, 563–570.
- Reid, D.E., Ferguson, B.J., Gresshoff, P.M., 2011. Inoculation and nitrate-induced CLE peptides of soybean control NARK-dependent nodule formation. *Molecular Plant-Microbe Interaction* 24, 606–618.
- Rivera-Vargas, L.I., Cshmitthener, A.F., Graham, T.L., 1993. Soybean flavonoid effects on and metabolism by *Phytophthora sojae*. *Phytochemistry* 32, 851–857.
- Robinson, M.F., Véry, A.A., Sanders, D., Mansfield, T.A., 1997. How can stomata contribute to salt tolerance? *Annual Botany* 80, 387–393.
- Rout, N.P., Shaw, B.P., 2001. Salt tolerance in aquatic macrophytes: Ionic relation and interaction. *Biologia Plantarum* 44, 91–95.
- Rudolph, A.S., Crowe, J.H., Crowe, L.M., 1986. Effects of three stabilizing agents proline, betaine and trehalose on membrane phospholipids. *Archives of Biochemistry and Biophysics* 245, 134–143.
- Santoro, M.M., Liu, Y., Khan, S.M.A., Hou, L.X., Bolen, D.W., 1992. Increased thermal stability of proteins in the presence of naturally occurring osmolytes. *Biochemistry* 31, 5278–5283.
- Schröppel-Meier, G., Kaiser, W.M., 1988. Ion homeostasis in chloroplasts under salinity and mineral deficiency. I. Solute concentrations in leaves and chloroplasts from spinach plants under NaCl or NaNO₃ salinity. *Plant Physiology* 87, 822–827.
- Seeman, J.R., Critchley, C., 1985. Effects of salt stress on the growth, ion content, stomatal behaviour and photosynthetic dro capacity of salt-sensitive species, *Phaseolus vulgaris* (L). *Planta* 164, 151–162.
- Serraj, R., Vazquez-Diaz, H., Drevon, J.J., 1998. Effects of salt stress on nitrogen fixation, oxygen diffusion, and ion distribution in soybean, common bean, and alfalfa. *Journal of Plant Nutrition* 21, 475–488.
- Singh, A.K., Singh, R.A., Sharma, S.G., 2001. Salt stress induced changes in certain organic metabolites during seedling growth of chickpea. *Legume Research* 24, 11–15.
- Siroka, B., Huttova, J., Tamas, L., Simonoviva, M., Mistrik, I., 2004. Effect of cadmium on hydrolytic enzymes in maize root and coleoptile. *Biologia* 59, 513–517.
- Smimoff, N., Cumbes, Q.J., 1989. Hydroxyl radical scavenging activity of compatible solutes. *Phytochemistry* 28, 1057–1060.
- Smith, D.L., Hume, D.J., 1987. Comparison of assay methods for N₂ fixation utilizing white bean and soybean. *Canadian Journal of Plant Science* 67, 11–19.
- Smith, R., Prichard, T., 2003. Using a Pressure Chamber in Winegrapes. UC Cooperative Extension Available at <http://cesonoma.ucdavis.edu/?search=yes2003>.
- Soussi, M., Lluch, C., Ocana, A., 1999. Comparative study of nitrogen fixation and carbon metabolism in two chickpea (*Cicer arietinum* L.) cultivars under salt stress. *Journal of Experimental Botany* 50, 1701–1708.
- Spaink, H.P., 1996. Regulation of plant morphogenesis by lipo-chitin oligosaccharides. *CRC Critical Reviews in Plant Science* 15, 559–582.
- Srinivas, V., Balasubramanian, D., 1995. Proline is a protein-compatible hydro-trope. *Langmuir* 11, 2830–2833.
- Sudhakar, C., Ramanjulu, S., Reddy, P.S., Veeranjanyulu, K., 1997. Response of some calvin cycle enzymes subjected to salinity shock in vitro. *Indian Journal of Experimental Botany* 35, 665–667.
- Sugiyama, A., Shitan, N., Kyazaki, K., 2008. Signalling from soybean roots to *Rhizobium*. *Plant Signaling & Behavior* 3, 38–40.
- Supanjani, S., Habib, A., Mabood, F., Lee, K.D., Donnelly, D., Smith, D.L., 2006. Nod factor enhances calcium uptake by soybean. *Plant Physiology and Biochemistry* 44, 866–872.
- Sutherland, T.D., Bassam, B.J., Schuller, L.J., Gresshoff, P.M., 1990. Early nodulation signals of the wild type and symbiotic mutants of soybean (*Glycine max*). *Molecular Plant-Microbe Interactions* 3, 122–128.
- Sutton, M.A., Erisman, W., Van Grinsven, H., Winiwarter, W., 2011. Too much of a good thing. *Nature* 472, 159–161.
- Tucker, M., 2004. Primary Nutrients and Plant Growth. In: SCRIBD (Ed.), *Essential Plant Nutrients*. North Carolina Department of Agriculture.
- Udvardi, M.K., Price, G.D., Gresshoff, P.M., Day, D.A., 1988. A dicarboxylate transporter on the peribacteroid membrane of soybean nodules. *Federation of European Biochemical Societies* 231, 36–40.
- Vessey, J.K., 1994. Measurement of nitrogenase activity in legume root nodules: In defence of the acetylene reduction assay. *Plant and Soil* 158, 151–162.
- Walters, R.G., Horton, P., 1991. Resolution of components on nonphotochemical chlorophyll fluorescence quenching in barley leaves. *Photosynthesis Research* 27, 121–133.
- Zhang, F., Lynch, D.H., Smith, D.L., 1996. Inoculation of soybean [*Glycine max* (L.) Merr.] with genistein pre-incubated *Bradyrhizobium japonicum* or genistein directly applied into soil increases soybean protein and dry matter yield under short season conditions. *Plant and Soil* 179, 233–241.
- Zhang, F., Smith, D.L., 1995. Pre-incubation of *Bradyrhizobium japonicum* with genistein accelerates nodule development of soybean [*Glycine max* (L.) Merr.] at sub-optimal root zone temperatures. *Plant Physiology* 108, 961–968.
- Zhang, F., Smith, D.L., 1997. Application of genistein to inocula and soil to overcome low spring soil temperature inhibition of soybean nodulation and nitrogen fixation. *Plant and Soil* 192, 141–151.
- Zhang, H., Prithiviraj, B., Souleimanov, A., Aoust, F.D., Charles, T.C., Driscoll, B.T., Smith, D.L., 2002. The effect of temperature and genistein concentration on lipo-chitooligosaccharide (LCO) production by wild-type and mutant strains of *Bradyrhizobium japonicum*. *Soil Biology and Biochemistry* 34, 1175–1180.