

1 **Depth of the source of the tectonic tremor in the**
2 **eastern Olympic Peninsula, Washington**

3 **Ariane Ducellier¹**

4 ¹University of Washington

5 **Key Points:**

- 6 • enter point 1 here
7 • enter point 2 here
8 • enter point 3 here

Abstract

enter abstract here

1 Introduction

Tremor is a long (several seconds to many minutes), low amplitude seismic signal, with emergent onsets, and an absence of clear impulsive phases. Tectonic tremor have been explained as a swarm of small, low frequency earthquakes (?), that is small magnitude earthquakes ($M \sim 1$) which frequency content (1-10 Hz) is lower than for ordinary earthquakes (up to 20 Hz). The source of the low-frequency earthquakes is located on the plate boundary, and their focal mechanisms represent shear slip on a low-angle thrust fault dipping in the same direction as the plate interface (?). Due to the lack of clear impulsive phases in the tremor signal, it is difficult to determine the depth of the tremor source with the same precision, and it is assumed to be also located close to the plate boundary. In subduction zones such as Nankai and Cascadia, tectonic tremor observations are spatially and temporally correlated with slow slip observations (?); (?). Due to this correlation, these paired phenomena have been called Episodic Tremor and Slip (ETS).

Specific question - Depth of the low-frequency earthquake on the plate boundary. More difficult for the tremor because no phase onset.

Explain here about the link between tremor and quartz vein. Problem: LFE on a plane / zone with quartz vein is several km thick. Interested in location and thickness of the tremor zone.

References on quartz veins

Audet and Bürgmann (2014 (?)) studied the relationship between the ratio between P-wave velocity and S-wave velocity in the subducted oceanic crust and the forearc and the periodicity of slow earthquakes. They computed the V_P/V_S ratio from receiver functions and data from the literature. They noticed that slow earthquakes are associated with a high V_P/V_S ratio in the subducted oceanic crust, but without relationship with recurrence time. However, they pointed out that the recurrence time of slow earthquakes increases linearly with the V_P/V_S ratio of the forearc. Moreover, along a margin-perpendicular profile from northern Cascadia, the V_P/V_S ratio of the forearc, and the recurrence time of Episodic Tremor and Slip (ETS) events, decrease with increasing depth. The authors explained the low V_P/V_S ratio in the forearc by the enrichment of forearc minerals in fluid-dissolved silica derived from the dehydration of the downgoing slab. However, they estimated that the fluid flux required for the formation of quartz veins was two orders of magnitude greater than the fluid production rates estimated from the dehydration of the slab. They hypothesized that silica-saturated fluids may originate from the complete serpentinization of the mantle near the wedge corner. They suggested that higher temperature and quartz content at depth may lead to faster dissolution - precipitation processes and more frequent slip events. Their model could also explain the global variation in recurrence time, with mafic silica-poor regions having longer ETS recurrence times than felsic silica-rich regions.

Hyndman *et al.* (2015 (?)) investigated the processes that control the position of Episodic Tremor and Slip (ETS) in the Cascadia subduction zone. They noticed that the high temperatures in the young subducting oceanic plate, the geodetic data, and the recordings of coseismic subsidence in buried coastal marshes during past great earthquakes, all point out to a downdip limit of the seismogenic zone located offshore. The position of the slow slip and the tremor is well known, although the depths have some uncertainty. The slip may extend seaward of the tremor, but there is a clear separation between the seismogenic zone and the ETS zone, with the ETS zone being located about 70 km east

of the downdip of the seismogenic zone, and the volcanic arc being located about 100 km east of the ETS zone. A previous study showed that the position of the subduction zone ETS does not coincide with a specific temperature or dehydration reaction. The authors pointed out that ETS has been related to high pore fluid pressures close to the plate boundary. They argued that the bending of the subducting plate at the ocean trench may introduce a large amount of water in the upper oceanic mantle, resulting in extensive serpentinization. Moreover, the serpentinization of the fore-arc mantle corner may increase its vertical impermeability, while keeping a high permeability parallel to the fault, thus channelling all the fluid updip in the subducting oceanic crust. The dehydration of the serpentinite from the upper oceanic mantle, and the focusing of rising fluids along the plate boundary should result in large amounts of fluids available at the fore-arc mantle corner. Additionally, there seems to be a good coincidence between the location of the fore-arc mantle corner, and the location of ETS. The authors then observed that the deep fore-arc crust has a very low Poisson's ratio (less than 0.22), and that the only mineral with a very low Poisson's ratio is quartz (about 0.1), which led them to conclude that there may be a significant amount of quartz (about 10 % in volume) in the deep fore-arc crust above the fore-arc mantle. Moreover, as the solubility of silica increases with temperature, fluids generated at depth and rising up the subduction channel should be rich in silica. The authors concluded that there may be a relation between quartz veins formation in the deep fore-arc crust and ETS. However, several constraints as the magnitude and mechanism of the low-frequency earthquakes, and the vertical extent of the tremor should be explained.

+ See papers by Fagereng

Review of past methods.

La Rocca et al. (2009)) stacked seismograms over all stations of the array for each component, and for three arrays in Cascadia. They then computed the cross-correlation between the horizontal and the vertical component, and found a distinct and persistent peak at a positive lag time, corresponding to the time between P-wave and S-wave arrivals. Using a standard layered Earth model, and horizontal slowness estimated from array analysis, they computed the depths of the tremor sources. They located the sources near or at the plate interface, with a much better depth resolution than previous methods based on seismic signal envelopes, source scanning algorithm, or small-aperture arrays. They concluded that at least some of the tremor consisted in the repetition of low-frequency events as was the case in Shikoku. A drawback of the method was that it could be applied only to tremor located beneath an array, and coming from only one place for an extended period of time. Part of the specific question - Just on the plate boundary or overlying layer and how thick the layer is?

If the source is on the plate boundary, you should have a constant time lag between P-wave and S-wave arrivals, and a peak in the cross correlation between the vertical component and the horizontal component.

Explain here why it is going to work (No problem with tremor streaks)

2 Data

The data were collected during the Array of Arrays experiment. Eight small-aperture arrays were installed in the northeastern part of the Olympic Peninsula, Washington. The aperture of the arrays was about 1 km, and station spacing was a few hundred meters. The arrays were around 5 to 10 km apart from each other (Figure 1). Most of the arrays were installed for more than a year, between June 2009 to September 2010, and were able to record the main August 2010 ETS event. Some of the arrays were also recording during the August 2011 ETS event. Ghosh et al. (2012) used a multibeam-backprojection (MBBP) technique to detect and locate tremor. They bandpass filtered the seismic data




Figure 1. Map showing the location of the eight arrays (red dots) used in this study. Blue dots are the locations of the source of the tremor recorded by the arrays. Inset shows the study area with the box marking the area covered in the main map. Contour lines represent a model of the depth of the plate interface (McCrory et al., 2006).

between 5 and 9 Hz. They divided the data into one-minute-long sliding independent (no overlap) time windows. They performed beam forming in the frequency domain at each array to determine the slowness vectors, and backprojected the slownesses in the 3-D space to locate the source of the tremor for each time window. We thus have two catalogs of tremors. The first one is a catalog of 28902 one-minute-long time windows during which tremor was detected between June 20th 2009 and September 30th 2010. For each time window, we have the beginning time, the end time, and the location (latitude and longitude) of the source of the tremor. The second one is a catalog of 5600 one-minute-long time windows between August 10th 2011 and September 6th 2011.

3 Method

We take a 5 km by 5 km grid cell located not too far (less than 25 km) from a given array. We then take all the one-minute-long time windows when tremor was detected and the source of the tremor was located inside this cell. For each one-minute-long time window, we download the seismic data for each seismic station of the array. Then, for each seismic station and each channel, we detrend the data, taper the first and last 5 seconds of the data with a Hann window, remove the instrument response, bandpass filter

between 2 and 8 Hz, and resample the data to 20 Hz. All these preprocessing operations are done with the Python package `obspy`. For each seismic station and each one-minute-long time window, we cross correlate the vertical component with the East-West horizontal component and the North-South horizontal component. Then, we stack the cross correlation functions over all the seismic stations. We experiment with a linear stack, a power stack, and a phase-weighted stack. Figure 2 shows an example of the cross correlation functions as function of time for the Big Skidder array for the 82 one-minute long time windows when tremor was detected in a 5 km by 5 km grid cell centered on the array. We can see that for about half of the tremor window, there is a peak in the cross correlation at about 4.7 s. As the energy of the P-waves is expected to be higher on the vertical component, and the energy of the S-waves to be higher on the horizontal components, we assume that this peak corresponds to the time lag between the arrival a direct P-wave and a direct S-wave. We then stack the cross correlation functions over all the one-minute-long time windows. Again, we experiment with a linear stack, a power stack, and a phase-weighted stack. We get the time of the maximum absolute value of the peak of the stack to be the time lag between direct P-wave and direct S-wave.

We notice that only about half of the cross correlation functions have a distinct peak that coincide with the peak in the stacked cross correlation functions. We want

Grid cells

Cross correlation HV Stack cross correlation over arrays

Stack cross correlation over time windows

peak about 5-6 s

Divide time windows into two clusters based on

New stack with only the good one

One image here to show the stacking

Time lag between direct P and direct S wave

Velocity model - λ tremor depth

Histogram of depths - λ Thickness of the tremor zone

4 Results

use all data

Quality of depth

Some sort of least square regression

One image here to show the depth + the thickness. Map the depth of LFE families too.

5 Discussion and Conclusion

Where are the other tremor? Not well located? Not on the plate boundary?

How analysis addresses the part of the specific question

How analysis addresses the specific question

How analysis addresses the general question

Acknowledgments

Abhijit Ghosh for catalog. NSF grant number. IGERT Big Data if computations ob AWS.

References

- Ghosh, A., Vidale, J., & Creager, K. (2012). Tremor asperities in the transition zone control evolution of slow earthquakes. *Journal of Geophysical Research*, *117*, B10301.
- La Rocca, M., Creager, K., Galluzzo, D., Malone, S., Vidale, J., Sweet, J., & Wech, A. (2009, January). Cascadia tremor located near plate interface constrained by S minus P wave times. *Science*, *323*, 620-623.
- McCorry, P., Blair, J., Oppenheimer, D., & Walter, S. (2006). *Depth to the Juan de Fuca slab beneath the Cascadia subduction margin - a 3-D model sorting earthquakes* (Tech. Rep. No. Series 91). U.S. Geological Survey.