

# 30 Years of Post-disturbance Recruitment in a Neotropical Forest

Ariane MIRABEL<sup>1\*</sup>

Eric MARCON<sup>1</sup>

Bruno HERAULT<sup>2</sup>

## Abstract

The role of tree diversity for tropical forests functioning and services makes it crucial tree diversity and composition fate in the global changing context. Community long-term response to disturbance rely on tree recruitment, long seen as following deterministic successional pathways. These pathways however might be altered in the hyper-diverse tropical forests and of slight but recurrent disturbances induced by global changes. Post-disturbance recruitment trajectories would (i) disentangle the determinants of tree recruitment between stochastic and deterministic processes that enhance a restricted pool of species, and (ii) elucidate tropical forests taxonomic and functional resilience. We examined the trajectories over 30 years of recruited trees taxonomic and functional diversity in 75 ha of forest following a disturbance gradient. We analyzed taxonomic richness, evenness, and turnover, and functional diversity and composition (regarding 7 leaf, stem and life-history functional traits). We highlighted a three-phased successional pathway defined by the interplay of stochastic and deterministic recruitment processes. The succession translated into (i) saplings growth mirroring pre-disturbance communities, (ii) light-demanding species enhanced recruitment entailing, above a disturbance intensity threshold, the dominance of pioneers and (iii) the recovery of pre-disturbance taxonomic and functional characteristics and of stochastic recruitment processes. Although tangible, community taxonomic and functional resilience was decades-long. Post-disturbance recruitment relied on deterministic competition processes for light balancing the stochastic processes ruling undisturbed communities. Although resilient, recruitment taxonomic and functional characteristics remained altered in the long-term, calling caution for forest management.

## Keywords

Disturbance Dynamics, Neotropical Forests, Recruitment, Resilience, Taxonomic and Functional Diversity, Tree Community

<sup>1</sup>UMR EcoFoG, AgroParistech, CNRS, Cirad, INRA, Université des Antilles, Université de Guyane.  
Campus Agronomique, 97310 Kourou, France.

<sup>2</sup>INPHB (Institut National Polytechnique Félix Houphouët Boigny)  
Yamoussoukro, Ivory Coast

\*Corresponding author: ariane.mirabel@ecofog.gf, <https://github.com/ArianeMirabel>

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Material and Methods</b>	<b>2</b>
2.1	Study Site . . . . .	2
2.2	Inventories Protocol and Dataset Collection . . . . .	2
2.3	Recruitment trajectories . . . . .	3
<b>3</b>	<b>Results</b>	<b>3</b>
3.1	Taxonomic richness and evenness and functional diversity	3
3.2	Functional composition . . . . .	3
3.3	Recruitment Turnover . . . . .	4
<b>4</b>	<b>Discussion</b>	<b>4</b>
4.1	A three-phased deterministic successional pathway	4
4.2	The achievement of communities recovery . . . . .	5
<b>5</b>	<b>Conclusion</b>	<b>6</b>
<b>6</b>	<b>Acknowledgement</b>	<b>6</b>
<b>7</b>	<b>Data availability</b>	<b>6</b>

## 1. Introduction

Determining the response of tropical forests to disturbance is key to predict their fate in the global changing context. In the last decades, tropical forests experienced a wide range of disturbance, from radical land-use changes for agriculture or mining (??) to more insidious changes following climatic changes or anthropogenic activities like selective logging (??). In that respect a vast literature successfully modeled community response to disturbance in terms of tree growth, tree height and fluxes of carbon, water and nutrients (????). Regarding tree community diversity and composition, the ecological theory of succession assumes predictable post-disturbance trajectories shaped by deterministic recruitment processes (??). However, the specific processes underlying these trajectories remain to be disentangled, which is essential to highlight community structuring mechanisms and improve exploitation and conservation guidelines (??).

In forest ecosystems, the successional framework translates first into the recruitment of saplings benefiting from the available resources and the low competition, then into the progressive exclusion of low-competitive species with the stand maturation and the increasing competition for re-

sources, and eventually into the senescence of early-successional and the emergence of late-successional species restoring community pre-disturbance composition and diversity (?). This deterministic succession involves a gradual recovery of pre-disturbance state through various recruitment processes depending on species resource use and competitive abilities (???). Observations, however, show that post-disturbance trajectories can result from the interplay between deterministic and stochastic processes. Stochastic processes would imply a random recruitment of species, depending on dispersal limitations, and no resilience of the pre-disturbance state (???). The balance between deterministic and stochastic processes remain specifically questioned in tropical forests, where the diversity is huge and the long-term monitoring are scarce (?). It is essential to understand the balance between deterministic and stochastic processes and its changes along time after disturbance, to understand and anticipate post-disturbance diversity patterns (?).

The underlying processes of post-disturbance trajectories affect both community taxonomic characteristics, that refer to neutral species assemblages, and functional characteristics, that account for species ecology and functioning (???). The different ecological processes can be assessed through the taxonomic and functional diversity and composition of recruited communities following disturbance (???). A deterministic succession thus imply predictable recruitment trajectories comprising a succession of recruitment processes based on species competitive ability (?). In tropical forests, light is the limiting resource: post-disturbance trajectories would be shaped by a shift from slow-growing, long-lived species with “conservative” resource use, to fast-growing species with “acquisitive” resource use (???). Key leaf, wood and life-history functional traits assessing species resources acquisition strategy and ecology would then grasp the competition processes at stake (???). The balance between deterministic and stochastic processes would translate in a random divergence of both community trajectories, contrasting with a deterministic convergence towards the recovery pre-disturbance state (?). The interplay between deterministic and stochastic processes might then translate into a divergence in the taxonomic space, with communities displaying different taxonomic composition and diversity, and a converge in the functional space (?).

In this paper we followed recruitment trajectories over 30 years of 75 ha of Neotropical forest plots set up on a gradient of disturbance intensity, from 10 to 60% of forest biomass removed. We examined the recruited trees (i) taxonomic composition, richness and evenness, (ii) taxonomic turnover compared to pre-disturbance community, and (iii) functional composition and diversity based on seven major leaf, stem and life-history traits. We compared the recruitment trajectories to neutral models corresponding to a stochastic recruitment and a randomization of species functional traits. Specifically, we (i) elucidated the successional pathway shaping community response to disturbance and the underlying ecological processes and (ii) clarified the extent of community taxonomic and functional resilience, in the sense of pre-disturbance characteristics recovery, and its consequences for tropical forest management.

## 2. Material and Methods

### 2.1 Study Site

The Paracou station is located in a lowland tropical rain-forest in French Guiana (518°N and 5253°W). Climate is tropical wet with mean annual precipitation averaging 2980 mm.yr<sup>-1</sup> (30-yr period) and a 3-months dry season (< 100 mm.mo<sup>-1</sup>) from mid-August to mid-November, and a one-month dry season in March (?). Elevation ranges from 5 to 50 m and mean annual temperature is 26°C. Soils are thin acrisols over a layer of transformed saprolite with low permeability generating lateral drainage during heavy rains. The experiment is a network of twelve 6.25 ha plots (Table 1) that underwent three disturbance treatments in 1987 according to a randomized plot design with three replicate blocks of four plots (?).

### 2.2 Inventories Protocol and Dataset Collection

Dominant families in the study site are Fabaceae, Chrysobalanaceae, Lecythidaceae and Sapotaceae. All trees above 10 cm DBH were mapped and measured annually since 1984. Trees are first identified with a vernacular name assigned by the forest worker team, and afterward with a scientific name assigned by botanists during regular botanical campaigns. Botanical campaigns have been carried out every five to six years from 2003 onwards but identification levels varied between campaigns.

These variability of protocols in time raised methodological issues as vernacular names usually correspond to different botanical species. It resulted in significant taxonomic uncertainties that had to be propagated to composition and diversity metrics. The uncertainty propagation was done through a Bayesian framework reconstituting complete inventories at genus level from real incomplete ones on the basis of vernacular/botanical names association. Vernacular names were replaced through multinomial trials based on the association probability  $[\alpha_1, \alpha_2, \dots, \alpha_N]$  observed across all inventories between each vernacular name  $v$  and the species  $[s_1, s_2, \dots, s_N]$ :

$$M_v([s_1, s_2, \dots, s_N], [\alpha_1, \alpha_2, \dots, \alpha_N])$$

See appendix 1 and ? for the detailed methodology.

To minimize the remaining identification uncertainties, the simulated botanical inventories were reported at genus level.

Six functional traits representing the leaf economics (leaves thickness, toughness, total chlorophyll content and specific leaf area) and stem economics spectra (wood specific gravity and bark thickness), and life-history traits (maximum specific height and seed mass) were considered. Traits were extracted from the BRIDGE project (<http://www.ecofog.gf/Bridge/>) where trait values were assessed from a selection of individuals located in nine permanent plots in French Guiana, including two in Paracou, and comprised 294 species pertaining to 157 genera. Missing trait values (10%) were filled by multivariate imputation by chained equation (?). Imputations were restricted within genus or family when samples were too scarce, in order to

**Table 1.** Intervention table, summary of the disturbance intensity for the 4 plot treatments in Paracou.

Treatment	Timber	Thinning	Fuelwood	%AGB lost
Control	-	-	-	0
T1	DBH $\geq$ 50 cm, commercial species, $\approx 10 \text{ trees.ha}^{-1}$	-	-	[12 – 33]
T2	DBH $\geq$ 50 cm, commercial species, $\approx 10 \text{ trees.ha}^{-1}$	DBH $\geq$ 40 cm, non-valuable species, $\approx 30 \text{ trees.ha}^{-1}$	-	[33 – 56]
T3	DBH $\geq$ 50 cm, commercial species, $\approx 10 \text{ trees.ha}^{-1}$	DBH $\geq$ 50 cm, non-valuable species, $\approx 15 \text{ trees.ha}^{-1}$	40 cm $\leq$ DBH $\leq$ 50 cm, non-valuable species, $\approx 15$ $\text{trees.ha}^{-1}$	[35 – 56]

account for the phylogenetic signal.

As seed mass information was classified into classes, no data filling process was applied and analyses were restricted to the 414 botanical species recorded.

All composition and diversity metrics were obtained after 50 iterations of the uncertainty propagation framework.

### 2.3 Recruitment trajectories

Communities were split into surviving trees of pre-disturbance communities and trees recruited afterward in 2-years intervals.

Taxonomic diversity trajectories were assessed through species richness and evenness (the Hill number translation of the Simpson index) (??).

The two diversities belong to the set of HCDT or generalized entropy, respectively corresponding to the zero and two order of diversity ( $q$ ), which grasps the balance between richness and evenness in the community through the value of  $q$  that emphasizes common species.

Functional diversity trajectories were assessed through the Rao index of quadratic entropy, which combines species abundance distribution and average pairwise dissimilarity based on all functional traits. Functional composition trajectories were assessed through the functional traits community weighted means (CWM), representing the average trait value in a community weighted by species relative abundance (?). Seed mass trajectories were reported by the proportion of each class recorded in the inventories.

The taxonomic similarity between recruited trees and pre-disturbance forest was measured with the turnover metrics detailed in ?.

The taxonomic and functional recruitment trajectories were compared to null trajectories obtained after 50 iterations of the null models. The taxonomic null model was a random sampling of recruited trees within the living communities, with the maintenance of species abundance and tree density. The functional null model was a reassignment of species trait values that randomized traits abundances but maintained communities abundance distribution (?). The null trajectories were similarly obtained after 50 iterations of the random sampling.

## 3. Results

### 3.1 Taxonomic richness and evenness and functional diversity


In undisturbed communities the recruitment taxonomic richness and evenness remained stable over the 30 years and with values equivalent to those of the taxonomic null model (Figure (1)).

In disturbed communities the taxonomic richness followed hump-shaped trajectories first increasing until a maximum reached after around 15 years and positively correlated to the disturbance intensity ( $\rho_{\text{Spearman}}^{\text{Richness}} = 0.93$ ). Afterward the taxonomic richness decreased and recovered the pre-disturbance values after 30 years. The observed taxonomic richness was increasingly lower than this of null model for 15 years, then the difference started to shrink but the observed richness remained negative remained negative until after 30 years. The taxonomic evenness decreased independently of the disturbance intensity over the 30 years ( $\rho_{\text{Spearman}}^{\text{Simpson}} = -0.35$ ). The observed taxonomic evenness was increasingly lower than this of the null model until 15 years after disturbance, when the difference stabilized.

The functional diversity in the undisturbed plots remained stable and equivalent to this of the functional null model over the 30 years. In the lowest disturbance plots the functional diversity remained stable or slightly increasing, and was higher than this of the null model for two of the T1 plots. In the disturbed plots of higher disturbance intensity (T2 and T3) the functional diversity decreased until 15 years after disturbance, when it started to recover towards initial values. The observed functional diversity remained lower than this of the null model over the 30 years.

### 3.2 Functional composition

In undisturbed plots functional traits values remained stable over the 30 years while it followed hump-shaped trajectories in all disturbed plots, to the exception of the leaf chlorophyll content. Trajectories of SLA and bark thickness first increased before decreasing towards initial values. Conversely, trajectories of leaf thickness, leaf toughness, wood specific gravity, and maximum height first decreased



**Figure 1.** Upper panels, trajectories over 30 years of taxonomic richness (a), taxonomic evenness (b) and functional diversity (c) of observed 2-years laps recruitment  $H_{obs}^q$ . Lower panels, diversity differences to null models  $H_{obs}^q - H_{null}^q$

and then started returning towards initial values but their recovery remained unachieved after 30 years (Figure 2).

### 3.3 Recruitment Turnover

Over the 30 years in control plots the turnover of recruited species compared to initial community remained low (Figure 3). In disturbed plots the recruited species turnover followed a marked hump-shaped trajectory, with a maximum reached around 15 years after disturbance. The maximum turnover was positively correlated to the disturbance intensity ( $\rho_{spearman} = 0.93$ ). Thirty years after disturbance the turnover had returned to low values.

## 4. Discussion

### 4.1 A three-phased deterministic successional pathway

Post-disturbance recruitment trajectories relied on a three-phased successional pathway defined by the emergence of deterministic competition processes for light gradually balancing the stochastic recruitment specific to undisturbed communities.


A first phase (0-8 years), corresponded to the recruitment of pre-disturbance surviving saplings (DBH < 10 cm) that immediately benefited from the increased enlightenment and alleviated competition induced by disturbance (?). The taxonomic and functional characteristics of recruited trees mirrored the pre-disturbance communities and recruitment processes matched the null stochastic recruitment model.

A second phase (8-15 years) was marked by a shift in community functional composition towards more “acquisitive” functional strategies and the dominance of a restricted


set of species. The recruitment then involved true recruits, *i.e.* trees germinating from the seeds bank, representing the main part of the whole post-disturbance recruitment (?). The recruitment was dominated by short-lived, fast growing hard pioneers displaying efficient light acquisition (???). As already demonstrated in temperate forests, the pool of recruited species was restricted by trait-based deterministic processes favoring species with efficient light acquisition (high SLA and leaf chlorophyll content) and inexpensive, short-lived tissues (low leaf thickness and toughness, small Hmax and low wood specific gravity and bark thickness)(?). This emergence of trait-based deterministic processes balanced the stochastic recruitment observed in the first place, and the relative importance of both processes was determined by the disturbance intensity. After low intensity disturbance (T1 plots) recruited species still mirrored pre-disturbance taxonomic composition, but included more long-lived pioneers and light-demanding species (?). For intense disturbance in contrast (T2 and T3 plots), the composition of recruited trees rapidly differed from pre-disturbance community and with the high dominance of hard pioneers, such as *Cecropia spp.* or *Vismia spp.*, likely entailing significant changes in communities functioning (?).

A third recruitment phase (15-30 years) corresponded to the recovery of pre-disturbance taxonomic and functional characteristics. Although the recruits remained mainly light-demanding species their functional diversity increased and they increasingly resembled the pre-disturbance taxonomic composition. The deterministic recruitment processes then gradually left room to stochastic recruitment processes spe-





**Figure 2.** Community weighted means (CWM) of the leaf, the two stem and specific maximum height. Shaded areas are the credibility intervals.



**Figure 3.** Trajectories over 30 years of the abundance-based turnover between 2-years laps recruited trees pre-disturbance communities.

cific to undisturbed forest (?).

#### 4.2 The achievement of communities recovery

After disturbance the stochastic recruitment specific to undisturbed communities was progressively restored and drove community taxonomic and functional recovery. This confirmed previous results from the Paracou experiment, conducted 10 years (?) and 20 years (?) after disturbance, where the early signs of the resilience of pre-disturbance taxonomic and functional composition recovery had been detected.

Recruitment taxonomic richness and evenness recovered pre-disturbance values and the taxonomic composition converged towards the pre-disturbance community, thus maintaining the initial differences among communities for all disturbance intensity. Community taxonomic convergence to the local pre-disturbance recruitment composition revealed the scarce recruitment of species that did not belong to pre-disturbance community, due to the commonness of dispersal limitation among tropical tree species (?).

Functional composition and diversity trajectories converged similarly in the functional space towards the recovery of pre-disturbance values, suggesting a common and resilient functioning despite communities' taxonomic divergence (?).

Trait-based enhancement processes made deterministic the community functional response to disturbance but dispersal limitation and steady-state stochastic recruitment made community taxonomic response historically contingent. Although resilient, the functional and taxonomic composition of recruited trees remained altered 30 years after disturbance by the dominance of light-demanding species.

This long-term impact specifically raises questions for the management of exploited forests, as most valuable species are late-successional and would thus require cutting cycles of more than 30 years (?).

## 5. Conclusion

The post-disturbance recruitment trajectories highlighted a three-phased deterministic successional pathway shaped by the emergence of niche processes enhancing light-acquisitive species and balancing the stochastic recruitment of undisturbed communities. The successional pathway first corresponded to the enhanced growth of pre-disturbance surviving saplings mirroring the taxonomic and functional characteristics of pre-disturbance communities. Second, recruitment trajectories were shaped by true recruits from the seeds bank selected through the emergence of competitive exclusion for light fostering pioneer species. Above a disturbance intensity threshold the second recruitment phase was dominated by short-lived hard pioneers that drastically changed community composition, diversity and likely functioning. A third phase eventually corresponded to the return towards pre-disturbance recruitment composition and taxonomic and functional diversity, through the recovery of stochastic recruitment processes specific to undisturbed communities. Besides, repeated disturbance might have increasingly strong impacts, as community recovery involved the seeds bank and probably altered the composition and diversity of the seeds stock (?).

## 6. Acknowledgement

We are in debt with all technicians and colleagues who helped setting up the plots and collecting data over years. Without their precious work, this study would have not been possible and they may be warmly thanked here.

## 7. Data availability

This article is based upon the dataset of the Paracou station, which is part of the Guyafor permanent plot network in French Guiana (Cirad-CNRS-ONF). The dataset is available upon request to the scientific director (<https://paracou.cirad.fr>).

---