

Appendix I

Ariane MIRABEL^{1*}

Eric MARCON¹

Bruno HERAULT²

Abstract

¹UMR EcoFoG, AgroParistech, CNRS, Cirad, INRA, Université des Antilles, Université de Guyane.
Campus Agronomique, 97310 Kourou, France.

²INPHB (Institut National Polytechnique Félix Houphouët Boigny)
Yamoussoukro, Ivory Coast

*: ariane.mirabel@ecofog.gf, <http://www.ecofog.gf/spip.php?article47>

Contents

1	Appendix figures	1
2	Figure A1	1
3	Figure A2	1
4	Figure A3	1

2. Figure A1

3. Figure A2

4. Figure A3

1. Appendix figures

The following information are available for this article:

Figure A1: Whenever the family or genus were recorded for a vernacular name, the pool of species sampled to replace the vernacular name was restricted to those pertaining to the same higher taxonomic level (i.e species of the same genus when the species is missing, species of the same family when the genus is missing) 1.

Figure A2: Trajectories of seed mass classes proportions over 30 years after disturbance, corresponding to the median (solid line) and 0.025 and 0.975 percentile (gray envelope) observed after 50 iteration of the taxonomic uncertainty propagation. No gap filling process was applied in this case. Initial treatments are represented by solid lines colors with green for control, blue for T1, orange for T2 and red for T3 2.

Figure A3: Trajectories of the Rao functional diversity over 30 years after disturbance. Trajectories correspond to the median (solid line) and 0.025 and 0.975 percentile (gray envelope) observed after 50 iteration of the taxonomic uncertainty propagation. Initial treatments are represented by solid lines colors with green for control, blue for T1, orange for T2 and red for T3. and the missing trait value filling processes. 3.




Figure 1. Scheme of the Bayesian process propagating the taxonomic uncertainty




Figure 2. Trajectories of seed mass classes proportions over 30 years after disturbance, corresponding to the median (solid line) and 0.025 and 0.975 percentile (gray envelope) observed after 50 iteration of the taxonomic uncertainty propagation. No gap filling process was applied in this case. Initial treatments are represented by solid lines colors with green for control, blue for T1, orange for T2 and red for T3.




Figure 3. Trajectories of the Rao functional diversity over 30 years after disturbance. Trajectories correspond to the median (solid line) and 0.025 and 0.975 percentile (gray envelope) observed after 50 iteration of the taxonomic uncertainty propagation. Initial treatments are represented by solid lines colors with green for control, blue for T1, orange for T2 and red for T3, and the missing trait value filling processes.