

1 Post-Disturbance Tree Community Trajectories in a 2 Neotropical Forest

3 Ariane MIRABEL^{1*}

4 Bruno Herault²

5 Eric Marcon¹

6 Abstract

7 In the current global change context, it is urgent to anticipate the fate of tropical forests. This means understanding tree
8 community response to disturbance and the underlying processes. In that respect, we aim here to clarify taxonomic and
9 functional post-disturbance trajectories, and determine the scope of the Intermediate Disturbance Hypothesis (IDH) that
10 remains debated in tropical forests. We analyzed community trajectories following a disturbance gradient in a Neotropical
11 forest over 30 years. We considered trajectories along time of community taxonomic and functional trajectories in
12 terms of richness, evenness, composition, and redundancy. We based on the annual botanical inventories of 75 ha
13 of a Neotropical forest and on large trait datasets comprising seven leaf, stem, and life-history traits. We identified a
14 decoupling between taxonomic composition, differing among communities, and functional composition, remaining similar
25 and convergent. The taxonomic diversity followed humped-shaped trajectories depending on the disturbance intensity,
16 which validated the IDH (Intermediate Disturbance Hypothesis). The functional diversity trajectories, however, were
17 homogeneous among all plots and dismissed the IDH. We explained this decoupling by the variations in community
18 functional redundancy that mitigated the functional impact of disturbance. Although consistent, the recovery of community
19 composition, diversity, and redundancy remained unachieved after 30 years. These results acknowledged the need of
20 decades-long cycles without disturbance to ensure community complete recovery, and questioned community resilience
21 after repeated disturbances.

22 Keywords

23 Community Ecology, Disturbance Trajectories, Intermediate Disturbance Hypothesis, Mid-term Resilience, Neotropical
24 Forests, Taxonomic and Functional Biodiversity

26 ¹UMR EcoFoG, AgroParistech, CNRS, Cirad, INRA, Université des Antilles, Université de Guyane.

27 Campus Agronomique, 97310 Kourou, France.

28 ²INPHB, Institut National Polytechnique Félix Houphouët-Boigny

29 Yamoussoukro, Ivory Coast.

30 *Corresponding author: ariane.mirabel@gmail.com, <https://github.com/ArianeMirabel>

32	Contents	
33	1 Introduction	2
34	2 Material and Methods	4
35	2.1 Study site	4
36	2.2 Inventories protocol and dataset collection	5
37	2.3 Composition and diversity metrics	7
38	3 Results	8
39	3.1 Community Composition	8
40	3.2 Community taxonomic and functional diversity	11
41	3.3 Functional redundancy	12
42	4 Discussion	12
43	4.1 Decoupled taxonomic and functional trajectories	14
44	4.2 The scope of the intermediate disturbance hypothesis	15
45	4.3 The functional redundancy, key of community resilience	15
46	5 Conclusion	16
47	6 Acknowledgement	16
48	7 Author's contributions	17
49	8 Data availability	17

50 1. Introduction

51 The large areas covered with tropical forests worldwide hold crucial environmental, economic, and social values. They
52 provide wood and multiple non-timber forest products, shelter a diversified fauna, and ensure cultural and human well-
53 being. They regulate local and regional climates, as well as carbon, water and nutrient cycles. However, the growing
54 demand in forests products together with current global changes increase the pressure on remaining undisturbed forests
55 (Morales-Hidalgo *et al.*, 2015). These threats affect the natural disturbance regime that defines and maintains the structure,
56 composition, and functioning of tree communities (Schnitzer & Carson, 2001; Anderson-Teixeira *et al.*, 2013; Sist *et al.*,
57 2015). To anticipate the fate of tropical forests, it is urgent to understand tree community response to disturbance, and
58 the underlying ecological processes. The forest cover is generally maintained following disturbance, but modifications
59 in the fluxes of light, heat, and water (Goulamoussène *et al.*, 2017) change community abiotic and biotic environments.

These changes translate into post-disturbance community trajectories that have been largely studied through trajectories of forest structural parameters such as aboveground biomass, tree height or stem density (Piponiot *et al.*, 2016; Rutishauser *et al.*, 2016). Some of the determinants of post-disturbance biomass trajectories are already identified, like the structure and composition of the pre-disturbance community, or the post-disturbance environmental parameters (Héroult & Piponiot, 2018). Community diversity and composition trajectories, however, have not been as thoroughly understood (Guitet *et al.*, 2018; Molino & Sabatier, 2001), and manifold biodiversity trajectories might emerge given the variety of species response to disturbance and the diversity of tropical forests (Lindenmayer *et al.*, 2012; Garcia Florez *et al.*, 2017).

An early conceptual basis of the linkage between biodiversity and disturbance is the Intermediate Disturbance Hypothesis (IDH). The IDH states a relationship between the community diversity and the intensity and frequency of disturbance events, and postulates a diversity peak at intermediate level of disturbance (Connell, 1978). This is based on the fluctuations of community environment following disturbance that foster both competitively superior species and fast colonizers, and prevents competitive exclusion (Shea *et al.*, 2004; Pulsford *et al.*, 2016). In tropical forests, however, observations of the IDH often diverge from theoretical expectations (Randall Hughes *et al.*, 2007; Sheil & Burslem, 2003; Norden *et al.*, 2017), and the underlying processes might be complicated by the diversity of tropical tree communities (Lindenmayer *et al.*, 2012; Garcia Florez *et al.*, 2017). In this context, the IDH is controversial in tropical forests and remains to be tested (Hubbell, 2001; Fox, 2013; Sheil & Burslem, 2013).

Analysing community response to disturbance and grasp all aspects of community changes requires a wide array of metrics (Sheil & Burslem, 2003; Shea *et al.*, 2004; Mayfield & Levine, 2010). The analysis should first consider community composition, which is crucial for conservation issues and reveals the pool of species fostered or hampered by disturbance (Lavorel & Garnier, 2002; Bellwood *et al.*, 2006). This should be completed with diversity metrics encompassing both community richness and evenness to assess the changes in community abundance distribution. Besides, functional approaches have been shown to usefully complement pure taxonomic approaches as they shed light on the species biological attributes directly linking community diversity, composition, and redundancy to ecosystem functioning (Violle *et al.*, 2007; Baraloto *et al.*, 2012). In that respect, a vast literature allowed recognizing major traits representing species ecological strategy and determining how they respond to changing conditions (Díaz *et al.*, 2005). Specifically, in tropical forests, the functional approach revealed the emergence of deterministic processes following disturbance. Such deterministic processes entailed a shift from a dominance of “conservative” slow-growing species dealing with scarce resources, to a dominance of “acquisitive” fast-growing species with rapid and efficient use of abundant resources (Rees *et al.*, 2001; Reich, 2014; Héroult *et al.*, 2011). This shift is translated into the trajectories of average community value of key functional traits related to resource acquisition, as leaf and stem traits, and life-history strategy, as seed mass and

90 maximum size (Wright *et al.*, 2004; ter Steege *et al.*, 2006; Westoby & Wright, 2006; Chave *et al.*, 2009).

91 The functional approach also encompasses the analysis of functional redundancy, that quantifies the amount of shared
92 trait values among species (Carmona *et al.*, 2016). The typical high functional redundancy of hyper-diverse tropical forests
93 (Bellwood *et al.*, 2006) mitigates the impacts of species removal on ecosystem functioning, and determines community
94 resilience after disturbance (Elmqvist *et al.*, 2003; Díaz *et al.*, 2005).

95 Here, we monitored over 30 years the response of 75 ha of Neotropical forest plots set up on a gradient of disturbance
96 intensity, from 10 to 60% of above-ground biomass (AGB) loss. We made use of a large functional traits database
97 encompassing major leaf, stem, and life-history traits in order to draw the taxonomic and functional trajectories in terms of
98 richness, evenness, composition, and redundancy. Specifically, (i) we drew taxonomic and functional post-disturbance
99 trajectories and examined the underlying ecological processes, (ii) we discussed the scope of the IDH regarding taxonomic
100 and functional facets of community diversity, and (iii) we analyzed community resilience and time to recovery.

101 2. Material and Methods

102 2.1 Study site

103 Paracou station in French Guiana (5° 18'N and 52° 53'W) is located in a lowland tropical rain forest in a tropical wet
104 climate with mean annual temperature of 26° C, and mean annual precipitation averaging 2980 mm.y⁻¹ (30-y period). The
105 climate comprises a 3-month dry season (< 100 mm.month⁻¹) from mid-August to mid-November, and a one-month dry
106 season in March (Wagner *et al.*, 2011). Across all plots, elevation ranges from 5 to 50 m, and the topography mainly
107 corresponds to hilltops or hillsides, while bottomlands cover less than 1 % of the area. Plots are shallow ferrallitic Acrisols
108 over a layer of transformed saprolite with low permeability and lateral drainage. Soil conditions are homogeneous, to the
109 exception of the highest hilltops where the thick surface allows a free vertical drainage (Gourlet-Fleury *et al.*, 2004).

110 The experiment is a network of twelve 6.25 ha plots (Table 1) that underwent three disturbance treatments in 1987
111 according to a randomized plot design (Gourlet-Fleury *et al.*, 2004).

112 The experiment comprised three replicates of three silvicultural treatments (hereafter plots T1, T2, and T3), and three
113 control plots (T0). All treatments T1, T2, and T3 comprised the logging of 10 trees/ha with 50 cm minimum DBH that
114 belonged to a set of 58 commercially exploited species (Gourlet-Fleury *et al.*, 2004). Treatment T2 additionally comprised
115 a thinning treatment by poison-girdling of non-commercially exploited species randomly selected with an average of 30
116 trees/ha with 40 cm minimum DBH. Treatment T3 additionally comprised the logging of 15 trees/ha with 40 cm minimum
117 DBH, and the poison-girdling of 20 trees/ha with a 50 cm minimum DBH, all belonging to non-commercially exploited
118 species. Considering the silvicultural treatments and the following damage, disturbance intensity was measured as the

percentage of aboveground biomass (%AGB) lost between the first inventory in 1984 and five years after disturbance (Piponiot *et al.*, 2016) estimated with the BIOMASS R package (Réjou-Méchain *et al.*, 2017). The three treatments were then transformed into a continuous disturbance intensity gradient with increasing above-ground biomass (AGB) loss.

2.2 Inventories protocol and dataset collection

The study site corresponds to a tropical rainforest typical of the Guiana Shield with a dominance of *Fabaceae*, *Chrysobalanaceae*, *Lecythidaceae*, and *Sapotaceae*. In the 12 experimental plots, all trees above 10 cm DBH have been mapped and measured annually since 1984. Trees are first identified with a vernacular name assigned by the forest worker team, and afterward with a scientific name assigned by botanists during regular botanical campaigns. In 1984, specific vernacular names were given to 62 common or commercially exploited species. More infrequent species were identified under general identifiers only distinguishing trees and palms. From 2003, botanical campaigns have been conducted every 5 to 6 years to identify all trees at the species level. In 2015, however, identification levels still varied among plots and campaigns.

This variability of protocols in time raised methodological issues as vernacular names usually correspond to various botanical species. This resulted in significant taxonomic uncertainty that had to be accounted for in the measure of composition and diversity metrics. Uncertainty propagation was implemented withing a Bayesian framework using vernacular/botanical names associations to reconstitute complete inventories at genus level from real incomplete ones. Vernacular names were replaced through multinomial trials based on the association probability $[\alpha_1, \alpha_2, \dots, \alpha_V]$ observed across all inventories between each vernacular name v and all species $[s_1, s_2, \dots, s_N]$:

$$M_v\left([s_1, s_2, \dots, s_N], [\alpha_1, \alpha_2, \dots, \alpha_V]\right)$$

See Supplementary Materials -Fig. A.1 and Aubry-Kientz *et al.* (2013) for the detailed methodology.

Six functional traits representing the leaf economics (*i.e.* leaf thickness, toughness, total chlorophyll content, and specific leaf area), and the stem economics (*i.e.* wood specific gravity and bark thickness) were obtained from the BRIDGE project ¹. Trait values were assessed from a selection of individuals located in nine permanent plots in French Guiana, including two in Paracou, and comprised 294 species belonging to 157 genera. Whenever a species was in the dataset but missed some trait values (10% of the species), missing values were filled using multivariate imputation by chained equation (van Buuren & Groothuis-Oudshoorn, 2011). To account for the phylogenetic signal in the filling process, imputations were based on samples of species from the same genus or from the same family. Whenever a species was not in the dataset, it was attributed a set of trait values randomly sampled among species of the next higher taxonomic level (same genus

¹<http://www.ecofog.gf/Bridge/>

Table 1. Intervention table, summary of the disturbance intensity for the 4 plot treatments in Paracou. Treatment intensities are defined by the minimum logging DBH (Diameter at Breast Height), the type of logged species (commercially exploited or not), the density of logged trees, and the total AGB (Above Ground Biomass) lost after treatment.

Treatment	Timber	Thinning	Fuelwood	%AGB lost
Control	-	-	-	0
T1, low	DBH ≥ 50 cm, commercially exploited species, ≈ 10 <i>trees.ha⁻¹</i>	-	-	[12 – 33]
T2, intermediate	DBH ≥ 50 cm, commercially exploited species, \approx 10 trees.ha^{-1}	DBH ≥ 40 cm, non-commercially exploited species, \approx 30 trees.ha^{-1}	-	[33 – 56]
T3, high	DBH ≥ 50 cm, commercially exploited species, \approx 10 trees.ha^{-1}	DBH ≥ 50 cm, non-commercially exploited species, \approx 15 trees.ha^{-1}	$40 \text{ cm} \leq \text{DBH} \leq 50$ cm, non-commercially exploited species, \approx 15 trees.ha^{-1}	[35 – 56]

or family). Two life-history traits, maximum specific height and seed mass, came from the Mariwenn database ². The database compiles information from a vast literature on the flora of French Guiana (Ollivier *et al.*, 2007) and comprises 362 species belonging to 188 genera. As seed mass information was classified into classes, no data filling process was applied and analyses were restricted to the botanical species recorded.

All composition and diversity metrics were obtained after 50 iterations of the uncertainty propagation framework.

2.3 Composition and diversity metrics

Because of the variable precision of botanical identification efforts, we had to conduct the taxonomic composition and diversity analysis at the genus level. Taxonomic and functional trajectories of community composition were drawn in a two-dimensional NMDS ordination plane. Two NMDS using abundance-based (*i.e.* Bray-Curtis) dissimilarity measures were conducted to map either taxonomic or functional composition, the latter based on the seven leaf, stem, and life history traits (without seed mass classes). Trajectories along time were reported through the Euclidean distance between the target inventories and the 1984 pre-disturbance inventories of reference. Univariate trajectories of the leaf, stem, and life-history traits were also visualized with the community weighted means (CWM) (Díaz *et al.*, 2007). Species seed mass were given in 5 mass classes, and seed mass trajectories were reported as the proportion of each class in the inventories (Supplementary materials).

The taxonomic and functional trajectories were analysed from the 1984 pre-disturbance inventories of reference. The taxonomic diversity was reported through species richness and the Hill number translation of the Simpson index (Hill, 1973). The comparison between these two metrics assesses community taxonomic richness and evenness: thereafter, results will be discussed directly in terms of taxonomic richness and evenness. Both indices are recommended for diversity studies (Marcon & Hérault, 2015), and belong to the set of HCDT or generalized entropy corresponding, respectively, to the 0 and 2 order of diversity (q). The functional diversity was reported using the functional richness and functional evenness, through the Rao index of quadratic entropy. The Rao index combines species abundance distribution, and the average pairwise functional dissimilarity between species computed by the Gower distance.

The impacts of the initial disturbance levels on the maximum gain or loss in richness and evenness were tested with the Spearman rank correlation tests. Richness and evenness trajectories were besides analyzed through polynomial regression between (i) taxonomic and functional richness and evenness, and (ii) the initial %AGB loss at 10, 20, and 30 years after disturbance.

Finally, functional redundancy was measured as the overlap among species in community functional space (Carmona *et al.*, 2016). First, the individuals of the trait database were mapped in the plane of the first two axes from a PCA analysis.

²<https://www.ecofog.gf/mariwenn/>

The PCA analysis lowered the weight of correlations among traits as the axes are combinations of the most decoupled traits. For each species, the traits probability density (TPD) were computed from the mapping of individuals through two-dimension kernel density estimators. Second, for each community, the TPD weighted by species abundance were summed across the functional space. Third, the functional space was divided into a 100 x 100 grid, and the number of species with a positive TPD was counted in each cell. The average count across cells minus 1 returned the Community Functional Redundancy, which was the average number of species in the community that share the same trait values.

3. Results

3.1 Community Composition

From 1984, the first pre-disturbance inventory, to 2015, 28 years after disturbance, 828 388 individual trees and 591 botanical species spanning 223 genera and 64 families were recorded.

In undisturbed plots, both taxonomic and functional composition remained stable (Fig. 1). In disturbed plots, both trajectories followed marked and consistent trajectories over time.

The functional composition trajectory resembled, in disturbed communities, to cyclic compositional changes with an incomplete recovery of the initial composition (Fig.1). The maximum dissimilarity with the initial state was positively correlated with the disturbance intensity for both taxonomic and functional composition ($\rho_{Spearman}^{Taxonomic} = 0.87$ and $\rho_{Spearman}^{Functional} = 0.90$, respectively). The maximum dissimilarity with the initial state was reached for taxonomic composition between 15 to 25 years, and between 20 to 25 years for functional composition.

In control plots, Community Weighted Means (CWM) of functional traits remained stable in time. In disturbed plots, they mostly followed unimodal trajectories, either stabilizing or returning towards their initial values, to the exception of leaf chlorophyll content, which continued to increase 30 years after disturbance for 4 out of 6 highly disturbed plots. Maximum height at adult stage (H_{max}), leaf toughness, and wood specific gravity (WSG) decreased in time and then slightly increased, but remained significantly lower than their initial value (Fig. 2). Bark thickness and specific leaf area (SLA) both increased along time. Bark thickness remained substantially high after 30 years, and SLA had almost recovered to its initial value. Whatever the functional traits, the maximum difference to initial value was highly correlated to the disturbance intensity. Positive correlations were observed for leaf thickness, chlorophyll content, SLA and bark thickness ($\rho_{Spearman}^{Leafthickness} = 0.76$, $\rho_{Spearman}^{Chlorophyllcontent} = 0.60$, $\rho_{Spearman}^{SLA} = 0.93$, $\rho_{Spearman}^{Barkthickness} = 0.71$). Negative correlation was observed for leaf toughness, WSG, and Hmax ($\rho_{Spearman}^{Leaftoughness} = -0.53$, $\rho_{Spearman}^{WSG} = -0.75$, $\rho_{Spearman}^{Hmax} = -0.40$) The proportions of the three lightest seed mass classes increased in all disturbed plots. After 30 years the proportion of lightest seed mass class decreased while it stabilized for the two other lightest seed mass classes (Supp. Mat. - Fig. A.2).




Figure 1. Plot trajectories in terms of taxonomic composition ((a) and (c)), and functional composition ((b) and (d)) in a two-dimensional NMDS plane. Lower panels ((c) and (d)) represent the Euclidean distance to initial condition along the 30 sampled years. Shaded areas are the credibility intervals.

Community Weighted Means




Figure 2. Trajectories of community weighted means over 30 years after disturbance of four leaf traits (leaf thickness, chlorophyll content, toughness, and specific area), two stem traits (wood specific gravity and bark thickness), and one life history trait (species maximum height at adult stage).




Figure 3. Trajectories of community taxonomic richness (a), Simpson diversity (b), functional richness (c), and Rao diversity (d). Values correspond to the difference over 30 years of community diversity with the values of 1984 pre-disturbance inventories of reference. Shaded areas are the credibility intervals

3.2 Community taxonomic and functional diversity

Taxonomic richness and evenness remained stable in control plots over the 30 years of monitoring. In disturbed communities, after low disturbance intensity the taxonomic richness increased, reaching a maximum gain of 14 botanical genera. After intense disturbance the taxonomic richness followed a more complex trajectory, decreasing for ten years after disturbance before recovering to pre-disturbance values. The maximum richness loss or gain after disturbance was positively correlated with the disturbance intensity ($\rho_{Spearman}^{Richness} = 0.50$).

In all disturbed plots the evenness first increased until a maximum reached after around 20 years. This maximum was positively correlated with the disturbance intensity ($\rho_{Spearman}^{Simpson} = 0.77$). The evenness then stabilized except for two intensively-disturbed plots (number 8 and 12) for which it kept increasing (Fig. 3).

Functional richness and evenness remained stable in control plots over the 30 years of monitoring. In disturbed communities, both trajectories depended on the disturbance intensity, with their maximum values in time being positively correlated to %AGB loss $\rho_{Spearman}^{Richness} = 0.76$ and




Figure 4. Relationship between the initial %AGB loss and community taxonomic richness (a), taxonomic evenness (b), functional richness (c), and functional evenness (d) at 10, 20, and 30 years after disturbance.

$\rho_{Spearman}^{Rao} = 0.60$. For low disturbance intensity, functional richness and evenness displayed a low but long-lasting increase up to a maximum reached after 20-25 years. For high disturbance intensity, they generally displayed a fast but short increase followed, after 10 years, by a slow decrease towards the initial values.

The second-degree polynomial regressions between (i) the percentage AGB loss, and (ii) the taxonomic and functional diversity showed various shapes depending on the diversity indices and on the time since disturbance (Fig. 4). Regarding taxonomic diversity, the relationship between disturbance intensity and diversity was more markedly hump-shaped for richness than for evenness, and peaked at 20% of initial AGB loss. Regarding functional diversity, the relationship was almost linear, and was similar between richness and evenness. All the relationships between disturbance intensity and diversity were stronger 20 or 30 years after disturbance than they were 10 years after disturbance.

3.3 Functional redundancy

Control plots displayed stable functional redundancy over time. All disturbed plots had lower functional redundancy than control plots and followed similar hump-shaped trajectories (Fig. 5). The maximum redundancy loss was positively correlated with the disturbance intensity ($\rho_{Spearman} = 0.47$) and the recovery trajectory had not attained initial values for any disturbed communities after 30 years.

4. Discussion

Our analysis revealed the decoupling between functional and taxonomic trajectories. While the initial differences in taxonomic composition among plots were maintained, the functional composition trajectories converged in the functional space. In terms of diversity, only the taxonomic trajectories validated the IDH that explained humped-shaped post-




Figure 5. Trajectories of the functional redundancy within the initial functional space over 30 years after disturbance.

Shaded areas are the credibility intervals.

disturbance trajectories with an amplitude depending on the disturbance intensity. The decoupling between taxonomic and functional response was explained by variations of community functional redundancy that mitigated the functional impact of disturbance, and appeared as a determinant of community recovery.

4.1 Decoupled taxonomic and functional trajectories

Community taxonomic composition substantially differed before disturbance, as materialized by their distinct location on the NMDS axis 2. These initial differences were maintained along the 30 years following disturbance, with the disturbance leading a displacement on the NMDS axis 1 only. Taxonomic composition changes were similar among plots and may correspond to the recruitment of a group of pioneers, like *Cecropia spp.* or *Miconia spp.*, common to all plots, whatever their initial taxonomic differences and the intensity of disturbance (Denslow & Guzman, 2000; Bongers *et al.*, 2009). Taxonomic trajectories initiated a recovery towards the initial composition. This recovery, although far from being achieved after 30 years, suggested the resilience of community taxonomic composition and the maintenance of initial composition differences (Folke, 2006). Such resilience suggested that species not belonging to the pre-disturbance community were rarely recruited in the long-term, probably because of species dispersal limitation that is common among tropical species (Svenning & Wright, 2005).

Community functional composition trajectories, in contrast, were similar in the functional space (Fukami *et al.*, 2005). The amplitude of the compositional changes depended on the disturbance intensity. Functional trajectories were probably driven by the recruitment of species infrequent or absent before disturbance, and belonging to a pool of species common to all plots. This common pool was composed of pioneers with “resource-acquisitive” strategies displaying low leaf toughness, wood specific gravity, maximum height, and high specific leaf area. The recruitment of these pioneers drove a displacement to the right along the first NMDS axis (Westoby, 1998; Wright *et al.*, 2004; Reich, 2014). Thereafter, the first recruited pioneers were progressively excluded by long-lived, more competitive, and shade-tolerant species. The recruitment of these late-successionals marked the recovery of the initial functional composition with more “resource-conservative” strategies. This recovery translated in the functional plane by a displacement left along the first axis and upward along the second axis (Fig. 1).

The decoupling between taxonomic and functional trajectories suggested that simultaneous operation of trait-based assembly rules and species-level priority effects shaped tree community assembly in Paracou forest. Tree community assembly would then be both deterministic in the functional space, and historically contingent in the taxonomic space.

4.2 The scope of the intermediate disturbance hypothesis

Trajectories of taxonomic richness and evenness differed markedly below and above an intensity threshold (Fig. 3). For low and intermediate disturbance, both taxonomic richness and evenness increased according to the disturbance intensity (Martin *et al.*, 2015; Chaudhary *et al.*, 2016). This suggested that the recruitment of pioneers, previously infrequent or absent, increased the taxonomic richness (Martin *et al.*, 2015; Chaudhary *et al.*, 2016), and that trees surviving after disturbance remained numerous enough to maintain the richness of the pre-disturbance community (Bongers *et al.*, 2009). The pioneers thus recruited became abundant, which balanced the usual hyper-dominance of tropical forests and increased the taxonomic evenness (Baraloto *et al.*, 2012). Above the intensity threshold, like for treatment 3 (high intensity), the taxonomic richness did not exceed the initial value in the first years following disturbance. No increase of the taxonomic richness suggested that the richness of surviving trees was lower than this of the pre-disturbance community, and that this difference was not offset by the recruitment of pioneers. In the Guiana shield, the pool of true pioneers specifically recruited after disturbance is restricted to a few common genera (e.g. *Cecropia* spp., *Miconia* spp., *Tapirira* spp.) (Guitet *et al.*, 2018). Considering the different times after disturbance, there was always a humped-shaped pattern linking the disturbance intensity with the post-disturbance increase in taxonomic richness and evenness (Fig. 4). Both taxonomic richness and evenness were maximized at an intermediate intensity, around 20-25% of AGB lost.

Regarding community functional trajectories, however, no marked differences were observed among post-disturbance trajectories (Fig. 3). Whenever the disturbance intensity, there was first an increase of both functional richness and evenness. Such increase suggested the recruitment of pioneers that were functionally highly different from the pre-disturbance community (Denslow, 1980; Molino & Sabatier, 2001). The recruitment of pioneers hampered the recruitment of other species in the first place. After 15 to 20 years, the first pioneer recruits declined and were replaced by species functionally more similar to the pre-disturbance community, which decreased the functional richness and evenness (Walker & del Moral, 2009).

4.3 The functional redundancy, key of community resilience

The loss of species following disturbance decreased community functional redundancy during the first 15 years. Progressively though, the functional redundancy was restored through the replacement of the species with “resource-acquisitive” strategies by more late-successional species, functionally closer to the pre-disturbance community. The replacement of pioneers by late-successionals followed the lottery recruitment rules. This rules mean an easy recruitment of the first species that becomes increasingly difficult, as the following species are hampered by the emergence of interspecific competition (Busing & Brokaw, 2002). The increasing competition among species explained why the recovery trajectory slowed down

20 years after disturbance. The recovery of the functional redundancy then relied upon the random process of species recruitment, and was increasingly slow and difficult to anticipate (Elmqvist *et al.*, 2003; Díaz *et al.*, 2005). This suggested a low resilience of the functional redundancy with the random recovery of infrequent species increasing the risks of losing keystone species, with unexpected ecological consequences (Jones *et al.*, 1994; Chazdon, 2003; Díaz *et al.*, 2005). Infrequent species might indeed have unique functional characteristics in the ecosystem, apart from the ones considered here, in the ecosystem or be a key resource for some of the fauna (Schleuning *et al.*, 2016).

5. Conclusion

Post-disturbance trajectories of tree community composition and diversity were driven by the recruitment of a determined pool of pioneers, identical among local communities, and independent of the disturbance intensity. The taxonomic composition trajectories maintained the initial differences among communities, while the functional trajectories were similar, and converged in the functional space towards the recovery of the initial composition. The diversity trajectories were contrasted as well. While the functional trajectories remained similar whatever the disturbance intensity, taxonomic trajectories were markedly different from a threshold of 20-25% AGB lost that maximized the taxonomic richness. The Intermediate Disturbance Hypothesis applied well to taxonomic diversity, but not to functional diversity. The decoupling between taxonomic and functional trajectories was mediated by the variations in functional redundancy, as the loss of a species does not necessarily entails the loss of its functional characteristics. Community resilience, in terms of recovery of the pre-disturbance state, was tangible but required several decades, and relied upon the random lottery recruitment of rare species. Given the long-term impacts of disturbance observed, we suggest that 30 years is not enough time for tropical communities to recover, even after relatively low intensity disturbance. Much of community response to disturbance rely on the processes of species recruitment. A more refined understanding of the post-disturbance trajectories would be gained by a closer analysis of the recruitment process.

6. Acknowledgement

We are in debt with all Paracou station technicians and colleagues who helped setting up the plots and collecting data over years. Without their precious work, this study would have not been possible and they may be warmly thanked here. Our work benefited from an “Investissement d’Avenir” grant managed by the Agence Nationale de la Recherche (LABEX CEBA, ref ANR-10-LBX-25). We thank Niklas Tysklind for the usefull help with english proofing.

7. Author's contributions

AM, EM & BH designed the study, developed the analysis framework, and interpreted the results. AM wrote the manuscript with contributions by EM & BH. All authors gave final approval for publication.

8. Data availability

This article is based upon the dataset of the Paracou station, which is part of the Guyafor permanent plot network in French Guiana (Cirad-CNRS-ONF). The dataset is available upon request to the scientific director (<https://paracou.cirad.fr>).

References

- Anderson-Teixeira, K.J., Miller, A.D., Mohan, J.E., Hudiburg, T.W., Duval, B.D. & DeLucia, E.H. (2013) Altered dynamics of forest recovery under a changing climate. *Global Change Biology*, **19**, 2001–2021.
- Aubry-Kientz, M., Hérault, B., Ayotte-Trépanier, C., Baraloto, C. & Rossi, V. (2013) Toward Trait-Based Mortality Models for Tropical Forests. *PLoS ONE*, **8**.
- Baraloto, C., Hérault, B., Paine, C.E.T., Massot, H., Blanc, L., Bonal, D., Molino, J.F.F., Nicolini, E.a. & Sabatier, D. (2012) Contrasting taxonomic and functional responses of a tropical tree community to selective logging. *Journal of Applied Ecology*, **49**, 861–870.
- Bellwood, D.R., Wainwright, P., Fulton, C. & Hoey, A. (2006) Functional versatility supports coral reef biodiversity. *Proceedings of the Royal Society B: Biological Sciences*, **273**, 101–107.
- Bongers, F., Poorter, L., Hawthorne, W.D. & Sheil, D. (2009) The intermediate disturbance hypothesis applies to tropical forests, but disturbance contributes little to tree diversity. *Ecology Letters*, **12**, 798–805.
- Busing, R.T. & Brokaw, N. (2002) Tree species diversity in temperate and tropical forest gaps: the role of lottery recruitment. *Folia geobotanica*, **37**, 33–43.
- Carmona, C.P., de Bello, F., Mason, N.W. & Lepš, J. (2016) Traits Without Borders: Integrating Functional Diversity Across Scales. *Trends in Ecology and Evolution*, **31**, 382–394.
- Chaudhary, A., Burivalova, Z., Koh, L.P. & Hellweg, S. (2016) Impact of Forest Management on Species Richness: Global Meta-Analysis and Economic Trade-Offs. *Scientific Reports*, **6**, 1–10.
- Chave, J., Coomes, D., Jansen, S., Lewis, S.L., Swenson, N.G. & Zanne, A.E. (2009) Towards a worldwide wood economics spectrum. *Ecology Letters*, **12**, 351–366.

- 341 Chazdon, R.L. (2003) Tropical forest recovery: legacies of human impact and natural disturbances. *Perspectives in Plant*
342 *Ecology, Evolution and Systematics*, **6**, 51–71.
- 343 Connell, J.H. (1978) Diversity in tropical rain forests and coral reefs. *Science*, **199**, 1302–1310.
- 344 Denslow, J.S. & Guzman, G.S. (2000) Variation in stand structure, light and seedling abundance across a tropical moist
345 forest chronosequence, Panama. *Journal of Vegetation Science*, **11**, 201–212.
- 346 Denslow, J.S. (1980) Gap Partitioning among Tropical Rainforest Trees. *Biotropica*, **12**, 47–55.
- 347 Díaz, S., Lavorel, S., de Bello, F., Quétier, F., Grigulis, K. & Robson, T.M. (2007) Incorporating plant functional diversity
348 effects in ecosystem service assessments. *Proceedings of the National Academy of Sciences*, **104**, 20684–20689.
- 349 Díaz, S., Tilman, D., Fargione, J., Chapin III, F.S., Dirzo, R., Kitzberger, T., Gemmill, B., Zobel, M., Vilà, M., Mitchell, C.,
350 Wilby, A., Daily, G.C., Galetti, M., Laurance, W.F., Pretty, J., Naylor, R., Power, A., Harvell, A., Potts, S., Kremen, C.,
351 Griswold, T. & Eardley, C. (2005) Biodiversity Regulation of Ecosystem Services. *Trends and conditions*, pp. 297–329.
- 352 Elmqvist, T., Folke, C., Nystrom, M., Peterson, G., Bengtsson, J., Walker, B. & Norberg, J. (2003) Response diversity,
353 ecosystem change, and resilience. *Frontiers in Ecology and the Environment*, **1**, 488–494.
- 354 Folke, C. (2006) Resilience: The emergence of a perspective for social–ecological systems analyses. *Global environmental*
355 *change*, **16**, 253–267.
- 356 Fox, J.W. (2013) The intermediate disturbance hypothesis should be abandoned. *Trends in ecology & evolution*, **28**, 86–92.
- 357 Fukami, T., Bezemer, T.M., Mortimer, S.R. & Van Der Putten, W.H. (2005) Species divergence and trait convergence in
358 experimental plant community assembly. *Ecology Letters*, **8**, 1283–1290.
- 359 Garcia Florez, L., Vanclay, J.K., Glencross, K. & Nichols, J.D. (2017) Understanding 48 years of changes in tree diversity,
360 dynamics and species responses since logging disturbance in a subtropical rainforest. *Forest ecology and management*,
361 **393**, 29–39.
- 362 Goulamoussène, Y., Bedeau, C., Descroix, L., Linguet, L. & Hérault, B. (2017) Environmental control of natural gap size
363 distribution in tropical forests. *Biogeosciences*, **14**, 353–364.
- 364 Gourlet-Fleury, S., Guehl, J.M. & Laroussinie, O. (2004) Ecology & management of a neotropical rainforest. Lessons
365 drawn from Paracou, a long-term experimental research site in French Guiana.

- 366 Guitet, S., Sabatier, D., Brunaux, O., Coutron, P., Denis, T., Freycon, V., Gonzalez, S., Hérault, B., Jaouen, G., Molino,
367 J.F., Péliissier, R., Richard-Hansen, C. & Vincent, G. (2018) Disturbance regimes drive the diversity of regional floristic
368 pools across Guianan rainforest landscapes. *Scientific Reports*, **8**, 3872.
- 369 Hérault, B., Bachelot, B., Poorter, L., Rossi, V., Bongers, F., Chave, J., Paine, C.E.T., Wagner, F. & Baraloto, C. (2011)
370 Functional traits shape ontogenetic growth trajectories of rain forest tree species. *Journal of Ecology*, **99**, 1431–1440.
- 371 Hérault, B. & Piponiot, C. (2018) Key drivers of ecosystem recovery after disturbance in a neotropical forest. *Forest*
372 *Ecosystems*, **5**, 2.
- 373 Hill, M.O. (1973) Diversity and Evenness : A Unifying Notation and Its Consequences. *Ecological Society of America*, **54**,
374 427–432.
- 375 Hubbell, S.P. (2001) *The Unified Neutral Theory of Biodiversity and Biogeography*. Princeton University Press.
- 376 Jones, C.G., Lawton, J.H. & Shachak, M. (1994) Organisms as ecosystem engineers. *Ecosystem management*, pp. 130–147.
377 Springer.
- 378 Lavorel, S. & Garnier, É. (2002) Predicting changes in community composition and ecosystem functioning from plant
379 traits:. *Functional Ecology*, **16**, 545–556.
- 380 Lindenmayer, D.B., Burton, P.J. & Franklin, J.F. (2012) *Salvage logging and its ecological consequences*. Island Press.
- 381 Marcon, E. & Hérault, B. (2015) entropart: An r package to measure and partition diversity. *Journal of Statistical Software*,
382 **67**, 1–26.
- 383 Martin, P.A., Newton, A.C., Pfeifer, M., Khoo, M. & Bullock, J.M. (2015) Impacts of tropical selective logging on carbon
384 storage and tree species richness: A meta-analysis. *Forest Ecology and Management*, **356**, 224–233.
- 385 Mayfield, M.M. & Levine, J.M. (2010) Opposing effects of competitive exclusion on the phylogenetic structure of
386 communities. *Ecology Letters*, **13**, 1085–1093.
- 387 Molino, J.F. & Sabatier, D. (2001) Tree diversity in tropical rain forests: a validation of the intermediate disturbance
388 hypothesis. *Science*, **294**, 1702–1704.
- 389 Morales-Hidalgo, D., Oswalt, S.N. & Somanathan, E. (2015) Status and trends in global primary forest, protected areas,
390 and areas designated for conservation of biodiversity from the Global Forest Resources Assessment 2015. *Forest Ecology*
391 *and Management*, **352**, 68–77.

- 392 Norden, N., Boukili, V., Chao, A., Ma, K., Letcher, S.G. & Chazdon, R.L. (2017) Opposing mechanisms affect taxonomic
393 convergence between tree assemblages during tropical forest succession. *Ecology letters*, **20**, 1448–1458.
- 394 Ollivier, M., Baraloto, C. & Marcon, E. (2007) A trait database for guianan rain forest trees permits intra-and inter-specific
395 contrasts. *Annals of forest science*, **64**, 781–786.
- 396 Piponiot, C., Sist, P., Mazzei, L., Peña-Claros, M., Putz, F.E., Rutishauser, E., Shenkin, A., Ascarrunz, N., de Azevedo,
397 C.P., Baraloto, C., França, M., Guedes, M.C., Coronado, E.N., D'Oliveira, M.V., Ruschel, A.R., da Silva, K.E., Sotta,
398 E.D., de Souza, C.R., Vidal, E., West, T.A. & Hérault, B. (2016) Carbon recovery dynamics following disturbance by
399 selective logging in amazonian forests. *ELife*, **5**, e21394.
- 400 Pulsford, S.A., Lindenmayer, D.B. & Driscoll, D.A. (2016) A succession of theories: purging redundancy from disturbance
401 theory. *Biological Reviews*, **91**, 148–167.
- 402 Randall Hughes, A., Byrnes, J.E., Kimbro, D.L. & Stachowicz, J.J. (2007) Reciprocal relationships and potential feedbacks
403 between biodiversity and disturbance. *Ecology letters*, **10**, 849–864.
- 404 Rees, M., Condit, R., Crawley, M., Pacala, S. & Tilman, D. (2001) Long-term studies of vegetation dynamics. *Science*,
405 **293**, 650–655.
- 406 Reich, P.B. (2014) The world-wide 'fast-slow' plant economics spectrum: A traits manifesto. *Journal of Ecology*, **102**,
407 275–301.
- 408 Réjou-Méchain, M., Tanguy, A., Piponiot, C., Chave, J. & Hérault, B. (2017) biomass: an r package for estimating
409 above-ground biomass and its uncertainty in tropical forests. *Methods in Ecology and Evolution*, **8**, 1163–1167.
- 410 Rutishauser, E., Hérault, B., Petronelli, P. & Sist, P. (2016) Tree Height Reduction After Selective Logging in a Tropical
411 Forest. *Biotropica*, **48**, 285–289.
- 412 Schleuning, M., Freund, J., Schweiger, O., Welk, E., Albrecht, J., Albrecht, M., Beil, M., Benadi, G., Bluethgen, N.,
413 Bruelheide, H. *et al.* (2016) Ecological networks are more sensitive to plant than to animal extinction under climate
414 change. *Nature communications*, **7**, 13965.
- 415 Schnitzer, S.A. & Carson, W.P. (2001) Treefall Gaps and the Maintenance of Species Diversity in a Tropical Forest.
416 *Ecology*, **82**, 913–919.
- 417 Shea, K., Roxburgh, S.H. & Rauschert, E.S. (2004) Moving from pattern to process: coexistence mechanisms under
418 intermediate disturbance regimes. *Ecology letters*, **7**, 491–508.

- 419 Sheil, D. & Burslem, D.F.R.P. (2003) Disturbing hypotheses in tropical forests. *Trends in Ecology and Evolution*, **18**,
420 18–26.
- 421 Sheil, D. & Burslem, D.F. (2013) Defining and defending connell’s intermediate disturbance hypothesis: a response to fox.
422 *Trends in ecology & evolution*, **28**, 571–572.
- 423 Sist, P., Pacheco, P., Nasi, R. & Blaser, J. (2015) Management of natural tropical forests for the future. Technical report,
424 IUFRO WFSE.
- 425 Svenning, J. & Wright, S. (2005) Seed Limitation in a Panamian Forest. *Journal of Ecology*, **93**, 853–562.
- 426 ter Steege, H., Pitman, N.C.A., Phillips, O.L., Chave, J., Sabatier, D., Duque, A., Molino, J.F., Prévost, M.F., Spichiger, R.,
427 Castellanos, H., von Hildebrand, P. & Vásquez, R. (2006) Continental-scale patterns of canopy tree composition and
428 function across Amazonia. *Nature*, **443**, 0–2.
- 429 van Buuren, S. & Groothuis-Oudshoorn, K. (2011) mice: Multivariate imputation by chained equations in r. *Journal of*
430 *Statistical Software*, **45**, 1–67.
- 431 Violle, C., Navas, M.L.L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I. & Garnier, E. (2007) Let the concept of trait be
432 functional! *Oikos*, **116**, 882–892.
- 433 Wagner, F., Hérault, B., Stahl, C., Bonal, D. & Rossi, V. (2011) Modeling water availability for trees in tropical forests.
434 *Agricultural and Forest Meteorology*, **151**, 1202–1213.
- 435 Walker, L.R. & del Moral, R. (2009) Transition dynamics in succession: implications for rates, trajectories and restoration.
436 *New models for ecosystem dynamics and restoration*, pp. 33–49.
- 437 Westoby, M. (1998) A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil*, **199**, 213–227.
- 438 Westoby, M. & Wright, I.J. (2006) Land-plant ecology on the basis of functional traits. *Trends in Ecology and Evolution*,
439 **21**, 261–268.
- 440 Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F., Cavender-Bares, J., Chapin, T., Cornelissen,
441 J.H.C., Diemer, M., Flexas, J., Garnier, E., Groom, P.K., Gulias, J., Hikosaka, K., Lamont, B.B., Lee, T., Lee, W., Lusk,
442 C., Midgley, J.J., Navas, M.L., Niinemets, &., Oleksyn, J., Osada, N., Poorter, H., Poot, P., Prior, L., Pyankov, V.I.,
443 Roumet, C., Thomas, S.C., Tjoelker, M.G., Veneklaas, E.J. & Villar, R. (2004) The worldwide leaf economics spectrum.
444 *Nature*, **428**, 821–827.