

Algorithmic Operation Research

Theodora Panagea - 1115201400135
Anna-Aikaterini Kavvada - 1115201500050

18-10-2019

1

Let $C \subseteq \mathbb{R}^n$ be a convex set with $x_1, \dots, x_k \in C$ and let $\theta_1, \dots, \theta_k \in \mathbb{R}$ satisfy $\theta_i \geq 0$ and $\theta_1 + \dots + \theta_k = 1$. Show that $\theta_1 x_1 + \dots + \theta_k x_k \in C$.

Solution:

We will start by proving the question for $i = 3$ which is trivial. So, suppose $i = 3$ and we also have $x_1, x_2, x_3 \in C$ and $\theta_1 + \theta_2 + \theta_3 = 1$ with $\theta_1, \theta_2, \theta_3 \geq 0$. We will show that

$$\theta_1 x_1 + \theta_2 x_2 + \theta_3 x_3 \in C \quad (1)$$

There is at least one of the θ_i that does not equal 1. Now, assume, without loss of generality that $\theta_1 \neq 1$, where

$$\theta_1 + \theta_2 + \theta_3 = 1 \iff$$

$$\theta_2 + \theta_3 = 1 - \theta_1 \iff, (1 - \theta_1 > 0)$$

$$\frac{\theta_2 + \theta_3}{1 - \theta_1} = 1 \Rightarrow$$

$$1 = m_1 + m_2 = \frac{\theta_2 + \theta_3}{1 - \theta_1} \Rightarrow$$

$$m_1 = \frac{\theta_2}{1 - \theta_1} \quad (2) \quad \text{and} \quad m_2 = \frac{\theta_3}{1 - \theta_1} \quad (3) \Rightarrow$$

$$\theta_2 = m_1(1 - \theta_1) \quad \text{and} \quad \theta_3 = m_2(1 - \theta_1)$$

Thus, from (1), (2) and (3), we have

$$\theta_1 x_1 + (1 - \theta_1)(m_1 x_1 + m_2 x_3) \quad (4)$$

Since C is convex and $x_2, x_3 \in C$, we conclude that $(m_1 x_1 + m_2 x_3) \in C$ as "extensions". Thus, since both points are in the convex C , then the function is in C as well.

Now that we have proven that the function belongs to set C for $i = 3$, we can prove with the same steps that it is also true for the general example $i = k$ where

$$\theta_1 x_1 + \theta_2 x_2 + \theta_3 x_3 + \dots + \theta_k \in C \quad (5)$$

. Following the method above, we find that

$$\begin{aligned} \theta_1 + \theta_2 + \dots + \theta_k &= 1 \iff \\ \theta_2 + \theta_3 + \dots + \theta_k &= 1 - \theta_1 \iff, (1 - \theta_1 > 0) \\ \frac{\theta_2 + \theta_3 + \dots + \theta_k}{1 - \theta_1} &= 1 \Rightarrow \\ m_1 = \frac{\theta_2}{1 - \theta_1} \quad (6) \quad \dots \quad m_{k-1} &= \frac{\theta_k}{1 - \theta_1} \quad (7) \end{aligned}$$

. As in the proof for $i = 3$ it is easy to see that from the (5), (6) and (7) we have

$$\theta_1 x_1 + (1 - \theta_1)(m_1 x_1 + m_2 x_3 + \dots + m_{k-1} x_k) \quad (8)$$

. Thus, we show that since C is convex and all points $x_1, x_2, \dots, x_k \in C$ then $(m_1 x_1 + m_2 x_3 + \dots + m_{k-1} x_k) \in C$. So the function given is proved to be convex and belong to set C .

2

Show that a set is convex if and only if its intersection with any line is convex.

Solution:

We have a convex set S . Suppose we have a convex set A .

\Rightarrow

Let's take two points $p_1, p_2 \in S \cup A$, and point p is on the line segment of $p_1 p_2$.

- $p \in S$ because S is convex
- $p \in A$ because A is convex

So $p \in S \cap A$

Since the two points p_1 and p_2 are arbitrary, we can assume that $S \cap A$ as well. (even in the case where $S \cap A$ is null it is convex)

\Leftarrow

Let $L \cap C$ which is convex for all lines in L . Assume that C is not convex, thus $\exists x, y \in C$, with $\lambda \in [0, 1]$, where

$$\lambda x + (1 - \lambda)y \notin C$$

. But, the line $ax + (1 - a)y, a \in \mathbb{R}$ intersects with C , thus, the intersection is convex.

The line segment between x, y belongs to the intersection $\lambda x + (1 - \lambda)y$, thus belonging to C as well, which negates our previous assumption, proving that C is indeed convex.

3

Show that a set is affine if and only if its intersection with any line is affine.

Solution:

As proved above in question (3), the intersection of convex set S with any line is convex.

Let's take two points x_1 and x_2 , where $x_1, x_2 \in S$. From the proof above, we know that the line between these two points is convex. Thus, the convex combination of x_1 and x_2 belongs to the intersection and expectedly to S . To conclude, since the line belongs to convex set S , the set is affine.

4

A set C is midpoint convex, if whenever two points $a, b \in C$, the average or midpoint $(a + b)/2$ is in C . Obviously, a convex set is midpoint convex. Prove that if C is closed and midpoint convex, then C is convex.

Solution:

5

Show that the convex hull of a set S is the intersection of all convex sets that contain S . (The same method can be used to show that the conic, or affine, or linear hull of a set S is the intersection of all conic sets, or affine sets, or subspaces that contain S .)

Solution:

6

What is the distance between two parallel hyperplanes $\{x \in \mathbb{R}^n : a^T x = b_1\}$ and $\{x \in \mathbb{R}^n : a^T x = b_2\}$?

Solution:

7

Let a and b be distinct points in \mathbb{R}^n . Show that the set of all points that are closer (in Euclidean norm) to a than b is a halfspace.

Solution: