Journal of Ecology



Mechanistic approaches to investigate soil microbemediated plant competition

Journal:	Journal of Ecology
Manuscript ID	JEcol-2023-0291.R1
Manuscript Type:	Mini Review
Date Submitted by the Author:	30-May-2023
Complete List of Authors:	Chung, Anny; University of Georgia, Ke, Po-Ju; National Taiwan University, ; Stanford University, Adler, Peter B; Utah State University, Wildland Resources
Key-words:	Coexistence, plant-microbe interactions, plant-soil feedbacks, competition, resource ratio theory, Plant population and community dynamics, Plant-soil (below-ground) interactions, Plant-plant interactions

SCHOLARONE™ Manuscripts

1	IIILE
2	Mechanistic approaches to investigate soil microbe-mediated plant competition
3	AUTHORS
4	Y. Anny Chung*, Po-Ju Ke, Peter B. Adler
5	
6	Y. Anny Chung: Departments of Plant Biology and Plant Pathology, University of Georgia, USA. ORC-ID:
7	https://orcid.org/0000-0001-5207-2872
8	Po-Ju Ke: Institute of Ecology and Evolutionary Biology, National Taiwan University, Taiwan. ORC-ID:
9	https://orcid.org/0000-0002-8371-7984
10	Peter B. Adler: Department of Wildland Resources and Ecology Center, Utah State University, USA. ORC
11	ID: https://orcid.org/0000-0002-4216-4009
12	*Corresponding author: Y. Anny Chung (yyachung@uga.edu)
13	
14	
15	
16	
17	

ABSTRAC [*]	Т
----------------------	---

- 1. Interactions between plants and soil microbes can influence plant population dynamics and diversity
 in plant communities. Traditional theoretical paradigms view the microbial community as a black box
 with net effects described by phenomenological models.
- 22 2. This approach struggles to quantify the importance of plant-microbe interactions relative to other
 23 competition and coexistence mechanisms and to explain context dependence in microbe effects.
- 3. We argue that a mechanistic framework focused on microbial functional groups will lead to
 conceptual and empirical advances, as demonstrated by extending resource ratio theory to plant microbe interactions.
- 4. We review the diverse pathways by which different microbial functional groups can influence plant
 resource competition. Finally, we suggest approaches to link theory with observations to measure the
 key parameters of our framework.
 - 5. *Synthesis.* Our review highlights recent experimental advancements for uncovering microbial mechanisms that alter plant host resource competition and coexistence. We synthesize these mechanisms into a conceptual model that provides a framework for future experiments to investigate the importance of plant-microbe interactions in structuring plant populations and communities.

KEYWORDS

37 Coexistence, plant-microbe interactions, plant-soil feedbacks, competition, resource ratio theory

38

30

31

32

33

34

35

Soil microbes mediate plant competition

Plant-soil microbe interactions are ubiquitous, and recognition of their role in mediating plant competition has advanced our understanding of plant coexistence (Bever 2003, Wagg et al. 2011, Chung and Rudgers 2016). In the past few decades, negative feedbacks between plant hosts and soil microbes (negative plant-soil feedbacks, negative PSFs) have become the dominant paradigm for understanding the role of soil microbes in maintaining plant diversity (Ke and Miki 2015, Crawford et al. 2019). This paradigm holds that plant species cultivate soil legacies via species-specific microbial assemblages that are more detrimental to themselves than co-occurring species; this self-limitation promotes coexistence.

Classic theory for PSFs is an extension of phenomenological models in which coexistence depends on the relative strength of intraspecific and interspecific interactions, without explicit reference to the mechanisms underlying those interactions (Bever et al. 1997, Revilla et al. 2013). This approach is experimentally tractable. The soil community can be treated as a single entity and evaluated based on its net effects on plant hosts. However, this tractability comes at the expense of mechanistic understanding. Recent reviews of PSFs have emphasized their importance (Gundale and Kardol 2021), but warned of high context-dependence and low predictability (Smith-Ramesh and Reynolds 2017, Bennett and Klironomos 2019, De Long et al. 2019). For example, PSF magnitude and direction can change depending on plant age and development stage, resource availability, abiotic environmental gradients, and host competitive context (Beals et al. 2020, Chung 2023). We should not be satisfied with demonstrations that PSFs occur, but should strive to quantify their contribution to the maintenance of diversity in nature.

To understand and predict the role of plant-soil microbe interactions in plant competition and coexistence, we must go beyond phenomenological description and consider the underlying plant-soil microbe interactions that drive the feedback. A mechanistic framework that explicitly considers microbial functions will help us predict microbial effects on plant coexistence in changing biotic and

Putten et al. 2016, Abbott et al. 2021, Semchenko et al. 2022). While microbial functions are hyperdiverse, they interact with plants primarily as mutualists, pathogens, or decomposers. Within the context of plant coexistence, PSF research has focused primarily on pathogens due to the emphasis on negative density (or frequency) dependence. However, strong evidence exists that coexistence-promoting feedbacks can also be facilitated by mutualists, such as nitrogen-fixing bacteria and mycorrhizal fungi, and decomposers (e.g. Bever 2002, Bennett et al. 2017, Whitaker et al. 2017, Siefert et al. 2018, Veen et al. 2019). Mechanistic approaches that open up the soil microbial black box may require more measurements than traditional phenomenological approaches (see *Future directions*), but are essential for predicting the consequences of rapid environmental change without the need to repeat experiments in new environments. We aim to synthesize recent evidence for microbial mechanisms that mediate resource competition between plants, and propose a resource-based mechanistic framework for understanding and predicting the role of soil microbes in plant competition and coexistence (Fig. 1).

Applying resource ratio theory to understand microbe-mediated plant competition

One of the most important alternative mechanisms for plant coexistence is soil resource partitioning. However, surprisingly few studies have investigated both the effects of resource addition and soil microbes on plant competition, and results have been equivocal due partly to a lack of clear theory (reviewed in Beals et al. 2020). Resource ratio theory (Tilman 1982) provides a way to directly link plant-soil microbe interactions to the key components of resource competition, namely the system's resource supply, the species' resource consumption, and the species' resource requirements (Box 1). These three components are inextricably linked with soil microbes (summarized in Table 1 and reviewed below). Since different microbial functional groups operate simultaneously and are linked by their impacts on resource flow (Fig. 1), a clear understanding of microbial function should help us to predict

how microbial effects vary with environmental context. Below, we highlight recent evidence for each of these mechanistic pathways and empirical approaches to quantify their effects.

Plant-soil microbe interactions and resource supply

One key component of the resource ratio theory is the resource supply (black point in Figure I in Box 1), defined as the resource availability for plants in the absence of plant consumption. Microbe-mediated decomposition alters resource supply for plants, sometimes in species-specific ways. Changes in resource supply caused by plant-soil microbe interactions could move competitors in and out of regions of coexistence (grey region of Figure I in Box 1). Studies in "home field advantage" in litter decomposition show that litter can decompose faster in its "home" location compared to other locations due to local adaptation of decomposers to the prevailing litter chemistry (Veen et al. 2015). This phenomenon can also operate at the scale of different plants within a community, where positive host-decomposer feedbacks favor competitive exclusion by moving resource supply points out of the coexistence region. For example, two Mediterranean oaks (*Quercus suber* and *Q. canariensis*) differed in their foliar nutrient content and associated saprotroph communities, generating soil conditions that favor the establishment of their own seedlings, which causes local aggregation (Aponte et al. 2013).

Among co-occurring European grassland species, grasses with more exploitative sulfur acquisition strategies also increased microbial arylsulphatase activities in their rhizospheres, thus promoting their own growth in a positive feedback loop (Legay et al. 2014).

Microbes can also alter resource availability by mediating nutrient losses. Theoretical models that include decomposition and nutrient recycling have demonstrated that coexistence is promoted when each plant species associates with microbes that accelerate loss of the nutrient that more limits its own growth compared to that of its competitor (Daufresne and Hedin 2005). On the contrary, host-specific decomposers that accelerate loss of nutrients that more limits its competitor's growth should

promote exclusion. In an example that promotes exclusion and leads to monodominance, some pines and poplars thrive in N-poor environments where their litter chemistry and decomposer associations continue to promote N losses. This maintains a favorable environment for these taxa, who are better competitors for N in such systems (Northup et al. 1995, Fierer et al. 2001).

Mutualistic plant growth promoting bacteria (PGPB) can also increase nutrient availability in the soil via specialized metabolic pathways (Ramakrishna et al. 2019). While the host-specificity of these taxa are less well known, plants likely differ in their ability to attract PGPB consortia, which could result in feedbacks. Finally, interactions between microbial groups could also moderate resource supply. For example, the Gadgil effect suggests that competition between ectomycorrhizal fungi and saprotrophs in forest ecosystems can reduce decomposition rates (Gadgil and Gadgil 1971, Fernandez and Kennedy 2016). However, whether these effects impact host coexistence may depend on host resource requirements and the N mining abilities of their respective ectomycorrhizal and saprotroph associates.

Plant-soil microbe interactions and resource consumption

The second component determining species coexistence is their *resource consumption* (consumption vectors in Figure I in Box 1), theoretically defined as the relative rates at which plants deplete resources. Differentiation in species' consumption vector stabilizes coexistence via self-limiting negative feedbacks (Letten et al. 2017). As long as each competitor consumes more of the resource most limiting its growth, then the greater the difference in their consumption vector, the wider the resource supply state-space available for coexistence. A species' consumption vector depends on its resource-specific consumption rates and its stoichiometry, which can be modified by resource mutualisms. For example, mycorrhizal associations increase concentrations of macro- and micronutrients in host tissues, implying increased consumption rates and altered stoichiometry (Smith and Smith 2011, Lehmann and Rillig 2015, Verzeaux et al. 2017, but see Nazeri et al. 2014). Rhizobia are

known to increase tissue N for legume hosts, and some hosts reap benefits when co-infected with rhizobia and mycorrhizal fungi (Primieri et al. 2022). Pathogens can also indirectly alter host consumption and stoichiometry via effects on host physiology. For example, the bacteria *Candidatus* Liberibacter spp. causes huanglongbing in citrus by disrupting phloem transport and feeder root development. This results in plant hosts with imbalanced photosynthate distribution and decreased micronutrient consumption (Mattos-Jr et al. 2020).

Evidence that plant-soil microbe interactions alter the host resource consumption is clear, but there is less consensus about whether such changes increase or decrease opportunities for host coexistence. Among hosts with similar mutualists, the evidence is mixed. For example, several studies have failed to find support for AMF-mediated partitioning of plant access to diverse phosphorus forms (Vogelsang et al. 2006, Reynolds et al. 2006, Ceulemans et al. 2017). However, hosts that associate with different mutualists (e.g. arbuscular mycorrhizal fungi (AMF), ectomycorrhizal (ECM) fungi, or rhizobia) may have access to different subsets of resources, thus altering their consumption vectors in a way that promotes coexistence (Steidinger and Bever 2014, Tedersoo et al. 2020). Furthermore, different resource mutualists respond in predictable ways to changing abiotic conditions. For example, plants benefit most from AMF association when phosphorus is limited (high tissue N:P) (Hoeksema et al. 2010). Thus, we can experimentally test the prediction that an increase in P supply will also decrease AMF effects on host resource consumption vectors (see *Future directions*).

Plant-soil microbe interactions and resource requirement

The final determinant of species coexistence in resource ratio theory is the species' resource requirement, reflected by the species' R* for resources. A species' R* is the minimum amount of a resource required to maintain zero population growth (i.e., where growth balances mortality at the population level). Thus, any effects of soil microbes on a host's mortality rate will alter the host's R*

values, as long as the effects of the microbes are independent of resource-supply rates. For example, when pathogens increase host mortality, the host's R* will also increase (Fig. IA). On the other hand, many belowground microbial mutualists benefit host survival, either via increased tolerance to environmental stress or protection from pathogens (Olanrewaju et al. 2017, Kumar and Verma 2018). There is also evidence of host-specific root endophytes decreasing the P requirement of legume *Ornithopus* species relative to *Trifolium*, potentially due to reduced C demand and P availability compared to the alternative treatment which was an AMF inoculation (Jeffery et al. 2018).

Microbe-mediated changes in R* for a given resource are empirically measurable (see *Future directions*), and could lead to changes in coexistence outcomes. For example, protective mycorrhizae interacting with an inferior competitor could decrease its R* due to lowered mortality rate, potentially shifting the outcome from exclusion to coexistence. How much a protective mutualism decreases mortality also depends on abiotic context, leading to testable hypotheses. For example, survival benefits of symbiosis with an *Epichloë* fungal endophyte for a native grass host was highest in locations with lowest precipitation (Afkhami et al. 2014). This interaction suggests that the relative decrease in R* for this native grass would be the strongest in contexts of low water supply.

Future directions to link theory and observation

Our synthesis identifies new research directions to understand the role of microbial functions in plant competition. First, how can we measure the components of resource ratio theory for plants?

Second, how can we isolate microbial contributions to those components? Past work has approximated R* by allowing plant monocultures to draw down a major resource until its concentration in the soil is constant (at R*) (Dybzinski and Tilman 2007). Supply rates, however, are more challenging to measure in a plant-soil system, and are often approximated by nutrient availability at the beginning of the experiment (Dybzinski and Tilman 2007), or more directly by measuring mineralization rates (Menge et

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

al. 2009). Finally, total resource consumption can be approximated by multiplying plant biomass and tissue resource concentration (Dybzinski and Tilman 2007). Or, it could be possible to use change in tissue stoichiometry to monitor the change in consumption vector angle. Given the number of moving parts, a good starting point for empiricists could be the single resource scenario, where dynamics are controlled by only R* and other components can be ignored (Fig. IA). This provides an entry point for experiments to verify model predictions before moving on to consider plants' consumption and requirements for other limiting resources

Empirical approaches exist to manipulate and quantify the influence of plant-soil microbe interactions on components of the resource ratio theory, with the ultimate goal of parameterizing a mechanistic model. For example, inoculation and exclusion experiments could isolate microbe effects on resource ratio components. While there is no perfect method to manipulate the presence or absence of microbial mutualists vs. pathogens vs. decomposers, established size-based and biocide methods can come close. To target the presence of AMF, researchers can inoculate whole AMF communities using spore extraction from field soils and pair it with a background microbial wash of a smaller size fraction as a control (Schnitzer et al. 2011). A gradient of increasingly small cell size restrictions can create a series of subset communities which can be mapped to known microbial traits (Zanne et al. 2020, Wagg et al. 2021). Size-based exclusion can be used in the greenhouse or field as colonization barriers using different-sized mesh (Chung et al. 2019). In addition to inoculating with specific microbial cultures, selective biocides are also available to remove target specific pathogen groups, such as fungal vs. oomycete pathogens, in competition and feedback experiments (Domínguez-Begines et al. 2021, Liu et al. 2022). While not as naturalistic, recent developments in "EcoFAB" flow-through root chambers allow potential for mechanistic experiments where plant-soil microbe interactions can be studied in close to chemostat conditions with constant resource supply (Gao et al. 2018). Finally, 'omics-based correlative approaches have potential, but rely on future improvements in quantitative microbiome profiling and

functional annotation. As annotations in microbial metagenomics and metatranscriptomics become more robust, it may be possible to link snapshots of microbiome composition to resource function (Knight et al. 2018, Shakya et al. 2019, Taş et al. 2021).

We have highlighted just a few ways that soil microbes affect plant coexistence, but future work should consider the full diversity of microbe-mediated impacts on plant niches. Microbes can modify niche elements for plant resource competition (Peay 2016), microbes can be limiting factors themselves (Koffel et al. 2021), or they can function as competitors of plants for nutrients (Orwin et al. 2011). Furthermore, one microbial functional group could simultaneously modify many elements of coexistence (Table 1). For example, pathogens can affect coexistence by changing host resource consumption, yet the literature has primarily focused on pathogen-mediated host coexistence via effects on mortality. A logical extension of our framework would consider how altered abiotic contexts, such as temperature or soil moisture, could change how multiple microbes alter not just one, but all components of the resource ratio theory. Finally, recognizing roles of specific microbial functional groups also makes it possible to study their interactions which, in theory, can lead to a wider range of possible dynamics than previously appreciated (Jiang et al. 2020).

Conclusions

The microbial black box has served us well. However, we now have the theory and technology to open it up using mechanistic approaches. Our framework generates specific, testable hypotheses that can guide future theory and experiments in a tractable way. Characterizing specific mechanisms by which microbial mutualists, decomposers, and pathogens affect plants is the critical next step in understanding how plant-soil microbe interactions mediate coexistence. Here, we used resource ratio theory as an example to demonstrate how a clear understanding of microbial function can help us predict microbial effects on plant communities under varying environmental contexts. Combining plant

and microbial ecology with plant physiological ecology and ecological stoichiometry could advance our
empirical understanding of mechanisms that underlie microbe-mediated plant competition and
coexistence. Ultimately, a shift away from phenomenological indices will promote better understanding
of plant-soil microbe interactions in the context of other coexistence mechanisms and environmental
change.
Acknowledgements
This work was supported by NSF DEB-1655522 to Adler and Chung, start-up funds from the University of
Georgia and NSF DEB-2105482 to Chung, and Taiwan MOE Yushan scholar program 111V1010-2 and
MOST 111-2621-B-002-001-MY3 to Ke.
Author contributions
All authors worked together to conceive and develop the ideas. Anny Chung took the lead in writing the
manuscript. All authors contributed critically to the drafts and gave final approval for publication.
Data availability
No novel data are associated with this manuscript

Figure 1: This conceptual diagram incorporates different plant-soil microbe interactions into a traditional resource partitioning framework with two plant competitors and two resources. Arrows represent the flow of energy and matter, not effects. The width of arrows indicates the strength of flow, which reflect hypothetical interaction specificity between plants, microbes, and resources (e.g. here each plant is attacked more by its specific enemy, and its specific decomposer processes more host litter). The dashed box represents the plant-enemy system, which contributes to litter via enemy-induced plant mortality. Blue arrows show external supply of resources. When microbes lack host specificity, this figure can be simplified by merging the microbial A and B components for each functional group.




Table 1: Examples of ways different microbial functional groups can affect coexistence through key components of resource ratio theory. Reference cited follow the specific examples discussed in the main text. Empty cells indicate knowledge gaps where the impact of the microbial functional group on the theoretical component requires further research.

	Resource supply	Resource consumption	Resource requirement (R*)
Decomposers	Plant litter stimulates specific		
	decomposer community to alter		
	resource supply rates and ratios.		
	Related to home field advantage		
	effects (Veen et al. 2019)		
Mutualists	The Gadgil effect suggests that	Resource mutualists such as	Resource mutualists can lower R*
	competition between ECM and	mycorrhizal fungi and rhizobia are	for hosts by reducing mortality via
	saprotrophs results in reduced	known to alter host consumption	stress tolerance or protection from
	litter decomposition rates in ECM	rates and stoichiometry (Smith	antagonists (Kumar and Verma
	forests (Gadgil and Gadgil 1971)	and Smith 2011, Primieri et al.	2018)
		2022)	
	Some plant growth promoting		Other changes to R* are also
	bacteria can solubilize		possible via changes to host
	phosphorus, calcium, iron, and		stoichiometry (Jeffery et al. 2018)
	increase their plant availability		
	(Olanrewaju et al. 2017)		
Pathogens	Indirectly contribute to litter	Pathogens can indirectly alter the	Pathogens directly increase host R*
	availability for decomposers via	host's consumption vector by	by increasing mortality.
	the increased production of	changing its stoichiometry	
	senesced host tissues.	(Verzeaux et al. 2017)	

BOX 1: Overview of resource ratio theory

The resource ratio theory (Tilman 1982), and its later generalization as contemporary niche theory (Chase and Leibold 2003, Koffel et al. 2021), provides a framework to study how different soil microbes affect plant-plant resource competition. We summarize the key components of the framework and the criteria for plant coexistence using a graphical approach.

When two hypothetical plants (green and gold colors in Fig. IA) compete for a single resource (e.g., nitrogen), the amount of resource where per capita growth rate equals zero (i.e., where growth balances mortality and the colored lines intercept the x-axis in Fig. IA) is their resource requirement, R*. In this scenario, the species with the lower R* (gold) will outcompete the other species (green) as it depletes resource to a level where the inferior species experiences negative growth.

When plants are competing for two resources, one needs to consider how the availability of the two resources jointly determine plant population growth and how plants differentially consume different resources. We consider two hypothetical plants competing for two essential resources that colimit plant growth according to Liebig's law of minimum, such as nitrogen and phosphorus (Fig. IB; but see Tilman 1982 for other types of resource pairs). We now need to take into account the resource supply point of the environment, which represents the resource availability of the two resources in the absence of plant consumption (black point in Fig. IB). Plant species are characterized by the relative rates at which they deplete the two resources (depicted as consumption vectors; dashed arrows in Fig. IB), and their resource requirement, the resource combinations at which growth balances mortality (depicted as zero net growth isoclines, ZNGIs; solid lines in Fig. IB). For purposes of simplicity, we consider essential resources, a plant's ZNGI is a right-angle corner positioned based on its R* values for the two resources, defined as the amount of a resource that causes growth to balance mortality when the other resource is not limiting. A plant is most limited by the resource with a higher R* value. This graphical framework can be extended to substitutable resources.

When the two plants compete for the two essential resources, coexistence is possible when the following three conditions are met. First, their ZNGIs must intersect, indicating a tradeoff where each plant is most limited by a different resource. Second, each plant must consume more of the resource it finds more limiting. In Fig. IB, these two conditions are fulfilled as the green plant has a higher R* for nitrogen and consumes more nitrogen relative to phosphorus, and vice versa for the gold plant. A final condition for coexistence is that the supply of the two resources must not disproportionately favor any particular plant; specifically, the supply point must be encompassed by the inverse of the consumption vectors (dotted lines in Fig. IB).




Figure I. (A) State-space diagram of two hypothetical plant species (colored as green and gold) competing for one resource. The colored lines represent a species' per capita growth rate under different resource level. (B) State-space diagram of two hypothetical plant species competing for two essential resources. Here, we depict the coexistence criteria in terms of the elements of niche theory, i.e., the resource requirements (solid lines), the resource consumptions (solid arrows as consumption vectors), and the resource supply (black point). See box text for more details.

303	References cited
304	Abbott, K. C., M. B. Eppinga, J. Umbanhowar, M. Baudena, and J. D. Bever. 2021. Microbiome
305	influence on host community dynamics: Conceptual integration of microbiome feedback
306	with classical host–microbe theory. Ecology Letters 24:2796–2811.
307	Afkhami, M. E., P. J. McIntyre, and S. Y. Strauss. 2014. Mutualist-mediated effects on species'
308	range limits across large geographic scales. Ecology Letters 17:1265–1273.
309	Aponte, C., L. V. García, and T. Marañón. 2013. Tree species effects on nutrient cycling and
310	soil biota: A feedback mechanism favouring species coexistence. Forest Ecology and
311	Management 309:36–46.
312	Beals, K. K., J. A. M. Moore, S. N. Kivlin, S. L. J. Bayliss, C. Y. Lumibao, L. C. Moorhead, M.
313	Patel, J. L. Summers, I. M. Ware, J. K. Bailey, and J. A. Schweitzer. 2020. Predicting
314	Plant-Soil Feedback in the Field: Meta-Analysis Reveals That Competition and
315	Environmental Stress Differentially Influence PSF. Frontiers in Ecology and Evolution
316	8:191.
317	Bennett, J. A., and J. Klironomos. 2019. Mechanisms of plant-soil feedback: interactions among
318	biotic and abiotic drivers. New Phytologist 222:91–96.
319	Bennett, J. A., H. Maherali, K. O. Reinhart, Y. Lekberg, M. M. Hart, and J. Klironomos. 2017.
320	Plant-soil feedbacks and mycorrhizal type influence temperate forest population
321	dynamics. Science 355:181–184.
322	Bever, J. D. 2002. Host-specificity of AM fungal population growth rates can generate feedback
323	on plant growth. Plant and Soil 244:281–290.
324	Bever, J. D. 2003. Soil community feedback and the coexistence of competitors: conceptual
325	frameworks and empirical tests. New Phytologist 157:465–473.

plant population dynamics: The utility of the feedback approach. Journal of Ecology 85:561–573.

Bever, J. D., K. M. Westover, and J. Antonovics. 1997. Incorporating the soil community into

329	Ceulemans, T., S. Bode, J. Bollyn, S. Harpole, K. Coorevits, G. Peeters, K. van Acker, E.
330	Smolders, P. Boeckx, and O. Honnay. 2017. Phosphorus resource partitioning shapes
331	phosphorus acquisition and plant species abundance in grasslands. Nature Plants 3:1-
332	7.
333	Chase, J. M., and M. A. Leibold. 2003. Ecological Niches: Linking Classical and Contemporary
334	Approaches. University of Chicago Press, Chicago, IL.
335	Chung, Y. A. 2023. The temporal and spatial dimensions of plant–soil feedbacks. New
336	Phytologist 237:2012–2019.
337	Chung, Y. A., S. L. Collins, and J. A. Rudgers. 2019. Connecting plant–soil feedbacks to long-
338	term stability in a desert grassland. Ecology 100:e02756.
339	Chung, Y. A., and J. A. Rudgers. 2016. Plant–soil feedbacks promote negative frequency
340	dependence in the coexistence of two aridland grasses. Proc. R. Soc. B 283:20160608.
341	Crawford, K. M., J. T. Bauer, L. S. Comita, M. B. Eppinga, D. J. Johnson, S. A. Mangan, S. A.
342	Queenborough, A. E. Strand, K. N. Suding, J. Umbanhowar, and J. D. Bever. 2019.
343	When and where plant-soil feedback may promote plant coexistence: a meta-analysis.
344	Ecology Letters 22:1274–1284.
345	Daufresne, T., and L. O. Hedin. 2005. Plant coexistence depends on ecosystem nutrient cycles:
346	Extension of the resource-ratio theory. Proceedings of the National Academy of
347	Sciences 102:9212–9217.
348	De Long, J. R., E. L. Fry, G. F. Veen, and P. Kardol. 2019. Why are plant–soil feedbacks so
349	unpredictable, and what to do about it? Functional Ecology 33:118–128.
350	Domínguez-Begines, J., J. M. Ávila, L. V. García, and L. Gómez-Aparicio. 2021. Disentangling
351	the role of oomycete soil pathogens as drivers of plant–soil feedbacks. Ecology
352	102:e03430.
353	Dybzinski, R., and D. Tilman. 2007. Resource Use Patterns Predict Long-Term Outcomes of
354	Plant Competition for Nutrients and Light. The American Naturalist 170:305–318.

355	Fernandez, C. W., and P. G. Kennedy. 2016. Revisiting the 'Gadgil effect': do interguild fungal
356	interactions control carbon cycling in forest soils? New Phytologist 209:1382–1394.
357	Fierer, N., J. P. Schimel, R. G. Cates, and J. Zou. 2001. Influence of balsam poplar tannin
358	fractions on carbon and nitrogen dynamics in Alaskan taiga floodplain soils. Soil
359	Biology:13.
360	Gadgil, R. L., and P. D. Gadgil. 1971. Mycorrhiza and Litter Decomposition. Nature 233:133–
361	133.
362	Gao, J., J. Sasse, K. M. Lewald, K. Zhalnina, L. T. Cornmesser, T. A. Duncombe, Y. Yoshikuni,
363	J. P. Vogel, M. K. Firestone, and T. R. Northen. 2018. Ecosystem Fabrication (EcoFAB)
364	Protocols for The Construction of Laboratory Ecosystems Designed to Study Plant-
365	microbe Interactions. JoVE (Journal of Visualized Experiments):e57170.
366	Gundale, M. J., and P. Kardol. 2021. Multi-dimensionality as a path forward in plant-soil
367	feedback research. Journal of Ecology 109:3446–3465.
368	Hoeksema, J. D., V. B. Chaudhary, C. A. Gehring, N. C. Johnson, J. Karst, R. T. Koide, A.
369	Pringle, C. Zabinski, J. D. Bever, J. C. Moore, G. W. T. Wilson, J. N. Klironomos, and J.
370	Umbanhowar. 2010. A meta-analysis of context-dependency in plant response to
371	inoculation with mycorrhizal fungi. Ecology Letters 13:394–407.
372	Jeffery, R. P., R. J. Simpson, H. Lambers, S. Orchard, D. R. Kidd, R. E. Haling, and M. H. Ryan
373	2018. Contrasting communities of arbuscule-forming root symbionts change external
374	critical phosphorus requirements of some annual pasture legumes. Applied Soil Ecology
375	126:88–97.
376	Jiang, J., K. C. Abbott, M. Baudena, M. B. Eppinga, J. A. Umbanhowar, and J. D. Bever. 2020.
377	Pathogens and Mutualists as Joint Drivers of Host Species Coexistence and Turnover:
378	Implications for Plant Competition and Succession. The American Naturalist 195:591-
379	602.

380	Ke, PJ., and T. Miki. 2015. Incorporating the soil environment and microbial community into
381	plant competition theory. Frontiers in Microbiology 6.
382	Knight, R., A. Vrbanac, B. C. Taylor, A. Aksenov, C. Callewaert, J. Debelius, A. Gonzalez, T.
383	Kosciolek, LI. McCall, D. McDonald, A. V. Melnik, J. T. Morton, J. Navas, R. A. Quinn,
384	J. G. Sanders, A. D. Swafford, L. R. Thompson, A. Tripathi, Z. Z. Xu, J. R. Zaneveld, Q.
385	Zhu, J. G. Caporaso, and P. C. Dorrestein. 2018. Best practices for analysing
386	microbiomes. Nature Reviews Microbiology 16:410–422.
387	Koffel, T., T. Daufresne, and C. A. Klausmeier. 2021. From competition to facilitation and
388	mutualism: a general theory of the niche. Ecological Monographs 91:e01458.
389	Kumar, A., and J. P. Verma. 2018. Does plant—Microbe interaction confer stress tolerance in
390	plants: A review? Microbiological Research 207:41–52.
391	Legay, N., E. Personeni, S. Slezack-Deschaumes, S. Piutti, and JB. Cliquet. 2014. Grassland
392	species show similar strategies for sulphur and nitrogen acquisition. Plant and Soil
393	375:113–126.
394	Lehmann, A., and M. C. Rillig. 2015. Arbuscular mycorrhizal contribution to copper, manganese
395	and iron nutrient concentrations in crops – A meta-analysis. Soil Biology and
396	Biochemistry 81:147–158.
397	Letten, A. D., PJ. Ke, and T. Fukami. 2017. Linking modern coexistence theory and
398	contemporary niche theory. Ecological Monographs 87:161–177.
399	Liu, X., I. M. Parker, G. S. Gilbert, Y. Lu, Y. Xiao, L. Zhang, M. Huang, Y. Cheng, Z. Zhang, and
400	S. Zhou. 2022. Coexistence is stabilized by conspecific negative density dependence via
401	fungal pathogens more than oomycete pathogens. Ecology 103:e3841.
402	Mattos-Jr, D., D. M. Kadyampakeni, J. R. da Silva, T. Vashisth, and R. M. Boaretto. 2020.
403	Reciprocal effects of huanglongbing infection and nutritional status of citrus trees: a
404	review. Tropical Plant Pathology 45:586–596.

405	Menge, D. N. L., S. W. Pacala, and L. O. Hedin. 2009. Emergence and Maintenance of Nutrient
406	Limitation over Multiple Timescales in Terrestrial Ecosystems. The American Naturalist
407	173:164–175.
408	Nazeri, N. K., H. Lambers, M. Tibbett, and M. H. Ryan. 2014. Moderating mycorrhizas:
409	arbuscular mycorrhizas modify rhizosphere chemistry and maintain plant phosphorus
410	status within narrow boundaries. Plant, Cell & Environment 37:911–921.
411	Northup, R. R., Z. Yu, R. A. Dahlgren, and K. A. Vogt. 1995. Polyphenol control of nitrogen
412	release from pine litter. Nature 377:227–229.
413	Olanrewaju, O. S., B. R. Glick, and O. O. Babalola. 2017. Mechanisms of action of plant growth
414	promoting bacteria. World Journal of Microbiology and Biotechnology 33:197.
415	Orwin, K. H., M. U. F. Kirschbaum, M. G. St John, and I. A. Dickie. 2011. Organic nutrient
416	uptake by mycorrhizal fungi enhances ecosystem carbon storage: a model-based
417	assessment. Ecology Letters 14:493–502.
418	Peay, K. G. 2016. The Mutualistic Niche: Mycorrhizal Symbiosis and Community Dynamics.
419	Annual Review of Ecology, Evolution, and Systematics 47:143–164.
420	Primieri, S., S. M. Magnoli, T. Koffel, S. L. Stürmer, and J. D. Bever. 2022. Perennial, but not
421	annual legumes synergistically benefit from infection with arbuscular mycorrhizal fungi
422	and rhizobia: a meta-analysis. New Phytologist 233:505–514.
423	van der Putten, W. H., M. A. Bradford, E. Pernilla Brinkman, T. F. J. van de Voorde, and G. F.
424	Veen. 2016. Where, when and how plant–soil feedback matters in a changing world.
425	Functional Ecology 30:1109–1121.
426	Ramakrishna, W., R. Yadav, and K. Li. 2019. Plant growth promoting bacteria in agriculture:
427	Two sides of a coin. Applied Soil Ecology 138:10–18.
428	Revilla, T. A., G. F. (Ciska) Veen, M. B. Eppinga, and F. J. Weissing. 2013. Plant–soil
429	feedbacks and the coexistence of competing plants. Theoretical Ecology 6:99–113.

430	Reynolds, H. L., K. M. Vogelsang, A. E. Hartley, J. D. Bever, and P. A. Schultz. 2006. Variable
431	responses of old-field perennials to arbuscular mycorrhizal fungi and phosphorus
432	source. Oecologia 147:348–358.
433	Schnitzer, S. A., J. N. Klironomos, J. HilleRisLambers, L. L. Kinkel, P. B. Reich, K. Xiao, M. C.
434	Rillig, B. A. Sikes, R. M. Callaway, S. A. Mangan, E. H. van Nes, and M. Scheffer. 2011
435	Soil microbes drive the classic plant diversity-productivity pattern. Ecology 92:296-303
436	Semchenko, M., K. E. Barry, F. T. de Vries, L. Mommer, M. Moora, and J. G. Maciá-Vicente.
437	2022. Deciphering the role of specialist and generalist plant-microbial interactions as
438	drivers of plant-soil feedback. New Phytologist in press.
439	Shakya, M., CC. Lo, and P. S. G. Chain. 2019. Advances and Challenges in
440	Metatranscriptomic Analysis. Frontiers in Genetics 10.
441	Siefert, A., K. W. Zillig, M. L. Friesen, and S. Y. Strauss. 2018. Mutualists stabilize the
442	coexistence of congeneric legumes. The American Naturalist 193.
443	Smith, S. E., and F. A. Smith. 2011. Roles of arbuscular mycorrhizas in plant nutrition and
444	growth: New paradigms from cellular to ecosystem scales. Annual Review of Plant
445	Biology 62:227–250.
446	Smith-Ramesh, L. M., and H. L. Reynolds. 2017. The next frontier of plant–soil feedback
447	research: unraveling context dependence across biotic and abiotic gradients. Journal or
448	Vegetation Science 28:484–494.
449	Steidinger, B. S., and J. D. Bever. 2014. The Coexistence of Hosts with Different Abilities to
450	Discriminate against Cheater Partners: An Evolutionary Game-Theory Approach. The
451	American Naturalist 183:762–770.
452	Taş, N., A. E. de Jong, Y. Li, G. Trubl, Y. Xue, and N. C. Dove. 2021. Metagenomic tools in
453	microbial ecology research. Current Opinion in Biotechnology 67:184–191.
454	Tedersoo, L., M. Bahram, and M. Zobel. 2020. How mycorrhizal associations drive plant
455	population and community biology. Science 367:eaba1223.

456 Tilman, D. 1982. Resource Competition and Community Structure. Princeton University Press. 457 Veen, G. F. (Ciska), G. T. Freschet, A. Ordonez, and D. A. Wardle. 2015. Litter quality and 458 environmental controls of home-field advantage effects on litter decomposition. Oikos 459 124:187-195. Veen, G. F. (Ciska), E. L. Fry, F. C. ten Hooven, P. Kardol, E. Morriën, and J. R. De Long. 2019. 460 461 The role of plant litter in driving plant-soil feedbacks. Frontiers in Environmental Science 462 7:168. 463 Verzeaux, J., B. Hirel, F. Dubois, P. J. Lea, and T. Tétu. 2017. Agricultural practices to improve 464 nitrogen use efficiency through the use of arbuscular mycorrhizae: Basic and agronomic 465 aspects. Plant Science 264:48-56. 466 Vogelsang, K. M., H. L. Reynolds, and J. D. Bever. 2006. Mycorrhizal fungal identity and 467 richness determine the diversity and productivity of a tallgrass prairie system. New 468 Phytologist 172:554–562. 469 Wagg, C., Y. Hautier, S. Pellkofer, S. Banerjee, B. Schmid, and M. G. van der Heijden. 2021. 470 Diversity and asynchrony in soil microbial communities stabilizes ecosystem functioning. 471 eLife 10:e62813. 472 Wagg, C., J. Jansa, M. Stadler, B. Schmid, and M. G. A. van der Heijden. 2011. Mycorrhizal 473 fungal identity and diversity relaxes plant-plant competition. Ecology 92:1303–1313. 474 Whitaker, B. K., J. T. Bauer, J. D. Bever, and K. Clay. 2017. Negative plant-phyllosphere 475 feedbacks in native Asteraceae hosts - a novel extension of the plant-soil feedback 476 framework. Ecology Letters 20:1064–1073. 477 Zanne, A. E., K. Abarenkov, M. E. Afkhami, C. A. Aguilar-Trigueros, S. Bates, J. M. Bhatnagar, 478 P. E. Busby, N. Christian, W. K. Cornwell, T. W. Crowther, H. Flores-Moreno, D. 479 Floudas, R. Gazis, D. Hibbett, P. Kennedy, D. L. Lindner, D. S. Maynard, A. M. Milo, R. 480 H. Nilsson, J. Powell, M. Schildhauer, J. Schilling, and K. K. Treseder. 2020. Fungal

functional ecology: bringing a trait-based approach to plant-associated fungi. Biological
Reviews 95:409–433.