

Observer-Patch Holography Cosmological Vacuum and Structure Formation: Finite Transfer Conditions, Fluctuation Ensembles, and Proto-Objects

B. Müller

September 21, 2026

Paper release: r2044 Released: September 21, 2026

Author affiliation: Bernhard Mueller, Pragma Research Inc.

Abstract

We specify finite quotient ensembles and transfer conditions for Observer-Patch Holography (OPH) cosmology. A declared quotient-intrinsic action and base measure determine a finite state, while reversible Euclidean data, reflection positivity and compatible refinement support its vacuum interpretation. On a separate supplied Whitney action, controlled real-field trajectories and a neutral interacting Hilbert space provide explicit matter and state constructions. We state the further source and transfer conditions that connect such objects to primordial seeds, observer-facing worldlines and large-scale structure observables. On the sourced causal order the expansion history is a record density; the capacity dictionary $\Lambda \ell_P^2 = 3\pi/N$ returns the cosmological constant $+0.62$ and $+0.38$ percent above the Planck central value from the two closure candidates, which sit -2.64 and -2.81 standard deviations below the DESI DR2 display of the same coordinate, and the fixed-capacity equation of state $w = -1$ sits 2.6 to 3.9 standard deviations from the DESI DR2 posterior means.

1 Scope

This technical companion states the vacuum and fluctuation boundary for finite-source cosmology, screen microphysics, the compact results, and finite-patch simulations. The simulator source is <https://github.com/muellerberndt/oph-physics-sim>, and its canonical public data archives are indexed at <https://github.com/FloatingPragma/observer-patch-holography/tree/main/evidence>. The large finite repair archive is a diagnostic whose physical-CMB and production-gravity gates are false; it does not supply the source and transfer receipts required below.

The [spacetime construction](#) gives a controlled flat $1 + 3$ causal-order and count-volume limit for conservative Fibonacci records with a specified population, complete local neighbour reads and layer duration. An observer with complete authenticated ancestry and interval anchors can read $(|I|/|J|)^{1/4}$ relative to a nondegenerate interval J in the same history. For interior timelike diamonds this tends to the proper-time ratio; the readout uses no coordinates, timestamps or density. Its reference fixes relative units, while the law underlying the limit remains supplied.

The vacuum term of the background has one dictionary on the fixed-capacity branch, $\Lambda \ell_P^2 = 3\pi/N$. The two closure candidates for the capacity, 3.2921×10^{122} and 3.3001×10^{122} , return

$\Lambda = 1.0959 \times 10^{-52}$ and $1.0933 \times 10^{-52} \text{ m}^{-2}$ against the Planck-central $1.0891 \times 10^{-52} \text{ m}^{-2}$, residuals of $+0.62$ and $+0.38$ percent, and sit 2.64 and 2.81 posterior standard deviations below the DESI DR2 with CMB display $(2.968 \pm 0.040) \times 10^{-122}$; both brackets are exact rational inequalities machine-checked in the Lean module `CosmologyLedgerBrackets`. Both candidates were exposed after the data and are target-informed declared maps, so the residual is a diagnostic of a frozen formula and carries no predictive weight; a relative capacity residual δ gives a relative constant residual $-\delta/(1+\delta)$, so the large capacity explains the small scale and not the closeness. On the same branch a constant capacity gives $w = -1$ exactly and a monotone capacity gives $w \geq -1$; the Gaussian moment diagnostic of the official DESI DR2 chains places $(-1, 0)$ between 2.6σ and 3.9σ from the four default posterior means, with monotone-branch posterior fractions between 0.007 and 0.11 percent, so the fixed branch is under tension on seen data and the pending freeze before DESI DR3 is a genuine risk commitment. The cosmology postdiction ledger released with these papers records these rows with their classes.

On the rank-three source Gram space with a real axis adjoined, a nonzero closed convex pointed cone is future or past Lorentzian if the proper icosahedral rotations and all rapidities of one boost axis preserve it. Time orientation selects the future cone. This premise must concern possible influence on histories, interventions and clock readouts, beyond an algebraic frame action. A cosmological vacuum requires the additional state, expanding geometry, physical clock calibration and screen-to-volume transfer conditions below. Smooth Einstein promotion requires same-family tensor-curvature convergence or the continuum small-ball/null-balance identification; scalar-curvature convergence alone is diagnostic.

2 Matter and States on a Supplied Spatial Domain

The Whitney construction on the fixed tetrahedral cone gives a controlled matter continuum in a specified sector of one charged scalar action. For $m^2 > 0$, $g \geq 0$, a supplied $u \in C^2([0, T]; H^2(\Omega))$ solving the real cubic Klein–Gordon equation with natural Neumann boundary, and Ritz-projected initial data, uniform conforming refinements give

$$\sup_{0 \leq t \leq T} (\|u_n - u\|_{H^1} + \|\dot{u}_n - \dot{u}_t\|_{L^2}) \leq C_T n^{-1}.$$

The real, zero-current solutions are invariant solutions of the full charged action. The proof uses a fixed shape-regular refinement family, the conserved nonlinear energy and a Ritz error estimate (`paper/tex_fragments/WHITNEY_REAL_CONTINUUM.tex`).

On the original finite mesh with nonzero coupling e , eliminating mean-zero gauge directions gives a complete kinetic metric γ on $\mathcal{Q} \cong \mathbb{R}^{30} \times \mathbb{C}^{13}$. With supplied $\hbar > 0$ and Laplace–Beltrami ordering, the neutral subspace of $L^2(\mathcal{Q}, d\text{vol}_\gamma)$ has a unique nonnegative self-adjoint Hamiltonian and unitary dynamics. Normalized Gaussian half-density states lie in its operator domain and have explicit initial matter moments (`paper/tex_fragments/WHITNEY_INTERACTING_QUANTUM.tex`). These are states of a finite spatial discretization, not a derived cosmological ground state or a continuum quantum field limit.

For the same autonomous action, the Jacobi duration

$$d\tau = \sqrt{\frac{G_q(dq, dq)}{2(E - V(q))}}$$

recovers action time on regular nonturning solutions at supplied energy E , using the full coupled kinetic metric or its gauge reduction (`paper/tex_fragments/WHITNEY_EPHEMERIS_CLOCK.tex`). Its

configuration-based reconstruction supplies an internal model clock. A cosmological interpretation additionally requires observer-selected geometry, physical preparation and units, and a compatible expanding background. Those attachments are separate from these matter and state theorems.

The golden source population of the spacetime construction carries a second matter and state construction on the same population that the causal order uses. Its positive source-metric scalar action has a finite-time continuum limit with an explicit detector error bound, a canonical quantization whose quadratic Hamiltonian has a unique normalized ground state, the original action vacuum, and a bounded probability comparison between the split evolution of that vacuum with a coherent displacement and the continuous evolution of the same finite action, with all sixty-four modes retained at $q = 5$. The joint spatial and temporal refinement of one compact detector is certified at $q = 233$, with total error below 0.0501 against a resolved response above 0.1377. This is a vacuum state of a finite scalar action on the source population, with the population, action and tick supplied. It is neither a reflection-positive Euclidean vacuum of a cosmological release law nor a source-instantiated triple $(J_r, V_r, a_{t,r})$ in the sense of the next section, and it carries no expanding background.

On the same family the expansion history has an exact finite form. The layered read law in comoving coordinates is conformally invariant, so the flat causal order is the causal order of every spatially flat expanding history; the scale factor enters only as the record-density weight a^4 per comoving cell and conformal tick, the count clock reads proper time, and the redshift between two epochs is the fourth root of a record-count ratio. The finite-source cosmology paper states and proves these statements, with their Lean module. The record-production profile is a supplied datum; the vacuum and fluctuation conditions below concern the field on that background and do not select the profile.

3 Finite Quotient Ensemble Theorem

Finite quotient ensemble theorem. Fix a finite regulator r . The physical presentation space is Σ_r , the presentation redundancy groupoid is Γ_r , and the finite physical quotient is

$$Q_r = \Sigma_r / \Gamma_r, \quad \pi_r : \Sigma_r \rightarrow Q_r.$$

The quotient removes nonphysical presentation data: gauge representatives, port relabelings, mesh labels, shard or worker identifiers, queue order, repair schedule identifiers, retry counters, timestamps unless declared semantic, hidden carrier coordinates, and inert ancillary labels. If only settled configurations carry probability, the probability space is the normal-form subset

$$N_r = n_r(Q_r).$$

The map n_r is a normal-form map, not a probability law. Any physical branch inherits this firewall: it must declare the quotient-intrinsic source law or action before a normal form can be read as a selection or prediction claim.

Observable algebras and reference states. In the finite classical case the quotient observable algebra is

$$\mathcal{O}_r = \ell^\infty(Q_r).$$

In the finite quantum case the physical algebra is a declared quotient algebra $\mathcal{A}_r^{\text{phys}}$ with state

$$\omega_r(A) = \text{Tr}(\rho_r A).$$

When the reference object is obtained from a finite lifted carrier, the load-bearing data are not an abstract groupoid cardinality alone. They are a tracially pointed quotient

$$\left(\mathcal{A}_{r,b}^{\text{phys}}, \tau_{r,b}^0\right), \quad \mathcal{A}_{r,b}^{\text{phys}} = z_{r,b} B(\tilde{\mathcal{H}}_r)^{G_r} z_{r,b},$$

where $U_r : G_r \rightarrow U(\tilde{\mathcal{H}}_r)$ is the compact gauge action and $z_{r,b}$ is the central projection for the declared boundary or superselection sector. The reference trace is

$$\tau_{r,b}^0(A) = \frac{\text{Tr}_{\tilde{\mathcal{H}}_r}(A)}{\text{Tr}_{\tilde{\mathcal{H}}_r}(z_{r,b})}.$$

If

$$z_{r,b} \tilde{\mathcal{H}}_r \cong \bigoplus_{\alpha} V_{\alpha} \otimes M_{\alpha},$$

with $d_{\alpha} = \dim V_{\alpha}$ and $m_{\alpha} = \dim M_{\alpha}$, then

$$\mathcal{A}_{r,b}^{\text{phys}} \cong \bigoplus_{\alpha} I_{V_{\alpha}} \otimes B(M_{\alpha}), \quad p_{r,\alpha} = \frac{d_{\alpha} m_{\alpha}}{\sum_{\beta} d_{\beta} m_{\beta}}.$$

These are the induced central-sector weights only after the carrier representation and boundary sector have been fixed.

OPH quotient ensemble. An OPH quotient ensemble is specified by a quotient-intrinsic base weight and action

$$m_r : Q_r \rightarrow \mathbb{R}_{>0}, \quad S_r : Q_r \rightarrow \mathbb{R} \cup \{+\infty\},$$

and

$$w_r(q) = m_r(q) e^{-S_r(q)}, \quad Z_r = \sum_{q \in Q_r} w_r(q), \quad \mu_r(q) = Z_r^{-1} w_r(q).$$

Equivalently, one may state an intrinsic projective prior ν_r on Q_r and set $\mu_r = (n_r)_{\#} \nu_r$. Uniform quotient counting, uniform representative counting pushed to the quotient, groupoid weights, and tracial central-sector weights are different physical claims. The paper must declare which one is being used.

Normal-form projector non-selection. For any retraction $N : Q \rightarrow Q_{\text{nf}}$ onto a subset $Q_{\text{nf}} \subseteq Q$, that is, any map whose restriction to Q_{nf} is the identity, the induced map on laws

$$\mathcal{C}_Q(\mu) = N_{\#} \mu$$

is idempotent:

$$\mathcal{C}_Q^2 = \mathcal{C}_Q.$$

Every law supported on Q_{nf} is fixed; both statements use the retraction property. Therefore settlement or canonicalization never selects a unique physical probability law by itself.

Selection-gap corollary. Let $X \subseteq Q_{\text{nf}}$ be a finite set of quotient-normal candidates distinguished by visible invariants. Normal-form data determine X and its quotient-visible invariants, but they do not choose a member of X . If two laws μ, ν are supported on X and concentrate on different candidates, both are fixed by $\mathcal{C}_Q$. Unique sector selection therefore requires source data: an intrinsic action with a unique minimizer, a declared physical ensemble, or a refinement-stable gap certificate. A defect or holonomy classification can classify possible sectors without choosing the physical sector, and a contraction or repair generator can certify convergence toward a declared target without creating the target law.

Finite MaxEnt quotient ensemble. For finite Q , positive m , and quotient observables $F_1, \dots, F_k$, maximizing

$$\mathcal{H}_m(\nu) = - \sum_q \nu(q) \log \frac{\nu(q)}{m(q)}$$

subject to

$$\sum_q \nu(q) = 1, \quad \sum_q \nu(q) F_a(q) = c_a$$

has the full-support solution, when the feasible full-support surface is nonempty,

$$\mu(q) = \frac{m(q) \exp[-\sum_a \theta_a F_a(q)]}{Z(\theta)}.$$

Boundary optima obey the same formula after restricting to their support. On a finite noncommutative quotient algebra with faithful reference state σ_r ,

$$\rho_r = \frac{\exp(\log \sigma_r - \sum_a \theta_a F_{r,a})}{\text{Tr} \exp(\log \sigma_r - \sum_a \theta_a F_{r,a})}.$$

The finite constraint ledger must name every $F_{r,a}$, its units and support, the target expectation and source, sector or zero-mode treatment, refinement transformation, and proof that no run output or observational output entered the source definition.

Refinement compatibility and RG closure. For $s \succeq r$, let $c_{sr} : Q_s \rightarrow Q_r$ be the physical coarse map. Exact compatibility of weighted ensembles is equivalent to the fiber-sum identity

$$\sum_{q': c_{sr}(q')=q} m_s(q') e^{-S_s(q')} = \alpha_{sr} m_r(q) e^{-S_r(q)}$$

for a constant $\alpha_{sr} > 0$ independent of q . Then

$$(c_{sr})_{\#} \mu_s = \mu_r.$$

If the one-step defects are

$$\delta_{k+1,k} = \|(c_{k+1,k})_{\#} \mu_{k+1} - \mu_k\|_{\text{TV}},$$

then

$$\|(c_{nr})_{\#} \mu_n - \mu_r\|_{\text{TV}} \leq \sum_{k=r}^{n-1} \delta_{k+1,k}.$$

For exponential-family refinement, exact closure requires the fine conditional free energy

$$G_{sr,\theta}(q_r) = -\log \mathbb{E}_{m_s^0} [\exp[-\theta \cdot F_s(Q_s)] \mid c_{sr}(Q_s) = q_r]$$

to equal $\kappa_{sr}(\theta) + R_{sr}(\theta) \cdot F_r(q_r)$. If the residual is uniformly bounded by ε , the induced total-variation defect is bounded by $\tanh \varepsilon$.

Implementation invariance and representative lifting. If implementations A, B have quotient bijections $h_r : Q_r^A \rightarrow Q_r^B$ satisfying

$$m_r^B(h_r q) = m_r^A(q), \quad S_r^B(h_r q) = S_r^A(q), \quad h_r \circ c_{sr}^A = c_{sr}^B \circ h_s,$$

then

$$(h_r)_\# \mu_r^A = \mu_r^B.$$

For tracially pointed quantum quotients the corresponding equivalence is a trace-preserving quotient equivalence. It is invariant under unitary intertwiners preserving the gauge action and sector, and under inert trivial ancillas $A \mapsto A \otimes I_{\text{anc}}$. It is not invariant under arbitrary changes of gauge-representation multiplicities.

If an implementation stores representatives, a representative-level law must be a conditional lift

$$\tilde{\mu}_r(x) = \mu_r(\pi_r x) \kappa_r(x \mid \pi_r x), \quad \sum_{x: \pi_r(x)=q} \kappa_r(x \mid q) = 1.$$

Then $(\pi_r)_\# \tilde{\mu}_r = \mu_r$. Uniform representative sampling yields orbit-size weights and is physical only if representative counting is the declared base measure.

Quotient-lumpable kernels and sampler correctness. A representative kernel $\tilde{P}(x, y)$ descends to Q_r only when

$$P_Q(q, q') = \sum_{y: \pi(y)=q'} \tilde{P}(x, y)$$

is independent of the chosen representative $x \in \pi^{-1}(q)$. For $w(q) = m(q)e^{-S(q)}$ and proposal $R(q, q')$ with reciprocal support, the Metropolis–Hastings acceptance rule

$$a(q, q') = \min \left\{ 1, \frac{w(q')R(q', q)}{w(q)R(q, q')} \right\}$$

gives detailed balance

$$\mu(q)R(q, q')a(q, q') = \mu(q')R(q', q)a(q', q).$$

Repair-informed proposals must include the Hastings asymmetry term; otherwise the stationary law is generically changed.

Repair generators are not selectors. A repair generator of the form

$$L_{\text{rep}} = \sum_C c_C (I - E_C)$$

is a relaxation or sampling object after a law has been selected. Conditional expectations E_C are defined on $L^2(X_r, \pi_r)$, so the reference law π_r is input. On overlapping collars the expectations need not commute. The correct finite gap certificate is the Poincare constant

$$\kappa_r = \inf_{f \perp 1} \frac{\sum_C \|(I - E_C)f\|^2}{\|f\|^2}.$$

If local fiber rates have a positive lower bound γ_* , then

$$L_{\text{rep}} \geq \gamma_* \kappa_r (I - P_0).$$

Finite repair completeness gives $\kappa_r > 0$ at fixed regulator. A uniform refinement lower bound $\inf_r \kappa_r > 0$ is a separate theorem or receipt.

Finite evidence accuracy. For bounded coarse observables O , if

$$\|\hat{\mu}_s - \mu_s\|_{\text{TV}} \leq \epsilon_{\text{samp}}$$

and the refinement defects sum to ϵ_{ref} , then

$$\left| \mathbb{E}_{\hat{\mu}_s}[O \circ c_{sr}] - \mathbb{E}_{\mu_r}[O] \right| \leq 2\|O\|_{\infty}(\epsilon_{\text{samp}} + \epsilon_{\text{ref}}).$$

Continuum-facing observables require a realization map and correlation Cauchy bound in addition to a finite histogram.

Physical-vacuum requirement. A stationary sampler is not a physical vacuum. For any faithful target law one can build a positive transfer operator with that law as ground state, so positivity alone is not a selector. A physical-vacuum interpretation requires source Euclidean slab data

$$\mathfrak{S}_r^E = (Q_r, m_r^0, J_r, V_r, a_{t,r})$$

whose conductance $J_r(q, q') = J_r(q', q) \geq 0$, local potential V_r , and slab thickness $a_{t,r}$ are derived without using the target law or sampler output. With connected event graph,

$$(H_r^E f)(q) = \frac{1}{m_r^0(q)} \sum_{q'} J_r(q, q') (f(q) - f(q')) + V_r(q) f(q)$$

is self-adjoint and bounded below on $L^2(Q_r, m_r^0)$; its finite Feynman–Kac semigroup is positivity improving. Perron–Frobenius gives a unique positive normalized ground state Ω_r , and the finite vacuum law is

$$\mu_r^{\text{vac}}(q) = |\Omega_r(q)|^2 m_r^0(q).$$

For $T_r = e^{-a_{t,r}(H_r^E - E_{0,r})}$, the Doob kernel is stochastic and detailed-balanced with μ_r^{vac} . A continuum interpretation additionally requires reflection positivity or equivalent reconstruction plus refinement compatibility of the transfer family.

Primordial and cosmological prediction firewall. A screen covariance contains incomplete radial information. The complete one-shell map

$$C_\ell = 4\pi \int_0^\infty \frac{dk}{k} \Delta_\zeta^2(k) j_\ell^2(k\chi_\star)$$

has an infinite-dimensional kernel that persists under positivity. OPH physical primordial interpretation requires the source-only stress, single-clock, entropy-repair, curvature-evolution, adiabatic-mode, phase-coherence, physical mode, radial-null-space, and forward-projection receipts together with a scale-natural physical dilation intertwiner or complete radial cross-covariance tomography. A finite radial prior produces a conditional continuation. Observable CMB comparison also requires declared source, solver, dataset, covariance, nuisance, data-use, and pooled-reducer provenance.

Interpretation classes and required receipts. Every ensemble-facing run records its ensemble id, interpretation class, regulator, representative schema, gauge action, canonicalizer, base measure, action coefficients, coarse maps, zero-mode projector, amplitude convention, sampler, smoothing

policy, source provenance, and explicit nonclaims. The seed belongs to the run receipt rather than the ensemble definition. The interpretation classes are

seed, proposal, or repair noise,
 conventional reference ensemble,
 OPH-native quotient ensemble,
 OPH physical vacuum,
 OPH primordial field,
 observable cosmological prediction.

The evidence bundle must keep separate receipts for stationary-law schedule invariance, detailed balance of the aggregate kernel, and pathwise partition invariance. Deterministic replay of semantic random streams or a canonical serial chain is useful, but it is not pathwise partition invariance. Smoothing must preserve raw coefficients, raw spectra, smoothing kernels, smoothed coefficients, smoothed spectra, and hashes of each stage; it is not part of S_r unless explicitly declared.

4 Vacuum Claim Boundary

A simulated field is classified as seed noise, repair jitter, or a conventional reference ensemble unless it satisfies the finite quotient ensemble conditions. An OPH-native vacuum requires source data of the form

$$\mathfrak{S}_r^E = (Q_r, m_r^0, J_r, V_r, a_{t,r})$$

with J_r , V_r , and $a_{t,r}$ derived from OPH source structure rather than from a target law or a desired visual spectrum. Accepted repair moves are generally directed and cannot simply be reused as reversible Euclidean conductances.

5 Structure-Formation Requirements

Object	Required source-side label
Fluctuating screen field	Seed noise or a conventional reference ensemble unless the quotient ensemble conditions hold.
OPH vacuum field	Source Euclidean slab, ground-state, Doob, reflection-positivity, and transfer-refinement conditions.
Proto-object worldlines	Diagnostic until production matter/particle and observer-facing object population conditions hold.
Growth seeds	Primordial bridge, physical mode basis, radial-null, positivity, forward-projection, and source-dilation or tomography conditions.
Large-scale structure comparison	Diagnostic; source, transfer, and official likelihood/data contracts are not supplied.

6 Numerical Acceptance Conditions

The finite-patch calculation reports the following five conditions separately:

- a tracially pointed quotient;
- a source-derived Euclidean slab;
- transfer stability under refinement;
- schedule invariance of the stationary law;
- pathwise invariance under partition changes.

A failed condition is part of the scientific result and excludes the physical vacuum classification for a reference baseline or rendering field.

7 Unsupplied Premises

The cosmological source construction does not supply the reversible conductances J_r , potential V_r , or slab thickness $a_{t,r}$ required above. A source-instantiated reflection-positive vacuum, or an equivalent reconstruction on this cosmological branch, is not supplied. The bridge from a source-derived vacuum to primordial growth seeds without cosmic microwave background or large-scale structure data is not supplied. A physical proto-object or worldline requires conditions distinct from those used for production particle claims.

AI Assistance Disclosure

This research project used research-grade commercial models, including Anthropic’s Fable and OpenAI’s GPT-6 Astra, for research support, software development, editing, and synthesis. The authors are responsible for the paper’s claims, methods, and final text.