```
root@pragma:~# ./init_oph --book
Fetching observer consistency protocols...
Connecting to floatingpragma.io...
```

REVERSE ENGINEERING REALITY

$$\mathcal{P} = \varphi + \alpha_{em}(\mathcal{P})\sqrt{\pi}$$

// WRITTEN BY
BERNHARD MUELLER

HTTPS://OPH-BOOK.FLOATINGPRAGMA.IO

Contents

Front matter 1
No Source Code, No Documentation 3
1 Why Is the Universe So Hard to Reverse Engineer? 9
2 Why Is There Anything Rather Than Nothing? 19
3 Why Would a Universe Need Anybody In It? 29
4 Why Do Two Descriptions Ever Agree? 36
5 Why Do You and I See the Same Table? 43
6 Why Isn't Agreeing With Your Neighbors Enough? 52
7 Why Is There No Such Thing As Now? 60
8 Why Can Strangers Agree Without a Boss? 68
9 What Move Does Reality Actually Make? 76
10 Why Doesn't It Matter Who Goes First? 86
11 What Does Everybody Actually End Up Holding? 95
12 Why Is the World Made of Relations and Not of Things? 104
13 What Is One Piece of Reality Actually Made Of? 111
14 Why Twelve? 119
15 Why Does Space Have Three Directions? 127
16 Why Does Time Run, and Only One Way? 137
17 Why Is There a Speed Limit? 145
18 Why Does Anything Have Energy? 159
19 Why Can't You Ask Everything At Once? 166
20 Why Does Nature Cheat, But Only By Exactly $2\sqrt{2}$? 179
21 Why Does Time Feel Like It's Moving? 186
22 Why Do All the Clocks Agree? 196
23 Why Does Writing Something Down Cost Heat? 203

24 Why Does a Region's Memory Scale With Its Surface? 212
25 Why Does Gravity Look Like Geometry? 219
26 Why Exactly These Forces? 229
27 Why This Exact List of Matter, and Why Three Times Over? 238
28 What Is a Particle, What Is a Wave, and Why Does It Look Like Both? 249
29 Why 1/137, and Why Is There a Limit to What the Universe Can Remember? 261
30 Why Do the Masses Land Where They Do? 270
31 What Is a Lagrangian, and Why Is It the Last Thing? 277
32 Why Does Anything Move? 289
33 Why Was Every Open Problem the Same Problem? 298
34 Why Was Newton Right for Two Hundred Years? 308
35 What Is Anything Made Of? 317
36 Why These Laws and Not Others? 326
37 Why Does the Loop Have No Beginning? 334
38 Why Does Anything Try to Persist? 345
39 Why Is a Bacterium Freer Than an Atom? 352
40 Does It Feel Like Anything, and What Is Left to Decide? 359
41 What Does a Repaired World Look Like? 370
Epilogue: The Last Rung 378
Appendix A: Eighteen Questions 384
Appendix B: Glossary 395
Appendix C: Sources and Further Reading 405
Appendix D: Run It Yourself 410

Front matter

Dedication

For my wife Noon.

Epigraph

> There is a theory which states that if ever anyone discovers exactly what the Universe is for and why it is here, it will instantly disappear and be replaced by something even more bizarre and inexplicable.
>
> There is another theory which states that this has already happened.

Douglas Adams, *The Restaurant at the End of the Universe*, 1980

How to read the claims

This book tells one story about what the world is. A complete self-consistent history and the computation carried by its own local interactions are one reality. No external computer and no extra time play that history. Events, changing local states and records belong to the structure itself. A complete quantum history assigns no simultaneous outcomes to incompatible measurements nobody made, and a closure condition on the whole history says nothing about its inhabitants approaching a final instantaneous state.

Space, durations, fields and matter are readouts of that structure, and the chapters build each of them out of observers repairing their disagreements. Chapter nine derives the move every repair makes and names the interference experiments that read how much quantum coherence a repair keeps. Where a number in these pages was measured before the arithmetic that lands on it, the book says so.

The finite constructions establish conditional charge representations, interaction tensors and common field descriptions. Identifying those representations with the particles in a laboratory requires their dynamics and measured spectral response. The observed mass spectrum and a repair-selected responsive geometry are not derived by the finite identities alone. A condition used by a theorem and an executable example satisfying that condition have different scopes, and the chapters distinguish them.

Locality forbids one thing: information reaching a local update from anywhere but its own neighborhood. Work counters, conventional orderings and global calculations used to check a construction are tools in the hands of whoever does the checking. There is no universal clock. An update count, a causal order, a coordinate, a modular parameter and a clock reading are different things, and moving between any two of them takes a map with a name on it. An observer need not be conscious. The last chapters, on experience and on ethics, take further steps, and they say where each step is taken.

Appendix B gives the working definitions, and the OPH interpretation and working terms use the same vocabulary alongside the papers.

No Source Code, No Documentation

In 1632 Galileo published a book in the form of a conversation between three men and gave the best passage in it to the one who agreed with him. Shut yourself below decks on a large ship, Salviati says. Bring flies and butterflies. Bring a bowl with fish in it. Hang a bottle so that it drips into a wide vessel underneath. Watch how everything behaves while the ship is tied up at the dock, and take your time about it.

Then have the ship put to sea at whatever speed you like, provided the motion is smooth and does not rock you about.

The flies do not pile up against the stern wall. The fish swim as easily toward the front of the bowl as toward the back, without having to work harder in either direction. The drops fall into the vessel below and not one of them lands behind it. You can jump the same distance forward as back. Throwing something to a friend takes the same effort whichever way you face. And from none of these effects, Galileo writes, can you determine whether the ship is moving or standing at rest.

That experiment is harder than it looks, and different in kind from what everybody means by testing something. To test a thing from outside you at least get to stand next to it and poke it. Galileo's man is inside the thing he wants to measure. He is a part of it. What he is asking is whether a container gives any sign of itself to what it contains. The answer he gets back is that this one does not.

Negative results of that kind are the most informative things in physics. He has not found that the measurement is delicate. He has found that there is nothing to measure: the information does not exist inside the cabin, no quantity of butterflies will bring it into existence, and the four centuries since have consisted largely of people discovering that the same is true of the cabin they are actually in.

The book was on the Index of forbidden works within a year and its author spent the rest of his life under house arrest.

Everything that follows is that experiment, run for longer with better equipment.

Six symptoms

What follows is six behaviors of the machine, for now without explanations which take the rest of the book.

Galileo's cabin has never been beaten. Four hundred years of trying, with instruments he could not have imagined. No experiment performed inside a smoothly moving laboratory has ever revealed that it was moving. The nineteenth century made a particular effort, looking for the laboratory's motion through the medium that light was supposed to be waving in, and came up empty to high precision. The information is missing so thoroughly that better instruments are unlikely to supply it.

Clocks disagree. The disagreement is engineered around. The satellites your phone uses to find itself carry atomic clocks, wrong in two directions at once. They move quickly, which makes them tick slow by about seven microseconds a day. They sit high up where gravity is weaker, which makes them tick fast by about forty-five. The two effects do not cancel. Without a correction the position your phone reports would drift off by something like ten kilometers a day.

When the first atomic clock of this kind went into orbit in 1977, the correction was built with a switch on it, because not everyone involved was convinced the physicists were right. It launched switched off. They watched the clock run fast by very close to the predicted amount for about three weeks, and then turned it on.

Asking a question changes the answer. In 1909 a twenty-three-year-old at Cambridge named Geoffrey Taylor set a needle in front of a gas flame and photographed the pattern of light and dark bands around its shadow. Then he began dimming the flame with sheets of smoked glass. He dimmed it until the light crossing his apparatus was so feeble that particles arrived essentially one at a time, with nothing else present for them to interfere with. To collect enough of them he had to leave the plate in place for three months. The bands came out looking exactly as they had at full brightness. Whatever each particle was interfering with, it was not the others, because there were no others.

The second half of that result is even stranger. Put a detector in the apparatus that can tell you which side of the needle each particle went, and the bands disappear. Delicacy does not rescue them. Any detector that succeeds in telling you the route destroys the pattern. Any detector that fails to tell you leaves the pattern intact, and how gently it fails makes no difference at all.

Two things far apart agree too often. In 2015 a team at Delft trapped a single electron in each of two laboratories 1.3 kilometers apart and measured both. The answers matched more often than they could have if each electron had been carrying its answer around with it. The laboratories were separated widely enough that no signal travelling at the

speed of light could have crossed between them while the measurements were happening. That experiment closed the last of the escape routes physicists had been finding in this result for fifty years. The correlation is real, it cannot be used to send a message, and no account in which the particles had properties before anyone looked survives it.

Falling into a black hole appears to destroy the record of what fell. On 6 February 1997 Stephen Hawking and Kip Thorne signed a wager with John Preskill. Hawking and Thorne held that information swallowed by a black hole is erased from the world. Preskill held that it survives. The stake was an encyclopedia of the winner's choice, from which, as the bet specifies, information can be recovered at will. In July 2004, at a conference in Dublin, Hawking conceded and handed Preskill a baseball encyclopedia. Thorne did not concede. Hawking remarked afterwards that burning the encyclopedia and handing over the ashes would have made the losing case rather better than the book did.

The bet was worth making because both positions are unacceptable. If the record is erased, quantum mechanics breaks. If it survives, it has to escape from the one place in the universe that nothing escapes from. Three of the most capable physicists alive spent seven years choosing between two impossibilities. At the end of it Hawking changed his mind and Thorne did not.

Gravity puts an area ceiling on what a region can hold. This last one is arithmetic rather than anecdote. Ask how much information you can pack into a region of space. Storage scales with volume, obviously: a warehouse twice as wide in every direction holds eight times as much. A black-hole horizon behaves differently. Double its radius and its entropy grows by four. Nobody put this in. It falls out of the thermodynamics of black holes. That the same ceiling holds over regions of empty space that contain no black hole at all is the holographic bound, a conjecture in the literature that this theory takes as its own ceiling; no counterexample to it is known.

A universe without a manual

People who take apart hardware for a living meet this problem constantly: a sealed device, no documentation, no source, nobody to ask, and no way to open the case without destroying what is inside. There is a discipline for it. It comes down to a few habits.

Feed it inputs and record what comes out. Change one thing at a time. Watch the timing, because timing leaks structure that the outputs conceal: a system that answers one question quickly and a similar question slowly is telling you that those questions take different routes. Above all, watch what makes it fail. A device gives away more about its architecture in the ten seconds around a crash than in a week of ordinary operation, which

is why the six items above are worth more than a century of things that worked.

And a third habit. Never assume a component you have not found on the board. If you have not located the clock, you do not get to reason about what the clock does. If you have not located the memory, you do not get to assume there is a place where the state is being kept.

This book runs that discipline under one condition, which Galileo's cabin introduced and which never goes away: the engineer is a part. There is no bench to put the machine on.

Every theory anyone has written down begins by assuming a great deal: a space for things to be in, a time for them to happen along, a list of fields, a symmetry group, and a page of numbers measured in laboratories and entered by hand. Those assumptions are not idle. They are what makes the equations work. But a component you assumed is a component you will never find. The six symptoms above are all complaints about the arena rather than about the things moving around in it.

The everyday picture of the world is that there is a three-dimensional container, time runs inside it at one rate for everybody, objects occupy positions in it, and physics describes how the objects move.

In 2002 the programmer Joel Spolsky wrote down a rule about descriptions of that shape. All non-trivial abstractions, to some degree, are leaky. An abstraction is a simplified account that lets you work without knowing what is underneath it. It holds until the thing underneath does something the simplification has no words for, at which point it leaks. The container picture of space and time is a **leaky abstraction** and it leaks in at least six places, which you have just finished reading. What comes back through each leak is not noise. It is a wrong answer, delivered confidently, and reproducible to as many decimal places as you care to measure.

One equation at a time

The framework is called Observer Patch Holography. It starts with bounded systems that read their boundaries, keep records and repair disagreements with their neighbors, and it treats agreement between them as a mathematical constraint. The algebra of questions such a system can read carries the rules of quantum probability. Complete reversible response at the boundary, together with transport inside it, fixes the local symmetries of the forces. A matter law on the same patches moves a real field through a continuum and, at fixed resolution, gives it an interacting quantum state space, with density and magnetic readings coming out of one field reconstruction in both descriptions. Objects that ordinary calculations introduce separately become parts of one connected argument.

The full version lives outside this book, in papers that carry the theorems and in machine-checked proofs for the parts that bear weight. Appendix C says where all of it sits. None of that is reproduced here, because reproducing it would produce a different book. What follows is the argument without the apparatus, down to the points at which a number falls out with nothing fed in to produce it. Where a step carries real load you will be told what it rests on and where the checkable version can be found.

There are equations in this book, not many, and every one of them is doing work. Each one is introduced by a sentence saying what it is about to say and followed by a sentence saying what it said. Every symbol in it is explained in words on the same page.

An equation is compressed prose. It decompresses under four questions.

What does it count? Every quantity in physics is a count of something: how many, how much, how often, how far. Find out what is being counted and you have read half the sentence.

What is being held fixed? An equation states that certain things change together. Nothing changes together unless something else is pinned in place. The pinned quantity is usually the interesting one, and usually not written down.

What is allowed to vary, and what happens at the edges? Set a term to zero and see what the equation collapses into. The collapsed version is generally a law you learned at school. Watching your old law fall out of the new one is how you check that you have understood either of them.

What do the units say? If one side of the equals sign is a length and the other is a time, then a speed is hiding inside the equals sign, or somebody has made a mistake. In 1999 NASA lost the Mars Climate Orbiter because one team's software reported thrust in pounds and the other team's software read it as newtons. The spacecraft went into the Martian atmosphere instead of around it. Checking units is the cheapest way to catch an error of that kind.

One line

There is a single line that working physicists write down when somebody asks what the world is made of. It is short enough to fit on a coffee mug, and it is often printed on one. Four or five terms, depending on how you group them. Each term is a compression: unfold one and out comes a force, or a family of particles, or the reason anything has mass.

Nobody derives that line. It is written down and then adjusted until it matches experiment. The symmetry group inside it was selected because it matched what the detectors reported. The number of particle families in it is three because experiments have found three. Somewhere between nineteen and twenty-six of the numbers in it were measured in laboratories

and typed in. The line is a transcription of those results, and the most successful description of anything human beings have ever produced.

By the last third of this book you will be able to read that line and say where every symbol in it came from.

That is the promise. This prologue makes no other.

Chapter 1

Why Is the Universe So Hard to Reverse Engineer?

Two sentences held up the whole of physics for two hundred and eighteen years, and the man who wrote them had announced, a few lines earlier, that he saw no need to write them.

They sit in the opening pages of the *Principia*, the book that founded the subject, which Isaac Newton published in the summer of 1687. He had spent the previous section defining everything else, carefully. Mass, momentum, force, centripetal force, all set out with their units and their edges. Then he stopped and wrote that he would not define time, space, place and motion, on the grounds that these were well known to everybody.

Then he defined them anyway.

Absolute, true and mathematical time, he wrote, of itself and from its own nature, flows equably without regard to anything external. Absolute space, in its own nature and without regard to anything external, remains always similar and immovable. Two sentences, offered as clarifications of the obvious, and neither of them derived from anything. They are the load-bearing assumptions of the *Principia* and they were put in by hand.

They held until 1905.

They did not hold unopposed. Within thirty years Gottfried Leibniz was writing furious letters about them to Samuel Clarke, who was answering on Newton's behalf and, most historians think, with Newton reading over his shoulder. Leibniz's objection was not technical. It was that absolute space never does anything you could catch it at. Space, he said, is nothing but the order of things that coexist, and time nothing but the order of things that follow one another. Suppose God had built the entire universe three feet to the left of where it is, everything shifted together. What would be different? Nothing measurable, nothing observable, nothing at all. So the position of the universe in absolute space is a fact about a picture somebody drew around it, and about nothing else.

Clarke replied. Leibniz replied to the reply. The correspondence ran to five long letters each and stopped in November 1716, when Leibniz died.

He was right, and it did not matter. Newton's mechanics worked, Leibniz's objection predicted nothing anybody could measure, and absolute space went into the textbooks for two centuries. The complaint survived in a thin line of people who were mostly ignored: George Berkeley restated it in 1721. Ernst Mach restated it in 1883, in a history of mechanics arguing that the inertia of a body has to come from its relation to all the other matter in the universe rather than from its relation to empty space. A young Einstein read Mach and said later that it had shaken him.

That line, the one saying that a quantity nobody can measure from inside is not a quantity, loses every argument it enters for two hundred years, and then wins.

It also leads to an awkward question. Three and a half centuries of work, by some of the most capable people who have ever lived, with instruments of extreme precision and large budgets. The job is not finished. There is no theory of everything. There are two theories, they do not fit together, and the seam between them is where all six of the symptoms in the prologue live.

Why not?

The answer is unflattering. Nobody's cleverness is the problem. Every attempt, including the ones that worked, began by assuming most of the answer.

1.1 Four parameters

In the spring of 1953 a young Freeman Dyson took a train to Chicago to show Enrico Fermi some calculations. Dyson's group had spent months computing how mesons scatter off protons. The numbers agreed with Fermi's measurements. Fermi was polite for about two minutes and then asked how many free parameters were in the calculation. Four, said Dyson. Fermi said that his friend Johnny von Neumann used to say that with four parameters he could fit an elephant, and with five he could make it wiggle its trunk. With that, Dyson wrote later, the conversation was over. He counted it among the luckiest things that ever happened to him, because it saved his group several more years of computing something that could not have meant anything.

Four parameters, and Fermi thought the answer was worthless.

Fermi's objection is arithmetic. There is a difference between a description that fits the data and a description that predicts it. The difference is countable. Suppose you have ten measurements, taken at ten different settings, and you draw a curve through them. If the curve has ten adjustable knobs, as a polynomial with ten coefficients does, the fact that it passes through all ten points tells you nothing, because such a curve passes through any ten points you like. If it has one knob and goes through all ten anyway, you have found something.

The knobs are called free parameters. Counting them is the single most useful thing you can do to an unfamiliar theory.

1.2 The log

What follows is a reading of the last three centuries as a crash log: what was assumed, what broke, and what got fixed. There are five entries.

Absolute space and time. Newton, 1687, by definition, as above. Everything in mechanics rests on it, everything in mechanics works, and for two centuries the assumption is invisible because it is also correct to within any measurement anybody can make.

A medium for light. The nineteenth century established that light is a wave. Waves wave in something: sound waves in air, water waves in water, and light waves in a substance nobody could detect but everyone was confident about, because the alternative was a wave in nothing, which was absurd. The ether had to be stiffer than steel to carry light at that speed, and thin enough that the planets swept through it without slowing. Physicists accepted both properties at once rather than give up the assumption.

In 1887, in a basement in Cleveland, Albert Michelson and Edward Morley floated a sandstone slab on a pool of mercury so it could be turned without vibrating, mounted an interferometer on it, and looked for the Earth's motion through the ether. The apparatus was sensitive enough to detect a fraction of the expected effect. They found nothing. Michelson described the result as decidedly negative and spent much of the next forty years repeating the experiment with better equipment, looking for the drift he was certain had to be there. He received the Nobel Prize in 1907, the first American to win one in the sciences, for his precision optical instruments and the measurements made with them.

Physicists at the end of the nineteenth century are supposed to have thought their subject was finished. Lord Kelvin did not, and said so at the Royal Institution in London on 27 April 1900, speaking on what he called the two clouds hanging over the dynamical theory of heat and light. Cloud one was the Michelson-Morley result: nobody could find the Earth's motion through the ether. Cloud two was that the accepted theory of how energy distributes itself among the parts of a system gave manifestly wrong answers for radiation and for the heat capacity of gases. Kelvin was seventy-five and he had put his finger on the two loose threads in the building. Cloud one became relativity within five years. Cloud two became quantum mechanics within twenty-five. Between them they replaced the foundations of classical physics.

Absolute time, withdrawn. 1905. Einstein's move was to stop treating the null result as a failure and start treating it as data. If no experiment can find the ether, take that as the law, and see what else has to change.

What had to change was simultaneity. Two events that happen at the same moment for you happen at different moments for somebody moving past you. There is no fact about which of you is right.

Notice what survived. Space and time stopped being separate and stopped being absolute, and stayed a stage. There is a four-dimensional arena, events have locations in it, and physics consists of saying what happens at those locations. Ten years later general relativity made the arena bend, which is a far more radical move than it is usually given credit for, because a bending stage is a stage that participates. But it is a stage. Every equation Einstein wrote is a statement about a set of points. The set of points is put in at the start.

Leibniz would have recognized the situation exactly. The quantity that got deleted in 1905, absolute simultaneity, was deleted for his reason: nobody could measure it from inside. The arena survived because nobody had a version of the objection that reached that far.

Definite values, withdrawn. 1926. Quantum mechanics arrived in about eighteen months of contentious argument. It removed the assumption that a thing has a value before you measure it. The rule for converting the theory's central object into odds you can bet on entered physics as a footnote. Max Born wrote a paper on collision processes that summer, said the probability went as the wave function, and then added a correction in proof: it goes as the square of the wave function. That footnote is the Born rule, it is the only place in quantum mechanics where probabilities come from, and it won him the Nobel Prize twenty-eight years later.

Look at what survived. The theory arrived with a space of states, a rule for how they evolve, and a rule for turning them into probabilities. All three were written down as axioms, and Born's has never been wrong. OPH gets it from the agreement that overlapping observers owe each other.

The Standard Model. Assembled between about 1954 and 1973, with a record of successful predictions nothing else in science comes near. Its structure arrived from outside, item by item. The symmetry group is a copy of what the detectors were reporting. The three families are a headcount of what those detectors found. Nineteen of its numbers are laboratory readings, or about twenty-six once neutrino masses are included, plus Newton's constant and the cosmological constant from the other theory entirely.

That is not the elephant problem. Fermi's complaint was that four knobs let you fit anything. The Standard Model's parameters are not knobs in that sense: they are measured quantities, each pinned down by many independent experiments, and turning any of them breaks a hundred other predictions. The Standard Model is not overfitted. Every one of those twenty-six numbers is an input. No part of the theory has an opinion about why any of them has the value it does. Ask why the electron weighs

what it weighs and the correct answer from inside the Standard Model is that somebody measured it.

Nineteen numbers is few for a theory covering every non-gravitational phenomenon ever observed, so the trouble is not the count. The trouble is direction. Numbers go into the Standard Model and predictions come out. There is no arrangement of the theory in which any of those numbers comes out instead. A framework that produced even one of them, from structure, with nothing fed in, would be doing something categorically different from fitting.

The seam. The two theories do not fit, and the way they fail is one you have just been given the equipment to read. It is a parameter-counting failure.

Quantum field theory needs a fixed spacetime to be defined on. That requirement is structural: the definition of a particle in the theory is a statement about the symmetries of the background, so if you do not know the background you cannot say what a particle is. General relativity says the background is dynamical and is determined by what the particles are doing. Each theory requires as an input the thing the other one produces as an output.

Try to force it anyway, quantizing gravity the way everything else was quantized, and you get a theory that produces infinities. Infinities on their own are survivable. The Standard Model has them and disposes of them by absorbing each one into a measured parameter, which is what those nineteen numbers are partly doing. For gravity the absorption never terminates. Every order of the calculation produces a fresh infinity requiring a fresh parameter. At low energies the extra terms are tiny, and Donoghue's calculation shows that the long-distance quantum corrections can be computed anyway. Push toward high energies and the list of parameters never closes. A theory with infinitely many free parameters fits anything and forbids nothing. Fermi's objection to four, taken to its limit.

Every revolution in that log worked by removing an assumption rather than by adding a mechanism. Einstein did not add a component to Newton's machine; he deleted absolute time and watched the rest rearrange itself. Born did not add a mechanism to make particles have definite positions; he deleted the requirement and wrote a footnote about odds.

And every deletion was resisted, always for the same reason. An assumption you have been making since you were four years old does not present itself as an assumption. It presents itself as the world. The ether was not a wild idea somebody talked physicists into; it was the obvious consequence of the obvious fact that waves wave in something. Giving it up meant accepting a sentence that sounded like gibberish. Absolute simultaneity was not a theory. It was what the word "now" meant.

1.3 Four approaches to the problem

Then, in the 1970s, everybody set out to finish the job. The next fifty years produced four major programs. Each one is clever, and each one surrendered something real.

Grand unification, 1974. Howard Georgi and Sheldon Glashow noticed that the three forces of the Standard Model would fit tidily inside a single larger symmetry, and published a paper called "Unity of All Elementary-Particle Forces". The idea was tidy in a way that felt like evidence. Quarks carry charges in thirds, which had never made sense and which suddenly did, because inside the larger symmetry a quark and a lepton are two faces of one object and the thirds are forced. The particles of one generation, which in the Standard Model are a list you write out, became a structure. And the three force strengths, which drift slowly with the energy you probe at, appeared to converge on a single value at an energy around ten to the sixteenth times that of a proton at rest. It was the most beautiful idea in particle physics and it made a prediction: if the forces are aspects of one symmetry, then quarks can turn into leptons, and the proton, which everybody had assumed was stable forever, decays.

The predicted lifetime was around ten to the thirty-first years. The universe is about fourteen billion years old, so you do not sit and watch one proton. You watch ten to the thirty-third of them for a year and wait for one to go. Which is what Super-Kamiokande is: fifty thousand tons of ultrapure water in a mine under a mountain in Japan, lined with photomultiplier tubes, watching since 1996 for a single flash of the right shape.

Nothing has happened. The floor on the proton's lifetime has been pushed past ten to the thirty-fourth years, three orders of magnitude beyond the original prediction. The simplest version of the theory has been dead for decades. The program did exactly what a scientific program should do. It made a specific claim, a detector was built under a mountain to check it, and the answer was no.

Lattice quantum chromodynamics, also 1974. In the same year, Kenneth Wilson made spacetime discrete, as a way of computing rather than as a claim about the world. The equations of the strong force cannot be solved with pen and paper, so put the world on a grid of points, let a computer grind through it, and shrink the spacing at the end. It works. It is how we know the proton's mass from first principles. The calculations run on some of the largest machines ever built.

This is the closest thing mainstream physics has to a granular world. It repays a careful look, because a finite world and a finite calculation are different animals.

The lattice is scaffolding. Its spacing is varied from run to run and taken to zero at the end, and the whole art of the subject is estimating and

controlling the error a finite spacing makes. The action and its parameters, quark masses included, go in as inputs. Lattice quantum chromodynamics is a discrete method for computing the consequences of a continuous theory in a continuous spacetime. It is a triumph, and it surrendered nothing.

String theory. Stop treating particles as points and treat them as extended objects, and you get a different framework for quantum gravity. In 1984 Michael Green and John Schwarz showed that the gauge and gravitational anomalies of a specific superstring theory cancel, and the field changed direction almost overnight. The price was extra dimensions, six or seven of them, curled up small enough to have escaped notice. In 1995 Edward Witten pointed out that the five competing versions of the theory that people had been arguing about were one theory viewed from five angles, in eleven dimensions, and named the eleven-dimensional thing M-theory without saying what the M stood for.

Then came the counting problem. The extra dimensions have to be curled up. There turned out to be an enormous number of ways to curl them, each giving a different set of particles and constants at ordinary energies. Estimates of how many run to ten to the five hundredth, a number with no useful comparison in the physical world. Somewhere in that collection is a version that looks like our universe. The difficulty is that somewhere in it is a version that looks like almost anything.

The program has real derivations to its name. Andrew Strominger and Cumrun Vafa counted quantum states and recovered the area-entropy relation for a class of extremal black holes. Juan Maldacena's correspondence relates specified quantum field theories to strings and gravity in anti-de Sitter settings, so that a gravitational bulk is reconstructed through a different description. That other description is a field theory on a fixed boundary spacetime.

What string theory gave up was the idea that the fundamental objects are points. What it kept was the arena they move in. A string is extended, and it is extended *in* something, and the something has to be specified before the string can be written down.

Background-independent quantum gravity. This program took on the hardest thing on the list, and it deserves to be named properly for that. General relativity says the geometry of spacetime is dynamical: it bends, it responds, it is part of the physics rather than the setting for it. So a quantum theory of gravity should not begin by fixing a geometry and doing quantum mechanics on top. Starting in the 1980s, loop quantum gravity set out to quantize gravity without ever choosing a background metric, and its quantum geometry comes with discrete spectra for area and volume. Abhay Ashtekar and Jerzy Lewandowski's account explains the construction and its starting structures. It succeeded in giving up the fixed metric, which nothing else on this list managed.

It kept a continuum of events to be geometric about. Points, and a smooth manifold of them, and the assumption that the world has a location structure that geometry describes.

Four programs, and a tally. One surrendered points. One surrendered the continuum for the duration of a calculation and handed it back at the end. One surrendered the fixed geometry, which is the hardest item anyone has managed to give up. One enlarged the symmetry group and surrendered the proton's immortality, and was ruled out by experiment.

Every one of them gave up something that had looked non-negotiable. Every one of them defended everything else.

None of them gave up the arena, the set of events with a place for things to happen in. Read the log again and the first half is a sequence of assumptions being removed, one per revolution, at forty-year intervals. The second half is a sequence of structures being added on top of an assumption that nobody touched.

1.4 The rule

There is a fifth program. It sits outside that tally because it did the one thing the other four would not.

On 14 April 2020 Stephen Wolfram announced a project that had been running quietly for years, with much of the technical work carried by a young mathematician named Jonathan Gorard. Its starting object is a **hypergraph**, a collection of relations among elements that are nothing in particular, carrying no positions, distances or dimension. One rewriting rule says that wherever some pattern of relations appears, replace it with another pattern. Apply it wherever it matches, and keep applying it.

Everything else is supposed to come out of that. Space is what the graph looks like from far enough away. The number of dimensions is a property you measure on the graph rather than a number entered in advance. Time is the updating. What an observer can know is the causal graph, the record of which updates fed which. The sharpest part of the project's technical account concerns order. If the causal graph comes out the same whatever order the updates are applied in, a condition called causal invariance, then two observers who disagree about the order still agree about the causes, and special relativity is what that agreement looks like from inside. Keep the orders running in parallel rather than collapsing them and the branching has the shape of quantum mechanics. Its treatment of observers places them inside the evolving system, with bounded computational resources.

Put that beside the four. Every program in the previous section defended the arena. This one deleted it. Every argument in this book is downstream of that move being available at all. The debt is not a courtesy.

Then look at what went onto the board in its place.

The rule. One rule, drawn from a space of candidates that is not small, wanted because it produces this universe rather than one of the others. The method for finding it is to search: enumerate simple rules, run them, inspect what comes out. Nothing inside the framework prefers any candidate to any other. The later proposal, that every rule runs and an observer samples the whole collection of them, is a way of living with the difficulty rather than a way of ending it.

Which is Fermi's objection arriving one floor up. Four parameters will fit an elephant. A free rule carries more freedom than any list of parameters does, because the parameters are what a rule has once somebody has chosen one. The arena came off the board and the largest adjustable thing anybody has written down went on in its place.

So the tally closes on a question rather than a fifth entry. Delete the arena, and what selects the rule? It cannot be a search run from outside, because the outside is the thing that was just deleted. Whatever does the selecting has to be inside, and has to be something the world does to itself.

1.5 What is left

Every mainstream framework in physics begins by assuming four things: a spacetime, a collection of fields living on it, a symmetry group chosen to match what has been observed, and a page of numbers copied off instruments. Every item on that list is load-bearing and each was hard-won. Take any of them away and the equations stop.

Absolute space was never going to come out of Newton's mechanics, because Newton put it in with the definitions. If you write down a spacetime at the top of the page, you will never derive one, and never be able to say why it has three large dimensions instead of two or seventeen. If you write down a symmetry group, you will never explain why that group. If you type in twenty-six numbers, the theory will never have an opinion about any of them. Whatever is going on underneath, you have ruled it out of scope on line one.

So try the other thing, and be strict about it, in the way the prologue's third habit demands. Assume none of it. No arena, no fields, no group, no numbers, and no place for events to be. Then go through the standard inventory of fundamental objects and see what can be kept.

A particle is a thing at a location, and there are no locations, so it goes. A field is worse: a field is an assignment of a value to every point, which presupposes the points and then adds bookkeeping on top. A wave is a disturbance in something. A string is extended, which means extended in something. A geometry is a statement about a set of points and how far apart they are. Even a spacetime event, the humblest object on the list, is a location with nothing at it.

The entire vocabulary of fundamental physics is defined by reference to where things are. The difficulty is that there is no where. Cross out every item that fails and the page is blank.

Almost blank. One object survives. It is a strange thing to find at the bottom of physics, because it is neither matter nor geometry. It is finite. It can read a part of itself and a part of whatever it is next to. It keeps a record of what it read. And it has a way of settling the disagreement when its record and its neighbor's record fail to match.

Note that nothing in this description mentions a location. Reading a neighbor does not require knowing where the neighbor is, only that there is one. Keeping a record does not require a place to keep it, only that the record can be read back. Disagreeing does not require a background, only two accounts of the same thing. This is the first object in the inventory that survives the deletion. It survives because it is described entirely by what it does rather than by where it sits.

That is a peculiar object to build a universe out of. Whether it works is the only question left. It has one property that recommends it immediately and one that ought to worry you. It assumes no arena, which is the entire point. And it has no obvious way of getting started, because to read a neighbor you need a neighbor, and to have a neighbor you need a world, and we have just thrown the world away.

Which is the next question, older than physics. If you assume nothing at all, not even a place for anything to happen, why is there anything here?

Chapter 2

Why Is There Anything Rather Than Nothing?

Alfred North Whitehead and Bertrand Russell spent ten years trying to build arithmetic out of nothing, and they meant it literally: no numbers assumed, no quantities taken as given, no facts about counting smuggled in from the outside world. They began with a handful of logical marks and rules for shuffling them. The plan was to show that everything true about numbers followed from the shuffling alone.

The result, *Principia Mathematica*, ran to three volumes. The proposition that one plus one equals two turns up in volume two, at proposition 110.643, after several hundred pages of preparation. Underneath it the authors added a line of commentary, which reads in full: "The above proposition is occasionally useful."

Russell wrote later that his intellect never quite recovered from the effort, and that he had been definitely less capable of difficult abstraction ever since. He was twenty-eight when they started. Cambridge University Press estimated the work would lose money and asked the authors to contribute fifty pounds each toward publishing it, which they did. The book brought them no income.

It is easy to read that as an anecdote about eccentric mathematicians and miss what the project was. Whitehead and Russell were not being pedantic about two. They were testing whether a body of knowledge could stand on itself: whether you could put down marks and rules with no borrowed meaning, turn the handle, and have arithmetic come out. If it worked, then nothing about numbers had to be assumed, because everything about numbers would be a consequence. Ten years, three volumes, and fifty pounds each was what it cost to find out for arithmetic alone.

What they were building has a name. A **formal system** is a set of marks you are allowed to write, a starting position, and a set of moves that turn one arrangement of marks into another. Meaning is absent by construction. The marks stand for nothing, the moves have no reasons, and the system has no idea what it is about. Hand the rules to somebody who shares no language with you and they will apply them correctly.

Which makes it the only kind of object anybody can examine without assuming something first. Chapter one ended by throwing away everything: no space, no fields, no symmetry group, no numbers, no place for events to

be. The question is whether anything at all survives that. It has a famous form. Leibniz put it in 1714, in a short piece written for a prince: why is there something rather than nothing? Nothing, he pointed out, is simpler and easier.

2.1 Two answers that do not work

The traditional replies come in two shapes.

The first says: something outside made it. Something caused the world. That something is the reason there is anything at all. This is the oldest answer on record and a serious one. Aristotle got to it by asking what moves a thing, and then what moves that, and noticing that the question does not stop on its own.

The trouble it runs into is arithmetic. Whatever you nominate as the cause, you can ask the same question about it. There are two ways out, both bad. Either the chain runs backward forever, in which case you never arrive at a reason and have only postponed the problem an infinite number of times, or you stop at some point and declare that this one thing needs no cause. But if one thing can need no cause, the question is why that particular thing gets the exemption. No answer to that is available except that we stopped there. In practice the item nominated as needing no cause is usually the one the arguer started with.

A first cause is also an assumed component, put on the board because the argument needs it rather than because anybody found it, which is the one rule a reverse engineer cannot break.

There is a second objection, the one that does the damage here, because it does not depend on the regress at all.

Something outside. Something before.

Outside is a spatial word. It means separated by a distance, on the far side of a boundary, in a different region. Before is a temporal word. It means earlier along an ordering that both things sit in. Chapter one did not suggest that space and time might turn out to be built from something else. It took them off the starting inventory altogether, along with the arena and the parameter that things happen along. So there is no distance for a cause to sit at and no earlier for it to have acted in.

Which makes “something outside made it” a sentence assembled out of the two components a reverse engineer has not found on the board. The regress was never the real complaint. A regress is what turns up when a question has quietly helped itself to a place to stand and a moment to stand in, and then goes looking for somewhere to put the answer.

The second reply says it is just here. A brute fact, with nothing behind it, and asking why is a mistake. This has the virtue of stopping the regress immediately and it has one crippling defect. It explains nothing and it also forbids nothing. If the world is here for no reason, then a world with

four spatial dimensions would also be here for no reason, and so would a world with none, and so would a world where the electron weighs twice what it does. You have not explained the world, you have declared that no explanation exists, and in the same motion you have thrown away any hope of understanding why it has the specific shape it has rather than one of the others. The twenty-six measured numbers of chapter one stop being a debt and become the natural state of affairs.

There is a way to feel the size of what that concedes. The mass of the electron is known to about eleven significant figures. Treat it as a brute fact and you have accepted that eleven digits arrived for no reason, and that the same is true of the other twenty-five numbers, and that the total quantity of unexplained precision in the world runs to a couple of hundred digits that simply are what they are. That is a great deal of precision to leave unexplained.

Both replies point away from the thing they are explaining, one to a cause and one to a shrug. Pointing away is the move that has just been withdrawn. With no outside and no before, there is nowhere for an account to come from except the thing itself. So what is left is not a third option chosen over two others. It is the only one still standing. It stands by elimination: a structure accounts for itself, holding together in a way that requires nothing to be added to it, so that asking what put it there has the same status as asking what makes a triangle have three sides.

That also does something to the question in the title. “Why is there anything” wants a because, and a because wants somewhere to come from. Take away the outside and the before and the why has no room to operate in, while the question underneath it survives intact and turns out to be a question about structure: what does a thing have to be like, to be the kind of thing that can hold itself up? What follows is the beginning of that answer. The beginning is small.

Stated like that it is a phrase rather than a proposal. Phrases of that shape have a long history of sounding like wisdom and meaning nothing. It gets edges from three things: what it means for a system to have consequences nobody chose, what it costs to contradict yourself, and what happens when you demand that something survive its own description.

2.2 Two rules and a pencil

Here is a formal system small enough to run by hand. You will need about ninety seconds.

The marks are the letters a and b. The starting position is the string `ab`. There are two moves:

Move one. Take any string you have and wrap it: put an a on the front and a b on the back. So `ab` becomes `aabb`.

Move two. Take any string you have and write it twice. So `ab` becomes `abab`.

That is the whole system. Start from `ab` and apply the moves in any order you like, as many times as you like. From `ab` you can reach `aabb` and `abab`. From `aabb` you can reach `aaabbb` and `aabbaabb`. From `abab` you can reach `aababb` and `abababab`. Go on for a while and you will generate a large and reasonably complicated family of strings, as many as you have patience for.

Here is a question about that family. Can you reach `aab`?

Try it for a minute and you will fail. By itself, failing to find something is not the same as showing it is not there. But look at the counts. The starting string has one a and one b. Move one adds one a and one b. Move two doubles both. So every string you can possibly reach has exactly as many a's as b's, in every case, forever, without anybody having written that down as a rule. `aab` has two a's and one b. It is unreachable. You know this without searching. You have proved something about an infinite collection of strings by looking at four lines of rules. The proof was sitting there before you began searching.

The decisive quantity is the difference between the number of a's and the number of b's. It starts at zero. Neither move can change it: one move adds one to each side of the subtraction, the other doubles both. A quantity the rules cannot touch, however long you run them, is the most useful thing anybody can know about a system.

There is a second thing about that little system that's easy to overlook. The strings it makes are not about anything. `aabb` is not a description of some other, realer `aabb` sitting elsewhere in a warehouse of genuine strings. Ask whether the one in front of you is the authentic article or a copy and the question dies on the way out, because there is nothing available for it to be a copy of. You ran the rules, and what came out is what there is.

That is the first useful property of a system with no meaning in it. Its consequences are not limited to what its author had in mind. Whitehead and Russell wrote down their marks and their moves, and then spent a decade finding out what they had committed themselves to. The answer took three volumes because the answer was not up to them. Nothing was consulted. Nothing was measured. The counts were fixed the moment the rules were written. The ten years were spent reading them off.

2.3 The price of contradictions

Suppose a bank's ledger says that an account holds nothing and also says that the same account holds a hundred pounds. Both entries, on the same page, at the same time.

A clerk looking at that does something sensible. They decide one entry is a mistake, or they pay out the hundred and open an investigation, or they

go and find the deposit slip. What a clerk does not do is conclude that the account holds nine million pounds. People reason around contradictions all day without much difficulty: two entries that cannot both hold get quarantined, while the rest of the ledger goes on working.

A formal system has no clerk. It has marks and moves, it applies every move that applies, and it has no faculty for deciding which of its own lines is the suspicious one. Applied to the letter, the moves take it somewhere no clerk would follow.

Watch. The ledger says the account holds a hundred pounds. From that alone I may write down a weaker sentence: either the account holds a hundred pounds, or the moon is a biscuit. That step costs nothing, because attaching an alternative to a sentence you already have cannot make it less true. Anybody who accepts the hundred pounds is obliged to accept it.

But the ledger also says the account holds nothing, so the first half of my weaker sentence is out. An either-or with its first half ruled out leaves the second half standing.

The moon is a biscuit.

No step in that is a trick. A clerk would sign off on each one taken by itself. Run it again with a different second half and the account holds nine million pounds. Run it again and you get whatever you please, because the second half was never constrained by anything: it was chosen freely at the first step and it survived to the last. Two entries on one page, and the ledger certifies every sentence in the language.

An inconsistent description asserts every world at once, and a description that permits everything distinguishes nothing. It is not a false theory. It is not a theory.

Gottlob Frege found this out in June 1902. The letter that did it came from Russell. Frege had spent two decades building arithmetic out of logic. The second volume of his *Grundgesetze der Arithmetik* was at the printer when Russell wrote to him on the sixteenth with one short observation about the class of all classes that are not members of themselves. Frege replied six days later. He had seen it at once. What he had seen was not that some theorem of his was wrong. It was that his system now proved everything, which is the same as proving nothing. He added an appendix while the volume was in press. It opens by saying that hardly anything more unwelcome can befall a scientific writer than to have one of the foundations of his edifice shaken after the work is finished.

Which is why *Principia Mathematica* runs to three volumes. The several hundred pages of preparation, and one plus one arriving at proposition 110.643, are not eccentricity. They are the price of a system that Russell's letter could not be written to.

The defense did not hold. Russell's letter worked by getting a system to talk about itself, and *Principia Mathematica* was built to make that impossible. In 1931 Kurt Gödel showed that it cannot be made impossible. Any

system strong enough to carry out ordinary arithmetic is strong enough to describe its own formulas, and once a system can describe its own formulas, sentences about itself are among the things it can build. What Gödel assembled out of one was milder than Frege's disaster and worse to live with. Frege's system proved everything, which killed it outright. Gödel's sentence is true, the system it is written in cannot prove it, and no patch closes the gap, because the repaired system has a sentence of its own. Whitehead and Russell had bought a system that could not be attacked the way Frege's had been. What was not for sale was a system with nothing to say about itself, because a system with nothing to say about itself is too weak to be the foundation of anything. Chapter thirty-seven is where the construction gets taken apart.

The result was read as a demolition, and as a demolition it was decisive: the programme of standing mathematics on logic did not recover. Read the same theorem as a specification and it says something the demolition reading discards. A structure rich enough to describe arithmetic is rich enough to describe itself. No arrangement of the axioms takes that capability away. It arrives with the strength.

Chapter one gave you the tool for seeing why that matters here. A theory with infinitely many free parameters fits anything and forbids nothing, which was the fatal problem with quantizing gravity by brute force. A contradiction is the same disease in its terminal form: not many knobs but all of them, not a poor prediction but every prediction at once. Consistency is what buys a description the right to rule anything out.

So the demand this book starts from can be stated. **A structure counts as existing exactly insofar as it accounts for itself with nothing left over.** No external cause is allowed, because a cause is one more thing standing in need of an account. No leftover assumptions are allowed, because an assumption is a thing you would have to explain. And it must not contradict itself, because a contradiction proves everything and therefore distinguishes nothing.

The obvious objection is that this demand is so severe that nothing could satisfy it. The answer to that objection is a number.

2.4 One demand, and a calculator

Take a calculator, or a phone, or a piece of paper. Pick any positive number at all. Any at all: your age, the year, seven.

Then do this. Divide one by your number, add one, and keep the answer. Then do it again to the answer. Then again.

Start with seven and you get 1.142857, then 1.875, then 1.533333, then 1.652174, then 1.605263, then 1.622951, then 1.616162, then 1.618750, then 1.617761. The numbers are hunting, overshooting one way and then the other, and closing in. Keep going and they settle on 1.6180339887.

Start with a thousand instead. Or with 0.001. Or with the number of stairs in your building. Same destination, every time, to as many decimal places as you have patience for.

There is a small bonus in the arithmetic if you look at the fractions rather than the decimals. Starting from seven you get 8/7, then 15/8, then 23/15, then 38/23, then 61/38, then 99/61, then 160/99. Read the numbers down the page: 7, 8, 15, 23, 38, 61, 99, 160. Each one is the sum of the two before it. You did not ask for that and the rule you were applying says nothing about addition.

What is 1.6180339887, and why does everything land on it?

It is the number that the operation leaves alone. If you feed in a number and get the same number back, then that number satisfies x equals one plus one over x. Only one positive number does. Multiply through by x and it reads x squared equals x plus one, which is a quadratic. Its positive root is one plus the square root of five, all over two. Which is 1.6180339887, and which people have called the golden ratio since long before anybody could write it down that way.

The hunting has a mechanism. When your number is too big, one over it is too small, so the answer comes out below the target. When your number is too small, one over it is large, so the answer overshoots above. Every step lands on the far side of the destination from where you started, and lands closer. The error does not merely shrink, it changes sign each time and shrinks, which is why the decimals settle from both directions at once.

A number that a process leaves alone is called a **fixed point**. This is the first of many in this book. Where it came from is the part that matters. Nothing was measured. No experiment was performed and no data existed at any point in the procedure. A single demand, that the operation return what it is given, reached into the continuum of real numbers and picked out exactly one.

Look at the shape of that demand rather than at the number it produced. The equation reads x equals one plus one over x. The unknown stands on both sides of the equals sign. What is being solved for appears inside its own specification, which in most arguments is the move that gets called circular and thrown out. Here it is the entire mechanism. Take the self-reference out and no equation is left, only a continuum with nothing in it to prefer.

That is the shape of the answer to the question in the title. A requirement severe enough to have exactly one solution.

That quadratic has two roots. The other is minus 0.618034. Check it: one divided by minus 0.618034 is minus 1.618034, and adding one gives you minus 0.618034 back. It is a fixed point, exactly as much as the other one is. But start your calculator anywhere near it and the numbers do not settle there, they flee. Being a point that a process leaves alone is one property. Being a point that a process moves toward is a different property.

A structure can perfectly well have fixed points that nothing ever arrives at.

2.5 What one requirement is worth

Look at what the demand bought. Before it there was a continuum of positive numbers available, uncountably many, with no reason to prefer any one of them. After it, there was one. One number, to as many decimal places as anybody cares to compute.

That exchange rate between what goes in and what comes out is the engine of this book. It fails in both directions.

Require nothing and you get nothing. This is the part that sounds wrong, because requiring nothing sounds like permitting everything, and permitting everything sounds generous. It is empty. A structure with no requirements on it does not have many possible forms; it has no form, because a form is exactly a set of things that are not allowed. If every arrangement is as good as every other, then nothing distinguishes anything, and there is no fact about what the structure is.

Which is the same failure the bank ledger had, arrived at through the opposite door. A contradiction permits every conclusion and therefore asserts nothing. A structure with no constraints permits every configuration and therefore is nothing. Total inconsistency and total freedom are emptiness. A great deal of confused metaphysics comes from noticing only one of them.

Between those two failures is a window, far narrower than anybody expects. Put a small number of requirements on a structure and what survives is not a comfortable family of possible worlds to choose among. It is very nearly one world, with a definite number of directions in it, a definite list of things it is allowed to contain, and definite numbers attached to them, none of which anybody selected. The golden ratio is that phenomenon at the smallest scale it can occur at: one requirement, one survivor, out of a continuum.

This cuts both ways and the second way is the uncomfortable one. If adding a requirement collapses a continuum to a point, then a requirement that slips in without being noticed produces a world that arrives without being earned. Chapter one is three centuries of exactly that going wrong. So the constraints have to be counted, they have to be few, and every one of them has to be visible.

2.6 Three requirements

Everything here comes out of three requirements, taken up in the chapters that follow. In the theorem library they are the premises the theorems

carry; the loop that holds the whole account together comes in chapter thirty-seven.

The first says that whatever the world is made of, each piece of it sees only a part and keeps a record of what it saw. The second says that where two pieces overlap, their records have to agree, and that any relabeling of a piece has to be a move the system can carry out on itself, with no hand reaching in. The third says that beyond what those two force, nothing is assumed: whatever is not pinned down stays as unconstrained as it is possible to be.

There is very little in there. Nothing about space, because there is none yet. Nothing about time, particles, forces or numbers. No mention of what the pieces are made of, how many there are, or what the records say. The first two are constraints on bookkeeping and the third is a refusal to add anything.

Three, and not more, and if that sounds too few to build a universe from, the history of the last comparable attempt is instructive.

Euclid began his geometry with five postulates. Four of them are the sort of thing nobody argues with: you can draw a line between two points, you can extend a line, you can draw a circle with any center and radius, all right angles are equal. The fifth is about parallel lines and it is longer and clumsier than the other four put together. Euclid avoids using it for as long as he possibly can. For two thousand years mathematicians tried to prove it from the other four and get rid of it, and failed, and assumed they had merely failed rather than attempted something impossible.

In 1733 an Italian Jesuit named Giovanni Saccheri published a book titled *Euclid Freed of Every Flaw*, in which he tried the last remaining tactic. Assume the fifth postulate is false, grind out the consequences, and wait for a contradiction. He ground out a great many consequences. Triangles whose angles sum to less than two right angles. Lines that approach each other forever without meeting. An entire coherent body of theorems, dozens of them, all following perfectly from four postulates and the denial of the fifth, and not one contradiction anywhere in it.

What Saccheri had in his hands was a complete and consistent geometry of curved space, a century before anybody else found one. What he concluded was that the results were repugnant to the nature of a straight line, which he offered as the contradiction he had spent the book hunting for. He published, and died that year, having discovered a new kind of space and rejected it.

A hundred and eighty years later Einstein went looking for a mathematics of gravity and found he needed geometry without the fifth postulate: space whose triangles do not add up to two right angles, whose parallel lines do not stay parallel, and whose shape is settled by what is sitting in it. The postulate Saccheri was trying to rescue is false in the universe he wrote the book in.

Two things come out of that. Four postulates and a decision about a fifth had been quietly fixing the shape of space for two thousand years. Nobody found out by inspection, because you cannot see a postulate by staring at it. They found out by working: by taking the requirements seriously and grinding out what they force, which took twenty centuries and was being got wrong as late as 1733. And the consequences of a small set of requirements are not up for negotiation. Saccheri rejected his, and they held regardless.

That is the whole starting position. There is a hole in it.

Look at what the three tools above have in common.

A formal system produces consequences. Those consequences are strings: arrangements of marks that stand one way rather than another and can be read back. Consistency is a property only certain objects can fail to have. A rock cannot contradict itself. No arrangement of atoms is a contradiction, because contradicting yourself means holding two claims about the same thing and a rock holds no claims. And a fixed point is whatever satisfies a condition, where a condition is a demand, and a demand has to be made of something and stated somewhere before it can do any demanding.

Every one of those is a property of a description rather than of a lump of world.

Which puts a clause into the founding demand that nobody wrote there. "Accounts for itself" has no application to a structure containing no accounts. Ask whether a universe with nothing in it that keeps a record is consistent, and the question slides off: there is nothing in such a place to be right or wrong about anything, so there is nothing for consistency to be a property of. The word does not become false. It stops reaching.

So the founding demand turns out to require something nobody put into it and nobody would have wanted. Somewhere in that structure there has to be a thing that holds a description, sets it against something, and is capable of getting it wrong. Otherwise the demand has nothing to bite on, and a structure the demand cannot bite on fails it, and by its own terms it does not exist.

One more thing follows. It does not get settled here. If the structure is everything, then whatever its record-keepers describe is inside it, so some of what they hold is a description of the thing they are part of. Gödel's theorem says a structure that strong can carry such a description. The demand above says it has to.

A calculator, two letters of the alphabet, and a bank ledger with two entries on it. What comes out at the bottom of them is that the universe has to contain something capable of being wrong.

Chapter 3

Why Would a Universe Need Anybody In It?

In 1883 a science teacher at the state normal school in Whitewater, Wisconsin, set out to keep janitors out of his classrooms. The heating in the building ran off a coal boiler in the basement. The only way to find out whether a classroom was too cold was for somebody to walk into it and look at the thermometer on the wall, which happened during lessons, repeatedly, all winter. Warren Johnson's solution was a device that would notice the temperature itself and ring a bell in the boiler room. He patented it, then built a company to sell it, and that company outlived him by more than a century.

The thing he built is on your wall. It has a sensor, a number you set, and a switch. It does one thing forever: it looks at what the temperature is, looks at what you asked for, and if those two disagree it does something about it.

Chapter two ended by discovering that consistency is a property of descriptions, and that descriptions have to be held by something. The obvious next move is to reach for a person. It is the wrong move, because reaching for a person means assuming a brain, a body, a planet, several billion years of chemistry and a spacetime for all of it to happen in, which is every assumption chapter one deleted arriving back through the side door.

The right move is to ask what the job actually requires, and then find the cheapest object in the universe that can do it. The answer is the thermostat on the wall. Nothing has to be added to it.

3.1 What the job requires

Take the job seriously. Something has to hold a description of part of the world. What does it need, and what can be removed?

It has to see something, and not everything. This one runs the opposite way from intuition, so take it slowly. You might suppose that a better describer is one that sees more, and that a perfect describer would see everything.

Lewis Carroll got there first. In *Sylvie and Bruno Concluded*, published in 1893, a character called Mein Herr explains that his country's cartographers began with maps at six inches to the mile, then tried six yards to

the mile, then a hundred yards, and finally arrived at the grandest idea of all, a map on the scale of a mile to the mile. It has never been spread out, he admits: the farmers objected that it would cover the whole country and shut out the sunlight. So we now use the country itself, as its own map, and I assure you it does nearly as well.

That is the whole argument. A description that leaves nothing out is a duplicate. A duplicate tells you precisely as much as the original does and adds nothing to it. Description begins where something gets left out. A boundary is what makes a thing a describer rather than a copy. It belongs at the top of the list rather than in a footnote about limitations.

It has to write. A description exists where it is recorded and nowhere else, because an unrecorded one leaves the world in exactly the state it would have been in had nobody formed it. So there has to be a somewhere: a state, a mark, a position, a magnetisation, anything at all that can be in one configuration rather than another and stay that way.

It has to read what it holds, and the reading has to make a difference. Seal the thermostat's set number behind glass where the switch cannot get at it. Every part still works. The sensor senses, the number sits there at twenty degrees, the room is cold, and the boiler stays off, because nothing in the device ever consults the number for any purpose. What is left on the wall works as a thermometer and nothing more.

So the requirement is not that a record be accurate. It is that something downstream depends on it. A mark nothing consults is a mark the world runs identically without. If the world runs identically without it then it is not describing anything, whatever it happens to say.

This is also the place where one thing gets to matter to another, worth being slow about, because everything later rests on it. Two systems can sit beside each other forever, each in whatever state it is in, with their states correlating as tightly as you like, and nothing whatever passing between them. What makes the second one matter to the first is that the first reads it and does something different for having read it. Remove that and there is no sense in which anything affects anything: there are states, and there is a tally of how often they coincide, and there is nothing else.

It is also where being wrong becomes possible, which chapter two asked for and did not have to. The mark could have been otherwise, so there is more than one state the thing might be in. It gets consulted, so what the thing does depends on which. And the view is bounded, so the mark does not contain the world, and there are always two ways the world can be that leave it saying the same thing. Two worlds, one mark, one response, and in at least one of them the response is not the one that world called for. Nothing was added anywhere to arrange that. It is what a bounded thing that writes, and then consults what it wrote, cannot avoid.

That is the list. Three items, and the third is doing most of the work.

3.2 Three for three

Score the thermostat.

A bounded view. It senses temperature, at one point, in one room. It knows nothing about the pressure, the humidity, the light level, the temperature two floors up, or the weather in the next country. Its window on the world is one number wide. Passes.

Somewhere to write. Two places, in fact: the current reading and the number you set. Passes.

A reading that makes a difference. This is all it does. The whole device exists to hold two numbers next to each other, consult the pair, and throw a switch on what it finds. Nothing in it is decorative. Passes.

Three for three. It is screwed to the wall, so it holds each of its numbers from the writing of it to the reading of it, which is all the lasting the third item asks for.

Then score the thermometer beside it, because a list of requirements is worth nothing until something fails one. A mercury thermometer has a bounded view: one number, one place, nothing else. It writes, in the sense that matters, since the height of the column is a physical configuration that can be one thing rather than another, and rewrites whenever the room changes. It persists, being made largely of glass and screwed to the same wall.

Two for three. Nothing in the glass consults the column. The mercury stands where it stands and no part of the instrument is downstream of where it stands, so the height makes no difference to anything the thermometer does, there being nothing the thermometer does. Engrave a second number on the case and nothing changes, because a number on a case is not consulted either. A thermometer has a state. A thermostat has a state and an opinion about it. The gap between those two sentences is the gap between a thing and an observer.

That is a useful failure, because it shows which of the three is load-bearing. Take away the boundary and you get a duplicate. Take away the writing and nothing is retained. But take away the reading and you are left with something that looks entirely functional, sits on the wall for decades, correlates with the world to a fraction of a degree, and observes nothing whatsoever.

The thermostat passes. So does a person, who has bounded access, retained state and feedback along with a great deal more organization. Passing the same test makes both of them observers and says nothing about whether either one feels anything; chapter forty takes that question up on its own.

A sensor, a number you set and a switch meet all three in full. That is the whole of what anything needs to be an observer, thermostats and people included.

3.3 What nobody has to add

The list is short enough that the omissions need saying out loud, because people reliably try to put things back on it.

Understanding. The thermostat has no idea what temperature is, what a room is, or what it is doing. Nothing in the three mentions comprehension. Adding it would exclude every instrument in every laboratory on Earth from a theory of measurement.

Intention. The thermostat does not want the room to be warm. Something wants that, and it is not the thermostat, and the device works identically whether anybody wants anything.

A model of itself. Nothing on the list requires the record-keeper to have any representation of its own state, its own boundary, or its own existence.

Complexity. Two numbers and a comparison. That is the entire specification. Any system that meets it is on the list regardless of what else it can do.

Life. Obviously not, and this cuts both ways, because it also means that the arrival of living things in the universe did not introduce a new kind of physical entity. It introduced a much better one of a kind that had been there all along.

The list has three things on it. Anything else that gets proposed describes how good an observer is rather than whether it is one.

3.4 The worst word in physics

Which brings up the vocabulary problem. It has to be handled here because the word has been widely misread.

The word “observer” arrived in physics through relativity, where it never meant a person and was never intended to. An observer in Einstein’s sense is a coordinate frame: a convention for assigning three numbers and a time to an event. Textbooks anthropomorphise them for readability, and then have to explain that these observers are equipped with rulers laid out across all of space and clocks at every point, synchronised in advance. Nobody was confused, because everybody knew the word was shorthand for bookkeeping.

Quantum mechanics then borrowed the same word for something that had to do the measuring. The shorthand stopped being harmless. If the theory says that an observation produces a definite outcome, and “observer” sounds like it means a person, then it sounds like the theory says people produce definite outcomes by paying attention. Which is not what any version of the mathematics says, and is nonetheless what a great many people have taken away from it.

Eugene Wigner is the honourable case, because he thought it through and then changed his mind. In a 1961 essay he argued seriously that

consciousness has to enter the physical description, on the grounds that a measurement apparatus can be described as a quantum system and so can the physicist looking at it, and something has to break the chain. He later abandoned the position. What did it was solipsism: if it takes a mind to make an outcome definite, whose mind, and what were the other minds looking at meanwhile.

The confusion has a way of surviving its own refutation. Schrödinger invented the cat in 1935 as an attack. He was arguing that a theory which lets a radioactive atom smear a live animal across two states is producing something ridiculous. He used the word. The example was a weapon aimed at a position he rejected. It is the single most widely reproduced illustration of that position, printed on t-shirts by people explaining what quantum mechanics says. A joke about a theory has become the theory's best-known exhibit.

So, unambiguously, because the whole structure rests on it:

An observer here is any bounded system that keeps records. A thermostat is one. A photographic plate is one. A rock with a scratch on it is not one, and fails on the requirement people would think of last: nothing reads the scratch. Put the rock where the scratch works a lever and it joins the list. Consciousness has nothing to do with any of it. Conscious observers are a special case that appears very late, arriving without anything being added to the list of three.

3.5 Constraint-first

There is a second misreading, subtler than the first and more persistent, and the one that makes people suspicious of observer-centered physics for good reasons.

Putting observers at the bottom sounds like putting *us* at the bottom. That sounds like a demotion of the universe in favour of the people looking at it. It sounds like the world is shaped by our attention. An anthropic selection argument conditions on observers existing; it does not require their preferences to arrange the universe. The observer architecture here likewise concerns what systems can record and compare.

That is the opposite of what is being said. An observer in this book does not get what it wants. It gets what it can consistently record. The second is far more restrictive than the first. The thermostat cannot decide the room is warm. It can hold a number, and the number can be wrong, and being wrong has consequences it does not choose. Everything in the next thirty chapters comes from what a bounded record-keeper is *forbidden* to do: it cannot see everything, it cannot write without limit, it cannot compare what it has no access to, and it cannot disagree with its neighbors indefinitely without something giving way.

There is a cheap way to feel the asymmetry. Set the thermostat to twenty-five degrees in January and switch the boiler off at the wall. The device will hold the number twenty-five for as long as you care to leave it, and the room will be freezing, and nothing anywhere in the arrangement will bend. A record-keeper may write whatever it likes. What it cannot do is make the comparison come out in its favour. That gap, between what a thing may write and what it finds when it checks, is where every result in this book comes from.

Observer-first is constraint-first. Nothing anywhere in this book turns on anybody wanting anything.

3.6 What the starting point explains

A theory that starts with a spacetime can explain how its geometry changes and what matter does within it. Why that starting structure is there at all is a question it cannot reach from inside. Chapter one's examples show several ways to move the boundary between what is assumed and what is derived.

OPH begins on the observer side of that boundary. Consider the sentence: *these two records agree*. It specifies a relation between bounded record-keepers without assigning them coordinates or a distance. The patch architecture makes such comparisons precise through its states, ports and repair laws.

That gives a vocabulary for reconstruction. Agreement is the common constraint, and the chapters that follow build distance, duration and curvature out of records, their relations and their dynamics, one bridge at a time.

3.7 The bill

One thing has been quietly assumed for six pages and it comes due later.

A physical record uses memory and must be carried by a physical process. Its energy cost depends on that process. Copying a readable value into a prepared register can retain the input; erasing an unknown value removes information from the memory being reset. Under the thermal conditions developed later, that reset has a mean heat bound.

Finite memory and thermodynamic reset connect the observer description to physical resources. Counting records and counting erased bits are different operations.

That is the whole starting position: bounded record-keepers, comparing what they hold against what they see, at a cost.

Which produces the difficulty immediately. If every one of these things sees a bounded part, and there are many of them, and none of them can see what the others hold, then there is nothing in the setup that makes

their records descriptions of the same thing. Each has its own numbers. Each is right about its own window. And no two of them have ever been in a position to check.

So what makes it one world?

Chapter 4

Why Do Two Descriptions Ever Agree?

Alice saw the food truck. Bob did not.

They were walking the same street in Barcelona, on opposite pavements, thirty meters apart, conducting the standard argument about where to eat. A delivery van had stopped on the corner of Carrer de Sepúlveda with its back doors open and a man unloading crates of bottled water. From where Bob was walking the van filled the gap between two buildings completely. From where Alice was walking the van was a van, and behind it, in the gap, was a truck selling grilled sandwiches with two people queueing and a third buying two at once.

At the corner they compared accounts. Alice said there had been a food truck. Bob said there had been no food truck, an empty stretch of pavement, and a man with a great many crates of water.

Both accounts were accurate. That is the difficulty.

Alice and Bob have been doing this since February 1978, when Rivest, Shamir and Adleman needed two people to exchange a message in a paper on public-key cryptography and preferred names to letters. They have since been assigned to nearly every problem in science that turns on two parties who cannot see what the other one knows, a category that takes in more of physics than anybody expected in 1978, and in close to fifty years of continuous employment neither of them has ever been told what the message says.

The comfortable picture of what agreement means says that there is one street, that the street has a definite content, that each of them writes down what the street contains, and that if both of them do the job properly the two descriptions come out the same. Two people did the job properly and the descriptions came out different. No amount of care by either of them would have fixed it, because the van is opaque and Bob was on the wrong side of it.

The question is harder than it looks. Take two record-keepers of the kind chapter three described, each seeing a bounded part, neither able to inspect the other's records, with no third party anywhere holding both. What would it even mean for their two accounts to agree? Something has to be demanded of them, or there is no shared world. Whatever is demanded has to be something they can check from inside their own windows.

4.1 What can be demanded

Nothing can require Alice and Bob to match on the food truck. Only one of them was in a position to see it. A rule that penalised them for that would be a rule against having a point of view. Every observer in this book has one by construction.

What can be required is that they match where they both looked. Both of them saw the van. Both of them saw the corner, the traffic, the time of day, the number of lanes. That shared region is called the **overlap**, the only place where a demand can land. Outside it, two accounts differing is two windows differing, which carries no information about anybody's competence.

Even inside the overlap the two accounts will fail to be identical, worth being precise about, because it is where the naive picture does its second piece of damage. Alice wrote down that the van was on her left. Bob wrote down that the van was on his right. Those are different sentences. They are not competing claims about the van. Nothing is wrong with either of them.

What holds between them is a rule. To turn Alice's account into Bob's, swap left and right, because they were facing opposite ways. Apply the rule to Alice's whole record and you get Bob's record for everything they both saw. So the demand that survives contact with two points of view asks for a rule, and asks two things of it. That it carry one account onto the other everywhere both of them looked. And that each of them be able to work out what it is without being told.

Where the rule fails to line up, there is a leftover. If Alice has the van two car-lengths from the corner and Bob has it four, no relabeling of left and right repairs that. The leftover is a quantity. You can measure it, you can add up all the leftovers across everything two parties both looked at, and you get a single number that is zero exactly when their accounts are compatible and positive when they are not.

That number is the most important quantity in the book. Physics is what happens when something tries to make it smaller.

4.2 Two detectors

Physics takes this problem seriously enough to spend money on it. The money is the argument.

On 4 July 2012, two collaborations at CERN announced that they had found a new particle. ATLAS gave its mass as 126.0 billion electron-volts, with a statistical uncertainty of 0.4 and a systematic uncertainty of 0.4, at a significance of 5.9 standard deviations. CMS gave 125.3 billion electron-volts, give or take 0.6, at five standard deviations. Five is the bar particle

physics sets before anybody uses the word discovery. It means the odds of a fluctuation doing it are about one in three and a half million.

Those are different numbers. 126.0 is not 125.3. A reader who was told that physics had converged on a value might reasonably ask which one it converged on. Neither, is the answer. What was announced was that two intervals overlapped, and overlapping is what agreement between measurements has always meant. A measurement is a claim about a range. Two measurements agree when the ranges have points in common. The sharper the measurements, the more that agreement is worth, because the ranges are smaller and there is less room to overlap by luck.

The reason there were two of them is the whole point. ATLAS and CMS sit at different places on the same ring and were built by different people to different designs. The differences are not cosmetic. ATLAS bends particle tracks with a thin two-tesla solenoid and then bends them again in three enormous air-core toroids. CMS does it with one superconducting solenoid six meters across running at 3.8 tesla. Different magnets, different detector materials, different trigger electronics, different reconstruction software and different teams.

Each collaboration also blinded itself. The region of the data where the signal was expected to sit was masked while the analysis was being built, and unmasked only after the collaboration had committed to its procedures, so that nobody could adjust the method until the answer came out pleasing. Neither experiment tuned its analysis to the other's number.

Building the second detector cost roughly what building the first one cost, which is a remarkable sum of money to spend on a second opinion. It bought the one thing a single detector cannot produce at any price. One detector reporting 126 produces a number. Two detectors that share no components, no code and no staff, reporting 126.0 and 125.3 on the same afternoon, produce something that a number by itself is not: a fact about the world rather than a fact about an apparatus.

That is the street corner again, with a budget. Alice and Bob compared accounts on the overlap and found them compatible. The overlap of two collaborations is the particle, and the compatibility is the discovery.

4.3 A clock face

It is quarter past three in the afternoon. The clock on the wall says three. A railway timetable says 15:15. Nobody is confused, and nobody thinks the clock and the timetable disagree about what time it is, because everybody knows the rule: fifteen and three are the same thing on a clock face.

Look at what that sentence does. Fifteen and three are different numbers. They stay different numbers. The clock declares that for its purposes, two numbers count as one thing when they differ by twelve, and having made that declaration it stops carrying the difference. A clock is a machine for

forgetting the difference between three and fifteen, and forgetting it is the entire service the clock provides.

Do that systematically and you get the tool. Start with a collection of things. Choose a rule that says which of them count as the same. The rule has to be sane, in a way that takes ten seconds to check: everything counts as the same as itself; if this counts as that, then that counts as this; and if the first counts as the second and the second as the third, then the first counts as the third. Sweep up each family of things that count as one another into a single bundle. Each bundle is called an equivalence class, the collection of bundles is called a **quotient**, and the operation of passing from the original things to the bundles is the single most useful operation in the book.

Twelve bundles come out of the clock. One of them contains 3 and 15 and 27 and 39. That bundle is what "three o'clock" names. There is no such thing as the true underlying number that three o'clock secretly is. There is a bundle, and 3 and 15 are two ways of writing it down.

Which brings the two friends back. Alice's record and Bob's record differ. Where they differ by something a permitted relabeling accounts for, they are two ways of writing down one thing, in the same sense that 3 and 15 are. Where they differ by more than that, the leftover is real and somebody's records have to change.

Everything hangs on the word permitted. The clock's rule is not a free choice made for convenience. If you declared that any two numbers count as the same you would get one bundle, which forgets everything, and a clock that says the same thing at every moment. The permitted relabelings are exactly the ones that no measurement made inside the system can detect, and no others. Working out which those are is a large part of the job of physics.

4.4 The rate has to be computable in the room

One clause on all of this carries the rest of the book. It is easiest to see with money.

Two firms keep books in different currencies and share an account. To settle it they need an exchange rate. In ordinary commercial life they look one up, which works because there is a market outside both firms that publishes rates all day.

Chapter three closed off that move. There is no outside. There is no venue where the correct description is kept, no referee holding both sets of books, and any procedure that requires one has smuggled in exactly the stage that chapter said cannot be assumed. The exchange rate has to be computable from inside the room, out of things both parties hold.

It can be. The two firms have both recorded the same transaction, one of them in each currency. The ratio of those two entries is the rate. It came out of the overlap, which is the one place both parties have data. Neither firm had to be told it by anybody.

Two properties come free with a rate obtained that way. It can be undone, because if the rate one way is known then the rate the other way is its reciprocal. And two of them can be chained, because a rate from the first currency to the second, followed by a rate from the second to the third, is a rate from the first to the third.

Chaining is where it stops being bookkeeping. Take three firms, each pair of which shares an account, so there are three rates. Convert from the first currency to the second, from the second to the third, and from the third back to the first. You have gone in a circle and arrived back where you started. The amount you are holding should be the amount you set out with.

If it is not, somebody can go round the loop repeatedly and end up with more money than they started with, having bought and sold nothing. Currency traders call this triangular arbitrage, they have machines watching for it, and when it opens it closes in milliseconds. The condition that closes it is the requirement that going around a loop of translations brings you back to yourself.

That condition is the whole content of what physicists call gauge structure. Accountants make it obvious and field theory makes it look hard. It is doing the work behind electromagnetism twenty-two chapters from here. Descriptions related by a permitted relabeling say the same thing. The relabelings can be undone and chained. And every closed loop of them has to come back to the identity, since otherwise a round trip would produce something from nothing.

4.5 Ninety meters

In June 1792 two French astronomers left Paris in opposite directions to measure the world.

France wanted a unit of length that no king had defined. The Academy of Sciences had settled on one ten-millionth of the distance from the north pole to the equator. To get it, somebody had to survey the meridian arc from Dunkirk to Barcelona by triangulation and scale up. Jean-Baptiste Delambre took the northern leg, Dunkirk down to Rodez. Pierre Méchain took the south, Rodez to Barcelona. The plan allowed a year. The survey took seven, most of them conducted through a revolution and a war, with both men repeatedly arrested as suspected spies.

Méchain finished his southern latitude work at the fortress on Montjuïc, the hill above Barcelona harbour, and had earlier measured the latitude at his lodgings in the city. Both were careful measurements of the same

quantity by the same excellent astronomer with the same instrument. In March 1794 he compared them and they disagreed by three seconds of arc.

Three seconds of arc is about a six-hundredth of the width of the full moon. On the ground, it is a little over ninety meters. It is a very small number, and a fatal one, because the two figures were both his and there was no third figure to adjudicate. War with Spain had closed the border behind him and he could not get back to the hill to remeasure.

He concealed it. When the international commission met in Paris in 1799 to fix the meter, the numbers Méchain supplied had been adjusted to remove the discrepancy. He spent the rest of his life trying to make the disagreement go away, went back into Spain in 1803 to redo the work, contracted yellow fever, and died at Castellón de la Plana on 20 September 1804 at the age of sixty. Delambre, preparing the official record of the survey afterwards, worked through the notebooks and found the alterations.

Set aside the concealment, which belongs to Méchain. The structure underneath it is the one the food truck had. Two records of the same quantity failed to agree on their overlap, the leftover would not go to zero, and the one move that could have driven it to zero, going back to the hill with the instrument, had been made unavailable. He was in the position of Alice and Bob with the van permanently in the way. A mismatch that cannot be repaired does not resolve itself and does not fade.

4.6 What the meter is

The meter they produced is wrong.

The quadrant they were measuring, from pole to equator, is 10,002,290 meters. They were trying to make it exactly ten million. Their meter is therefore short of its own definition by about a fifth of a millimeter, roughly the thickness of two sheets of paper. The error has nothing to do with Méchain's three seconds of arc. It comes from the flattening of the Earth, which they had to estimate and which was not then known well enough.

Nothing whatever has gone wrong.

Every meter stick in the world was made to match the standard, and every meter stick in the world therefore matches every other one. The definition was replaced in 1889 with a platinum-iridium bar, replaced in 1960 with a count of 1,650,763.73 wavelengths of the orange-red light of krypton-86, and replaced again in 1983 with the distance light travels in one 299,792,458th of a second. Through three replacements the meter has kept the same length.

It has never changed length because the meter was never the bar in the vault. The meter is the bundle. It is the equivalence class of every stick, wavelength and timing that counts as agreeing with every other one. A definition is a way of pointing at the bundle by holding up one member of

it. Point at it with a bar, point at it with krypton, point at it with the speed of light and a clock, and you have pointed at the same bundle three times. A better representative buys precision. It has no power to change what is being represented, because it was never carrying the identity in the first place.

This is why the survey's error is not embarrassing and why French commerce did not collapse in 1799. The unit's job is to let two parties translate their measurements into each other's terms. It does that job through what everybody adopts, which fixes the translation. Nobody needs to go back to the pole.

4.7 One world

The naive picture of agreement wanted one street with a definite content and two matching descriptions of it. A delivery van was enough to take it apart. There is no master copy that anybody holds a copy of. There are records held by parties who cannot see each other's, and there are overlaps, and on the overlaps there are translations, and the translations are constrained: they must be computable from inside, they must be undoable, they must chain, and every closed loop of them must come back to where it started.

What makes it one world is that the leftovers on the overlaps go to zero. That is the only sense in which Alice and Bob live in the same city, the only sense in which ATLAS and CMS found the same particle, and the sense that failed for Pierre Méchain on a hill above Barcelona.

Which raises the question the next chapter exists to answer. Alice and Bob compared notes and their accounts were compatible. Compatible accounts are two labelings of one thing, in the way that 3 and 15 are two labelings of one bundle. The labeling is thrown away by the comparison.

So something survived the comparison. It was not either of their descriptions.

They agree. On what?

Chapter 5

Why Do You and I See the Same Table?

Bob's kitchen table is oak, scratched, and a hundred and twelve centimeters along its long edge. At two in the morning he crosses the kitchen in the dark to get a glass of water. He does it by memory and by hand: four steps from the doorway, the cold edge of the worktop under his left palm, then the corner of the table, which he takes wide because he has caught his hip on it before. He never sees it. Nothing reaches his eyes at all.

Alice, coming in behind him, turns on the light and sees the whole thing at once.

Two accounts of the table. The physical signals behind them have nothing in common. Alice receives photons, scattered off the oak in a band of wavelengths a few hundred nanometers wide, focused onto a patch of light-sensitive cells at the back of each eye. Bob receives mechanical deformation of the skin of his palm, transduced by pressure receptors that would not respond to a laser shone directly at them. No signal either of them receives is available to the other. The two channels do not overlap anywhere in the physics.

They agree about the table instantly and without discussion. Neither of them experiences the agreement as an achievement.

Chapter four left them agreeing and did not say what they were agreeing about. The comparison threw away both descriptions. Something came through it anyway.

5.1 What survives

Start with what does not survive.

The oak is brown. That belongs to Alice. Bob has no version of it, because brown is what her visual system calls a certain distribution of wavelengths, and no amount of running your hand along a table delivers it. The oak is slightly waxy, and cooler near the window. That belongs to Bob. Neither of them is wrong, and neither of them can hand their version across.

What both of them have is that the edge runs straight for a hundred and twelve centimeters and then turns through a right angle. That the far corner stands a hundred and twelve by seventy from the near one. That

the surface is rigid, so the distance between any two points on it stays what it was while you walk around it. That there is a hard flat obstruction between the doorway and the window, and that walking into the space it occupies will hurt.

Every item on that list is a relation between one part of the table and another. None of them is a property of the table on its own. The rigidity is a statement about pairs of points, the right angle is a statement about two edges, and the hundred and twelve centimeters is a comparison with a metal bar in a vault outside Paris, or with whatever has since taken over that bar's job. Two channels with no physics in common delivered the relations and dropped everything else.

A quantity that comes out the same under a change in how you look is called an **invariant** of that change.

The order matters here. It is the order most treatments get backwards. The invariant comes first. You find out what survives, and only then do you have any business asking what it survived.

5.2 A square and a pencil

Take a sheet of paper and draw a square on it, big enough to handle, five or six centimeters. Mark one corner with a pencil dot, and mark the same corner on the reverse side so you can find it after the square has been face down.

Cut it out. Then find every way to pick it up and put it back down so that the square occupies exactly the same patch of paper it did before. The outline has to land on the outline. The pencil dot may end up anywhere.

Most people find four and stop. The four are the rotations: leave it alone, turn it a quarter turn, turn it a half turn, turn it three quarters. Then they remember they are allowed to turn it over. There are four more: one flip about the vertical line through the middle, one about the horizontal line, and one about each of the two diagonals.

Eight. There are exactly eight. Three things about those eight can be checked with the paper in your hand.

Do one move, then do another. Whatever you did, the result is one of the eight. A quarter turn counterclockwise followed by a flip about the diagonal that runs up to the right is a flip about the horizontal line. You can watch the dot arrive to confirm it. There was nothing else it could have been, because the square is back on its outline, with only eight ways to be back on your outline. The set is closed. Nothing you can do with these moves takes you outside them.

Every move can be undone by another one on the list. The quarter turn is undone by the three-quarter turn. Each of the four flips is undone by itself. No move in the set strands you.

Leaving it alone is on the list, too. It looks like a formality, but without it, the result of "undo the quarter turn" has no name.

Checking the closure by hand means checking sixty-four pairs of moves. An eight-by-eight grid holds them all at once, the eight moves along the top and the same eight down the left edge. Each cell answers one question: do the move on the left, then the move along the top, and write down which of the eight you have ended up with. Sixty of the sixty-four are filled in below.

Eight short names first, to avoid writing "the flip about the diagonal that runs up to the right" sixty-four times. Each rotation gets the number of quarter turns it is, counted counterclockwise: 0 for leaving the square alone, 1 for the quarter turn, 2 for the half turn, 3 for three quarters. Each flip gets the line it is a flip about: V for the vertical line down the middle, H for the horizontal line across it, S for the diagonal that leans like a slash, running from the bottom left corner up to the top right, and B for the diagonal leaning the other way. The four lines belong to the table rather than to the square, so they stay put while the square moves. Rows are the first move, columns the second.

then	0	1	2	3	V	H	S	B
0	0	1	2	3	V	H	S	B
1	1	2	3	0	S		H	V
2	2	3	0	1	H	V	B	S
3	3	0	1		B	S	V	H
V	V	B	H	S	0		3	1
H	H		V	B	2	0	1	3
S	S	V	B	H	1	3	0	2
B	B	H	S	V	3	1	2	0

Four cells are empty. Pick the square up and fill them in.

Two of the four hold the same pair of moves in opposite orders. Row 1 under column H is a quarter turn and then a flip about the horizontal line. Row H under column 1 is the flip first and the quarter turn after. Both answers are diagonal flips. They are flips about different diagonals, and the pencil dot settles which is which: one order leaves the dot in the corner it started from, and the other sends it to the corner diagonally opposite.

The other two blanks are quieter. Row 3 under column 3 is a three-quarter turn done twice: six quarter turns, which is a half turn. Row V under column H is a flip about the vertical line followed by a flip about the horizontal. The square, turned over twice and lying face up again, comes back rotated through a half turn. Two mirrors make a rotation.

Count along any row of the finished grid. Each of the eight moves appears exactly once, and the same holds down every column. The row

for the quarter turn shows why. Two different second moves cannot land the square in the same place there, because a three-quarter turn stuck in front of both sequences cancels the quarter turn and leaves the second move standing on its own, so those two second moves would have to have been one move. Eight second moves, eight different answers, and only eight moves available for the answers to be. All eight are in the row, once each, and nothing anywhere in the grid produces a ninth thing.

Closed, undoable, and containing the do-nothing move. A collection of changes with those properties is called a **group**. Mathematicians have been calling it that since the 1830s. The word arrives last on purpose. The object was in your hands before it had a name. It would have been there if nobody had ever named it.

The three properties fit on one line of symbols. Write G for the collection of the eight moves, g and h for any two moves in it, and a dot between them for doing g and then doing h. Write e for leaving the square alone, and g with a raised minus one for the move that undoes g. The rounded E is read as "is one of". The conditions come in the order closure, do-nothing move, undo, because the undo is defined by what it gets you back to.

$$g \cdot h \in G \qquad e \cdot g = g \cdot e = g \qquad g \cdot g^{-1} = g^{-1} \cdot g = e$$

The first says the answer to any pair of moves is on the list. The second says that e changes nothing whichever side of another move it sits on, and the third that every move has an undo which works from either side. The do-nothing move is called the **identity**, the undo is the **inverse**, and one further condition is usually listed alongside them and costs nothing here: with three moves in a row, grouping the first two or the last two makes no difference, because the square gets picked up three times in the same order either way.

A group is a set with one operation obeying those conditions. Nothing in them mentions paper. The eight moves of the square are the **dihedral group of order eight**, order eight for the number of moves in it and dihedral from the Greek for two-faced, which here is a square with a front and a back. The rotations of an ammonia molecule satisfy the same three conditions, and so does every way of reordering a deck of cards, so anything proved from those conditions alone holds for the molecule, the deck and the paper square without being proved three times.

One more thing to see with the paper. The dot moves. Under the eight moves it visits all four corners, every corner reachable from every other. The set of places a thing can be sent by the moves in a group is its **orbit**, so the orbit of a corner is the four corners. The orbit of the midpoint of an edge is the four edge midpoints, a different set of four. No move carries a corner to an edge midpoint.

The center of the square has an orbit of one. It sits where it sits under every one of the eight. Nothing you can do to the square touches it.

That is an invariant, in your hand, on a scrap of paper. Something the moves leave alone.

5.3 Chapter four, from the other side

Look again at the three conditions.

Chapter four asked what has to hold between two record-keepers before they can be said to share a world. It came back with conditions on the translations between their records. A translation had to be computable from inside, using only what the observer holds. It had to be undoable, so that going from Alice's account to Bob's and back returns Alice's account. Translations had to chain, so that a route through a third party gives the same answer as the direct one. And every closed loop of them had to come back where it started.

Those are the conditions on the paper square. Undoable is the inverse. Chaining is closure. Coming back where you started is the do-nothing move sitting at the end of every loop.

The overlap rule of chapter four and the symmetry of a square are one thing seen from two directions. The translations between observers form a group, in the way the moves of a square do. The question "what do Alice and Bob agree about" is the question "what does that group leave alone".

Which gives the sentence the rest of this book depends on.

A symmetry is a change that leaves a named invariant alone, so a symmetry is the claim that part of your description was never physical.

Brown is not physical in this exact sense: it is what changes when you switch from Alice's channel to Bob's. The world does not change when you make that switch. The distance from corner to corner does not change. That one is the table.

5.4 Nought degrees

The clearest case of a quantity everybody treated as a fact about the world, until it was subtracted, is the one printed on every map.

Latitude has an origin the planet supplies. The equator is where the spin axis says it is. Any surveyor anywhere can find it without asking permission. Longitude has no such thing. The Earth is very close to symmetric about its spin axis, which means that turning the whole planet through some angle about that axis leaves every measurable relation between places exactly as it was. Rotating everything is a symmetry, so the number you attach to a meridian is a labeling. No experiment can read it off the ground.

For three centuries this was treated as an unsettled question rather than an empty one. The French measured from Paris, the Spanish from Cádiz

or Tenerife, the Americans from Washington, and captains carried tables for converting between them. In October 1884 forty-one delegates from twenty-six nations sat down in Washington to fix it, and on the thirteenth they voted: twenty-two in favour of Greenwich, one against, which was San Domingo, and two abstentions, which were France and Brazil. France went on printing charts from Paris for another quarter of a century.

What the conference settled was a convention, and everybody in the room knew it. The tell is in the mechanism, because there is no other way to settle it. You cannot measure which meridian is the real zero, and you cannot be wrong about it either. After the vote, two charts could be compared without conversion arithmetic.

Distances between places survived the vote unchanged. Distance is the invariant, and the labeling was never physical.

Every symmetry anybody has found in physics is that same operation run on a bigger description. Here is a change you can make to how the world is written down. Here is what survives it. Whatever did not survive was never a fact about the world.

5.5 Vienna, orbit, and the Cretaceous

There is a symmetry so large that almost nobody notices they are assuming it.

A physicist in Vienna measures how fast a ball falls. A physicist on the International Space Station, four hundred kilometers up and moving at seven and a half kilometers a second, does the same experiment in a way that accounts for the local conditions. They get different readings, because the conditions differ, and the same law relating the readings to the conditions.

Do it again a hundred and fifty kilometers to the west. Same law. Do it in the Cretaceous: same law, and a version of this can actually be checked, because light arriving from a quasar eleven billion years old carries the spectral fingerprint of hydrogen behaving exactly as hydrogen behaves in a lamp in a basement in Vienna.

The laws do not depend on where you are. They do not depend on when you are. Those are two symmetries. Shifting everything sideways in space leaves the laws alone, and shifting everything along in time leaves the laws alone. Galileo's cabin is a third change of the same kind: put the whole laboratory into smooth motion and the laws do not notice that either, which is why the flies stayed where they were.

Strip them out and watch what happens to the idea of an experiment. If the law of falling depended on where you stood, a result obtained in Vienna would say nothing about a ball in Graz, and the Graz result would say nothing about the next street. If it depended on when, a measurement made on Tuesday would say nothing about Wednesday, which is to say it

would say nothing at all, because a measurement is finished before anybody can use it. Repeating an experiment would be pointless. The word “law” would have no work to do.

Every experimental result anybody has ever quoted is leaning on those two symmetries. They are the reason a number obtained once is worth writing down.

And there’s more to it. Emmy Noether proved in 1918 that a continuous symmetry of the laws always comes with a conserved quantity, exactly one, determined by the symmetry. Shifting in space gives you a quantity that never changes: momentum. Shifting in time gives you another: energy.

Momentum is conserved because the laws do not care where you are. Energy is conserved because they do not care when. Those are two statements of one fact, about two directions. The procedure that turns a symmetry into its conserved quantity is exact and mechanical. Running it takes more mathematics than a paper square supplies.

5.6 The letter to Chevalier

The paper square is the whole of what Évariste Galois was working on, run on something harder.

He was asking which polynomial equations can be solved by a formula built from the coefficients using addition, multiplication and the extraction of roots. The quadratic has such a formula, which every schoolchild is made to memorize. The cubic and the quartic have them, found in Bologna in the 1500s, ugly and real. The fifth degree had resisted for two hundred and fifty years.

Galois stopped looking at the equation and looked instead at the symmetries of its solutions: the ways you can permute the roots among themselves without disturbing any relation between them that the coefficients can see. Those permutations form a group, in exactly the sense the square’s eight moves do. He then showed that whether a formula exists is a property of that group, and that from the fifth degree onward the group is the wrong shape.

The submissions went badly. Cauchy took the first memoir in 1829 and it never appeared. Fourier took the second in 1830, for the Academy’s Grand Prix, and died in May of that year with the manuscript among his papers, where it was not found. Poisson took the third in 1831 and reported in July that the reasoning was neither sufficiently clear nor sufficiently developed for the Academy to judge it.

On the night of 29 May 1832 Galois wrote to his friend Auguste Chevalier, setting out the results and asking him to make a public request: that Jacobi or Gauss be asked to give an opinion, not on whether the theorems were true, but on whether they were important. In the margin of one page,

beside an argument he had not finished, he wrote that there was something to complete in the demonstration and that he did not have the time.

He was shot in a duel the following morning and died on 31 May, aged twenty. The letter was published in 1846. What was in it is taught to undergraduates in their first year. It is why the eight moves of a piece of paper have a name.

5.7 The size of an atom

A notion that applies to everything explains nothing. Symmetry has a boundary, findable from where you are sitting.

Shift everything in space: symmetry. Shift everything in time: symmetry. Rotate everything: symmetry. Change the size of everything, so that every length in the universe is doubled and nothing else about the arrangement is altered: the world notices.

The evidence is in the room. Atoms have a definite size. A hydrogen atom is about one ångström across, a ten-billionth of a meter. That number is the same in Vienna, in orbit and in the Cretaceous. It is a length built into the physics. A world without scale symmetry is exactly a world in which such a length can exist. If doubling every length left the laws alone, no size could be preferred, and nothing would distinguish the atom you have from one twice as big. Atoms would come in every size, and chemistry, which depends on atoms of a given element being interchangeable, would not be available.

The consequences are structural rather than decorative. Galileo worked one of them out in 1638, in the *Two New Sciences*, and illustrated it with a drawing of a bone. Scale an animal up by a factor of ten in every direction and its weight goes up by a factor of a thousand, while the cross-section of the bones that carry the weight goes up by a hundred. The scaled animal is ten times worse off than the original, and its legs snap. For the same reason a bridge cannot be designed by scaling up a small one. Every engineer who has tested a model in a wind tunnel has had to do arithmetic to correct for the fact that the model is not a small copy of the world. That arithmetic is a bill for a symmetry the world does not have.

So symmetry is a claim with content. The way you can tell is that it is false about scale.

5.8 What the table was

Alice and Bob agree about the table because there is a change, from her channel to his, that leaves something alone, and what it leaves alone is the geometry: the edges, the right angle, the hundred and twelve centimeters, the rigidity. Everything else in either account belongs to the observer that produced it.

That is the same operation as the paper square, where the eight moves leave the center alone. It is the same operation as chapter four's overlaps, where the translations between records leave the shared content alone. It is the operation physics has been running since the 1830s, and each time it has been run, something everybody had taken for a property of the world has turned out to be a property of the description, and has been subtracted.

There is a question left over, and Alice and Bob cannot answer it between them.

Put a third person in the room. Alice agrees with Bob. Bob agrees with Charlie. Charlie agrees with Alice. Every pair compares records, finds a translation, and comes away satisfied. Each pair shares one invariant, and each pair is content.

That has the sound of a settled matter. Three coins in three separate rooms are enough to take it apart.

Chapter 6

Why Isn't Agreeing With Your Neighbors Enough?

Give Alice, Bob and Charlie a coin each and put them in three separate rooms. Each of them looks at their own coin and sees heads or tails. Nobody can see anybody else's.

They are allowed to meet in the corridor two at a time. When a pair meets they compare coins and write down one word: **same**, if the two coins match, or **different**, if they do not. The only restriction is that the pair must not write down which way either coin is facing. What gets recorded is the relation between two coins, which is the only thing either of them could check together.

Three meetings happen. Alice and Bob meet, and record different. Bob and Charlie meet, and record different. Charlie and Alice meet, and record different.

Set the coins.

Suppose Alice's is heads. Bob's has to be tails, because they recorded different. Charlie's has to be heads, because Bob and Charlie recorded different. And Charlie and Alice recorded different, so Charlie's has to be tails. That contradicts the step before.

Try the other branch. Suppose Alice's is tails. Bob's is heads, Charlie's is tails, and Charlie and Alice are both tails, having recorded that they differ.

That is all the branches there are. There are eight ways to set three coins. Every one of them contradicts at least one of the three records.

The obvious response is that somebody is wrong. Three coins have three faces. Records that no arrangement of faces can satisfy are records with a mistake in them, so the thing to do is find out whose. Look again at what the eight cases ruled out. Not the three records by themselves: the three records together with the assumption that there were three settled faces for them to be records of. The check kills the pair. It does not say which half to drop.

Nobody lied. No coin was misread. Each of the three meetings produced a record that is perfectly satisfiable on its own. Any two of the three can be satisfied together without the slightest difficulty. It is the third one that

has nowhere to go. Which of the three you call the third one is your choice, because they are symmetric and none of them is the culprit.

There is nothing the three of them can do about it. They can meet in pairs for the rest of their lives. Every meeting will confirm what the previous meeting found. No pairwise comparison anywhere in the arrangement contains the information that the three records are jointly impossible, because that information is not held by any pair. The three of them can go on being individually correct indefinitely.

Chapter five put a third person in the room and left the three of them agreeing. This is the bill.

The trap is older than the corridor. In 1785 the Marquis de Condorcet published an essay on applying analysis to decisions reached by majority vote, and buried in it is an observation about committees.

Take three people and three options. The first person ranks them A, B, C. The second ranks them B, C, A. The third ranks them C, A, B. Ask the committee to choose between A and B, and A wins by two votes to one. Ask it to choose between B and C, and B wins two to one. Ask it to choose between C and A, and C wins two to one.

Every one of those votes is a real majority. Nobody was manipulated, nobody abstained, and each vote can be repeated as often as you like with the same result. There is no winner, and there is no winner in a way that no amount of further voting will fix, because the information that the three results are jointly impossible is not present in any of them.

Condorcet's committee was the first warning that pairwise agreement is a weaker thing than it sounds. Local data can be flawless, checkable and reproducible, and refuse to be assembled into anything. The voting version has no arithmetic attached to it that helps much. The version with coins comes with arithmetic.

6.1 "Same same but different" as arithmetic

The coin argument is short enough to check by hand, but doing things by hand doesn't scale very well here. Twelve people generate sixty-six possible pairings. What is needed is a way of computing the answer.

Write **0** for same and **1** for different.

Then chain two records together. If Alice and Bob are the same, and Bob and Charlie are the same, Alice and Charlie are the same: 0 and 0 give 0. If Alice and Bob are the same and Bob and Charlie differ, then Alice and Charlie differ: 0 and 1 give 1. And if Alice and Bob differ and Bob and Charlie differ, then Alice and Charlie are back to being the same, because two flips return you to where you started: 1 and 1 give 0.

Ordinary addition says one plus one is two. Here there is no two, because there are only two states in the world being described. The count that

matters is whether the number of flips is even or odd. So one plus one is zero.

This is called **arithmetic modulo 2**. It is the arithmetic of light switches: flick the switch twice and the room is as you found it. It is the arithmetic of turning a sock inside out. Every fact in it fits on two lines, the whole of what the coins require.

Take the three records again. Different, different, different: 1, 1, 1. Add them. One and one is zero, and zero and one is one.

The three records around the loop sum to 1.

For the coins to be settable at all, that sum has to be 0, because going all the way around the loop returns you to Alice's own coin, and Alice's coin does not differ from itself. An odd number of flips around a closed path is a demand that something be different from itself. The coins cannot arrange that.

Eight cases checked by hand, replaced by adding three ones.

6.2 Loops

The picture generalizes as soon as it is drawn.

Put a dot for each observer and a line between two dots whenever those two have compared records. A collection of dots and lines like that is a **graph**, one of the few pieces of mathematics that looks exactly like the thing it is about. The dots are called vertices, the lines edges, and two dots joined by a line are neighbors.

Walk along the edges from dot to dot and you have a **path**. Come back to the dot you started from without retracing your steps and you have a **cycle**. The triangle of Alice, Bob and Charlie is the smallest cycle there is.

Every edge carries a record, 0 or 1. Every cycle therefore carries a total, which is the sum of the records around it, computed in the arithmetic that has no two in it. The quantity you pick up by going all the way around a loop and arriving back where you started is called the **holonomy** of that loop.

The condition on the whole arrangement fits in one sentence.

A consistent global assignment exists exactly when every cycle has holonomy zero.

One direction of that is the coin argument again: if the coins can be set, then walking around any loop and coming home must find the coin you left, so the flips have to cancel.

The other direction is constructive: Suppose every loop sums to zero. Pick any dot and set its coin however you like. Walk outward. Every time you cross an edge you know whether to keep the value or flip it, because the edge says same or different, so every dot you can reach gets a value. The only way this can go wrong is if two different routes to the same dot

disagree. Two routes to the same dot form a loop. Every loop sums to zero. So they cannot disagree. The assignment is forced, everywhere, from one free choice at the start.

A route through the dots that reaches all of them without ever closing a loop is called a **spanning tree**, the skeleton of that construction. For any connected graph, a spanning tree has exactly one edge fewer than it has dots. Every remaining edge, the ones the tree did not use, closes exactly one independent loop when you put it back.

Which gives you the count of conditions, from nothing but arithmetic. Take the number of edges, subtract the number of dots, add one. That is how many independent loops the graph has, and therefore how many separate sums have to come out zero. The triangle has three edges and three dots: three minus three plus one is one. One loop, one condition, and it failed.

6.3 Ascending and descending

Roger Penrose went to the International Congress of Mathematicians in Amsterdam in 1954, where there was an exhibition of prints by M. C. Escher, and came away preoccupied. He and his father Lionel, a psychiatrist, spent some time afterward constructing figures that behave locally like ordinary drawings and cannot be assembled into an object, and published them in the *British Journal of Psychology* in 1958. One of them was a staircase.

Every flight goes up. Take any corner of the drawing, cover the rest, and there is nothing to object to: a step, then another step, each one higher than the last. Follow all four flights around and you arrive back at the landing you started from, having climbed the whole way.

The Penroses sent a copy to Escher, who put the staircase into a lithograph in March 1960 and populated it with a file of monks trudging around the roof of a monastery forever.

The staircase is the coin problem drawn. Each edge of the loop carries a record, in this case "up", every local record is fine, and the sum around the loop is not zero. What is impossible is not any of the steps. It is the closing.

6.4 A magnet that cannot decide

Physics met this in the 1950s and gave it a name.

A magnet is many small magnetic moments, each able to point one of two ways, each pushing its neighbors around. In an antiferromagnet, every pair wants to be opposite: if this one points up, the one beside it should point down. On a square grid that is easy to arrange: the result is the tidy alternating pattern of a chessboard.

Put the same rule on a triangular grid and it comes apart at the first triangle. Set the first moment up and the second down, which satisfies that pair. The third has to be opposite to both. There is no third direction. Whatever it does, one of its two bonds is unhappy, with no principle available to say which one.

Gregory Wannier worked out the consequences for a whole triangular lattice in 1950 and found something unexpected: the system keeps a finite amount of disorder as the temperature goes to absolute zero. Ordinarily, cooling something to zero freezes it into its one best arrangement. Here there is no one best arrangement, only an enormous number of equally good ones. The system goes on having a choice after the energy has been taken away.

Gerard Toulouse gave the phenomenon its name in 1977, while working on spin glasses, and gave it the same test the coins gave it. Go around a closed loop of bonds and multiply out what each one demands. If the loop comes back consistent, the region can settle. If it does not, the loop is **frustrated**. No arrangement of the moments inside it satisfies every bond, however long you wait.

Frustration is the coin problem in a material, and a measurable one. It changes the heat capacity, it shows up in neutron scattering, and it is the reason a class of magnets never orders properly no matter how carefully they are cooled. The obstruction lives in the loops, exactly where the arithmetic said it would.

6.5 Twelve patches

The triangle is small enough to be dismissed as a puzzle. So take the smallest thing that deserves to be called a universe and check every state of it.

Twelve observers. Wire them as the corners of an icosahedron, the twenty-sided solid, so that each observer has exactly five neighbors and there are thirty edges in all. Each edge is an overlap in chapter four's sense: two observers with something they can both check, and no way to reach each other except by checking it. Alice and Bob's overlap was the stretch of Barcelona street they could both see. Here it is one comparison, which is as small as an overlap can get and still be one. Twelve corners, thirty edges. The count above says the graph has thirty minus twelve plus one, which is nineteen independent loops, so nineteen conditions have to hold at once for the thing to be settable.

Each observer holds one bit. That is four thousand and ninety-six arrangements. Flipping all twelve at once changes no comparison anywhere, so there are two thousand and forty-eight distinct states.

Each of the thirty overlaps holds a record of the kind the corridor produced: same or different, the relation between two bits and nothing about

either one. The thirty used here are a set whose nineteen loop sums all come out zero, so by the argument above the bits can satisfy them.

Two thousand and forty-eight is a number you can exhaust. So exhaust it.

First the machine needs a move. The one used here is the spanning tree of two sections ago, run as a procedure: take eleven of the thirty edges, enough to reach all twelve observers without closing a loop, and root them at one observer. An observer repairs by copying its parent's bit through the record on the edge between them. That is the whole move. The remaining nineteen edges are never repaired against. They are the conditions being checked.

That move is a demonstration engine rather than a law. Nothing so far says repair has to work this way, and chapter nine is where the move an observer actually makes gets derived from what a repair is allowed to cost. What this one buys is a machine small enough to enumerate.

It also stops. Repairs run outward from the root. An observer that has just been fixed can be unsettled again if its parent changes later, but its parent sits nearer the root, and nothing unsettles the root at all, so agreement sets from the root outward and every run runs out of work. A run can only stop when every observer agrees with its parent, and with the root pinned that leaves exactly one assignment.

Start in every one of the two thousand and forty-eight states and repair until nothing is left to do. That is one **run**: one starting state, one order of working through the observers, carried to a stop. Do it in every order. Sixteen schedules apiece, which for observers with no clock, no leader and no queue are sixteen different worlds as far as anyone inside can tell. Counted over every state, the machine has eleven thousand two hundred and sixty-four repairs available to it.

Two thousand and forty-eight states, sixteen schedules apiece, and eleven thousand two hundred and sixty-four repairs is not a sample. It is the whole space. Nothing here rests on a well-chosen starting configuration or a lucky order, because there was no choosing and no luck: every configuration this universe admits was started, and every order these observers could repair in was run.

Two results come out. The first is the boring one that has to be true before the second means anything.

Exactly one of the two thousand and forty-eight states is globally consistent. That one is arithmetic. The spanning tree argument already settled it: the nineteen loop sums are zero, so fix one observer's bit and every other bit is forced, which leaves one free choice, two assignments, and one distinct state once the all-over flip is divided out.

The second result is the machine's. Every single run lands on that state. Every starting point, every schedule, every order of repair. The answer does not depend on who went first. This machine's indifference is bought

by the score, which falls whichever order you work in. Whether a repair rule has to behave that way in general is chapter ten's subject. The leftover at the end is zero, which is to say the observers end up in complete agreement and no observer's position decided it.

6.6 One edge

Flip a single edge, one of the nineteen the tree did not use. Change one record, out of thirty, from same to different. Change nothing else. Every observer has five neighbors, every overlap works, and the machine is untouched.

Some loop now sums to 1. How many of the nineteen do depends on which nineteen you drew, since there is more than one way to pick them. What the choice does not affect is that they no longer all close.

Of the two thousand and forty-eight states, the number that are globally consistent is zero. There is no way to set twelve bits that satisfies all thirty records. No amount of repair produces one, because the obstruction is a property of the loops rather than of the bits.

What happens instead is the interesting part. The machine does not thrash, fail to halt, or give different answers on different days. It converges, exactly as before, from every one of the two thousand and forty-eight starting states and under every one of the sixteen schedules, onto one terminal arrangement.

That arrangement has a leftover of exactly one.

Every repair the observers can perform has been performed. Every disagreement that could be pushed somewhere else has been pushed somewhere else. And when the pushing stops, one unit of disagreement sits in the system. It cannot be removed by working harder, it cannot be removed by choosing a better order, and it cannot be pushed off the edge of the world, because a system of twelve observers wired into a closed surface has no edge to push it off.

You can move it. Repair in a different order and it sits somewhere else. What you cannot do is get rid of it. The total is one wherever it happens to be sitting.

6.7 The leftover

Look at what that leftover does.

It has a definite quantity, exactly one, not approximately one and not one on average. It has a location, in the sense that the loops carrying holonomy tell you where it is. It survives every process the world can apply to it. It is not a property of any single observer, because every observer is behaving correctly and repairing whatever it finds. And it is not a property of any pair, because every pair is satisfiable and always was.

It is a property of the way the whole thing is wired. It is as real as anything else in the arrangement, in the only sense of real that has been available since chapter four: it is what survives comparison between every point of view.

A conserved quantity, indivisible, located, indifferent to how you approach it, that no amount of local work will destroy.

The word for that costs twenty more chapters to earn, so it stays unsaid.

6.8 The order of the meetings

Alice, Bob and Charlie met in a corridor, two at a time, and the account above knows which meetings happened before which. The twelve observers ran sixteen schedules. Calling them schedules is a claim that there is an order to run things in. Every one of those sentences assumed that comparisons are events, that events happen one after another, and that there is a fact about which.

Take that away.

Three observers with no clock, no shared present, and no way of telling a neighbor who has gone quiet from one that has stopped existing.

They have to agree anyway.

Chapter 7

Why Is There No Such Thing As Now?

On 4 October 1971 two men boarded an eastbound commercial flight out of Washington with tickets made out for a party of four. Two of the four were listed as Mr. Clock.

Mr. Clock was four cesium-beam atomic clocks, Hewlett-Packard model 5061A, each about the size of a large suitcase and too delicate for the hold. Joseph Hafele was a physicist at Washington University in St. Louis who had worked out, while preparing a lecture in 1969, that an ordinary airliner and an ordinary atomic clock were enough between them to measure something people had been arguing about since 1905. He spent the following year failing to find anybody who would pay for it. After a talk he gave in 1970, Richard Keating, an astronomer at the United States Naval Observatory, came up and mentioned that he worked with atomic clocks for a living and that his employer owned several. The whole thing cost eight thousand dollars, of which seven thousand six hundred went on economy fares.

They flew east around the world, came back, and flew west from 13 to 17 October. Then they carried the clocks back into the Naval Observatory and compared them against the clocks that had never left the bench.

Every clock in the building had been running the entire time. None of them agreed.

Going east, the flying clocks had lost 59 nanoseconds against the ones on the ground, give or take 10, where the prediction had been a loss of 40 give or take 23. Going west they had gained 273 nanoseconds, give or take 7, against a prediction of 275 give or take 21. Both papers went into *Science* in July 1972, one titled "Around-the-World Atomic Clocks: Predicted Relativistic Time Gains" and the other, published immediately after it by the same two authors, "Around-the-World Atomic Clocks: Observed Relativistic Time Gains".

Look at what those numbers are attached to. Each came out of subtracting one reading from another in one room, at the end, with the clocks side by side.

That is not the only comparison available, and why the others do not help is the whole of the point. Nothing stops you comparing clocks at a distance. Radio does it. A navigation satellite does nothing else all day:

the plane could have sent its reading down and the observatory could have written down what arrived.

The trouble is that the message takes time. The reading arrives stale, and correcting for that means knowing how long the trip took. You can time a round trip without difficulty: send a pulse up, wait for it to come back, read your own clock twice. Splitting that total into the trip out and the trip back is where it stops. Calling the two halves equal is the usual move and a perfectly good working rule, but nothing measures it, and checking it would mean comparing two clocks that are far apart, which is the thing you were trying to do.

What did the flying clock read at the instant the bench clock read noon on the second day out? Every route to an answer goes through an assumption somebody made rather than a measurement somebody took.

Everyone carries a picture around without inspecting it. There is a single present moment, everything in the universe is doing something in it, and clocks are devices for finding out what. On that picture the question about noon on the second day has an answer that Hafele and Keating merely failed to measure. It has none. The single present moment is a part of the description rather than a part of the world. The evidence is that no instrument reads it and no procedure produces one.

Agreement between observers has to be built out of messages: one observer sends, another receives, both write. Do messages need a shared present to happen in?

7.1 The messenger

Take away the clock and look at what is left, which is two observers and a channel.

Two armies sit on hills either side of a valley. The enemy is in the valley. Either army alone loses; both together win, provided they attack at the same time. The only way to communicate is to send a messenger down through the valley, where the messenger may be captured.

The first general sends a rider: attack at dawn. The rider may not arrive, so the first general cannot attack on the strength of having sent him. Suppose the rider does arrive. The second general reads the message and cannot attack either, because he knows that the first general does not know the message got through. A general who suspects he may be attacking alone stays put. So the second general sends the rider back with an acknowledgment. That rider may be captured, so the second general cannot rely on the acknowledgment having landed. The first general receives it, and cannot attack, because he knows the second general does not know the acknowledgment arrived.

Acknowledging the acknowledgment moves the problem one rider further down the road and changes nothing else.

E. A. Akkoyunlu, K. Ekanadham and R. V. Huber published the impossibility proof in 1975, in a paper called “Some Constraints and Tradeoffs in the Design of Network Communications”. The argument is three lines long. Any protocol that solves the problem delivers some finite number of messages. Consider the last one delivered. Its sender cannot know it arrived, so the sender’s decision cannot have depended on it, and the receiver’s decision cannot depend on a message that a working protocol has to be prepared to lose. Delete it. Both generals behave exactly as before. Delete the last message of what is left, and keep going, and you reduce any protocol whatever to one that sends no riders at all. Jim Gray named it the Two Generals Paradox in 1978, in a set of lecture notes on database operating systems.

The proof kills certainty, for any number of exchanges, on any channel that can lose anything. Anyone who has operated a network knows what that leaves. A participant that has gone quiet and a participant that has stopped existing send you exactly the same thing, which is nothing, and no amount of listening tells the two cases apart. You can wait longer, as every networked system does, which is what a timeout is: a decision to call a slow neighbor dead at a moment somebody chose in advance and wrote into a configuration file.

The channel does something else to messages besides losing them. Send two messages one after the other across a network with more than one route through it, and the second can land first, because the routes are of different lengths and one of them had a queue on it. The receiver writes them down as they arrive. That is a real order, it belongs to the receiver, and the sending order it fails to match is held by somebody else on another hill. There is a fix, and everyone who builds these systems uses it: put a number in each message before sending it. The number works because both parties agreed on the numbering in advance, which makes it a piece of freight the message is carrying rather than a fact the receiver discovered.

Every requirement chapter three put on an observer survives that. A bounded view, somewhere to write, and a reading that makes a difference, with the record lasting from the writing of it to the reading of it. Read the fifth one again. It asks that the thing outlive its own operations, which is a statement about order. It never mentions a duration, a rate or a second.

7.2 Water, pasta, cheese

A recipe carries the whole distinction. Most people have run one a thousand times without needing a name for it.

Five steps. Fill the pot with water. Bring it to the boil. Put the pasta in. Drain the pasta. Grate the cheese.

Four of those steps are wired together and one is loose. Boiling has to come after filling, since there is nothing to boil otherwise. The pasta goes

in after the water boils, and putting it into cold water is a different act with a different result. Draining comes after the pasta goes in. The cheese can be grated at any point, before the pot comes out or while the pasta cooks or the previous evening. No arrangement of the kitchen makes that choice matter.

In how many different orders can this be cooked? The four wired steps have to happen in their one sequence. Grating can be dropped into any of five positions: before filling, between filling and boiling, between boiling and the pasta, between the pasta and draining, or after draining. Five orders, one dinner, and nothing distinguishes them afterward.

Draw it. Four dots in a row with arrows running along them, filling to boiling to pasta to draining, and a fifth dot sitting off to one side with no arrow touching it at all. There is no left-to-right axis in that picture and no place to put one, because putting one in means deciding where the fifth dot goes. Nothing in the kitchen decides that.

The structure underneath is a relation that says, of some pairs of steps, that one must come after the other. The relation has to behave: no step comes after itself, and if grating comes after boiling and boiling comes after filling then grating comes after filling. Beyond that it is permitted to be silent, and about grating and boiling it says nothing at all. A relation like that is a **partial order**. When it is silent about no pair, when every two items are related one way or the other, it is a **total order**.

A printed recipe is a total order. Step one, step two, step three, down the card, and somewhere between the cooking and the cardboard the one choice that never mattered got made and printed as though it had. A card is read in some order, so an order was invented for it.

Leslie Lamport wrote the distributed-systems version in July 1978, in the *Communications of the ACM*, in a paper called "Time, Clocks, and the Ordering of Events in a Distributed System" that runs from page 558 to page 565. He defined one event as happening before another when the first could have influenced the second, showed that this relation is a partial order and nothing stronger, and then gave an algorithm for extending it to a total order so that a computer system could number things. The algorithm needs an arbitrary rule for breaking ties between events that the partial order leaves unrelated. Any rule will serve. Choose a different one and you get a different total order, equally correct, over the same events.

Causality is the partial order. A timeline is a total order. You can have the first without the second. The first is the one the records hand out, built from which read depended on which write. The second is available to anybody willing to break ties, which can be done more than one way.

There is a physics program built around the same shape. In 1987 Luca Bombelli, Joohan Lee, David Meyer and Rafael Sorkin proposed that spacetime, looked at finely enough, is a partial order in which any two related elements have only finitely many others between them. They called it a

causal set. Sorkin later put what the program asks of one into five words: order plus number equals geometry.

The order half rests on two theorems from the 1970s, one by Stephen Hawking, Alan King and Patrick McCarthy in 1976 and one by David Malament in 1977. Take a spacetime in which no chain of causes loops back on itself and no two distinct events share both their past and their future, and hand somebody nothing but the list of which events can influence which. From that list alone they can rebuild the light cones at every point, and with the light cones every angle and every path a ray of light can take. What the list cannot tell them is how big anything is. A region and a copy of it blown up to twice the size carry the same list. The number half supplies the size: count the elements in a region and the count is its volume.

The hard part runs the other way. Almost no partial order looks like a spacetime. Daniel Kleitman and Bruce Rothschild proved in 1975 that as the number of elements grows, almost every partial order has three layers, a bottom, a middle and a top, with no chain longer than three. A world like that would be enormously wide and last three ticks. Causal-set theory gets orders that look like spacetime by scattering points at random through a spacetime given in advance, which tests the idea and takes the spacetime as given. A theory that produces its own partial order has to say why the order it produces is one of the rare ones with a continuum in it.

The observers build an order of this kind, and the next section spells out how. Each event gets a **source height**, the number of steps in the longest chain of dependencies ending at it. Height goes up along every dependency and comes from who read what, never from anybody's position in an execution list. It counts steps and carries no unit, so it orders events without measuring any duration between them.

The record grammar is generous. Every finite partial order can be written as a pattern of authenticated reads, so records alone force neither single-file events nor a spacetime. The spacetime comes from what the reads are allowed to reach, and chapters fifteen and seventeen build it.

7.3 Caused, without earlier

Putting the pasta into unboiled water is a different event from putting it into boiling water. The arrow from boiling to pasta records that dependency and nothing else. It says the second act consumes what the first produced. Nobody consulted a clock to establish it. The arrow would be there in a kitchen with no clock in it.

Carry that across to observers. One record depends on another when its accepted commit certifies a read of a version the other commit actually wrote. The writer mark must name that parent, and the value read must equal the value the parent left. Assemble the authenticated dependencies

and their chains and you have the partial order, which is the entire causal structure available. Two records with no chain between them have no fact about which came first. An executor that happened to process one of them first adds nothing, and no clock, retry counter or queue position can supply the missing edge afterwards.

7.4 Eight things called time

Values in a program have types, meaning a declaration of what kind of thing each value is, and compilers convert quietly between types they consider close enough. Write a whole number where a decimal is expected and most languages hand you the decimal without comment. The convenience is real, and so is the class of defect where two quantities that were never the same kind of thing get added together because the conversion happened silently.

Eight typed objects enter this discussion, held apart by exactly that mechanism, run in reverse. They are not eight clocks: closure of a complete history and a map from records into spacetime are not clocks at all.

There is closure of the complete history under its self-reference map. There is the repair order, a list of solver moves and the positions they occupy. There is an observer's record order, which says which records depend on which. There is a modular parameter attached to an algebra-state pair, carrying no unit or duration by itself. There is a worldline, a map from records to events preserving the relevant order. There is a clock readout, a real number attached to each record that increases along that order. There is proper time, the calibrated duration along a physical worldline. And there is a global time function increasing along the model's causal order. Host wall time measures a computation's resource use and belongs to a further level of description.

The eighth needs a distinction. Every finite dependency order admits a numbering that increases along its arrows, and Lamport's construction supplies useful logical clocks. Spacetimes admit global time coordinates too. None of them is a preferred clock, a universal simultaneity or one rate all observers share, and none of them is forbidden either: a global ordering is a perfectly good tool for a calculation.

"That repair took about a nanosecond" reads a position in a list as a duration. Accept it once and the positions can be added up, so a stretch of the world's history acquires an elapsed time in units, with no clock anywhere in the reasoning and no calibration performed by anybody. The sentence is humdrum, which is exactly why it gets through. Nothing about it sounds like a mistake.

Crossing between two of the eight costs an explicit map with a name on it, doing the work in the open. Chapter seventeen builds one: the count of records inside a diamond measures its volume, and the fourth root of a ratio of counts is a ratio of durations. The ratio of counts taken straight

is a ratio of volumes. Going from a clock readout to a proper time takes a calibration as well.

Take any clock readout and add seventeen to every reading. Every record reads later than the records it depends on, exactly as before, so the shifted readout is a perfectly legal clock for the same history. It is a different clock. Multiply every reading by any positive number instead and the same thing happens. The record order alone fixes neither the origin nor the rate.

7.5 Two meetings

Einstein got to the same place from the opposite direction in 1905. He started from the speed of light being the same for every observer, which is a strong thing to be handed, and derived that two events simultaneous for one observer are not simultaneous for another. The demand was made by a light signal, and simultaneity came apart under it.

Coming the other way, an observer begins with dependencies among records it can check, and those dependencies select no shared present. A logical numbering or a global coordinate can be built on top of them with further choices, and building one makes nothing simultaneous. What is missing is a preferred present, and every common ordering anybody constructs leaves it missing.

Einstein's route needs the speed of light to come out the same for inertial observers, a measured input to his argument. The records' route gets the same fact from the sky: chapter seventeen finds one invariant speed in the transformations that keep an observer's sky intact. Independent records need no canonical interleaving, even when an executor or a coordinate convention chooses one.

Which is why the only numbers Hafele and Keating could bring home were differences taken at a meeting. Two clocks compare where they meet. Each trip gave the flying clocks two meetings with the bench clocks, one on the way out of the Naval Observatory and one on the way back in. Everything between those two events belongs to a route rather than to a moment. The two routes, east and west, gave answers about a third of a microsecond apart.

7.6 Sixteen schedules

Go back to the twelve observers of chapter six, wired as an icosahedron, thirty overlaps between them, and every one of the two thousand and forty-eight states run under sixteen different repair orders.

Calling them schedules was generous. The sixteen are sixteen lists, each naming which repair move occupies which position, with no duration attached to any position and no rate at which positions go by. Two of those lists that differ only by swapping a pair of moves touching no overlap in

common are not two histories of the world. They are one partial order, written down twice, and the ordering between those two moves is the numbering on the recipe card.

That is why all sixteen had to be run. Taking turns properly has nothing behind it here, so the agreement the twelve reach cannot be an artifact of a well-behaved order. The same answer comes out of all sixteen lists.

Which leaves the twelve of them in the position the generals were in. Each sees five neighbors and nothing else. Each holds an order over its own records and over nobody else's. A message from a neighbor lands after an unbounded number of intervening moves, out of the sequence it was sent in, and sometimes it does not land at all. A leader would have to be recognized as one by observers with no way of comparing what they hold, and a vote would have to be taken in a present that none of them share.

They converge anyway. Every state, every order, one answer, arrived at under exactly the conditions that a branch of engineering has been designing systems to survive since 1975.

Chapter 8

Why Can Strangers Agree Without a Boss?

At 22:52 UTC on 21 October 2018, maintenance work on a failing piece of 100-gigabit optical equipment cut the link between GitHub's East Coast network hub and its primary East Coast data center. The link came back forty-three seconds later.

During that forty-three-second break, GitHub's database-topology software did what it was built to do. It runs a leader-election protocol called Raft. It could not reach the East Coast machines, and Raft carries no test that separates a machine which has stopped answering from one that has stopped. What it does instead is count. The nodes that could still hear one another formed a majority, agreed among themselves that the East Coast primary was gone, and promoted a cluster on the West Coast to accept writes in its place.

The East Coast primary had lost nothing except the ability to talk to the other coast. Applications in its own building went on sending it work, and it went on taking it. One of the busiest clusters accepted 954 writes during the window.

Then the link came back. Each coast was holding writes the other had never seen. Neither copy contained the other, so neither could be made the primary without throwing away data that had been acknowledged to somebody. GitHub ran for 24 hours and 11 minutes in a degraded state, showing information that was out of date and inconsistent, unable to publish pages or deliver webhooks. The divergent writes had to be reconciled by hand.

There is no bug in that story. Every component behaved as specified, on the information available to it. Forty-three seconds of missing fiber cost a day of the largest code host on the planet. The problem was that two groups of machines, neither able to see the other, both concluded they were in charge.

So how does a crowd of parties who cannot see each other's internal state, who share no clock, and who answer to nobody end up holding the same answer, and what does it cost? Engineers buy that agreement with hardware, messages and delay. Alice, Bob and Charlie get theirs for free.

8.1 Two ways to be broken

In 1982 Leslie Lamport, Robert Shostak and Marshall Pease published a paper in *ACM Transactions on Programming Languages and Systems* called "The Byzantine Generals Problem", about a group of commanders surrounding a city who must agree on whether to attack, and some of whom are traitors.

Lamport wanted a nationality for the traitors that no reader would object to, and settled on Albania, which at the time was sealed tightly enough that he doubted any Albanian would ever see the paper. His colleague Jack Goldberg pointed out that there were Albanians outside Albania and that Albania might not stay sealed forever. Byzantium, gone since 1453, raised neither problem.

A **crash fault** stops. The machine dies, the process is killed, the power goes, and what everybody else sees is silence.

A **Byzantine fault** keeps talking. It sends messages that are wrong, or that are inconsistent with each other, or that say one thing to one neighbor and something else to another, all of it with no outward sign that anything has happened. Malice is one way to get there and by far the least common. A memory cell that flips, a disk returning data from before the last write, a program that applied half of an update and then fell over: each of these produces a participant confidently answering questions with garbage. The confidence is the whole of the problem, because the other participants have no way to weigh it.

On 18 May 2003 a vote-counting machine in Schaerbeek, in Brussels, credited one candidate in the Belgian general election with 4,096 votes that nobody had cast. No fault was ever found in the software. The official finding was a spontaneous inversion of a single bit in memory at the thirteenth position, a position worth two to the twelfth, which is 4,096. The machine reported the result the way it reported every other result, with no error and no warning. The reason anyone caught it was that the municipality had returned more votes than it had voters.

Crash faults are cheap to survive. You wait, you notice the silence, you carry on without the missing party. Byzantine faults are expensive. The exact price is a short piece of arithmetic.

8.2 Four dots

Draw four dots. They are four participants. You would like the group to survive one of them being Byzantine, without knowing which one.

Decide that any three of them, agreeing, may commit a decision on behalf of the group. Call a set that large a **quorum**.

Take any two quorums. Each has three members drawn from four, so the two of them together name six memberships among four participants,

which means they share at least two members: three plus three minus four is two. At most one participant of the four is faulty. Therefore at least one member of the shared pair is sound. A sound participant does not vote for two different answers to the same question. So no two quorums can ratify conflicting decisions, whatever the faulty participant says to whom.

That is the entire safety argument. The same subtraction runs at any size. Surviving three broken participants out of ten takes quorums of seven. Any two quorums of seven out of ten share four members, at most three of them faulty. Surviving four out of thirteen takes quorums of nine, which share five, at most four of them faulty. Either way at least one sound participant sits in both. The property is called **quorum intersection**, the reason the ratio is two thirds rather than one half. Ordinary majority voting survives crashes. Surviving liars costs an extra third of the hardware.

Read the ratio as a purchase order. To survive one broken participant you buy four machines and use them to do the work of one. Every decision costs a round of messages from three of them. To survive two you buy seven. The bill is linear in the number of liars you are willing to tolerate and it never gets cheaper, because the subtraction that makes the argument work is doing so at the tightest possible margin: two quorums of three out of four share two members, and if either quorum shrank to two, they could share one, and that one could be the liar.

Notice also what the four dots require you to have before any of this starts: a list of who the participants are, a way to reach any of them, and a fixed count of how many there are. Take away the roster and the arithmetic has nothing left to subtract.

8.3 What Fischer, Lynch and Paterson proved

In April 1985 the *Journal of the ACM* published a paper by Michael Fischer, Nancy Lynch and Michael Paterson titled “Impossibility of Distributed Consensus with One Faulty Process”.

Suppose messages always arrive eventually, but with no bound on how long they take. Suppose one single participant may crash, and only crash: no lying, no corruption, one machine that stops. Then there is no deterministic protocol that guarantees both that everybody who decides decides the same thing and that everybody decides at all.

The result does not say that no efficient protocol exists. It says that no protocol exists. The shape of the proof is that any such system passes through configurations from which both answers are reachable, and that a schedule delivering one message at a time can always steer it out of one such configuration and into another. Every message gets delivered. Nobody is starved. The system is kept poised forever by nothing more than a choice about the order in which messages are delivered.

Faster hardware does not fix this. Better engineering does not fix it. What the result attacks is the combination of three things: asynchrony with no bound on delay, a deterministic rule, and the demand that agreement and termination both hold always. Give up any one of them and consensus comes back.

The industry actually made two of those surrenders.

The first gives up asynchrony. Assume that after some unknown point, message delays are bounded by some unknown constant, an assumption Cynthia Dwork, Nancy Lynch and Larry Stockmeyer formalized in the *Journal of the ACM* in 1988 and called partial synchrony. The protocol never violates agreement, whatever the network does; it promises to finish only during the stretches when the network behaves. Every leader-based system in production is standing on this hatch, which is why every one of them has a timeout in it, and why the timeout is the parameter that operators argue about.

The second gives up determinism. Let participants flip coins, which Michael Ben-Or showed in 1983 is enough: the protocol terminates with probability one, meaning that the runs in which it goes on forever have a total weight of zero, while carrying no promise about any particular deadline.

A leader with a timeout, or a coin. Every consensus protocol running in production is one of those two, or both.

The failover on 21 October 2018 was the first hatch working as designed. The nodes that could see each other formed a quorum, elected a new topology and never disagreed with one another about it, which is exactly what a protocol standing on partial synchrony promises: agreement always, progress when the network behaves. What diverged was the data underneath, because the databases accepting writes on the East Coast were not participants in the election and had made nobody any promises at all. Agreement held, about which machine was in charge rather than about the 954 writes.

8.4 A leader and a clock

Since September 2022 Ethereum has run its consensus on a schedule. Time is cut into slots of 12 seconds, one validator is drawn for each slot to propose a block, and a committee attests to what the proposer produced. Thirty-two slots make an epoch of 6.4 minutes. A checkpoint at an epoch boundary is finalized once the following epoch has been agreed, which puts final on the record 12.8 minutes after the fact and backs it with stakes that can be destroyed.

What that machinery buys is a single history that every participant agrees on, in one order, with a definite moment after which nothing in it can move. What it costs is on the face of the design. There is a leader for

every slot. There is a wall clock that everybody reads, because a 12-second slot means nothing to a participant with no idea what time it is. Every full node holds the entire state, because checking the next block means having the previous one in full. And every transaction is placed in the one order relative to every other transaction, whether or not the two have anything whatsoever to do with each other. The system spends real work establishing which of them came first. Bitcoin buys the same single order with a lottery in place of a schedule, a race to find a number whose hash falls below a target that is retuned every 2,016 blocks to hold those 2,016 blocks to 1,209,600 seconds. Ten minutes a block is what that works out to, and 1,209,600 seconds is two weeks. Whoever wins a round is that round's leader. The chain the winner extends is one line.

8.5 Twenty strangers

In May 2018 a pseudonymous group calling itself Team Rocket posted a paper to a file sharing network, "Snowflake to Avalanche: A Novel Metastable Consensus Protocol Family for Cryptocurrencies", describing a way of reaching agreement in which nobody is in charge and nobody holds the tally.

Each participant keeps two things: its current preference, and a counter. That is the whole of its state. There is no vote count, no view of the network, no record of who said what.

Each round, a participant picks twenty others at random and asks each one what it prefers. If fifteen of the twenty give the same answer, the asker adopts that answer and increases its confidence in it. If twenty rounds in a row come back the same way, it decides, and the decision does not reverse. Twenty as the sample size, fifteen as the threshold, twenty as the number of consecutive rounds: those are the documented defaults on Avalanche's primary network, settings rather than laws. The protocol built out of the sample and the counter is called Snowball.

If a bag holds a hundred marbles and thirty of them are red, the probability of drawing a red one is thirty out of a hundred: the fraction of the cases that come out red, which is the fraction a long run of draws settles toward. That is all the sampling needs. When a sample of twenty comes back fifteen to five, that fraction is evidence about the fraction in the whole population. It can also mislead, because a sample of twenty drawn from a population split sixty to forty will occasionally come back fifteen for the minority, in the way that a fair coin occasionally lands heads six times running. The counter is what handles this. Being misled twenty rounds consecutively is enormously less likely than being misled once. The decision threshold is the dial that pushes the chance of a wrong decision below one in a billion, a bound the published parameters hold with a fifth of the network lying.

Each participant sends twenty questions per round regardless of how large the network is. Twenty when there are a hundred participants, twenty when there are a hundred thousand. The number of rounds needed grows like the logarithm of the population, which is the slowest growth on offer. There is no proposer, no miner, no round leader and no view change, so there is no single machine an attacker can take off the air to stop the thing.

And the dynamics are metastable. Start the network at an even split and it sits on a knife edge, where the sampling itself is the only thing capable of breaking the tie. As soon as one answer runs slightly ahead by accident, participants sampling the network get that answer slightly more often, adopt it, and become part of the majority that the next participant samples. The tilt feeds itself. Past a certain point the outcome is fixed. The only thing that broke the tie was the order in which twenty names came out of a hat.

8.6 Where the analogy inverts

A crowd of parties, none of them in charge, none of them holding the global picture, converging on one answer: that is what a network of overlapping observers does. The engineered version of it settles real payments in production.

Watch an Avalanche node take its sample. It picks twenty participants uniformly at random from the whole validator set, which requires it to hold a roster of every participant in the network and to be able to open a connection to any of them. That roster is a global object, and holding one is the price of the sample being uniform. The locality in that protocol lives in the size of the sample, twenty out of however many, rather than in who the twenty are allowed to be.

An observer in this world has neighbors, fixed, and communicates through those and through nothing else. The twelve of chapter six each had five neighbors. The five came with the wiring: the same five in every round of every schedule, with no round of repair ever producing a sixth. To reach a non-neighbor an observer has to go through a neighbor. There is no address book and no directory. A channel that bypassed the wiring would itself be an overlap, at which point it would be in the wiring.

So the engineered protocol is the loose version. What an observer can be affected by is settled by the diagram of who overlaps with whom. That diagram is finite and fixed.

Snowball's safety is probabilistic. It holds except with an error probability that the parameters drive down. One in a billion is a small number rather than zero. The order-independence of a repairing network of observers is of a different kind. Where it settles is fixed by the disagreements and the wiring, and running the repairs in a different order lands on the same place, in every run rather than in overwhelmingly many of them.

The twelve-observer sweep of chapter six is what that looks like from below, where every state and every schedule was enumerated rather than sampled. The count of exceptions was zero because there was nowhere for an exception to hide.

An engineer would take that guarantee over Snowball's in an instant and cannot have it, because getting it requires the thing engineers are never given, which is control over how the participants are wired to each other.

8.7 What a fixed neighbor list buys

Fix an observer and ask a question that the engineered protocols cannot even pose: what could the reading it holds possibly depend on?

Before any update, its state depends on its own inputs. Each repair touches one overlap, so dependence crosses at most one edge per repair event. A round in which every overlap is repaired once adds at most one edge of depth; after two such rounds, neighbors of neighbors can contribute. A sweep that repairs a path of overlaps in order can carry dependence several edges at once, one event per edge, so the bound counts repair events along a chain of dependence, whatever round counter an executor keeps.

On the twelve-observer graph, five vertices are one edge away, five are two edges away, and the opposite vertex is three edges away. Those counts give possible source sets by dependency depth: six vertices including itself at depth one, eleven at depth two and twelve at depth three. They say who can be heard from and leave open how long a hop takes.

Call the set the **dependency cone**: the observers within reach, where reach is counted in steps along overlaps rather than in kilometers.

A value outside the dependency cone has not influenced that reading. At most one overlap per repair event is a speed limit, and chapter seventeen puts meters and seconds on it.

Two distant readings in this world can be correlated, and chapter six built an arrangement where they are correlated by force, since the twelve observers had exactly one consistent state available to them and every one of them was pinned by it. Correlation is allowed and is ordinary. What no arrangement permits is a distant party choosing what you read. There is no dial they can turn, because the only channel between you is the wiring and the wiring has a pace, so whatever they do arrives as something that walked, or does not arrive.

Correlation without communication is the shape of the Delft result from the prologue, where two laboratories 1.3 kilometers apart got answers that matched too often and could not use the matching to send each other a single bit. In an arrangement whose only channels are its overlaps, that combination is arithmetic about which observers are within how many steps

of which. The name for it is **no-signalling**, arriving here before there are any particles to attach it to.

8.8 Only the conflicts

One feature of the engineered protocol carries over exactly. It looks at first like an efficiency trick.

Avalanche does not order transactions. It organizes them into conflict sets, where a conflict set is the group of transactions that try to spend the same coin, and runs its sampling only within those sets. Two transactions that touch nothing in common are never compared, never ordered, and never made to wait for each other. There is no fact in the system about which of them came first, so no work is done to manufacture one.

The leader family does the opposite, and pays for it in the queue. Every transaction goes into one line whether or not it interacts with anything else in the line, which is why a payment in Lisbon waits behind a payment in Osaka that has nothing to do with it.

A network of observers repairing their overlaps is in the first camp, there by construction rather than by choice. A repair happens where a comparison disagrees. Two disagreements in unrelated parts of the arrangement are not competing for anything, do not touch a common quantity, and have no order between them, which is chapter seven's partial order doing work in this world. One repair comes after another when it read a value the other one wrote. Repairs with no such read between them stand in no order at all. Nothing else in the arrangement assigns them an order.

The local rule needs no leader, no synchronized clock and no roster beyond a patch's own neighbors. A patch proposes and checks its update from the data on its own boundary. Whether all of those local checks add up to one global agreement is the question chapter ten answers.

Alice and Bob share an overlap. Alice reads twenty across it, Bob reads twenty-two, in whatever units that boundary is measured in. Both are careful, neither has malfunctioned, and both readings are of the same thing, which is why two numbers where there should be one is a problem rather than a curiosity. Something has to move. The protocol says nothing about which number moves, or by how much, or why that answer rather than one of the infinitely many others that would also leave the two of them matching.

Chapter 9

What Move Does Reality Actually Make?

David Kinnebrook arrived at the Royal Observatory at Greenwich on 23 May 1794 as assistant to the Astronomer Royal, which meant that most nights of the week he sat at the transit instrument and timed stars.

The method was eye and ear. You listen to the clock beating out seconds, you watch a star cross a wire in the eyepiece, and you split the interval between the last beat and the crossing into tenths by judgment. Nevil Maskelyne had been doing it for thirty years. Both men used the same instrument on the same stars. For a year their numbers matched.

From the beginning of August 1795, Maskelyne found that his assistant was setting the transits down half a second later than he should. He told him. Kinnebrook tried to correct it, and the gap widened. By January 1796 it stood at eight-tenths of a second. Maskelyne wrote that Kinnebrook did not seem to him likely ever to get over it, noted for the record that the man had been diligent and useful, and dismissed him. Kinnebrook left the observatory on 12 February 1796.

A star crossing a wire issues no receipt. No instrument at Greenwich held the true time of the transit, and none could have, because the true time of a transit was the thing the transit instrument existed to produce. Two men, one event, two numbers, and the single measurable quantity in the place was the difference between them: eight-tenths of a second, attached to nobody. Maskelyne resolved it by appointing one of the two readings correct. The reading he appointed was his own.

Twenty-seven years later, at Königsberg, Friedrich Bessel worked out what had happened. In 1823 he published the finding that these differences are involuntary, constant, and present between any two competent observers of the same event. He measured himself against his colleagues and published the results as pairs, because a pair was all there was to publish. Against Argelander he ran 1.223 seconds early. The corrections never attached to a person. They attached to a pair of people. No transit time belonging to nobody was available.

Every observer in this world sits where Maskelyne and Kinnebrook sat. Two of them share a boundary, each holds a reading of it, the readings disagree, no third reading exists, and nobody has been appointed correct.

The question is arithmetic. What does one observer physically do to its own state, and by how much?

9.1 Two readings and a seam

Take chapter six's wiring back out: twelve observers, each with exactly five neighbors, thirty shared boundaries between them. Call a shared boundary a **seam**.

In that universe each observer held one bit and each seam recorded one word, same or different, which is the coarsest reading a seam can carry. Widen it. Let each observer hold a quantity, so that what a seam compares is two accounts of one quantity and the record on a seam is a number rather than a word: how far apart the two ends are, and which way round. A seam is satisfied when the two readings at its ends are equal. The simplest such quantity is a count of units held.

A rule that acts on a small piece of a state and leaves the rest alone, applied over and over in no particular order, is a **rewriting system**. Chapter two had one with two rules and an alphabet of two letters. A state that no rule applies to is where the process stops. A stopping state is a **normal form**. Whether a state is one is a local question: an observer looks at its own five seams, finds nothing to do, and has its answer without knowing anything about the other six observers or about what any of them is doing at that moment.

So what is a repair allowed to be? Three requirements fix it.

First, it changes nothing outside the seam it repairs. The two readings at that seam's ends and the record between them are all it can touch, because an observer talks through its ports and nowhere else, and a move that reached further would need a fact about a place it cannot see.

Second, it leaves the total of those two readings alone. What the two of them are disagreeing about is one shared quantity described twice, so a repair re-attributes between two accounts rather than adding to them. The readings have no absolute level for a repair to correct them toward.

Third, when it is finished the two readings are equal. A move that leaves them disagreeing has repaired nothing.

Solve it. Call the two readings a and b. The third requirement says both end at one value, so call that value v. The second says twice v equals a plus b. Therefore v is half of a plus b. There is nothing further to determine, because two conditions have fixed two unknowns.

Both parties move to the midpoint. Split the difference.

The move carries no adjustable constant. It has no rate, no step size and no fraction of the gap to be closed per event, so there is nothing in it for anybody to have chosen or fitted to a measurement.

Three consequences arrive free. If a and b are equal to start with, the move does nothing, so the states where the law idles are exactly the con-

sistent ones. Neither observer is corrected by the other: both move, by the same amount, in opposite directions. And Maskelyne's move, which was to overwrite the assistant's reading with the Astronomer Royal's, fails the second requirement, because it changes the total of the two numbers it touches, and needs somebody appointed before it can be carried out at all.

9.2 Why does the universe keep books?

Remove the second requirement and the answer disperses: both readings going to a, both going to b, both going to zero, and every weighted blend of the two are all legal moves. No information available at the seam picks between them. Restore it and one move survives.

The requirement belongs there because a rule that changed the total would have to say by how much, and every answer to that question is a fact about a quantity nothing in this world holds. The two observers can add their own two readings together. There is no register anywhere carrying the total of all twelve, no clearing house, and nobody to ask. A universe that repairs locally cannot help conserving things, because a local repair that changed a total would have to know the total. Nothing here knows a total.

The readings have no absolute level. Add the same amount to every observer's books and every seam in the arrangement reads exactly what it read before, which makes the level a convention of the kind chapter five found at Greenwich: unmeasurable from inside, impossible to be wrong about. The gap between the two numbers is the whole of what the pair carries. Now look again at the moves the requirement threw out. Both readings to zero, or to nine tenths of the midpoint, or to any weighted blend of a and b: every one of them needs a zero to move toward, and a level fixed by convention supplies none. Shift every book by the same amount and the midpoint shifts with it, while each of the alternatives lands somewhere else. Conservation is what survives when a move is forbidden to know where the origin is. Chapter twelve takes that freedom seriously enough to get a force out of it.

No admissible move alters the total of the two readings it touches, and reaches no reading beyond those two, so the total across all twelve observers is the same after one repair, after a thousand, in any order, forever. The total is an invariant, a quantity that comes out the same under a change, and the change it survives is every repair the world can perform.

A conserved quantity, in this world, is a quantity that no admissible local move can alter. What makes a move admissible is that it touches one seam and reduces disagreement. Newtonian mechanics conserves momentum and energy too, and nothing inside it says why the universe should keep accounts; the conservation goes in by hand and is then observed to hold. Here it falls out of the only move available.

Chapter five gave the standard correspondence, that momentum is conserved because the laws do not care where you are and energy because they do not care when. The correspondence is exact and it says which quantity goes with which symmetry. It takes the existence of the books for granted before it starts. The books come from the second requirement, which is there because no seam and no observer holds a total. Charge conservation, energy conservation, and the conservation of the quantities that turn out to label particles are all this same argument run on a different reading of a seam.

9.3 Take three numbers

Write three numbers at the corners of a triangle: 6 at the top, 0 at the bottom left, 3 at the bottom right. Every corner is joined to the other two. Take a pencil and work out, for each corner in turn, its own number minus the average of the two numbers it is joined to, and then double the answer, because each corner has two neighbors. Three subtractions and three doublings. Write the results beside the corners before reading the next paragraph.

The top corner sits at 6 and its two neighbors average 1.5, so it gets 9. The bottom left sits at 0 against an average of 4.5, so it gets minus 9. The bottom right sits at 3 against an average of 3, so it gets 0.

Three numbers in, three numbers out. Each output says how far above or below its own neighborhood that corner sits, scaled by how many neighbors it has. The three outputs sum to zero, because every gap in the list appears twice with opposite signs. The operation is called the **graph Laplacian**, it is defined on any graph whatever, and the sentence carried out three times with the pencil is all of it:

$$(Lx)_{\text{me}} \;=\; n_{\text{me}} \times (x_{\text{me}} - \text{average of } x \text{ over my neighbors})$$

Here x is the list of readings, one number per corner, x with the subscript me is the reading at my own corner, n with the subscript me is how many neighbors my corner has, and L is the name of the operation that turns one such list into another. Nothing on the right-hand side looks past the corners a corner is joined to, which is why an observer can work out its own entry without leaving its patch.

Repair a seam by hand. Take the 6 and the 3, whose midpoint is 4.5, and send both there: the triangle reads 4.5, 0, 4.5. Add them. Start over and repair the 0 and the 3 instead, which puts both at 1.5: the triangle reads 6, 1.5, 1.5. Add them. Three seams, six orders of working through them, and the running total is 9 at every point in every one of those histories, because a move that sends two numbers to their midpoint puts back exactly what

it took out of them. No corner holds the 9, and no move available on the triangle can spend it.

The midpoint move is the graph Laplacian of a single seam, halved and subtracted: each reading moves by half the gap to its partner, toward the partner. Apply it to all thirty seams and weight the seams equally, and what acts on the twelve readings is one operator:

$$T = I - \frac{1}{60}L$$

Here T is the average effect of one repair on the twelve readings, I is the operation that leaves all of them alone, L is the graph Laplacian of the wiring, and the sixty is the thirty seams doubled, because each endpoint moves by half the gap, closing the full pairwise gap. So a reading moves toward the average of the five readings around it by a twelfth of the distance: the Laplacian counts that distance once per neighbor, and five sixtieths is a twelfth.

That operator is the one in the heat equation, written out on a graph instead of on a line. Heat in an iron bar obeys exactly this rule: the temperature at a point changes at a rate set by how far that point sits from the average of the temperatures next to it, which is why a bar heated at one end finishes up uniformly warm and why the warmth arrives at the far end late. The hot side and the cold side of a bar land on their midpoint for the reason two observers do. Repair is diffusion.

This gives a descent calculation. Score a state by the sum of the squares of the twelve readings. Replacing two readings by their mean lowers that score by exactly half the squared gap. For real-valued states, a decreasing score bounded below can approach a limit through infinitely many moves; finite termination needs a stronger, well-founded progress argument.

Check it on the triangle. Square 6, 0 and 3 and add them: 45. Repair the seam between the 6 and the 0, which sends both of them to 3, then square and add again: 27. The level fell by 18. The gap that was closed was 6, and half of 6 squared is 18. The average was 3 throughout. A single move on the widest seam finished the triangle.

When the readings are counts, the level is a whole number. A whole number that strictly drops and cannot go below zero runs out of room after finitely many steps. That is the general recipe for showing a rewriting system terminates: find a quantity every rule strictly lowers, which cannot be lowered indefinitely.

A repair event touches one seam, so dependence crosses at most one seam per repair event. Whatever a reading depends on reached it along a chain of repairs, one seam to a link, and some chains attain that limit exactly. A round in which every seam is repaired once extends the reach by at most one seam. A sweep that repairs the seams along a path in order can carry it several seams in one sweep, one event per seam. Counted in

repair events, that is a speed limit. Chapter seventeen puts meters and seconds on it.

9.4 The rule that got stuck in 302 places

Let the readings be counts, and give the twelve observers twelve units to hold between them. Since the total is untouchable, the entire history of any run is confined to the arrangements of twelve units over twelve observers. There are 1,352,078 of those.

A space that size can be checked all the way through, so check it all the way through.

The obvious rule goes first. If a neighbor and I differ by two units or more, hand one unit across the seam. It reduces the gap, it preserves the total, it touches nothing but the two of us, and it is wrong.

Run every one of the 1,352,078 arrangements until the rule has nothing left to do. Three hundred and three of them have nothing to do at the outset. One is the balanced arrangement, everybody holding one unit, which is the answer. The other 302 are lopsided and the rule cannot move them.

They are simple to describe once you have seen one. Give one observer two units, give another observer none, place those two so they are not neighbors, and give everybody else one. Every seam in that arrangement has a gap of one or zero, so the rule has no move available anywhere. The arrangement is not the balanced one. Count them yourself. An observer has five neighbors out of the other eleven, which leaves six it is not joined to, and twelve places to put the full observer times six places to put the empty one comes to seventy-two. Two hundred and ten more have two observers holding two units and two holding none. Twenty have three holding two and three holding none. The three counts add to 302.

The defect is at a gap of exactly one. Handing that unit across reverses which observer is ahead without narrowing anything. A rule that accepts only moves which improve the score refuses it. So the process halts on a plateau where no move lowers the score, one observer keeps a spare unit and another keeps none.

The repair is to complete the move instead of nibbling at it. On a seam whose two readings add to an even number, the midpoint is a whole number and both sides go to it. On a seam whose readings add to an odd number, no pair of equal whole numbers has that total. The closest the two readings can come is half a unit either side of the midpoint, a pair that can be placed two ways round. Both placements are legal completions, including the one that reverses which observer is ahead. That one is legal because it changes nothing measurable.

Allowing that one move changes the audit. Under the completed law, exactly one arrangement out of 1,352,078 rests. It is the balanced one.

From every other arrangement there is a path of at most three events to a strictly lower level. Most of the stuck ones need two.

Watch one of them get out. An observer holding two units sits opposite an observer holding none, with the ten others holding one each. Every gap in the arrangement is one or zero, so nothing improves anywhere. The full observer and one of its neighbors hold two and one, which totals three, so the midpoint is one and a half, the two readings land half a unit either side of it, and they may land either way round: the surplus unit walks to the neighbor, and the level has not moved. It walks again, to a neighbor of that neighbor, and the level has not moved on that event either. On the third event the observer holding the surplus is finally adjacent to the observer holding none, their readings total two, the midpoint is a whole number, and both go to one. The level drops. Nothing was gained by the first two events except position, which is exactly what the one-unit rule refused to spend anything on.

Three hundred and two failures out of 1,352,078 arrangements is a failure rate of two in ten thousand, which is why the rule had to be run over every arrangement rather than over a sample of them.

9.5 What a repair writes down

One freedom is left in the completed move. When the two readings add to an odd number, somebody has to end up with the extra unit, and nothing at the seam prefers either end. So the choice is a fair coin. That finishes the rule: nothing lost, the job done in one move, a tie broken by a coin that looks at nothing, and no favorite end. Take the coin away and let the fuller side always keep the spare unit, and the 302 stuck arrangements come straight back. The coin is what lets the surplus walk.

In a quantum world a repair does one more thing. It writes something down. The requirements, the completed move and the coin fix where the units go. They do not fix what the event records, and the record decides how much of the world's quantum coherence survives the repair. A repair can record only how many units crossed the seam. It can record the coin as well. It can add whether the seam's total was odd, or the full count at both ends. Each extra entry is one more fact the world has committed to, and each one erases interference that a leaner record would have kept.

All of these records agree about everything you can learn by counting units. Prepare definite arrangements, repair, read how many units crossed, and the statistics come out the same whichever record the world keeps. The difference is in the coherence, and the cleanest place to see it is between an empty seam and a seam holding a single unit, which the coin may leave where it is. Put those two arrangements into superposition, repair once, and ask whether the superposition came back. The chance is three eighths plus half the coherence that survived. A record of the crossing alone gives

about 0.73. Adding the coin gives five eighths. Adding the parity of the total gives three eighths, flat, at every phase.

The parity record and the full count share a property the leaner records lack. After either of them, repairing the same seam a second time changes nothing at all: one repair finishes the job, the way a projection does. Every record that keeps an empty seam coherent with a full one ends at the parity record if the same seam is repaired often enough.

Which record the world keeps is a question for interference experiments, and the one above answers only part of it. It reads a single number, the coherence a repair leaves between an empty seam and a waiting unit, and no record lets that number exceed one over the square root of two, about seven tenths. The parity record and the full count both erase it, so both give three eighths and the experiment cannot tell them apart. A second experiment can. Put an empty seam and a full one into superposition instead, repair once, and ask again: the parity record brings the superposition back every time, and the full count brings it back half the time. The whole family of possible records has nine numbers in it, and nine such measurements, each a superposition of two seam arrangements read after one repair, fix all nine.

9.6 Two tellers, one account

An account holds 1,000 and two tellers open it at the same moment. The first takes a deposit of 100, adds it to the 1,000 in front of her, and writes 1,100. The second pays out 200, subtracts it from the 1,000 in front of him, and writes 800. Both did correct arithmetic on the number they were given. The account ends at 800, the depositor’s 100 is gone, and no entry anywhere records its disappearance, because every individual entry is right.

Two operations that touch nothing in common can be done in either order. No record afterwards can tell which happened first. Two operations that touch the same item cannot, because the order is part of the answer. Operations that conflict are operations whose order matters, which is the whole content of the word conflict.

In June 1981, H. T. Kung and John Robinson published “On Optimistic Methods for Concurrency Control” in ACM Transactions on Database Systems, proposing that a database should stop asking permission. Their transaction runs in three phases. During the read phase it works from the copy of the data it took when it started, keeping a list of everything it looked at, its **read set**, and a list of everything it intends to change, its **write set**, and writing nothing anybody else can see. During the validation phase it checks whether anything in its read set was changed by a transaction that committed while it was working. Only if that check passes does it reach the write phase and make its changes visible. A transaction that fails validation is discarded and run again from the start.

Locking says: nobody touches this while I work. Optimistic control says: work, then prove nobody touched it. Where conflicts are rare the second is cheaper, because nothing waits for anything. Where conflicts are common it is expensive, since every collision costs the whole transaction and the work is done twice, which is the trade the designer of any such system is making whether or not anybody writes it down.

The acceptance layer of the repair law is that mechanism, item for item. A repair is proposed rather than performed. Its read set is the pair of readings at the ends of one seam, its write set is the same pair. It commits only if the fit on every overlap it touches is at least as good afterwards and strictly better on one of them. Otherwise it is discarded. The observer reads again. Nothing in that loop is a rule handed to the observer from outside. The three requirements say what a proposal has to satisfy; working out a proposal that satisfies them is the observer's own computation, and the check is what decides whether the proposal becomes the world. Because read set and write set are confined to a single seam, two repairs on seams with no observer in common cannot conflict, which is why nothing has to put them in an order. The check has a second effect that the tellers make obvious. A repair cannot buy agreement on one seam by wrecking it on another, because it is validated against every overlap it touches and refused if any of them comes out worse.

Put a dot for each transaction and draw an arrow between two dots whenever those transactions touched the same item in a way whose order matters. That is a **conflict graph**. A batch of interleaved transactions can be rearranged into a run where they went one at a time exactly when that graph contains no cycle. Whether local records assemble into one global account is settled around loops, which is the condition Alice, Bob and Charlie's coins failed, turning up here in a banking system.

9.7 Two budgets

The three requirements fix what the two readings end up holding. They say nothing about which register the repair gets written into, and a seam has two: the readings in the observers' own patches, and the record sitting between them. So a repair comes in two kinds. The difference is who pays for it.

In the first, an observer rewrites its own patch so that its reading matches what the seam says. The capacity spent is the observer's own, nothing outside its patch changes, and the boundary between it and its neighbor says afterwards what it said before.

In the second, the boundary record itself is rewritten, so that both sides read something new. The capacity spent is held jointly by the two observers on the overlap. Neither of them owns it alone.

The two budgets are not interchangeable. An observer with capacity to spend on itself can bring its own reading into line and leave the boundary saying what it always said. Nothing outside that patch registers that anything happened. Rewriting the boundary takes both of them, and afterwards both are holding something neither one held before.

9.8 Sixty ways to go next

Chapter six ran a twelve-observer universe under sixteen schedules and got the same answer every time, on a system holding one bit per observer and 2,048 distinguishable states. The machine with counts in it is larger: 1,352,078 arrangements, and at every point in a run there are sixty directed completions to choose from, one for each seam in each direction.

The rule says which moves are allowed. A run of the machine picks them in some order, and nothing in the rule needs a synchronized clock or a leader to do the picking. Different orders pass through different intermediate readings, and the records of which repair read which keep part of that history.

They land on the same arrangement. The three requirements do not say they have to: two rules on two letters are enough to build a rewriting system where the answer depends on which rule you reach for first, and where both answers are places at which nothing further can be done. So every schedule landing in the same place is either a lucky property of this particular wiring or a theorem. If it is luck, then what the twelve of them end up holding depends on which seam happened to be repaired first. There is no objective answer anywhere in the machine.

Chapter 10

Why Doesn't It Matter Who Goes First?

Hand the same sudoku to two people on the same train. One of them works the sevens first, because four of them are showing and they constrain each other. The other goes box by box from the bottom left. They do not speak, they do not look at each other's paper, and neither has the faintest idea what the other has written down. Forty minutes later both grids are full and all eighty-one squares match.

Nothing arranged that. Two different procedures ran in two different orders on different parts of the grid. The agreement at the end was neither negotiated nor checked.

The puzzle in their laps is older than its name. Among the crosswords in the May 1979 issue of *Dell Pencil Puzzles and Word Games* there was a nine by nine grid with some of its squares filled in and the rest blank. It was called Number Place. Every row, every column and each of the nine three by three boxes had to end up holding each digit from one to nine exactly once. The man who devised it was Howard Garns, a retired architect in Indianapolis, seventy-four years old, and Dell did not print his name on it. He died in October 1989. Five years before that, the Japanese publisher Nikoli had picked the puzzle up and given it the name it travels under, which is sudoku, and in 1997 a retired Hong Kong judge named Wayne Gould found one in a Tokyo bookshop, spent six years writing a program that could make them by the thousand, and talked The Times of London into printing one on 12 November 2004. Within weeks every British newspaper had a grid on its puzzle page.

The two grids on the train match because the finished grid was fixed by the clues before either of them picked up a pen. There are 6,670,903,752,021,072,936,960 ways to fill a nine by nine grid legally, a figure Bertram Felgenhauer and Frazer Jarvis computed in 2005. A printed puzzle is a set of clues that cuts that number to one. In 2012 Gary McGuire, Bastian Tugemann and Gilles Civario finished an exhaustive search and established how few clues can manage it. Seventeen. Sixteen never suffices, on any grid, in any arrangement. Below seventeen the clues stop picking out a single completion. The two people on the train can hand in different grids and both be right.

The clues do half of the work and the pencils do the other half. Every move either of them makes is forced. A square gets a five because five is the only digit its row, its column and its box have left over for it, and once that is true it goes on being true whatever else gets filled in, because filling in another square removes possibilities and never restores one. A forced entry cannot be un-forced by a later forced entry. So the order the two of them work in decides which square gets filled at which minute. It has no way to reach the digit that square ends up with.

Repairs in the network of the last chapter happen when a mismatch is noticed. Which mismatch gets noticed first is whatever the world happens to hand over. There is no schedule, in the sense that there is no fact anywhere about the order. If the endpoint depends on that order, then the world an observer finds is a fact about scheduling. No observer inside can see a schedule.

10.1 Two librarians

Take a shelf of books that ought to be in order and are not. The rule is one line: if you find two neighboring books out of order, swap them. Any pair, whenever you like.

Photograph the shelf. Let one librarian sort it. Then put the books back exactly as the photograph shows and hand the shelf to a second librarian who has been out of the room. Both finish with the shelf in order, which is no surprise, because there is one way for a shelf to be in order. Count their swaps. Each of them will have made exactly the same number. Neither of them chose it.

Count, at any moment, every pair of books in the wrong relative order, wherever on the shelf the two of them happen to sit. A book that belongs before another one and stands to its right is one such pair, whether the two are touching or at opposite ends of the run. Call each of those a wrong pair. Swapping two neighbors that are out of order fixes exactly one wrong pair, because it reverses the relative order of those two books and leaves the relative order of every other pair untouched. So the count drops by exactly one at every swap. It drops to zero. The number of swaps is the number of wrong pairs the shelf started with, which is a property of the shelf and not of the librarian. This is the descent argument of the last chapter, applied to a shelf: a quantity that strictly falls and cannot fall below zero cannot go on falling.

The shelf does something else on the way down. Take a shelf where two swaps are available: books three and four are out of order, and so are books seven and eight. Do the first one. Do the second one instead. The two shelves you get are different from each other. From the first, swap seven and eight; from the second, swap three and four. Both land on the same shelf.

Draw those four shelves with arrows between them and the picture is a diamond: one at the top, two down the sides, one at the bottom where the paths meet. The **diamond property** is the name for that shape. The property it names is that any two moves out of a state can each be followed by one move that brings them back together. Allow the branches any number of moves to find each other again and the condition is called **local confluence**. Allow the branches themselves to be any length, so that two long histories out of one starting state have to be reconcilable, and the name for that is **confluence**.

Three books show what the looser condition buys. Put Chesterton, Cervantes and Bunyan in positions three, four and five. The pair three and four is out of order, the pair four and five is out of order, and the two swaps share a book, so they are not independent the way the first example's were. Swap three and four and the shelf reads Cervantes, Chesterton, Bunyan. Swap four and five instead and it reads Chesterton, Bunyan, Cervantes. Those two shelves are further apart than the first example's pair. No single swap carries either one to the other. From the first, swap four and five, then three and four. From the second, swap three and four, then four and five. Both arrive at Bunyan, Cervantes, Chesterton. Two moves on each branch rather than one, and three swaps down each route, which is the number of wrong pairs those three books started with.

Those last two conditions are separate statements. Local confluence says any two first steps can be reconciled. Confluence says any two histories can be. A history is a hundred moves of drift, and knowing that its opening move was harmless says nothing about where the other ninety-nine took it.

10.2 A snowflake on Triple Divide Peak

Triple Divide Peak stands 8,020 feet up in Glacier National Park in Montana, on the one summit where two continental divides cross. Water leaving that summit on one face reaches the Pacific by way of the Columbia. On another face it reaches the Gulf of Mexico by way of the Missouri and the Mississippi. On the third it reaches Hudson Bay by way of the Saskatchewan. Two snowflakes landing a hand's width apart on the summit finish in different oceans.

Inside any one of those three basins the outcome stops depending on the route. A raindrop falling anywhere in the Columbia basin can take any of a billion routes down, through gullies and side creeks and a reservoir or two, and arrives at the same Pacific as every other drop in the basin. Different paths, one sea. That is confluence, drawn at the scale of half a continent.

Water is also guaranteed to stop. It goes downhill, its height is a quantity that strictly decreases, there is a floor, and every drop that lands on that mountain reaches sea level and stays. Termination promises that the

process ends. It promises nothing about ending in one place. A mountain with three basins meeting on its summit is what the failure looks like.

The failure runs the other way as well. Loosen the librarians' rule so that any two neighboring books may be swapped whether or not they are out of order, and the branches go on being reconcilable, since enough swapping turns any shelf into any other shelf. What goes is the stopping. The two librarians can push Bunyan past Cervantes and back again forever, able to agree at every step and never arriving at a shelf to compare.

The mountain misleads in one place. Water runs downhill because there is a downhill, and it stops at the sea because the sea is there before it arrives. The network has neither. Its settled configuration is the arrangement that satisfies every seam at once, picked out by the constraints in the way seventeen printed clues pick out one grid, before a single repair happens. Repairs delete the arrangements that fail. Nothing travels anywhere. Nobody who finishes a sudoku believes the grid was assembled by the pencil. The same restraint is owed to a network that ends up consistent.

Max Newman supplied the missing piece in 1942, in the *Annals of Mathematics*, in a paper on theories with a combinatorial definition of equivalence. He was a Cambridge topologist whose lectures had sent Alan Turing off to write "On Computable Numbers", and in the year that paper appeared he arrived at Bletchley Park, where he was given a section that everyone called the Newmanry and told to break the German high command's teleprinter cipher with machines. The first Colossus was delivered to him on 18 January 1944.

His result fits in a sentence. If a process always stops, and if every pair of single steps out of a state can be brought back together, then every pair of histories can be brought back together, and every starting state has exactly one endpoint. That is Newman's lemma.

The argument runs from the bottom of the descent upward. Take a state, and suppose that everything reachable below it has a single endpoint. That supposition is safe, because the process descends and cannot descend forever, so you can start at the bottom and work upward. Two moves lead out of your state. Local confluence brings those two moves back together somewhere below. The two states they led to both sit below the state you began at, so each of them has a single endpoint, and so does the state where they rejoined, which forces all three endpoints to be the same one. Your state inherits it. The induction runs over every level of the descent. The descent is finite because the disagreement it counts down is finite.

10.3 Two things to check

Newman's first condition, that the process stops, was settled in the last chapter. Every accepted repair strictly lowers the total disagreement across the network, that total is a sum of terms none of which can go negative,

and so it has a floor at zero. A quantity that drops by a positive amount every time and cannot pass zero runs out. The network stops.

The second condition is the one that does the work. Two repairs happen at two different seams. If those seams share no patch, each repair reads and writes numbers the other never touches, and the two orders give identical results, because the repairs are not interacting in the first place. That is the shelf's first example, where books three and four had nothing to do with books seven and eight.

The case that has to be checked is the shelf's second one. Draw three patches in a row. There is a seam between the first and the second, another between the second and the third, and the middle patch sits on both of them. A repair on the left seam moves the middle patch's reading. A repair on the right seam moves the same reading. The middle patch has only the one.

A repair does one thing. It takes what the two sides of a seam expose to each other and moves them together. What each side exposes is overlap data, information about a region both patches can see, and assembling overlapping regions is associative: glue the first two and then the third, or glue the last two and then the first, and the object you finish holding is the same object. Two repairs sharing a patch are two such gluings. The order in which overlapping regions were glued is not recorded anywhere in the result. Whichever repair went first, one further move on each branch brings both branches to the same configuration.

The network has both of Newman's conditions. Every configuration of it settles into exactly one state. That state carries no trace of the order in which the repairs happened.

Schedule independence is what objectivity means here. The twelve-observer universe of chapter six settled onto one arrangement from all 2,048 of its states, under sixteen orders of repair apiece. Newman's two conditions cover the configurations nobody enumerated, and past twelve observers that is all of them.

In most of engineering the order does change the answer. It changes the answer here too, on the smallest arrangement that can show it, which takes two observers and one seam.

Give each of them a single bit and one rule: when you disagree with your neighbor, adopt your neighbor's value. Nothing about that is unreasonable. It is the politest repair rule available, it strictly reduces disagreement, and it touches only the seam it is applied at.

Start them disagreeing. The left one holds false and the right one holds true.

Let the left one move. It takes the right one's value, both hold true, and since they agree neither will ever move again. Go back and let the right one move instead. It takes the left one's value, both hold false, and neither will ever move again.

Two endings, both final, out of one start. Whoever moves first wins, permanently. The two worlds are easy to tell apart, and that is the trouble rather than the consolation: both holding true means the left one moved first, both holding false means the right one did, so the contents of the world are a transcript of a schedule that no rule chose and nothing required.

Confluence belongs to the repair rule rather than to locality. Leave those two patches wired exactly as they are and change only the rule, from adopt your neighbor's value to something that defers to a fixed criterion, say that whichever of them holds true gives way. From the same disagreeing start there is one move available instead of two, both orders arrive at false, and the fork is gone. Courtesy was the whole problem: a rule in which each party defers to the other leaves nothing to settle the case.

The twelve-observer network has confluence, and chapter six settled it by starting every configuration the arrangement admits, rather than by arguing that the alternative sounded unphysical.

The engineered protocols of chapter eight buy order-independence with sampling and probability. The guarantee they buy is a guarantee about how often. What the observers get instead is two conditions and an induction over a finite descent, which is why their version has no failure rate attached to it.

10.4 What the network is made of

The arrangement in this chapter is one the OPH research team runs as a program, and the program is called OPH-FPE. It is not a numerical model of a continuous world. Every quantity inside it is a whole number or one of the six ways of permuting three labels, every move is exact, and no step anywhere rounds anything off. A run either meets a condition or fails to, and it returns the same verdict on every machine that executes it.

Each published run is archived entire: the configuration it was given, the seeds it drew from, the arrays it started with, the arrays it finished with, the full trace, and the state of all twelve ports on every patch. Beside them sits a verifier that borrows none of the simulator's own code. It builds the six permutations from scratch, recomputes the starting disagreements from the raw arrays, works out the finishing state, and recomputes every fingerprint. The run read here was archived on 27 August 2026. Its archive has since left the repository's working tree but stays in the repository's history, and Appendix C gives the commit. A reader with Python who checks out that commit runs one command and either gets the same answer or finds out that they do not.

The run read here is the level-six rung of a tower built by subdividing an icosahedron over and over. It holds 81,920 patches and 122,880 seams between them, and every patch carries twelve local port slots, three of

which route seams while nine stay exposed. At the start, 102,415 of those seams disagree, which is 83.3 percent of them.

10.5 Two cycles

The trace is kept in cycles. A cycle is a round in which every patch with something to repair gets a turn. That count belongs to whoever reads the trace afterward. Nothing in the network counts cycles, nothing waits for the next one, and there is no tick anywhere in the machinery.

A run of a process that failed to be confluent would be an anecdote: what happened to one arrangement on one pass, with no warrant to say anything about the next pass, which would take a different order and could finish somewhere else entirely. Newman's two conditions are what make the last line of this trace worth more than the run that produced it. The lines in between are the accidents of one order. The line at cycle 1 would have read zero under any of them.

Cycle 0 accepts 81,920 repairs and leaves 20,495 seams in disagreement. Cycle 1 accepts the remaining 20,495 and leaves nothing. Of 122,880 seams, every single one agrees, and the trace runs on to cycle 15 without another mismatch posted against it.

Each repair removes exactly one disagreement, which is the librarians' shelf at the scale of a network, and the number of moves the run accepts is 102,415, the number of disagreements it began with. Neither figure has anything in it about the order the repairs came in.

10.6 Sixteen shuffles

One trace records one order. What settles the question is running the same arrangement sixteen times and shuffling the order before every pass.

All sixteen accept 102,415 moves. All sixteen finish with no seam in disagreement. All sixteen arrive at the same finishing state, and the check on that last point is a fingerprint: the whole terminal state is boiled down to one line of characters, and the sixteen lines are the same line.

Sixteen orders. One destination.

The same replay takes 512 places where two repairs share a patch and confirms that the two ways around the diamond meet, takes 512 places where two repairs have nothing in common and confirms that they commute, and relabels the local frames sixteen times over to confirm that nothing in the outcome was resting on the labels. None of those checks turns up a violation.

The records arrive after the arguing stops. Nothing is committed through cycle 6. Cycle 7 posts 47,412 of them, a little under three in five. Cycle 8 posts the rest, 81,920, one for every patch in the network, and from there to the end of the run nothing changes at all.

The spread of those records at the finish is 11.3134. The natural logarithm of 81,920 is 11.3135, and the whole of the gap between those two figures is the price of four coincidences: four pairs of patches hold matching readings, and each of the other 81,912 holds a reading that no other patch in the network holds. Spread counts how many different things a collection holds, on a scale where doubling the number of distinct items adds a fixed amount instead of doubling the figure.

10.7 Agreement first

Records lag agreement by six cycles, and they lag it because of what a record is. A record is something you can go back to and find unchanged. Write one into a neighborhood whose seams disagree, and a repair comes through and revises what you wrote, which makes what you wrote a draft.

The run puts a number on that. A patch's reading becomes a record when it has held still for eight cycles together. Anything in the middle of being repaired fails that test, because being repaired is what failing to hold still consists of.

Read the rule closely and notice what it does not say. It does not tell a patch to look at its neighborhood, or to wait for anybody, or to ask whether the argument nearby has finished. It asks one question, and the question is about the patch itself: has this reading stopped changing. The neighborhood enters only through the back door, because a patch whose seams are in dispute is a patch whose reading keeps getting revised out from under it.

Agreement comes first and memory second. The order is forced by what a record is rather than by any instruction to observe it, and it runs the opposite way to the picture most people carry, in which observers write down what they see and then compare notes and argue about the differences. Through six cycles of this run, every patch in the network is in the middle of a disagreement or fresh out of one, and not one of them has written anything down, and there is nothing anywhere to compare.

10.8 The last line

The last line of the trace reads like this. One hundred and twenty-two thousand eight hundred and eighty seams, every one of them in agreement. Eighty-one thousand nine hundred and twenty patches, each holding a record of its own. Whatever order the repairs ran in, that is where the network lands, and Newman's two conditions are the reason.

Everything settled. Nothing that settled is a number any two patches share. Each patch holds its own reading, taken through its own ports, of the piece of the world it can see. Walk up to two neighboring patches at

the end of that run and ask each of them what the world is like, and you get two different answers from two observers in complete agreement.

The network has a settled state. It is written down in none of the eighty-one thousand nine hundred and twenty places where things get written down.

Chapter 11

What Does Everybody Actually End Up Holding?

What is not in the record is not in the world. Continental procedure keeps the sentence as a maxim, in Latin, quod non est in actis, non est in mundo, and a courtroom is a machine for living by it.

A trial does not ask what is inside a witness's head, partly because there is no instrument for getting at it and partly because the contents turn out to depend on how you ask. What a trial produces instead is a record: the words actually spoken in the room, written down as they are spoken by a court reporter. Exhibits go into it. Private recollection does not, and neither does anything a juror happens to know from outside.

The record is a physical object made by a person under load. The Registered Professional Reporter certification, from the National Court Reporters Association, requires taking down two speakers at 225 words a minute with 95 percent accuracy, on a keyboard with 22 keys, which is fewer keys than there are letters and is the reason the output is phonetic rather than alphabetical. Everything a court later does with the case runs through what came off those 22 keys.

No authority outside the room supplies the verdict. Twelve people who sat through the trial compute it from the record. When they cannot agree about what a witness said they ask for the transcript and it is read back to them.

How much the contents of a head depend on the asking was measured in 1974, when Elizabeth Loftus and John Palmer sat forty-five students down at the University of Washington and showed them seven short films of traffic accidents. Afterward each student was asked how fast the cars had been going. It was the same question every time except for one word. Some were asked how fast the cars were going when they smashed into each other, and the others got collided, bumped, hit, or contacted.

Everybody saw the same films. The smashed group put the cars at 40.8 miles an hour on average, the hit group at 34.0, and the contacted group at 31.8.

A second experiment used a hundred and fifty students and one film of a multiple-car crash. Fifty of them were asked about smashing, fifty about hitting, and fifty were asked nothing about speed at all. A week later, all of them were asked whether they had seen any broken glass. Sixteen of the fifty in the smashed group said they had, against seven in the hit group and six in the group that had never been asked about speed. There is no broken glass in the film.

Loftus and Palmer called the paper "Reconstruction of Automobile Destruction". Nobody in those rooms was lying, and nobody was playing back a stored recording. Each of them was assembling an account out of the material available, and the material available included the verb.

Chapter ten put eighty-two thousand patches through the repair process and the last of their 102,415 disagreements went at cycle 1, whatever order the repairs ran in. That settles the schedule. It says nothing about what any patch is holding at cycle 2. Each finished the run holding its own state, visible to nobody else, connected to the rest through what crosses its seams. So what do they end up holding in common, and how much of the world does it pin down?

11.1 The list everybody can check

When Alice and Bob met in the corridor with their coins, they wrote down one word, same or different, and never which way either coin was facing. That was the only restriction in the whole arrangement. Two observers cannot compare what only one of them holds; they can compare the relation between them. The relation is what gets written down.

Collect every one of those written comparisons into a single list. That list is what anybody can check, in the sense that any observer can walk up to the seam it names and find out whether the entry is right. Everything else in the network is somebody's private state.

The **record map** takes the full state of the network and returns the part of it that survives comparison: every seam entry, and nothing that only one observer could have known. Two observers who disagree about anything are disagreeing about an entry, because an entry is the whole of what either of them can put on the table.

What the record map deletes is real. Alice's coin is a feature of Alice, she can look at it whenever she likes, and it does not appear in the list because there is no procedure by which she and anybody else could jointly establish which way it faces. She has been deleting it at every meeting since the corridor, and so has everybody she has ever compared with.

A state of the network is **consistent** when every entry on that list is true of the patches it names. Twelve coins and thirty written records: consistent means all thirty records say what the pairs of coins they name actually do. The collection of consistent states is the **consistent set**.

Chapter six counted that set for the icosahedral net. Thirty records that pass the loop test, twelve coins, four thousand and ninety-six ways to set them, and exactly one setting satisfies every record, along with the setting you get by turning all twelve coins over at once.

11.2 Three verdicts

Fix a list. Ask which consistent states carry it.

The answer is a set: all the ways the world could be that would produce exactly this record and no other. In court it is the set of stories the testimony fits. The prosecution has one of them in mind, the defense usually has another, and the size of the set is what the argument is about. The set of consistent states carrying a given reading is called the **fiber** of that reading.

Wire three patches to each other and to nobody else. Three coins, three seams carrying one entry each, and eight ways to set the coins.

Read all three entries as same. Heads all round satisfies them and tails all round satisfies them, and those two differ by turning over every coin in the world, which no entry sees, so they count once. One world.

Read all three entries as different instead. Chapter six wrote different as 1, added the three of them in the arithmetic where one and one make zero, got one rather than zero, and no setting of three coins survives a loop that fails to close.

Let the third patch go quiet. Its two seams go unread, the record shrinks to the single entry between the first two, and that entry reads same. Count this one yourself. Four of the eight settings have the first two coins matching: heads heads heads, heads heads tails, tails tails heads, tails tails tails. All four satisfy the record, because the record has nothing to say about the third coin. Pair each setting with its all-over flip, the way the first reading paired heads all round with tails all round, and the four collapse to two: one world in which the third coin agrees with the other two, one in which it differs. The record names both and picks neither.

Three readings, one question put to each, and the fibers came back holding one world, none, and two. One recipe produced all three counts. It takes three letters to write down. Write r for a reading, s for a state of the network, and R for the record map that strips a state down to its entries. The fiber of r is then every consistent s whose R(s) comes out r: keep the states that satisfy the entries, discard the ones whose entries read differently, count what is left over. Fix a map, name one of its outputs, collect the inputs that land on it, and that operation is called taking a preimage. A fiber is a preimage.

A fiber can have three sizes, counting a state and its all-over flip as one. The three sizes are three different situations rather than three points on a scale.

It can be empty. No consistent state produces this record, so the testimony describes nothing that could have happened. No amount of further questioning will produce a story that fits. The reading is **unrealizable**.

It can hold exactly one. The record pins down the world completely. Any two observers who hold the record hold the same world whether or not they ever compare anything else. The reading is **reconstructing**.

It can hold several. Every state in it satisfies every record. Nothing anybody wrote distinguishes them. The reading is **ambiguous**. This is the fiber a jury is looking at when it acquits somebody it privately suspects. The acquittal is the correct output: an ambiguous fiber has more than one member and the record does not name one.

Advocates argue about which of the three they are in. The defense wants the fiber shown to hold more than one member, and needs only one alternative story that fits every piece of testimony. The prosecution wants it shown to hold exactly one. A witness caught in an unrealizable reading has produced an account that no arrangement of the world satisfies, which is a result rather than a doubt.

Only the middle case convicts.

The twelve-patch net supplies all three. Its consistent record list is reconstructing: two coin settings satisfy it, they differ by flipping every coin in the world, and no meeting anyone could arrange would tell them apart. Flip one edge of the record, changing a single entry out of thirty from same to different, and the reading goes unrealizable: two of the nineteen independent loops carry a residue, and the number of coin settings satisfying all thirty records drops from two to zero.

For the ambiguous case, take away a patch's ability to talk. Its five seams go unread, so the record shrinks to twenty-five entries. The eleven patches that go on comparing are pinned exactly as before, and the silent patch's coin is not pinned at all: four settings of the twelve coins satisfy every record that exists, in two pairs that differ by the all-over flip. The reading leaves one bit undetermined. The bit has an address.

Repair cannot fix that. Repair works by reducing disagreement. There is no disagreement anywhere: every record is satisfied. A rule that picked one of the two would be picking on grounds no record contains, which makes it a choice rather than a repair.

11.3 Four demands

Run the machine on the flipped-edge record anyway. Chapter six did, from all two thousand and forty-eight distinguishable starting states and under all sixteen schedules. It converged every time onto one terminal arrangement with a leftover of exactly one. The machine halts. It halts in the same place from everywhere. And the state it halts in violates a

record, which is to say that halting and being consistent are two separate properties. The second one has to be checked by somebody.

The check is a procedure with a four-line specification. Take any state at all, read off its record, and return the consistent state that record names. It has to meet four demands. What comes back carries the record it was handed, since a procedure that revised the testimony on its way past would be answering a different question. A state that was consistent to begin with comes back untouched. Two states with the same record come back with the same answer. And when no consistent state carries the record at all, what comes back is a failure value, written bottom, rather than an answer. There is exactly one procedure meeting all four. It exists precisely when no reading is ambiguous.

The third demand is the one doing the work, and the one an ordinary repair procedure fails. Returning the same answer for two states with the same record forbids the procedure from consulting anything the record does not contain, including which state it was handed and which repairs it happened to run. Hand it an ambiguous reading and no procedure meets all four, because the third requires one answer and the second requires two.

Bottom is a proof that the fiber is empty, which a search that stops without finding anything does not supply. On the twelve-patch net you can hold the proof in your hand. Add up the entries around each of the nineteen loops in the arithmetic where one and one make zero. Two of the sums come out one. A loop summing to one demands that some coin differ from itself. Nineteen sums settle four thousand and ninety-six settings without a coin being set, and they name the two loops the obstruction sits in, which no amount of running through the settings would.

A normal form in chapter nine’s sense was what a process stops at, which made it a statement about the process. This one is a statement about the reading: the settled state is determined by the record rather than by the path, so it is an **observation-determined normal form**. Two observers who agree on the record agree on the world. They can be handed the record by anybody, in any order, having watched none of the repairs.

Which makes the settling two functions run one after the other. Write N for the whole procedure, the one handed a state and returning the world it settles into. The third demand says N consults what it was handed only through the record, so it comes apart into two steps:

$$N(s) \;=\; \mathrm{world}(R(s))$$

Here world is the lookup that takes a reading to the consistent state that reading names, and returns bottom when the fiber is empty. The state s appears on the right inside the record map’s brackets. The left-hand side reaches it through nothing else. Two networks holding different coins and writing down the same thirty entries settle into one world, and whatever

separated them at the start is a difference neither can carry forward, because carrying it forward would take an entry, and the entries match.

Nineteen sums decided the twelve-patch case because its constraints are linear. For constraint systems at large, deciding whether any consistent state carries a given reading is NP-complete: hand somebody a proposed state and they can check it against the reading in seconds, and no known method of producing one beats searching. The ambiguity question comes out the same shape. Two consistent states sharing one record can be put in front of anybody in a moment, and ruling out every such pair is the expensive half. The record determines the world. Working out which world is a second problem, its cost set by the shape of the constraints rather than by the length of the record.

11.4 Flip all twelve coins

Take the settled state of the twelve-patch net and turn every coin over. Every same stays same. Every different stays different. All thirty records are untouched, so the record map returns exactly what it returned before, and the two states are indistinguishable to every observer in the net, to every observer who joins later, and to any sequence of meetings anybody cares to arrange. That is why chapter six counted two thousand and forty-eight states rather than four thousand and ninety-six: the two members of each pair are one thing.

The silent patch's undetermined bit looks similar and is not. Flipping that coin also leaves every existing record alone, but the moment the patch's seams come back into use, somebody writes down a comparison that comes out differently. It is a difference this record fails to see. The all-over flip is a difference nothing could see, under any arrangement of comparisons, ever.

Changes nothing could see form a group, in chapter five's sense: doing two of them in a row is another one, doing nothing is one of them, and every one of them can be undone. The group has a job here that chapter five could only describe. **The symmetry group is exactly the set of changes the record map cannot see.** On the twelve-patch net that group has two members: leave the coins alone, or turn all twelve over.

That fixes what physics is allowed to talk about. Only quantities that survive a change of description may be called physical. Chapter five arrived at that sentence by watching what two accounts of one table had in common, and the settled answer enforces it: relabel every patch's internal state however you like, run the repair, and the result is the relabeled version of the result you would have got. The answer is a statement about whole families of descriptions rather than about any one of them, which is called **presentation invariance**. It means no internal labeling choice anybody makes can reach the physics.

So the word objective has an exact referent here. It is a state. Objective reality is the observable normal form of the network: the consistent state that the shared record picks out, up to changes nothing can see. There is no further fact about the world hiding behind it, because a further fact would be a difference no record distinguishes. Those are exactly what the group deletes.

11.5 Paths and fibers

Schedule independence and reconstruction sound alike. They are statements about different objects. The first does not give the second.

Confluence is a property of paths. It takes one starting state, lets two repair sequences run out of it, and says the branches come back together. Reconstruction is a property of fibers. It takes two starting states that happen to carry the same record and says they settle to the same place. The first says nothing whatever about the second, because the first never mentions a second starting state.

The silent patch is the counterexample, small enough to check. Everything about it is confluent: every repair sequence terminates, every schedule from a given start lands in the same place, and Newman’s lemma applies without amendment. Two observers who start from different coin settings and hold identical records land in different worlds anyway, because the silent coin is never touched by any repair and never recorded by anybody. Schedule independence in full, and a coin nobody can name.

Run the two together and you get a network that passes every confluence test there is and has no objective world in it. Any two observers can compare records forever without turning up a discrepancy. The thing they are converging on differs between them in a way no comparison reaches. The repairs are finished, the schedules agree, and there are two worlds.

The traffic runs the other way with a condition attached. A repair moves coins and leaves entries alone, so two branches out of one start carry that start’s record all the way down, and if both branches stop, a record that pins down one consistent state hands them the same world. Schedule independence falls out, provided every state has somewhere to settle at all. Three properties, then: branches out of one state rejoin, endpoints from different states with one record agree, and every state has an endpoint to reach. The first and third leave the second free to fail, which is the silent patch. The second and third force the first.

11.6 One over n

The trichotomy counts the members of a fiber and says nothing about how far apart they are. The distance is what decides whether the answer survives a closer look.

Instruments quote a number for exactly that, called sensitivity: how much of a difference in the world it takes to move the dial by a readable amount. A thermometer whose scale marks crowd together at the top of its range reads temperatures perfectly well and separates two nearby ones badly. A buyer who checks only that each mark has its own temperature has checked the wrong thing. A reconstruction is an instrument pointed backward, from the dial to the world, and needs its sensitivity quoted in that direction.

A reading can be refined. Resolve more seams and the record gets longer, and throwing away the extra entries hands back the shorter record you started with. Stack those refinements and you have a tower. The question is whether the answers at each level are converging on anything.

Build one. At level n there are two consistent states a full unit apart, and the record separates them: it reads 0 on the first and 1 over n on the second. Two distinct readings picking out two distinct states is reconstruction, so every level of the tower reconstructs. Then watch the levels go by. The readings converge on each other, the states sit a unit apart wherever you look, and the factor by which a difference in the record gets magnified into a difference in the state is n. At level ten thousand the record separates two worlds by one part in ten thousand. The limit has one reading and two worlds.

So being determined by the record is not the same as being determined stably by the record. What separates them is the largest factor by which a difference in the record can be magnified into a difference in the state, which is the conditioning of the reconstruction, and a tower of readings has a limit worth speaking of exactly when the conditioning is bounded all the way up rather than level by level. Chapter ten's curve went to zero across all 122,880 seams in the network, and 122,880 seams is one level. The bound is what says a finer one would have gone to the same place.

This is why a continuum is available at all. Space that looks smooth at the scale of a laboratory is a limit of finite readings. A limit taken through a tower with unbounded conditioning is an artifact of whichever level you stopped at. With the bound, different solvers, different refinements and different orders land on one limit. The smooth description is a thing about the world rather than about the grid.

11.7 Hold the boundary fixed

An accepted repair leaves the record it was handed exactly as it found it, so a settled state carries the record it started with, and what it settles into depends on that record from the first move to the last.

The settled state is unique given the boundary data, which is the part of the record fixed from outside the region doing the repairing. Every start carrying that boundary data lands in the same world. Change the boundary

data and the region settles into a different world, just as consistent, just as settled, and just as indifferent to who went first.

On the twelve-patch net the boundary data is easy to point at. Repair a region of the net and the entries on the seams that leave the region are handed to it from outside: the patches inside cannot change them, and every setting the region can settle into has to agree with them. Those entries are the region's boundary. The settled interior is a function of them. Two neighboring regions with different boundary entries settle into different interiors, and both are correct.

There is no single endpoint that the whole thing is heading toward. The terminal state is unique relative to its boundary record, and boundary records are not unique. Two different readings on the edge of a region produce two different settled interiors. Both of them are objective in the only sense the word has here, which is that every observer inside either one agrees with every other.

Which is the whole of what a jury does. It is handed a record it did not choose. The verdict follows from the record. Hand it a different record and a different verdict follows, with no less force. Twelve people, one transcript, and a verdict that belongs to the transcript rather than to any of them.

So look at what the settled world is made of. Thirty entries, each one a comparison between two patches, none of them a property of either patch on its own. The coins were the things in this world, in the sense that they were what the patches actually held, and the coins are the part that dropped out: two settings of them produce every entry identically, and the answer is stated about the pair rather than about either member.

Thirty entries survived the procedure. Twelve coins did not appear anywhere in the answer. The record map never once returned a coin.

Chapter 12

Why Is the World Made of Relations and Not of Things?

A box of gauge blocks holds 102 slabs of hardened steel, each finished to one exact length and so flat that two of them wrung face to face cling together. Stack the right handful and you have any length between 0.5 and 300 millimeters in steps of a hundredth of a millimeter. A Swedish machinist named Carl Edvard Johansson worked the set out in 1896 and patented it in 1901, and within twenty years it was what the world's engineering shops measured against. Every shop had its own box, and every box was checked against a better box somewhere else.

In the spring of 1918 Hermann Weyl proposed that the universe skips the last step: every point keeps its own box, and no box anywhere is the master. He sent Albert Einstein a paper arguing that the universe has no fixed unit of length. Einstein disagreed with it, presented it to the Prussian Academy of Sciences in Berlin anyway, and his objection was printed along with it.

Let every point of space set its own unit of length, the way every workshop sets its own gauge blocks, and comparing two lengths at different places then needs a rule for carrying a unit from one to the other. Weyl's rule was the electromagnetic potential. Electricity and magnetism would be the bookkeeping of a freedom that had been sitting unexamined inside geometry, the freedom to recalibrate the ruler at every point, and the German word for setting the scale on an instrument is Eichung, calibration. The word that came back through the translation is gauge.

Einstein's objection was short. If carrying a unit of length depends on the route it takes, two identical objects sent by different routes arrive with different sizes. Atoms are objects of that kind and the light they emit is their scale marking, so hydrogen that spent a billion years in the arm of a distant galaxy would emit at frequencies displaced from hydrogen in a Berlin discharge tube by an amount depending on where it had been. Astronomy consists in large part of recognizing elements in starlight by matching their lines against laboratory samples. The lines match.

Weyl argued back for years and gave the scale geometry up in the early 1920s, and the word outlived it. In 1927 Fritz London noticed that the factor Weyl had written works exactly as intended if it is made imaginary, at which point it stops rescaling the size of anything and starts rotating the phase of a quantum wave function, which no instrument reads off. Weyl rewrote the theory in those terms in 1929, in a paper on the electron and gravitation. The phrase gauge invariance came through the rewrite with its modern meaning attached. What had to change was the quantity chosen freely at each point, which has to be one that no measurement reaches. The length of a ruler fails that condition, as the spectral lines show.

Twelve observers wired into thirty seams have a freedom of exactly that kind, for reasons with no geometry in them. Each keeps its state in its own labels, and nothing outside the twelve fixes the labeling, because there is nothing outside the twelve. Watch what the settling then produces. The repairs run, the disagreement drains out of the network, every one of the thirty seams reports itself content, and if you walk round the twelve afterwards and read off what each of them is holding, nothing in the run has pushed those numbers together. The seams converged and the labels did not. What is it that the twelve of them agree about?

12.1 Three teams, one algorithm

Three teams are handed the same specification and told to implement it. One writes Python, one writes Java, one writes Rust. When the programs are done, the only comparison anybody can perform is to feed all three the same inputs and check the outputs, because that is the only place the three implementations touch. Everything else about them, variable names, memory layout, which loop runs backwards, is invisible to that test. Each team can choose it without consulting the others.

Chapter five made a symmetry the claim that part of the description was never physical. Here the choice gets made three times, independently, by parties who never see each other's source. A symmetry with one global choice is a claim about the whole world. A symmetry whose choice can be made separately in every place is a **gauge** symmetry, and twelve observers make twelve of them.

The picture has a client in it. The client is the part to delete. Somebody outside the three teams wrote the specification and knows what the program is supposed to do. Nothing plays that role here. The input-output behavior of the twelve observers is the only thing about them that exists. Each of them wrote itself. What is left when the client goes is three implementations whose agreement has nothing to appeal to but each other.

Local choice has a price and it is charged at the seams. Two patches sharing a seam compare port readings written in two private conventions. The comparison needs a translation. Chapter four supplied the shape of

it, with two firms settling one account in two currencies and reading their exchange rate off the transaction they both recorded. Put a dictionary of that sort on every seam. The consistency condition stops being "my reading equals yours" and becomes "my reading equals this seam's dictionary applied to yours."

A rule that lives on an edge and translates one end into the other is called a **connection**. Chain the dictionaries around a closed loop and you come back holding a translation from your own conventions into your own conventions, which is chapter six's holonomy, and the one thing it tells you is whether the trip returned you unchanged.

Notice where the information went. Each observer's own labels are its private business. The thirty dictionaries are the only content the arrangement holds that anybody can check. In the large runs the freedom each patch holds is one of the six ways of permuting three labels, and every seam's dictionary is one of the same six. If a patch decides to swap the names of two of its labels, the dictionary reading into it picks up that swap, the dictionary reading out of it picks up the reverse, and every round trip through the patch comes back with what it came back with before. Redenominate one city's currency and each rate into and out of it changes, while the round-trip factor does not move.

12.2 One number for the whole arrangement

Take the wiring itself, and give each observer the freedom London's rewrite left standing, an angle on a dial rather than one of six permutations. Twelve ports, thirty seams, five seams meeting at each port. A field on that graph is a number on each of the thirty seams. A relabeling is a number at each of the twelve ports, and what it does to a seam is add the difference between the numbers sitting at the seam's two ends.

Some fields can be manufactured that way out of nothing and some cannot. Every field splits into those two parts. The part no relabeling could have manufactured is the collection of loop readings. That part is the field strength, because it is the part a measurement can reach.

The split shows up in what two observers can argue about. A seam value is something each of them wrote in private. The gap between their two numbers closes if either one changes a convention. A loop reading is the result of a round trip. No convention either of them can change will move it. Relabel one port and all five seam values touching it move, while the round trips through it come back exactly as before.

Then a source. Charge moves and the source has to move with it, so the source is a current: one number on each seam, saying how much runs along it and in which direction.

Score the arrangement with a single number: the energy held in the field strength, less the thirty products of each seam's value with the current

running along it. A number scored over a whole arrangement like that is called the **action**.

Apply a relabeling and work out what the score does. The first term does nothing, since the field strength was built to ignore relabelings. The second term moves, and the amount it moves by is one line of arithmetic: for each of the twelve ports, take the number that port chose and multiply it by the imbalance in the current there, meaning what flows out along its five seams less what flows in, then add the twelve products.

Both directions of the theorem are sitting in that sum. If the current balances at all twelve ports, every product is zero and no choice of labels moves the score, whatever numbers the observers pick. If the current fails to balance at even one port, the observer at that port can pick a number that makes its product nonzero, and the score moves. Send three units out of one port along its five seams and two units back in. The imbalance there is one. Let that observer add seven to its own labels, and the score of the whole arrangement moves by seven while the eleven other products sit at zero.

The second direction is the one doing work. A current that fails to balance somewhere gives the arrangement a score that depends on conventions the twelve observers picked privately, and any one of the twelve could then move that score by rewriting its own labels, having changed nothing whatever about the world. An arrangement like that has no score at all.

Charge conservation and gauge symmetry are one statement read in two directions. Both directions came out of the same twelve products.

Look at what went into that and what came out. In went a wiring diagram of twelve ports and thirty seams, a rule saying which changes count as relabelings, and a source. Out came a restriction on what a source is allowed to be, with a conservation law as its content. Nothing about electricity was assumed anywhere in the argument. The only geometry in it is a list of which port is wired to which.

12.3 Nineteen loops

Count what the labels can reach. There are twelve dials to turn, one at each port, and turning all twelve by the same amount changes no seam, so eleven of the twelve directions do anything at all. Thirty seam values, eleven directions of pure convention, nineteen left over.

Nineteen turned up in chapter six as well, off this same wiring, out of a rule with three pieces of arithmetic in it: take the edges, subtract the dots, add one. Thirty seams, twelve ports, and thirty minus twelve plus one is nineteen.

Run the rule on two drawings small enough to check by eye. Sketch six observers in a row, which is five seams and six dots. Sketch two triangles

sharing an edge, which is five lines and four dots. Do both subtractions in your head and keep the second answer.

Count the loops in the second drawing and the eye finds three: the left triangle, the right triangle, and the four-sided circuit around the outside. Walk the left triangle clockwise and then the right one clockwise. The shared edge gets crossed once in each direction and cancels. What survives of the two trips is the circuit around the outside. The third loop is the first two performed in sequence, two of the three are independent, and the drawing gives no hint as to which two. The eye overcounted by one on a figure with five lines in it, which is the argument for doing the arithmetic on a figure with thirty.

Three drawings, one rule, and the arithmetic for all of them fits on a line.

$$5 - 6 + 1 = 0 \qquad 5 - 4 + 1 = 2 \qquad 30 - 12 + 1 = 19$$

The row of six has nothing to walk around, the two triangles carry two independent loops where the eye found three, and twelve observers wired into thirty seams carry nineteen.

Nineteen does two jobs. It counts the sums that have to come out zero before the twelve can settle on anything, and it counts how much of the wiring is physics rather than paperwork.

The two nineteens are the same nineteen. Strip every label choice out of a field and what survives is nineteen numbers, one for each independent loop, and nothing else about the thirty seam values survives at all. Two fields with the same nineteen loop readings are one field written down twice.

Those nineteen numbers are also the only things any two of the twelve can both compute and both check. A round trip begins and ends at one port, so the observer sitting there can work it out from the dictionaries alone, without being told what anybody else calls their own state and without taking anybody's word for anything. The dictionaries sit on the seams where both ends can read them, and the shortest of the round trips chains three.

12.4 The laziest field

Give the numbers on the seams a different job. They stop being anybody's private labeling and become the field itself. A field costs energy: square each seam's number and add the thirty of them. The law tying charge to that field reads at the ports: the net outflow along a port's five seams equals the charge sitting there. Give it a list of twelve charges and it answers with a family of fields rather than one, because adding a pure circulation around any loop leaves every port's outflow alone. The family is nineteen-dimensional, one dimension for each independent loop.

A law that answers with a nineteen-dimensional family has not finished the job. Two further conditions each pick out one member of it. They pick the same member. Exactly one field in the family has zero reading on every loop. Exactly one has the least energy. They are one field for a single reason: the energy of any solution equals the energy of that one plus the energy of the difference between the two, and the difference is pure circulation, which costs energy unless there is none of it.

The field around a charge is the solution that circulates nowhere. Every other member of the family is that field with a circulation laid on top of it, which costs energy and transports nothing. The operator that selects against it is chapter nine's repair rule, my value against the average of my neighbors, running on the twelve ports of this same graph, and the only reading that rule leaves alone is the one that is the same at every port.

On a twelve-port graph all of this is exact arithmetic rather than approximation. Put one unit of positive charge at a port and one unit of negative charge at a neighbor. Send the whole unit straight down the seam that joins them and the arrangement costs one, one squared being one. Let it settle into the field that circulates nowhere and the same two charges cost eleven thirtieths, and the nineteen thirtieths that went missing was circulation delivering nothing to either port. Eleven over thirty and nineteen over thirty, off the same wiring that gave eleven directions of convention and nineteen loops.

12.5 One hundred and two thousand three hundred and thirty-three seams

The same behavior is visible in the large runs, where nothing about it was arranged. In the run of chapter ten, 81,920 observers wired into 122,880 seams, the raw labels began by disagreeing across 102,519 seams and ended disagreeing across 102,333, while the transport-corrected disagreement began at 102,415 and ended at zero. The dictionaries themselves were never touched on the way: a repair rewrites the label at one end of a seam and leaves the translation between the ends exactly as it found it.

At the finish, 102,333 of the 122,880 seams have two ends holding different labels, and no seam in the network is in disagreement. Both counts came off the same run.

Count the other way at the end of that run. The seams whose two ends held identical labels numbered 20,547, and the seams whose dictionary was the permutation that does nothing numbered 20,547, and they were the same seams. Two ends of a seam carry the same label exactly when the translation between them happens to be the do-nothing one, which is a coincidence of conventions with no content in it. The other 102,333 seams were in perfect agreement and said so in different words.

What settles is the relation, never the values. It could not have been the values. The labels are each observer's private business, no procedure anywhere in the arrangement reads them, and a process that drove them together would be enforcing a convention that nothing can check.

12.6 The residue in the loop

Chapter six ended with twelve observers, one flipped record, and a leftover of exactly one unit of disagreement that no repair removed and no ordering avoided. Repairing in a different order moved it somewhere else and left the total at one wherever it came to rest. It sat in the loops.

The nineteen loop readings are where it lives. A relabeling cannot reach it, because relabelings leave round trips alone, the way redenominating a currency leaves the arbitrage factor where it was. Repair cannot flush it, because repair works one seam at a time and no single seam holds it. A loop that comes back carrying a residue is a loop with something inside it. No rewriting of anybody's conventions disposes of it, because the conventions have been divided out of the reading before the residue is read.

That residue is charge. The quantity a current has to conserve for the arrangement to have a score at all, the quantity the law at the ports reads as an outflow, and the quantity trapped in a loop that fails to close are one quantity. A loop carrying a residue cannot be talked away as somebody's bookkeeping, because the bookkeeping was divided out of the reading before the reading was taken, and what survives that division has to be paid for in energy and balanced at every port. Weyl proposed in 1918 that electromagnetism is the bookkeeping of a freedom chosen separately at every point. He had the wrong freedom, and the bookkeeping was right.

Thirty seams. Twelve ports, five seams at each. Eleven directions of pure convention, nineteen loops, and one unit of charge sitting in the loops where no relabeling reaches it. Every number in that list was read off a wiring diagram. Nobody has opened one of the twelve observers to see what a port is.

Chapter 13

What Is One Piece of Reality Actually Made Of?

The standard way to test a circuit board used to be a bed of nails: a fixture with a spring-loaded probe for every pad, which pressed the board down onto several hundred needles at once and read off which connections had taken solder. It worked while the connections were where a needle could touch them.

Then the boards got dense. Traces went down into inner layers, and chip packages arrived with their contacts in a grid underneath the body of the part instead of along its edges, so the pins of a soldered-down chip sat sealed in a sandwich a probe cannot enter. The nails had nowhere to land.

In 1985 a group of electronics firms formed the Joint Test Action Group to deal with it. Their answer gave up on reaching the pins and asked the chip to report itself. Put a small register cell beside every pin, chain those cells into one long shift register running around the boundary of the die, and add wires that let an outsider clock the chain out a bit at a time. In 1990 the arrangement became IEEE Standard 1149.1, titled Standard Test Access Port and Boundary-Scan Architecture. The port takes four wires: a test clock, a mode select, data in, data out, with an optional fifth for reset. Behind those four wires sit several million transistors, and the standard says nothing about them.

Hand somebody an unfamiliar board and the first thing they look for is the port: four or five pads in a row on a tenth-of-an-inch grid, often with no header soldered into them. Those pads say the board expects to be interrogated and on how many lines, before anybody knows what a single chip on it is.

Chapter twelve left twelve observers wired into thirty seams, nineteen independent loops, and one unit of charge sitting where no relabeling reached it. All of that came off a wiring diagram. A wiring diagram says which port is joined to which port and says nothing whatever about what sits behind either one.

Open one. The question is what a piece of the world is made of when you get all the way down. The answer comes back as six parts and six moves.

13.1 The parts list

Start with the thing that holds a state. A **patch** is a machine small enough to be described completely: a bounded amount of internal state, a set of interfaces, and a finite list of moves it is allowed to make. Bounded has been in force since the definition of an observer. A patch that could hold unlimited state would have an unlimited amount to compare. The comparison would never finish.

A **port** is one place where a patch exposes part of itself to whatever is outside it. What appears at a port is a **packet**: a reading, drawn from a finite set of possible readings, that anything on the other side can pick up. The packet is not the patch's state. It is what the state looks like from the outside at that one interface, in the way that the cell beside a pin holds what that pin is doing rather than the internal state of the die.

Suppose a port could show any value on a continuous dial. Reading it to a hundred decimal places takes a hundred digits of storage in the reader, reading it exactly takes an infinite amount, and a machine with bounded state cannot hold either. A neighbor who writes down the first four digits and drops the rest has not read that port. It has read a different, coarser port with a finite list of settings, which is the port the architecture had all along.

Two ports routed to each other make a **seam**. Chapter six drew these as edges and counted thirty of them. A seam is where two patches read the same thing twice, once from each side, the only place in the architecture where a disagreement can be detected at all.

Everything any patch can ever learn about its neighbor arrives through the seams they share. Collect all of it, every reading available at every interface an observer has, and what you are holding is that observer's **screen**. A screen is where an observer's accessible information lives, in the same sense that the boundary-scan chain is where a chip's pin states live. It shows nothing to anybody, it faces no direction, and there is no seat in front of it.

An observer assembled out of several patches has a screen too, the readings on the ports facing outward from the assembly. The ports that patches inside the assembly share with each other are internal wiring to everything beyond it. So the size of a screen is a count of readings rather than a width in meters, and asking what an observer's screen looks like is asking how many distinct things its outward ports can be doing at once.

Last, a **record**. A patch that completes a move writes the outcome into the part of its state that later moves are required to leave alone. Chapter

three priced this and chapter nine identified what gets written, which is the accepted repair. A record is a write that survives the next read, and survives it because the rules of the machine protect that region of the state rather than because writing is somehow permanent.

Six words, and every one of them names something you could point at on a board: the machine, the interface, the reading at the interface, the join between two interfaces, the set of all the readings, and the part of the state that stays put.

13.2 Six moves and no others

A patch does one thing, over and over.

It reads the packets exposed on its ports. It compares them against its own, through a rule saying which of its readings answers which of the neighbor's, since two patches keeping their state in private conventions have to be told what corresponds to what before a comparison means anything. That rule is chapter twelve's dictionary on the seam. The technical word for whether the comparison is possible at all is commensurability. Then, if the comparison came back unequal, the patch chooses a move from a finite menu of allowed local updates, one entry off a printed card. It writes a record of the move it made. It exposes the resulting packet at its ports, so the neighbor's next read sees the new state rather than the old one. And any claim about what it did is settled by an exact check run on the exposed data, by anything that has access to that data, and the patch's own account of its work does not count.

Read, compare, choose, write, expose, verify. That is the entire operational repertoire of a piece of reality. No move reaches a patch on the far side of the network, none of them consults a clock, and the menu they are drawn from has an end to it.

Chapter twelve's large arrangement was running that loop at 122,880 seams at once. A patch reads what its neighbor has put on their shared seam. It applies the dictionary sitting on that seam, which tells it which of the neighbor's three labels answers which of its own. Suppose they do not answer each other. The menu on that patch holds the six ways of permuting three labels, and the entry that gets taken is the one the dictionary names: the neighbor's label carried across the seam and written down at this end as its own. The dictionary is not on the menu, and no move in the run rewrites it. The patch takes one entry, writes down that it took it, and puts the resulting packet back on the port for the neighbor's next read. Nothing in that sequence required knowing anything at all about the far side of the network, which is the property that lets 81,920 of these run at once with no coordination between them.

The sixth move is the one engineers recognize on sight. A patch can be wrong. It can have run its comparison against a stale packet, or chosen a

move that was not available, or written a record that does not follow from what it read. What settles the matter is a check performed on the readings themselves, which either passes or does not, and which anybody holding the same data performs identically. Anything that runs that check is a verifier. The data it runs on is chapter eleven's list everybody can check, the seam entries and nothing private to either end of a seam, so a verifier needs no access to the inside of the patch whose work it settles.

13.3 Adding up the seams

Each seam can score itself. Take the two packets, apply the seam's dictionary to one of them, and produce a number that is zero when they answer each other and positive when they do not. That number is local: it involves the two ends of one seam and nothing else in the world.

Sum the seam scores and accept a move only when it lowers that sum. The sum takes finitely many values, and a quantity that strictly falls through finitely many values can fall only finitely many times, so the moves terminate.

Chapter ten watched that number come down. Of the 122,880 seams in that run, 102,415 were scoring above zero at the start, and at cycle 1 out of 16 none of them were. The settling curve counted the seams contributing anything at all. The quantity the machine was working against was their sum.

So every sequence of repairs stops. Wherever it stops, no patch has a move available: each one has read its ports, compared, and found nothing its menu can improve. That is a local normal form, exactly as much as the descent argument buys.

Everybody stopping and everybody stopping in the same place are two different claims. The score establishes only the first.

13.4 What the second claim costs

Three more conditions, and each is a property the interface either has or lacks.

The first is that the repair menu is complete: for every mismatch a seam is able to report, the menu contains a move that clears it. Chapter nine showed what a missing entry looks like. Twelve observers passing units across seams, and a rule that acts whenever two neighbors differ by two or more. In 302 of the 1,352,078 arrangements, no seam has a gap of more than one, no rule can fire, and the arrangement sits lopsided. Every patch content, every local check passing, and the world not settled. The gap of one had no entry on the card.

The second is about the inside of a patch rather than the seam. When a repair changes what a patch exposes at a port, the patch has to have an

interior state consistent with the new reading. If some admissible boundary value has no interior that produces it, then clearing the seam breaks something inside, repairing the inside changes what the port exposes, and the two repairs chase each other around the boundary forever. The condition that rules this out is a gluing condition: every value a port can legally be asked to display extends to a consistent state of the patch behind it.

A third requirement comes out of the relabeling freedom of chapter twelve. Two repairs leaving one state have to be reconcilable, which is chapter ten's diamond, and the diamond has to close in the description with private conventions divided out. Two patches that relabeled themselves differently have landed in the same place as far as anything checkable goes, so closing the diamond in the raw labels is the wrong test in both directions: it can fail where the world agrees, and pass where the world does not.

With the menu complete, the gluing condition holding, and the diamond closing where conventions have been divided out, the terminal state of the network is the same whatever order the repairs ran in. Chapter ten proved the shape of that argument on a shelf of books and a mountain with three basins. This is what it costs in parts: a scoring rule on every seam, a printed menu with no gaps in it, and interiors that can support every boundary reading their ports allow.

Two patches on one seam gave chapter ten two endings out of one start. The score in that arrangement runs down to zero, every move taken is an entry on the card applied to a mismatch that was there, and the sixth move's check passes on every reading either patch exposed. The descent supplies all of that and leaves both endings standing. A gap in the menu and an interior that cannot support its boundary are both printed into the card and the wiring before the first repair runs.

Those three requirements are the **synchronization contract**: obligations on the parts. Any assembly that meets them synchronizes. The descent gets the machine to stop. The contract is what makes the place it stops one place rather than a place per schedule, which is to say that whether there is a single world at the end of the repairs is a question about the connector.

13.5 Four things that have to travel

A host needs maintenance, so the live machine running on it is stopped, written down and started again on another host, with the programs inside carrying on through the move. Server operators do this on a schedule. What makes it work is a decision about which parts of the machine have to travel and which parts can be left behind and rebuilt, and getting the list wrong shows up as a program that resumes into a world it does not recognize.

A machine that is interrupted and resumed is the same machine when enough of it has been put back. Enough, for an observer, is four things. Its records, meaning the writes that later moves protect. Its accessible state, meaning the state of what it can actually reach. Its interfaces, meaning which ports it has and what is routed to them. And its future law, meaning which moves and which readings its machinery allows. Those four together are a **checkpoint**.

The fourth item is the one that gets forgotten. Restore an observer’s records and its accessible state into a machine whose menu of moves is different, and the restored thing reads the same history and takes different steps from it. The law that says which moves are available has to travel with the records, or the records land somewhere that treats them differently.

Two checkpoints that agree exactly on all four give the same probabilities to everything the observer subsequently sees, for every event it is able to register. If the two restored states differ slightly, the two futures differ by at most as much, which is what makes a rebuilt observer usable when the copy is imperfect.

A great deal of what a running system holds is outside that list. Worker identifiers. Queue positions. Retry counters, repair-cycle indices, timestamps, and the latency of individual packets. All of it is real, all of it is needed to replay the machine’s history, and none of it is part of the observer or of the observer’s time.

13.6 A code with no distance in it

Look at what the seams do to the space of configurations. Each one demands that its two packets answer each other under its dictionary. A configuration that satisfies every one of those demands at once is legal, and one that fails any of them is not. Chapter twelve’s arrangement had 81,920 patches holding one of six labelings apiece, which is finitely many possibilities, though writing out their number takes 63,747 digits, and that is before any seam dictionary is counted.

An object of that description has a standard name in engineering: it is a **constraint code**, the same kind of thing as the codes that protect a hard drive or a deep-space link, where a handful of parity checks pick the legal words out of all the possible words. This carrier is a finite constraint code.

What a code usually arrives with is a distance. Send every bit three times and the legal words are 000 and 111, which differ in three places. Corrupt one symbol and you hold 010, which is one step from 000 and two steps from 111, so the nearer legal word is the one that was sent and the error is corrected rather than merely noticed. Corrupt two symbols and 011 is nearer to 111, and the correction produces the wrong bit. The number three is the distance of that code, and the distance is what says how much the code can fix: a distance of three corrects one error, a distance of five

corrects two, and a distance of two catches an error without being able to say which symbol moved.

The architecture supplies the checks and does not supply the distance. Nor does it supply a bottleneck across which information has to squeeze, a rate at which disagreement mixes through the network, or a bound on how many seconds settling takes. Each of those is a property of a particular wiring diagram, and has to be computed for that diagram, from the diagram. The six moves say which configurations are legal. How far apart two legal configurations sit depends on which port is joined to which port.

13.7 What one boundary can hold

Take a patch and ask what anything outside it can determine about it.

Everything arrives through the ports. What arrives at a port is one packet from a finite set. So two interior states that expose the same packets on every port are the same state to every neighbor, to every neighbor's neighbor, and to the whole network. They can differ internally in any way you like. The difference has nowhere to be read.

Which turns the amount that can be known about a region into a count of arrangements of its boundary. The number of values one port can show, multiplied by itself once for every port the patch has. In the runs of chapter twelve the freedom each patch held was one of the six ways of permuting three labels, so a reading in that arrangement is one of six things, and the bound is six multiplied by itself once per port.

Assemble many patches into one observer and the same argument applies to the assembly. Adding a patch deep inside adds state and adds nothing to what an outsider can read, because every port it has is routed to another patch inside the assembly. The count moves only when a patch puts a port on the outside. Two assemblies with the same outward-facing ports and very different interiors are one object to everything beyond them, with the same list of possible readings.

The interior does not appear in that product anywhere. However much a patch holds inside, and however finely it holds it, what an outsider can distinguish is bounded by the readings on its boundary. That is half of the area law: the information associated with a region of space goes as the area of the surface around it rather than as the volume inside, and black holes are where that was first noticed. The other half is a number, how much information one unit of boundary area holds, and fixing that number takes a horizon and a thermodynamic argument. Getting this far took a machine with a bounded interior and finitely many ports, and nothing about gravity at all.

Six, multiplied by itself once per port. The six is an accident of one arrangement's three labels. The exponent is a count of holes in a connector. The six moves do not fix it.

The connector is a closed surface, its ports have to be arranged so that every one of them sees the same thing, and the seams have to pair up with nothing left over. Impose them all at once and the arithmetic permits one count and no other.

Count the ports.

Chapter 14

Why Twelve?

In the Egyptian galleries of the Metropolitan Museum of Art there is a lump of serpentinite the size of a plum, 3.2 centimeters tall, cut into twenty triangular faces with a Greek letter carved into each one. A missionary named Chauncey Murch picked it up in Egypt at some point between 1883 and 1906, the museum bought it from his family in 1910 with money Helen Miller Gould put up, and the label allows the object itself anywhere from the second century BC to the fourth century AD. Nobody knows what it was for. The museum's own entry on it offers an alphabet oracle, a text with one line written out against each letter, where the letter that lands face up picks the line that is yours.

Whoever cut it was working to one number. The number was twenty. Twenty faces, twenty letters, twenty answers.

Count its corners. There are twelve, and five faces close around every one of them. Nobody in Ptolemaic Egypt sat down and picked twelve. Twelve is what happens when twenty triangles close up with five meeting at each corner, and any craftsman in any century who set out to make that object would have arrived at the same twelve. The same object sits on shelves in a good many houses in injection-molded plastic with the numbers one to twenty on its faces. Its corners come to twelve as well.

Chapter thirteen opened one piece of the world and found a boundary with ports on it, packets going out through the ports, seams where two patches meet, and a verifier deciding claims. The question the boundary leaves standing is how many ports it has. That count is settled the same way the twelve corners in the museum case are settled, by four conditions on how the pieces meet and some arithmetic anyone can do on paper.

The conditions are these. The boundary is cut into triangular faces, since three is the smallest number of patches whose comparisons close into a loop. Every seam has exactly two faces along it, because a seam is where two patches meet and a third would have nothing to meet across. The faces around any port close into a single ring with no loose end anywhere, because there is no edge of the world for a loose end to hang off. That ring is five faces long. And any loop of comparisons you can walk on the boundary is the rim of some collection of faces, with none left over that the faces cannot reach.

14.1 Count it twice

A theater with thirty rows of twenty-four seats has seven hundred and twenty seats, and counting the same room column by column, twenty-four columns of thirty, gives seven hundred and twenty again. The two answers agree because they are answers about the same seats. Counting one collection two different ways and setting the two results equal is what settles the port count.

The collection to count is seam-ends. Every port has five seams running out of it, so the number of seam-ends is five times the number of ports. Every seam has two ends, so the number of seam-ends is twice the number of seams. Five times the ports equals twice the seams. On the object in the museum case that reads sixty seam-ends counted corner by corner, and thirty edges with two ends apiece, which is the same sixty.

The second collection is face-sides. Every face is a triangle and has three sides, so the number of face-sides is three times the number of faces. Every seam has a face on each side of it and supplies exactly two, so the number of face-sides is twice the number of seams. Three times the faces equals twice the seams.

Two equations, and feeding one into the other leaves everything standing on the port count. If the ports number P, the seams number five P over two and the faces number five P over three. Both of those have to come out as whole numbers, since the boundary cannot have half a seam, so P is even and P is divisible by three. Before any question of shape arises, the number of ports is a multiple of six.

14.2 Nineteen loops, twenty faces

Chapter six counted the independent loops in a wiring diagram of ports and seams. Lay a route that reaches every port and never closes on itself, which uses exactly one seam fewer than there are ports, and then put back the seams the route did not use: each one closes exactly one new loop. Seams, minus ports, plus one. On the twelve-port wiring that came to nineteen, and chapter twelve found the same nineteen again from the other end, as the nineteen loop readings that survive when every observer's private labeling is divided out.

Count the faces of the same object. Twenty.

Nineteen and twenty sit one apart. The reason they do is the entire content of the word spherical.

Take the twenty faces and add up their boundaries in chapter six's arithmetic, where two of anything cancel. Each seam lies along exactly two faces, so each seam gets counted twice and drops out, and the grand total is nothing at all. So the twenty face boundaries carry exactly one relation between them: any nineteen of them determine the twentieth, and

no smaller collection of faces adds up to nothing, since a collection that stopped short would have a seam with a face on one side of it and nothing on the other, and that seam has nothing to cancel against. Twenty faces supply nineteen independent loops.

The wiring has nineteen independent loops. The faces supply nineteen independent loops. Those two nineteens are the same nineteen. Every closed walk across the ports is a sum of triangles.

That is what being a sphere consists of, checkable from inside the object without stepping back to look at it. Cut the surface of a doughnut into triangles and two loops on it are the boundary of nothing at all: the loop that goes around the hole and the loop that goes through it. Take any collection of triangles on the doughnut and its boundary misses both of them. On a doughnut the faces fall two short of the loops. The shortfall runs two for every handle.

So for a boundary that closes into a sphere, faces minus one equals seams minus ports plus one. Move the terms about and the same statement reads: ports, minus seams, plus faces, equals two. Cut a sphere into a million triangles of wildly unequal size and the combination comes out two. Cut a doughnut any way at all and it comes out zero.

Each of the three counts on its own depends entirely on how the cutting was done. Drop a new port into the middle of one face and join it to that face's three corners: the ports go up by one, the seams by three, the faces by two, and one minus three plus two is nothing. The combination is unchanged. That is an invariant in chapter five's sense, a quantity left alone by a family of changes. The family here is every way of cutting a surface into triangles. What survives all of them is one small whole number that counts holes. The name it goes by is the Euler characteristic of the surface.

14.3 Twelve, thirty, twenty

Two independent relations on the same three counts, and they meet. Take the port counts the whole-number test left standing, fill in the seams and the faces from the two double counts, and do ports minus seams plus faces on paper. Six ports carry fifteen seams and ten faces, and six minus fifteen plus ten is one. Twelve ports carry thirty seams and twenty faces, and twelve minus thirty plus twenty is two. Eighteen carry forty-five and thirty, and eighteen minus forty-five plus thirty is three. One, two and three, against six, twelve and eighteen.

The letters say the same thing and say why it happens: ports minus seams plus faces is P minus five P over two plus five P over three, which over a common denominator of six is six P minus fifteen P plus ten P, all over six, and that is P over six. The counting alone, with nothing about

shape anywhere in it, ties the port count to the one combination a sphere pins down.

$$\text{ports} - \text{seams} + \text{faces} = \frac{\text{ports}}{6}$$

The left side is two for anything that closes into a sphere, so the right side is two, and the ports number twelve. Feed twelve back through the two double counts: the seams number thirty and the faces number twenty. That is the icosahedron chapters six through twelve have been running on, arrived at from four conditions and a page of arithmetic.

Run the argument backwards. Twelve ports with five faces around each of them make ports minus seams plus faces equal to a sixth of twelve, which is two, and among closed surfaces the value two belongs to the sphere alone. Two is the largest value any closed surface reaches. The only other positive value is one, belonging to a one-sided surface that cannot sit in ordinary space without cutting through itself. Twelve ports force a sphere, and a sphere forces twelve ports.

14.4 Four around a point, six around a point

Every condition in that argument does work. Three of the four can be broken and leave a boundary behind. The fourth cannot: put a third face along a seam and the seam has no two sides, so there is nothing left to run a count on. Break the other three one at a time and watch the twelve go.

Change the ring around a port from five faces to four. Run the identical double counting: four times the ports equals twice the seams, three times the faces equals twice the seams, and ports minus seams plus faces comes out at a third of the ports. A sphere sets that combination to two, so the ports number six. That object exists, has six ports, twelve seams and eight faces, and sits in every mineralogy display case as an octahedron, the shape a fluorite crystal cleaves into. Three faces around a port gives four ports, which is the shape of the smallest die in the box.

Six faces around a port drops the port count out of the arithmetic entirely. The double counting gives six times the ports equals twice the seams, and the combination collapses to zero times the ports, which is zero however many ports there are. Zero is the doughnut's number and the sphere's is two. Six triangles around every point is the flat triangular grid, which runs across a tabletop forever and closes up into a doughnut of any size, and never closes into a sphere.

Break the spherical condition instead. Take the twenty-faced object and glue every point of it to the point diametrically opposite, so that walking off one side brings you back on the other. Ports pair up and there are six of them left, seams pair up and there are fifteen, faces pair up and there are ten. Six minus fifteen plus ten is one. Five faces ring every port. The

surface has one side and cannot be a sphere. Its port count is six, and a sixth of six is one.

Break the triangles. Pick up the six-sided die from the same shelf and try to find a triangle in its wiring: start at a corner, walk along an edge, walk along another, and see whether a third edge takes you home. It never does. On a cube, no two neighbors of a corner are neighbors of each other, so there is no face anywhere in it whose three corners are pairwise joined. A cube does not fail the count. It fails to admit the structure at all. Relabeling its eight corners cannot repair that.

14.5 Sixty rotations

Chapter five had you cut a paper square and find every way of putting it back on its own outline, and eight came out. Do the same thing to the twenty-faced object, turning it without ever reflecting it, and count the ways it lands on itself.

Every one of those turns leaves the object standing where it stood and shuffles its parts underneath. A turn carries every port to a port, every seam to a seam and every face to a face, and a collection of moves that shuffles a set of things among themselves in that way is a **group action**. One collection of turns is acting on three different sets here. The same sixty turns are doing all three at once. Chapter five also left a word for the set of places a single thing can be sent to under such an action, which is its orbit. Every port here can be carried to every other, so the orbit of a port is all twelve. Ask the opposite question. Which turns leave one chosen port sitting exactly where it is? Only the ones that spin the object about the line through that port, and since five faces ring it, five such turns bring the ring onto itself, counting the do-nothing turn. The set of moves that leave a chosen thing where it is is its **stabilizer**. The stabilizer of a port has five members.

Every turn in the whole collection is accounted for exactly once: send the chosen port to one of its twelve destinations, then spin about the destination in one of five ways. Twelve times five is sixty. That multiplication is the general fact, and works from any starting point: the size of an orbit times the size of a stabilizer is the size of the group.

Which means the count can be run backwards through any feature of the object, and run with the object in your hand.

Pinch a twenty-sided die between finger and thumb at the centers of two opposite faces and turn it about that pinch. A third of the way round it fills its own outline again, because the triangle under your thumb has three corners and takes each of them in turn, which is three positions in a full turn counting the one you started in. Move the pinch to two opposite corners, where five triangles meet under each fingertip, and turn it again. Five positions. Take the awkward grip, at the midpoints of two opposite

edges, and there are two, the second being the half turn about the line through those midpoints.

Three, five and two, read off with a hand rather than argued. Every face can be carried to every other face by some turn in the collection, and every edge to every other edge, so each of those three numbers is a stabilizer sitting under a full orbit. Multiply each one by the number of things of its kind on the solid.

$$20 \times 3 = 12 \times 5 = 30 \times 2 = 60$$

Twenty faces held three ways apiece, twelve corners held five ways, thirty edges held two ways. Three counts that share nothing except the solid they were taken from, and one answer. The solid reports the size of its own symmetry three times over. The port count comes back out of it as sixty divided by five.

Nobody picked those sixty turns. Four conditions on how comparisons meet forced twelve ports, and the twelve came with sixty turns attached, as much a consequence of the conditions as the corner count is. The symmetry belongs to the boundary the way the corner count belongs to it, arrived at rather than assumed, and every rule that can be written on such a boundary has to come back to itself under all sixty turns. Most rules do not.

Sixty moves carrying twelve ports around means no port is distinguished from any other by anything the wiring holds. Any port can be sent to any other, and having sent it there, its ring of five neighbors can be spun into any of five positions, so a description of the boundary that starts at one port and works outward could have started anywhere and come out the same. Two patches comparing readings through a seam are comparing quantities of the same kind. No calibration step anywhere in the machine had to establish it.

Fifteen of the sixty are half turns, one for each pair of opposite seams. They fall into five families of three, each family being three half turns about three lines at right angles to each other, and every turn in the group shuffles those five families among themselves. Follow the shuffling and you get sixty permutations of five objects. That is the group behind chapter five's fifth-degree result, the one whose shape rules out a solution formula of the sort the quadratic has, and it is contained in the turns of a twenty-sided die.

14.6 Ten thousand three hundred and ninety-five ways to pair up twelve things

Twelve ports. Nothing in a list of twelve labels says which of them has anything to do with which.

Pair them up, six pairs with every port used once. Take the first port and choose its partner from the remaining eleven. Take the lowest unpaired port and choose from the nine left. Then seven, then five, then three, then one. Eleven times nine times seven times five times three is ten thousand three hundred and ninety-five. That is how many pairings twelve labeled things admit. Every one of them is as good as every other, so long as the twelve are only labels.

Put the wiring back and all but one of the pairings dies. Start at any port and count how far the others are, in seams, along the shortest route. Five ports are one seam away, which is the ring around it. Five more are two seams away. That accounts for eleven of the twelve, counting the port you started from. The twelfth is three seams off. Chapter nine leaned on that three when it put a ceiling on how far news has to travel, and every port has exactly one port at distance three, never two.

One pairing out of ten thousand three hundred and ninety-five survives that, the one matching each port with the single port farthest from it. It uses every port once and gives six axes.

The same six pairs can be found without counting a single distance. Take any two ports and count the ports adjacent to both of them. Sixty of the sixty-six pairs come back with two ports in common. The other six come back with none, the six axes again, each one a pair of ports that share no neighbor anywhere in the wiring.

Any relabeling that preserves the wiring preserves distances, so it carries the far port of a port to the far port of its image. None of the sixty turns can scramble the pairing. Count the group against the axes and the arithmetic closes again: six axes, and the moves holding one axis in place number ten.

The pairing itself is a move, sending every port to the port opposite. Include it and the moves that preserve the wiring number a hundred and twenty rather than sixty, the extra sixty being mirror moves that no amount of turning the solid in your hands reproduces. That single move commutes with every other one in the collection: whatever else happens to the ports, opposite stays opposite.

Pick up a modern twenty-sided die and check what the manufacturer did with the numbers. One is opposite twenty, two is opposite nineteen, three is opposite eighteen, and every pair of opposite faces sums to twenty-one. The reason is mechanical. A die pressed slightly flat on one pair of faces lands on those two more often than it should, and if the pair reads nineteen and twenty, the die runs high. Pairs summing to twenty-one cancel that bias. The numbering is a decision. The pairing it uses is the only one the solid offers.

14.7 Thirteen ports

A boundary with thirteen ports on it, each with five seams running out, would have sixty-five seam-ends. Seams have two ends. Sixty-five is odd. There is no such object. The obstruction is arithmetic. Fourteen ports pass that test and fail the other one: they give thirty-five seams, thirty-five seams give seventy face-sides, and seventy face-sides will not divide into triangles. Between them the two double counts dispose of every count that is not a multiple of six. The multiples of six need the surface to settle them. Eighteen ports would need ports minus seams plus faces to come out at three, twenty-four would need four, and no closed surface reaches either, since two is the largest value and only the sphere has it. Six ports would need one, and one belongs to the surface you get by gluing every point to the point opposite, where a port and its far partner are the same port and a patch carried once around comes back mirrored. Twelve is the whole of what the conditions permit.

The one line working physicists write down when somebody asks what the world is made of carries somewhere between nineteen and twenty-six numbers that were measured in laboratories and entered by hand. The number of particle families in it is three because three is how many the detectors reported. Twelve came by the other route. Four conditions on how comparisons meet, every one of them checkable by something sitting inside the boundary with no view of it, leave exactly one count standing, and the count is twelve, with thirty seams and twenty faces attached.

None of which puts a small solid anywhere. The object in the case in New York is 3.2 centimeters of serpentinite lying on a shelf in a gallery. A screen is where an observer's accessible information lives. Being spherical was a statement about loops closing. An observer checks it by counting its own comparisons without ever standing outside anything. The three directions in that gallery are built out of twelve readings like these. The count that produced the twelve used no direction anywhere in it.

Twelve readings, one per port, is everything one piece of the world exposes about itself.

Twelve numbers. The world an observer finds when it looks around has three directions in it, no more and no fewer. Physics has always taken that three as given.

Chapter 15

Why Does Space Have Three Directions?

You can walk forward, walk sideways, and climb a ladder, and any motion you are capable of is some combination of those three. Three numbers pin a body down and two will not: an air traffic controller is given latitude, longitude and altitude, and losing any one of the three leaves a whole line of places the aircraft could be sitting. A fourth independent number has never been available to anybody. General relativity generalizes the shape of space in every respect except the count of directions, which it takes as a starting datum. Nothing in its field equations prefers three to five or to ten. The same equations take it for granted that the three run smooth, with a place between any two places and no smallest step anywhere in them.

It comes out of a bell.

In September 1879 a man left his house in Essex carrying a rack of brass resonators and walked to the parish church to strike a bell.

The house was Terling Place, where John William Strutt, third Baron Rayleigh, had a wing fitted out as a laboratory. The bell was the second of the five in the village tower, counting down from the highest, cast by a founder named Gardiner in 1723. The brass was a set of Helmholtz resonators made in Paris by Rudolph Koenig, who had been apprenticed to the violin maker Vuillaume in 1851, set up on his own account in 1858, and spent the next forty years turning Helmholtz's ideas into apparatus a physicist could order by post. Each resonator is tuned to a single frequency. Hold one to your ear inside a noise and that frequency comes up out of the mess loud enough to name, while everything else drops away. Rayleigh's verdict on them, printed a decade later, was that without the security the resonators afford, deciding which octave a bell is sounding in is very uncertain.

He published in January 1890, in the *Philosophical Magazine*, under the title "On Bells". Much of the paper is a list of what he heard, bell by bell, with the founder and the casting date beside each one, going back to a Terling bell cast by Graye in 1623.

One entry is a hemispherical bell of about three hundredweight, a little over 150 kilograms of bronze, lent to him by the London founders Mears and Stainbank. Four tones came out of it plainly and he named all four. Then this: "The gravest tone has a long duration. When the bell is struck

by a hard body, the higher tones are at first predominant, but after a time they die away, and leave e-flat in possession of the field."

For a fraction of a second the bell has four voices. The loudest of them are the high ones. After that it has one. Which of those two descriptions is the bell depends on how fast the listener is. Give the job to an instrument that cannot register anything shorter than a few seconds, and the four voices never happen. It reports one note, the lowest, and it reports that note as the whole bell.

15.1 Twelve readings and a grid of weights

Chapter fourteen left the carrier with twelve ports, thirty seams, five seams meeting at each port, six axes of paired ports, and sixty rotations carrying the whole arrangement onto itself. Chapter nine left a rule running on it: each port's reading becomes itself, less a sixtieth of the total gap between it and the five readings around it. Sixty because there are thirty seams and a midpoint move closes half of a gap rather than all of it.

Write the twelve readings as a column of twelve numbers, top to bottom, one line per port. A column like that is a **vector**, and the word carries no more than that here: an ordered list, kept in order so that the third entry always means the third port.

The rule turns one such column into another. Look at what a single output line is made of. Port three's new reading is eleven twelfths of port three's old reading, plus a sixtieth of each of its five neighbors' old readings. Every output line has that shape, with different neighbors in it. Each new number is a weighted sum of the old numbers, and the weights can be laid out in a square grid: twelve rows for the twelve outputs, twelve columns for the twelve inputs, and in row three, the number eleven twelfths in column three, a sixtieth in the five columns belonging to its neighbors, and zero in the other six. A square grid of weights used this way is a **matrix**. This one has 144 entries in it, of which 72 are zero, and eleven twelfths plus five sixtieths is one, so every row of it adds to one and so does every column. That is the arithmetic statement of something chapter nine established by a different route: repair moves the total around the twelve observers without creating or destroying any of it.

Run it once on something. Put 12 units at one port and nothing anywhere else. That port keeps eleven twelfths of what it had, or 11, and each of its five neighbors picks up a sixtieth of 12, which is 0.2. Eleven plus five lots of 0.2 is 12, so the total sat where it was, and the gap between that port and each of its neighbors shrank from 12 to 10.8. That is one pass.

The question the bell asks is what the grid does after many passes. That means applying it a hundred times, which by hand means a hundred rounds of multiplying and adding across 144 entries, and after all of it a column of twelve numbers that has to be interpreted.

15.2 Two observers and a number raised to a power

Alice holds 10 and Bob holds 0. The only rule in operation is that each of them moves a tenth of the way toward the other.

One pass: Alice holds 9, Bob holds 1. Second pass: 8.2 and 1.8. Third: 7.56 and 2.44. Do the fourth on paper. Nine tenths of 7.56 plus a tenth of 2.44 gives Alice 7.048, and the same sum with the tenths swapped gives Bob 2.952. The numbers get messier with each pass, and their sum is 10 at every stage, and their difference goes 10, then 8, then 6.4, then 5.12, then the 4.096 you just produced.

The arithmetic behind that is two lines. Let Alice hold A and Bob hold B. After one pass Alice holds nine tenths of A plus a tenth of B, and Bob holds a tenth of A plus nine tenths of B. Add those two expressions together and the tenths cancel, leaving A plus B, which is what the pair started with. Subtract the second from the first and the tenths reinforce, leaving eight tenths of A minus B.

Look at what the rule did to those two combinations. It multiplied the sum by 1 and the difference by 0.8, exactly, and it will do the same on the next pass and on every pass after that, from any starting values whatever, because the arithmetic that does it never refers to what the numbers are. The whole two-by-two grid, read in those two combinations, is two ordinary numbers.

A direction that a grid of weights merely scales, without swinging it round to point somewhere else, is called an **eigenvector**, and the number it gets scaled by is its **eigenvalue**. The prefix is the German for “own”, and these are the grid’s own directions, the ones it is built out of.

What is left of the gap between Alice and Bob after a hundred passes is 0.8 raised to the hundredth power of what it was, which is one button on a calculator rather than a hundred rounds of arithmetic. Ten passes leave 0.8 to the tenth, which is 0.107, so a gap of 10 has closed to a little over 1. The same substitution works with twelve observers instead of two, on one condition.

The weight carrying port three into port seven has to equal the weight carrying port seven into port three. A grid where that holds everywhere is **symmetric**. When it holds, a full set of such directions always exists: as many of them as the grid is wide, sitting at right angles to one another, and every possible column of readings is a sum of pieces along them. That statement is the **spectral theorem**. It holds for every symmetric grid of numbers of any size. The repair grid qualifies, because a seam pushes both of its ends toward the middle by the same amount.

So the twelve-port rule amounts to twelve numbers. Getting them out of the 144 by force means solving an equation of the twelfth degree.

15.3 Sixty rotations and four numbers

The wiring hands them over instead.

The rule cannot tell one port from another. It is written entirely in terms of which port shares a seam with which, and a rotation of the arrangement carries seams to seams, so rotating the readings and then repairing gives the same twelve numbers as repairing and then rotating. Either order, same answer, all sixty rotations.

That is a heavy constraint. It turns into a count. The sixty rotations shuffle the twelve readings around among themselves, and under that shuffling the twelve directions of reading-space fall into pieces: collections of directions that the rotations mix internally and never mix with anything outside. Some such pieces can be cut into smaller pieces of the same kind. Some cannot. The ones that cannot are where the argument lands. For twelve readings on this wiring the uncuttable pieces have sizes one, three, three and five, and one plus three plus three plus five is twelve.

A rule that commutes with all sixty rotations acts on each uncuttable piece as multiplication by a single number. Suppose it did something else, treating one direction inside a piece differently from another. The rotations carry the first direction to the second, so the rule cannot treat them differently. That argument is **Schur's lemma**, which Issai Schur published in Berlin in 1905 to put the character theory of finite groups on a new footing. It is chapter five's symmetry doing arithmetic on 144 unknown weights.

So a rule respecting this wiring, however elaborate its 144 entries look, has at most four distinct answers in it. One for each uncuttable piece.

The two pieces of size three are not copies of each other. The sixty rotations move the directions inside one of them along a different pattern from the other, which is why the two get separate numbers and why the count is four rather than three. Two pieces of size three went into the wiring. The world is built on one of them.

15.4 The three rates

The piece of size one is the direction in which all twelve ports shift together. Nobody disagrees with anybody, there is nothing for a repair to close, and the rule leaves it exactly where it is: eigenvalue one, forever. It also carries no comparison, which means no observer inside the arrangement can read it, and it drops out. Eleven directions are left.

The other three numbers are the fraction of a disagreement that survives one pass of repair in each of the three remaining pieces. Two of the three are irrational.

$$\frac{55+\sqrt{5}}{60} \approx 0.9539 \qquad \frac{9}{10} = 0.9000 \qquad \frac{55-\sqrt{5}}{60} \approx 0.8794$$

Three directions keep 0.9539 of whatever they were holding, five keep exactly nine tenths, three keep 0.8794, and those three numbers, with the 1 that no observer can read, are the entire content of a twelve-by-twelve grid of weights. The 55 over 60 in the outer two is the eleven twelfths a port keeps of its own reading, and the square root of five over sixty is what its five neighbors put back or take away. Slow band, middle band and fast band, named for the rates rather than the sizes. The quantum repair keeps the same slow rate: with the records of each repair set aside, the slowest quantum pattern of the twelve ports also keeps 0.9539 per pass, and every coherence between different arrangements fades faster than that.

The square root of five is the same square root of five a reader met in chapter two, in one plus the square root of five all over two, which a calculator kept landing on from any starting number. It is here because the solid carrying the twelve ports is built out of it, and because a fraction cannot be its own reciprocal plus one. The distance between the top rate and the bottom one is the square root of five divided by thirty, which comes to 0.0745 per pass, and thirty is the number of seams. One of the two pieces of size three fades slower than the other by that margin.

Both of them can be written down. Set the arrangement on a table, choose any flat surface to measure heights above, and give each port a reading equal to its own height above that surface. Those twelve numbers fade at 0.9539 a pass and at no other rate. There are three independent versions of the pattern because there are three independent directions to measure a height along. The other three-dimensional piece is the same recipe run on a different solid. Chapter two's calculator settled on 1.618034 and the quadratic behind it had a second solution, minus 0.618034, a fixed point that nothing ever arrives at. Build the twelve corners using minus 0.618034 everywhere the golden ratio stood, read the heights off those, and the patterns that come back are the ones that fade at 0.8794.

15.5 Slow ears

Strike the arrangement, meaning disturb it away from agreement in no particular direction, and the disagreement lands across all eleven directions at once. Repair works on all three bands simultaneously, at the three rates above.

Follow it. A disturbance in the fastest band is down to half in 5.4 passes. The middle band takes 6.6 passes to halve, and the slow band takes 14.7. Spread the initial disagreement evenly across the eleven directions and three elevenths of it, 27 percent, sits in the slow band at the start. After ten passes the slow band holds 42 percent of what is left. After thirty passes, 73 percent. After sixty, 94 percent. After a hundred, 99.5 percent.

Those are shares of a total that is itself collapsing. After a hundred passes the slow band holds nearly nine parts in a thousand of what it was

handed, the fast band holds under three parts in a million, and the ratio between them is 3,414. Nothing was tuned to produce that. Change the rates and you would have to change the wiring. The wiring is what chapter fourteen showed there is only one of.

That is the behavior of Rayleigh's hemispherical bell. Four voices at the strike, one voice a moment later, and e-flat left in possession of the field.

Observers here are built out of repair records. A repair record is not made in one pass. Chapter nine's limit on how fast influence travels says one seam per repair, and the two most distant ports on this wiring sit three seams apart, so the quickest possible confirmation between that pair is three repairs out and three back. Six repairs is longer than the fast band's half-life of 5.4 passes, so more than half the fast content is gone before the quickest question this arrangement can ask about itself has its answer back. Any observer inside this arrangement is therefore reading it late, after the fast content has gone, which makes observers built out of repair records slow ears. Over those same six passes the slow band loses a quarter of what it holds and no more.

That settles which content can carry a position. A position is a quantity two separated observers can compare and both get the same answer for. The comparison takes rounds to run. Content that has faded to nothing before the comparison completes cannot anchor anything: by the time the second observer reports, the quantity it was reporting on has relaxed away. Only the slowest band lasts long enough. Its size is three.

The five-dimensional piece is the largest one on the wiring, and it fades faster. Its content halves in 6.6 passes against the slow band's 14.7, and after sixty passes the five directions are holding about a twentieth of what the three are holding. Whatever an observer writes into those five directions has drained most of the way out before a second observer can be told about it. A quantity like that describes nothing that has a place.

That is where three comes from. It is the number of independent directions in the one band that outlives every other band, on a wiring that has no alternatives.

15.6 The table of angles

The slow band is three directions wide. Look at what the twelve ports are doing inside it.

Each port leaves a shadow in the slow band, a direction with a length. To compare two directions, take the number that says how much they point the same way: one when they coincide, zero when they are at right angles, minus one when they point exactly opposite. It is the cosine of the angle between them, the only thing about two directions that survives forgetting where they are.

Do that for all 144 pairs of port shadows and the table has exactly four different numbers in it. A port against itself gives 1. Two ports sharing a seam give one divided by the square root of five, which is 0.4472, an angle of 63.43 degrees. Two ports that are two seams apart give minus that. A port and the port directly across the arrangement from it give minus 1.

Those are the angles the twelve ports of chapter fourteen's arrangement stand at, arriving here out of a rule about disagreement rather than out of anybody having measured anything. What went into the calculation was a list of which port shares a seam with which. What came back was a solid.

An observer inside the carrier can read the same table off its own records, with no ruler and no view from outside. Prepare the ports, let repair run, and count the units crossing each port, timing the counts by nothing but the observer's own local events. The counts fix the table of angles exactly, and every one of the thirty seams lands at the same shortest distance. The way the observer's local clocks fluctuate gives the same distances a second time.

The count of independent directions in a table like that is called its **rank**: how many of the twelve shadows you have to be given before the other shadows are forced. Two will not do it. Four is more than necessary. The rank of that table is three. Every entry in it is 1, or minus 1, or one over the square root of five with a sign in front, so the whole calculation runs in arithmetic built out of the square root of five and nothing anywhere in it is rounded.

That three is the slow band's size arriving a second time, since shadows lying in a three-directional band cannot supply more independent rows than the band has directions. A world with four directions in it would need a slowest band of size four. The sizes available on this wiring are one, three, three and five.

15.7 Six counters

A three-dimensional space in the abstract has nowhere in it to stand, and getting into it takes whole numbers.

What an observer holds about where it is, at any moment, is a tally. Every comparison it completes through a port adds one to a count. The twelve ports come in six pairs facing opposite ways, and a comparison recorded out through one port is the same event as a comparison recorded inward through the port facing the other way, so the two counts are one count with a sign on it. Twelve ports, six pairs, six running totals, and every one of them a whole number, because a comparison either completed or it did not. An observer that has completed forty comparisons out through one port and thirty-seven inward through the port facing it holds a 3 on that axis, and five more numbers like it, and that list is everything it has ever known about where it is.

Six whole numbers, dropped into a space with three directions in it. That looks like it should lose information, and on ordinary coordinates it does: six real coordinates fed through the table of angles collapse onto three. On whole numbers it loses nothing. For two different tallies to name the same point, some nonzero collection of whole numbers would have to be flattened by the table, which would make the square root of five rational. It is not. Distinct tallies name distinct points. Different histories can finish with the same tally.

What those tallies do instead is fill the space. Take any point of the three-dimensional band and any tolerance you care to name, and some finite tally lands inside that tolerance of it. The landing places can be numbered off one after another the way the fractions can, and they are packed everywhere. Some points are reached exactly. Every point can be approached as closely as anybody likes.

15.8 Nobody ever holds pi

Write down 3. Then 3.1. Then 3.14, then 3.141, then 3.1416.

Every one of those is a ratio of two whole numbers: the fifth is 31,416 over 10,000. The list crowds in on something, and the something it crowds in on is not on the list and never will be. Nobody has ever written pi down. Every circle any engineer has ever cut was cut to a finite number of digits, and the number the digits are converging on is not a thing anybody has held.

The argument that a list like that is a problem is older than the number line. Zeno of Elea made it in the fifth century BC, in a book that is lost. It survives because Aristotle set it out in the *Physics* in order to answer it: the first argument "asserts the non-existence of motion on the ground that that which is in locomotion must arrive at the half-way stage before it arrives at the goal." Cross a room and you cross half of it first. Then half of what is left, then half of what is left after that. The halves do not run out, so the crossing is a list of tasks with no last entry on it, and Zeno took that to mean the crossing never happens.

Add the list up by hand. A half. A half and a quarter is three quarters. Put in an eighth and you have seven eighths, and the gap left over is an eighth, the very piece you just put in. That happens at every stage: each total falls short of 1 by exactly the piece last added, which is also exactly what all the pieces after it come to. Writing n for the number of pieces added, the pattern is one line.

$$\frac{1}{2} + \frac{1}{4} + \cdots + \frac{1}{2^n} = 1 - \frac{1}{2^n}$$

At n equal to three the left side is the three pieces you just added, seven eighths, and the right side is 1 minus an eighth, which is the same

number. Name a tolerance and the line says which stage gets you inside it. A thousandth needs ten pieces, because 1 over 2 to the tenth is 1 over 1024, and every total after the tenth is closer than that one.

That is the whole of what "the sum is 1" says. The surprise is in what it does not say. The limit is not the last term. There is no last term. None of the totals is 1, no amount of adding produces one, and the statement is about where the totals pile up rather than about any total on the list. Calling it a trick is fair, provided the trick gets named accurately: it replaces an act nobody can perform with a test on finite quantities that anybody can run. For any tolerance somebody names, some finite stage gets inside it and every stage after that stays inside. That is what an infinite sum is, all of it. Cauchy had pushed the argument into inequalities in the *Cours d'analyse* of 1821 and stopped short of putting it that way; Weierstrass put it that way to a differential calculus class in Berlin in the summer of 1861, twenty-three centuries after Zeno.

The fix for pi is a single operation and it is the operation that built the number line. Take the ratios of whole numbers, which have gaps in them wherever a sequence crowds together without arriving, and add in the limit of every such sequence. What comes out has no gaps left. It is the real numbers: the fractions together with every destination the fractions were pointing at. The name for the operation is **completion**.

Run that operation on the record points inside the three-dimensional band, using the distances the table of angles supplies, and what comes out is a continuous three-dimensional space. The record points belong to it, just as fractions belong to the real line. Every other point is the destination of some sequence of ever more accurate records, in the sense that 3.1416 and its successors are a destination.

The sixty rotations come along. Each of them permutes the twelve ports, so each leaves every entry of the table of angles precisely where it was, so each preserves every distance in the completed space. A transformation that preserves all distances is a **rigid motion**: what a machinist does picking up a part and setting it down the other way round, with nothing stretched. The group chapter fourteen counted at sixty elements acts on the space it built as the turns of a rigid body.

A carrier is one piece of something larger. Carriers join into a sphere, twenty of them, then eighty, then three hundred and twenty, each level finer than the last, and repair running across the joins has a smooth limit of its own. That limit lives on the sphere the carriers tile. Count its modes below each frequency and the count grows the way it grows on a surface: the limit has dimension two, the dimension of a screen, the surface a hologram is written on. The three slow directions inside each carrier drop out of the screen's low-energy modes at that scale. Three is the number of directions in each carrier's slowest response, the screen the carriers make together is two-dimensional, and chapter twenty-four sets the two side by side.

15.9 What holds at a finite resolution

Both constructions have a limit in them, and nothing in the world sits at either one. The two limits answer different questions: how strongly the slow band dominates, and how finely records can sample its space.

Run the repair grid any finite number of passes and count the independent directions in what it does. The answer is eleven. Every time, at ten passes and at ten million, because all three bands are present, none of them has reached zero. Three appears when the overall size is divided out and the number of passes is allowed to grow: the fast and middle bands vanish against the slow one, and the rank of what survives is three. At any actual number of passes it is eleven.

Space gets sampled at a finite resolution. Choose a region and a minimum separation between records, and keep adding records until no more fit without breaking that separation. Because the tallies are packed everywhere, the records chosen leave no gap wider than a fixed multiple of the separation. Their number grows as the inverse cube of the separation, which is the three directions showing up in a count.

A field can live on a population like that. Give each record point a real value, weight it by the volume nearest to it, and couple nearby values with a positive interaction. One action, in chapter twelve's sense of the word, then fixes the motion. Refine the population quickly enough compared with the shrinking reach of the interaction and the field approaches an ordinary continuous wave. The error in any smooth reading of it goes to zero over a fixed stretch of time, before anything from the boundary arrives. No grid of little cubes was put in anywhere.

A continuum for practical purposes asks for an error small enough for the observer's question. It does not need a finite list to contain every real point. Having a record near a point and getting there quickly are different things: a tiny displacement can be the difference between two long runs of comparisons. Zeno's runner crosses the room because crossing it was never the same act as finishing a list with no last entry.

That leaves one word carrying much of the argument. Every rate here is per pass. Nine tenths per pass, half in 5.4 passes, 94 percent after sixty. A pass is an event that has finished happening. The shares only mean anything if the events fall into an order. What a pass is, why passes fall into an order at all, and why that order has a preferred way round are three questions a wiring diagram does not answer. Rayleigh's bell rang down and did not ring up.

Chapter 16

Why Does Time Run, and Only One Way?

A latex bead a little over six thousandths of a millimeter across sat in a dish of water in a laboratory in Canberra, held in place by a focused laser. A trap of that kind pulls the bead toward the brightest point with a force that grows the further out it strays, which is how a spring behaves. The dish was then dragged sideways past the trap at 1.25 micrometers per second while a quadrant photodiode read the bead's position a thousand times a second, for as long as ten seconds at a stretch.

Dragging a bead through water takes work. The work finishes up in the water as heat. Scored across a few seconds and longer, every one of the five hundred and forty recorded runs did what the second law of thermodynamics requires: the bead lagged behind the trap and the water took up the difference. Score the same recordings over a hundredth of a second instead, and the stretches that run backwards come out about as numerous as the stretches that run forwards. The water hands energy to the bead rather than taking it. The entropy of the arrangement goes down. Widen the window to two seconds and the backwards stretches thin out; widen it past a few seconds and they stop occurring.

G. M. Wang, E. M. Sevick, E. Mittag, D. J. Searles and D. J. Evans published that in *Physical Review Letters* on 29 July 2002, under the title "Experimental Demonstration of Violations of the Second Law of Thermodynamics for Small Systems and Short Time Scales".

A rule that fails on short recordings and holds on long ones is a statement about an average. So some quantity is being averaged, the average comes out with a sign, and something has to say where the sign comes from. That is one of the three questions hiding inside the word time, and the other two are older.

16.1 Three jobs, one word

Ask anybody what time is and you get a description with four separate claims folded into it: events fall in a single line, the line runs forward and cannot be run backward, the line is the same for everybody, and it goes at one rate. Chapter seven took away the third one. Two clocks in different places have no shared present in which to disagree.

The other three claims are three different objects with one word stretched over them.

The first is **order**: which of two things happened before which. Order is a relation between events. It has no units, no rate and no duration in it. Chapter seven's recipe carried the whole of it, four steps wired in a line and a fifth that floats.

The second is **direction**: why the relation runs one way round and never the other. That is a property of an operation rather than of a relation, and the one the bead in Canberra was testing.

The third is **flow**: a parameterized evolution, the thing a clock reading is laid against. An order alone supplies no durations. Chapter twenty-one builds the flow out of an algebra and a state, and finds a clock in it.

16.2 What gets ordered

Chapter seven taught partial orders on a kitchen counter, before this world contained anything to order, and chapter nine supplied the missing ingredient. The events are commits: committed repairs, whose reads and writes are recorded, together with the records, readbacks and feedback that observers write down.

Take the twelve observers of chapter six, wired as an icosahedron, thirty seams between them, every observer with five neighbors. A repair reads the two readings at the ends of one seam, writes the two readings at the ends of that same seam, and touches nothing else in the world.

That is enough to settle every question about order that has an answer. Two repairs on seams that share an observer both read and write that observer's register. The later one worked from the register after the earlier one had written it, and its certified read carries the version it used, so there is a fact about which came first. Following another repair in an executor's list creates no such fact. Two repairs on seams with no observer in common have no direct fact of that kind. There is no register anywhere holding it, neither observer at either seam was in a position to write it down, and no clock or queue position could supply it afterwards, because supplying it would mean constructing the shared present chapter seven found nothing to build out of.

Two pairs, concretely. A repair on the seam between the first observer and the second, then a repair on the seam between the second and the third: the later move worked from a number the earlier move wrote, so any observer holding both records can see the dependency. The writer's mark on that number and the read certificate on the later move establish it, and the order between them belongs to the arrangement rather than to a convention. Against that, a repair on the seam between the first and the second and a repair on the seam between the fourth and the fifth touch four different observers, four different readings and no common register. Ask

which of those two came first and neither record has anything to say about the other.

Count the pairs. Thirty seams make four hundred and thirty-five pairs of seams. A given seam has two ends, and at each end four other seams arrive, so eight seams share an observer with it and twenty-one do not. That gives one hundred and twenty pairs of seams whose repairs share a register and three hundred and fifteen that share nothing. Those counts belong to the wiring. The order in any one run comes from the versions its commits actually read, and a chain of repairs through other seams can order two repairs that share nothing.

That is a great deal less order than a line would carry. A line puts everything in single file and answers all four hundred and thirty-five pairwise questions before a single repair has run. The observers instead produce a web after the fact. Every strand is a certified dependency of a read on a write, and commits with no chain between them stay unordered. The wiring says where dependencies are possible. The records say which ones occurred.

The everyday picture flattens that web into a timeline. The difference shows up in what you are allowed to ask. “Which of these two happened first” has an answer for some pairs and none for others. Which pairs those are is fixed by the records themselves rather than by any observer’s ignorance.

16.3 Seven pairs go in, one comes out

Chapter nine derived the repair move from three requirements and got the midpoint. Two readings at a seam. Both of them go to the average.

Watch what that does to counts. A seam whose two ends hold 6 and 0 finishes with both ends holding 3. So does a seam holding 5 and 1. So does 4 and 2, and 2 and 4, and 1 and 5, and 0 and 6, and 3 and 3, which needs no change. Seven arrangements of two whole numbers go into that move and one arrangement comes out, and every one of the seven is consistent with what the seam says afterwards.

Run it backwards. You have two readings of 3. The question is what they were. The readings do not say. The observers do not say, because their state consists of what they hold, and what they hold is 3 and 3. The seam does not say, because a seam records the comparison and the comparison is satisfied. The information distinguishing those seven cases was in the arrangement and the arrangement was overwritten, so it is somewhere else or nowhere.

That is what a repair **commit** does: it leaves the total alone, satisfies the seam, and discards which disagreement it resolved. The acceptance test of chapter nine refuses any move that makes the fit worse anywhere,

which rules out the move that would put 6 and 0 back. Even without that rule, nothing at the seam records which of the seven to restore.

Those two facts fail differently. The acceptance test is a rule about which moves are available, and a rule can in principle be relaxed. The overwriting is arithmetic. Seven into one is a fact about averaging whole numbers, and no amount of relaxing anything recovers the input from the output.

A write can be made undoable, and the method is old: keep a copy. Chapter nine's repairs keep one as they go, the number of units that crossed the seam, and that number together with the result returns the input, so the seven cases stop being interchangeable. Copying onto a blank register erases nothing. The cost comes when memory runs out and a register has to be cleared for reuse, and clearing it is another write that discards what it held. Every route out of the collapse arrives at another commit, and chapter twenty-three prices it.

The quantum version of the same move loses in a second way. Put an empty seam and a seam holding one waiting unit into superposition and repair once, and the coherence between them comes back smaller than it went in, by an amount the record law fixes. No later repair restores it. Written as an operator, what one repair takes away, summed over all thirty seams, is positive, and it drains coherence the way friction drains motion. Repair generates a dissipative process, and no reversible rotation of the carrier's own states produces it.

Order is a relation between two records, checkable by whoever holds both and carrying no units. Direction is a property of a single move, showing up as an arithmetic loss inside that move. Drawing one axis and putting an arrow along it welds the two into a single object. The weld is why people go looking for the arrow in the first instant of the universe.

The standard account of the arrow puts it somewhere else entirely. Boltzmann's argument is that high-entropy arrangements vastly outnumber low-entropy ones, so a system wanders into them and stays, which explains why the film runs one way provided the film started in an arrangement of the rare kind. David Albert named that proviso the past hypothesis in *Time and Chance* in 2000, because the account depends on it: the reason your coffee cools becomes a claim about the state of the universe some fourteen billion years ago. The seven-into-one collapse asks for nothing of the kind. It is a property of the move. It holds on the first repair of a fresh arrangement exactly as it holds on the ten billionth, and a universe assembled in any condition whatever runs its repairs in the same direction, because the direction is in the operation.

16.4 Counting what got lost

The seven arrangements that collapse into one are a quantity of missing information. Ask how many yes-or-no questions it takes to pin down one item out of a list. Two items, one question. Four items, two questions, since the first splits the four into halves and the second splits a half. Eight items, three. Sixteen, four. Each doubling of the list costs one more question, which is the defining habit of the **logarithm**: it converts multiplying into adding. Counting questions this way makes the logarithm base two. The unit of the answer is the **bit**.

The list is rarely flat. If one item out of the four is likely and the other three are long shots, a well-chosen first question separates the likely one from the rest, and most of the time you are finished after one question rather than two. So the count wanted is an average over the odds. Claude Shannon wrote it down at Bell Labs and published it in the *Bell System Technical Journal* in July and October of 1948, in two halves of a paper called "A Mathematical Theory of Communication".

$$H = -\sum_{i} p_i \log_2 p_i$$

Here H is the average information in one outcome, measured in bits; the index i runs over the possible outcomes, and p sub i is the probability of outcome i. The minus sign compensates for the negative logarithms of probabilities below one.

Three cases, by hand. A fair coin has two equally likely outcomes, and the answer is exactly 1 bit. A coin biased nine to one comes out at 0.469 bits, and that one is the only one of the three with any arithmetic in it. Nine times in ten the answer is heads, worth 0.152 bits of surprise, contributing 0.137. One time in ten it is tails, worth 3.322 bits, contributing 0.332. The two add to 0.469, so a coin that lopsided carries less than half a bit. A fair eight-sided die gives 3 bits, one for each halving of eight. Give the seven arrangements at the seam equal odds and they come to log base two of seven, which is 2.807 bits. A commit on that seam removes 2.807 bits from what the seam can say about what it used to hold.

Shannon told Myron Tribus that he had gone to John von Neumann for a name, and that von Neumann told him to call it entropy, on the grounds that the same expression appeared in statistical mechanics and that nobody knows what entropy is, so in an argument he would have the advantage. Tribus and Edward McIrvine printed the story in *Scientific American* in September 1971.

The entropy belongs to the description rather than to the thing described. The seam holds 3 and 3 whatever anybody knows. The 2.807 bits are a fact about somebody's odds over the seven arrangements. A different set of odds gives a different number for the same seam. Chapter ten met this

quantity under another word: the run that finished with 81,920 records had a spread of 11.3134, which is the same count in a different unit, the way a length is the same length in inches or in centimeters. In bits it is 16.32.

Which is where the schoolroom gloss goes wrong. Repair makes the world tidier and more determinate, and a seam that goes from 6 and 0 to 3 and 3 has had its uncertainty cut, so anybody reading entropy as a measure of how messy the seam is will get the sign backwards. The seam is as messy after the repair as before it: two whole numbers, sitting at two ends of one overlap. What changed is how many questions it takes to say which two.

16.5 Two messages down one channel

The quantity that runs one way is a comparison rather than a count. Take a settled arrangement of the sort the repair law leaves alone, and call the odds it assigns to each state the reference. Take your own odds over states, which need not be the reference. Ask how many extra questions per state your odds cost you compared with the reference, and the answer is a second quantity, the **relative entropy** of your odds against the reference. It is zero exactly when the two agree, positive otherwise, and measures how far your description sits from the settled one.

Pass both descriptions through the same process, whatever the process is, and they get harder to tell apart. Send two distinguishable messages through the same noisy channel and the outputs are closer together than the inputs were; nothing you do to both of them afterwards can make them further apart again. Relative entropy is the number that quantifies that, and can only fall under further processing.

Repair is a process of that kind, and it leaves the reference where it found it. So the relative entropy of any description against the settled one can only fall as repair runs. That is the second law, as a property of the move.

16.6 The identity underneath the inequality

The falling is an average of something that does not always fall. Take one repair step. Before it, the world is in some state and your odds give that state a probability, and the reference gives it a probability too. The logarithm of the ratio between them is your surprise at finding that state. After the step, the world is in another state, and there is a second surprise, computed the same way against the same reference. The **entropy produced** by that transition is the first surprise minus the second.

Take the exponential of minus the entropy produced and average it over every transition the repair can make, weighted by how likely each one is.

When every state has some chance under both your odds and the reference, the average comes out at exactly one, whatever your odds were.

One further condition is checkable at the seam: measured against the reference, the weight the step carries from one state to a second matches the weight it carries back. A repair in that balance satisfies the following at every pair of states, exactly.

$$p(x)\,K(x,y) = e^{\sigma(x,y)}\,q(y)\,K(y,x)$$

Here p of x is the probability your description gives the state before, K of x y is the chance the repair takes that state to the state after, q of y is the probability of the state after, K of y x is the chance the same repair runs the other way, and sigma is the entropy produced. The equation says the forward weight of any transition exceeds the reverse weight of the same transition by exactly the exponential of the entropy produced, at every pair of states, with nothing approximated. Gavin Crooks published the general relation of this form in *Physical Review E* in 1999, the relation the Canberra bead was measured against.

Two consequences follow. Collect all the transitions that produce the same amount of entropy and add up their weights, and the forward total is that same exponential times the reverse total. Count the entropy produced in bits and the exponential turns into a doubling per bit, which is the same statement in the unit the coins were counted in. A transition producing ten bits runs forwards about a thousand times as often as backwards. One producing forty bits, about a million million times as often. The arrow is a ratio rather than a prohibition, which is why a small enough system watched for a short enough time can be caught running the wrong way, and why a cup of coffee is never seen doing it.

And the average of the entropy produced, over all transitions weighted by how likely they are, equals the fall in relative entropy exactly. The mean of the quantity that fluctuates is the descent itself, with no correction term. The second law then follows in one step. The exponential curve bends upward, so averaging it overshoots the exponential of the average, a fact called Jensen's inequality. The average was exactly one, and the only way both can hold is for the mean entropy production to be at least zero. Nothing anywhere in the argument is approximated.

One more relation drops out of the same bookkeeping. Take two quantities you can measure about the arrangement, evaluate the first before a step and the second after, and average over the settled state of a balanced repair. Swap which one you evaluate first. The two averages are equal. Lars Onsager published that symmetry in *Physical Review* on 15 February 1931, worked out its consequences for coupled processes such as the way a temperature difference across a metal junction drives an electric current and a current drives heat, and collected the Nobel Prize in Chemistry for it

in 1968. Transport coefficients come in pairs and the pairs are equal. Here it is a two-line consequence of the same relation that gave the arrow.

16.7 Faking a past

If a commit destroys what it resolved, a record of the past is the only remaining access to it. The question is whether a record can be forged.

Set the machine one task. A chain of records claims a history worth twenty bits, meaning twenty yes-or-no questions' worth of detail about what happened. A competing chain claims the same history and has no such history behind it, so the twenty bits it would need to produce are twenty bits it can only guess. A blind guess at twenty bits succeeds one time in two to the twentieth, which is one time in 1,048,576, or 9.5 times in ten million.

The machine ran a hundred thousand attempts. The expected number of successes at that rate is about a tenth of one success, and the number obtained was zero. Computing power buys more attempts and leaves the odds on each one exactly where they were. The bits were discarded by the commits that made the history, so there is nothing to copy and nothing to compute.

The past is the set of commits that have been made, together with the certified reads that tie them to one another. Its content is fixed because rewriting it would need information the writes destroyed. The future is the disagreements that have no commit yet. They are not being kept from anybody: their content does not exist in any register in the world, because the repairs that would fix it have not been made.

Chapter six left one unit of disagreement sitting in a world of twelve. One flipped record, thirty seams, nineteen loops, and a leftover of exactly one that no allowed repair removes. Repair in a different order and it sits somewhere else. Every move that put it where it sits was a commit, each one discarding the arrangement it came from. A defect that survives every repair the world can perform is a thing with a history. The history is the chain of writes that carried it, in an order that any observer holding two of those records can check.

Every strand in that web was laid by one commit reading what another commit wrote. Chapter nine counted how fast that can spread: one seam per event, no faster, so a commit can only depend on commits within a certain number of seams of it. The edge of the web advances by at most one seam per event. Turn that count into meters and seconds and it is a speed. Every laboratory that has ever measured that speed has come back with the same number, whatever its own motion.

Chapter 17

Why Is There a Speed Limit?

Stand in rain that falls straight down and it lands on the top of your head. Start walking and it seems to slant toward you, so that an umbrella held upright keeps your hair dry and leaves your shins wet, and it has to be tipped forward to cover them, further the faster you go. Nothing has happened to the rain. The slant is your speed set against the speed of the drops, and the angle the umbrella needs is set by the ratio of the two.

Starlight does the same, and the astronomer who caught it doing so was looking for something else.

On 3 December 1725, in a house on Kew Green, James Bradley pointed a telescope straight up through a hole in the ceiling and a second hole in the roof.

The instrument belonged to Samuel Molyneux, who had ordered it from the clockmaker George Graham. Twenty-four feet from lens to eyepiece, hung from an iron frame bolted to the chimney stack, because the chimney was the most rigid part of the house. Pointing it at the zenith and nowhere else was the design: a star directly overhead is seen through the least air, so any displacement left over belongs to the sky rather than the weather.

The star was gamma Draconis, which passes almost exactly overhead at London. What the two of them wanted from it was parallax. If the Earth goes around the Sun, a nearby star should shift against the distant ones, one way in June and the other in December, and nobody had ever caught it happening.

The star moved. It moved the wrong way, it moved too far, and it moved at the wrong time of year.

Parallax keeps a schedule set by where the Earth is. Gamma Draconis kept one set by which way the Earth is going, and those two run three months apart. By March 1726 the star sat about twenty seconds of arc south of its December position, by June it had come back, and by September it was twenty seconds north.

Every star Bradley checked answered to the same number, which settled it. Parallax has to shrink with distance, being a triangle with the Earth's orbit for a base and the star at the far vertex, and this displacement did not depend on distance.

The displacement is a ratio instead. Divide the speed of the Earth along its orbit by the speed of light, thirty kilometers a second against three hundred thousand, and you get one part in ten thousand, which as an angle is twenty and a half seconds of arc.

The effect is called **aberration**, what motion does to a direction. The number Bradley divided by has to be a speed light has, the same in June as in December, when he and his telescope were moving in opposite directions at sixty kilometers a second. Why is there a top speed at all, and why does everybody who measures it come back with the same figure?

17.1 Two mirrors

Build a clock out of light. Two mirrors facing each other one meter apart, and a pulse of light bouncing between them. One tick is one round trip: two meters of travel, about six and two-thirds billionths of a second.

Put the clock on a train and stand beside the track.

From the platform the pulse does not go straight up and straight back down, because the mirrors slide sideways while it is in the air. It leaves the bottom mirror, travels along a slant, and arrives at a top mirror that has moved along the track since it set out. Then it slants back to a bottom mirror that has moved again. The path from the platform is a pair of slants. A slant is longer than the gap.

The speed is the same on both paths, by the fact Bradley's ratio needs. A longer path at the same speed takes longer, so one tick of the clock on the train occupies more time as read from the platform than it does as read on the train.

How much more is a right-angled triangle and nothing else. The gap between the mirrors is one side. The distance the train carries the mirrors while the pulse is crossing is the second side, at right angles to the first. The slant the pulse actually travels is the hypotenuse, and Pythagoras fixes it: the square on the hypotenuse is the sum of the squares on the other two. Divide through by the gap and the stretch factor turns on one thing, the train's speed as a fraction of the pulse's, and on nothing about the mirrors or the clock. Put the train at eighty percent of light speed and the triangle comes out whole: call the slant five, the train covers eight tenths of it while the pulse is in the air, which is four, and twenty-five take away sixteen leaves nine, so the gap is three. The slant is five thirds of the gap, and so is the tick. At ninety-eight percent it is a shade over five.

Cosmic rays supply the check. A muon is a heavy relative of the electron, manufactured in quantity when a cosmic ray hits the upper atmosphere ten or fifteen kilometers up. Left alone it survives 2.2 millionths of a second on average, which at almost the speed of light carries it about 660 meters, so almost none of them should reach the ground. In 1941 Bruno Rossi and David Hall counted muons at Denver and again at Echo Lake, 1,624 meters

higher, sorted them by how fast they were traveling, and found the slower ones thinning out over that drop faster than the quicker ones did. Read backwards, their counts gave the muon a lifetime of its own of about 2.4 millionths of a second, against the 2.2 that later measurements settled on. The fast ones were not sturdier. They were running on longer ticks.

A clock built out of records shows the same slowing with no mirrors in it. Let a quantum walk carry two packets with slightly different masses and let a small detector read their interference. The beat between them is a clock, and when the packets move, the beat slows by exactly the relativistic factor for the walk's own top speed, which is a third of the fastest speed a record can travel at in it. A finite detector, a finite preparation and an explicit error bound leave the swing plainly visible through a complete moving cycle.

17.2 The train and the lightning

Lightning strikes both ends of the train at once. A guard stands on the platform at the midpoint between the two strike marks, and a passenger sits at the midpoint of the train, and the passenger is traveling toward the front.

Light from the two strikes reaches the guard together. Both flashes crossed equal distances at one speed, so for the guard the strikes happened at the same moment. The passenger is moving toward the flash from the front and away from the flash behind, so the front flash reaches the passenger first, and the passenger, doing the same arithmetic with the same speed, concludes that the front of the train was hit first.

There is no third party who can break the tie, because breaking it would take a comparison that is not the arrival of light at a place, and nobody has one. The two of them hold different facts about the order of two events. Both sets of records are correct.

Chapter seven reached the same absence from the other end, with no signals in the argument at all: no observer in the arrangement was in a position to assemble a shared present out of what it held. Here it fails for a narrower reason, that light has one speed, handed over as a measured fact about one kind of signal and left standing.

17.3 Two events, one subtraction

The guard and the passenger disagree about the order of two events, about how far apart they were, and about how long the gap between them lasted. Chapter four's overlap rule says something has to come through a comparison like that intact, or the two of them are not describing one world.

Take two events. Take the time between them on one observer's clock, multiply by the speed of light so that it becomes a length, and square

it. Then square each of the three distances between the events along that observer's three directions, and subtract all three.

$$s^2 = c^2t^2 - x^2 - y^2 - z^2$$

Here t is the time between the two events on one observer's clock, x, y and z are the distances between them along that observer's three directions, and c is the speed of light, present so that a duration and a length can appear in the same subtraction. Two observers in relative motion assign different values to every letter on the right and the same value to s^2, making the quantity a fact about the pair of events rather than about anybody's account of them.

That quantity is the **interval**. Its sign sorts every pair of events in the world into three classes.

When the interval is positive, the events are separated by more time than distance. One can influence the other, some observer sees them happening in the same place at different moments, and every observer agrees on which came first. When it is negative, they are separated by more distance than time. Influence runs in neither direction, some observer sees them happening at the same moment, and observers disagree about the order with no fact anywhere to settle it. When it is exactly zero, the separation is one that light can cross and nothing slower can.

Draw the zero case at a single event and it is a cone opening into the future and another into the past. That cone is the **light cone**. Every pair of events in the universe sits inside one, on one, or outside.

All of which runs on a single input, that light comes out at one speed for everybody. Einstein put that in by hand in 1905 and said he was putting it in by hand. A null result in a Cleveland basement eighteen years earlier had said the same thing, and promoting a null result to a law is a way of writing physics down. It is not a way of explaining it.

17.4 The sky is the data

So take the arrangement apart and ask what an observer physically holds.

An observer has a bounded interface. What reaches it reaches it through a port. Everything it can read is indexed by which way the thing came in. At the resolution of one carrier that is twelve directions, and refining the carrier fills them in, and the object the directions fill in is a sphere. That is the screen with its geometry showing: the sky, in the plain sense that the sky is where you see everything from where you are standing.

Two observers in relative motion look at the same stars and disagree about where they are. Bradley measured the disagreement at one part in ten thousand. It grows with speed. His stars did not swing round together the way the constellations swing round on a globe you turn. They were all

pulled toward one point, the point the Earth was heading for that month, and a pull toward a point is a different operation from a turn: stars near that point crowd together, stars out to the side barely change their spacing at all, and the amount depends on where on the sky you look. So angular separation between two stars does not survive: your sky and mine put the same pair of stars at different distances apart, and neither of us is wrong. Overall size does not survive either. There was never anything to measure it against, since a sphere of directions has no radius that anybody in the arrangement holds.

What does survive is the angle at a crossing. Two families of directions that cut across each other at right angles on my sky cut across each other at right angles on yours. A set of directions that closes into a circle on mine closes into a circle on yours, at whatever size it comes out. Those relations hold up under every comparison two observers can run. Chapter five called that an invariant.

Name the invariant and the group follows.

The maps of a sphere onto itself that preserve every crossing angle have a picture. The picture is a map projection. Set the sphere on a flat table so it touches at the south pole. From the north pole run a straight line through any other point of the sphere and carry it on until it hits the table. Every point except the north pole lands somewhere. The entire infinite table gets used. Navigators have bought angle-preserving charts for centuries and they all buy them for the same property: two roads crossing at forty degrees on the globe cross at forty degrees on the chart. What a chart like that destroys is size, which is why Greenland looks enormous on a sea chart. Navigators accept that, because nobody steers by area.

It does one further thing. Every circle on the sphere arrives on the table as a circle, except for circles through the north pole, which arrive as straight lines, a straight line being what a circle of unlimited size looks like.

Which turns the question into a question about the flat table. The table has an arithmetic on it.

Write a point two units right of the origin and three units up as the single number $2 + 3i$, where i is the instruction "turn a quarter turn counterclockwise". Turn twice and you are facing backwards, so i times i is -1, and that is the entire content of the symbol. Numbers of this shape are **complex numbers**. They add the way arrows add, tip to tail. The useful part is that they multiply. Multiply $2 + 3i$ by i using the ordinary rules of school algebra and you get $2i + 3i^2$, which is $-3 + 2i$: the point three left and two up, the arrow you started with swung a quarter turn to the left. Nothing was arranged to make that come out. It follows from two quarter turns making a half turn, and from the rule for expanding brackets. Multiplying the whole table by one complex number turns it through an angle and stretches it by a factor, both at once, everywhere at the same time.

Turning keeps angles. Stretching keeps angles. Sliding keeps angles, and so does the operation that undoes multiplying. Build a map of the table out of those and it keeps angles, and the most general such map that covers the whole table once and sends every circle to a circle sends a point w to $(aw+b)/(pw+q)$, with a, b, p and q four complex numbers. Maps of that shape are called Möbius transformations. Every angle-preserving map of the sphere onto itself is one of them.

Count the knobs. Four complex numbers is eight real ones. Multiplying all four by the same thing moves no point anywhere, which deletes two. Six knobs.

Turn them and watch the sky. Three of the six rotate it rigidly, the way you turn a globe: the constellations swing round and no star changes its distance from any other. The remaining three do something no rotation does. They gather the whole sky toward one point and thin it away from the opposite point, so the constellations crowd toward the direction you are heading and open out behind you, and each of the three is labeled by a speed. Bradley's twenty seconds of arc is that motion run at one part in ten thousand, and the speed shows through the fact that the twenty is not quite a constant: the displacement runs about 20.8 seconds of arc in January, when the Earth is nearest the Sun and traveling at 30.3 kilometers a second, and about 20.2 in July, when it has slowed to 29.3.

Those six, three rotations and three crowdings, are the Lorentz group, named for Hendrik Lorentz, who wrote the transformations down in 1904 as a way of keeping the ether. They are the transformations relating observers in relative motion, the ones Einstein reached from two postulates, and the sphere carries them as the only thing it can do to itself without disturbing an angle.

Textbooks from 1905 onward drew a fast-moving sphere as an ellipsoid squashed along its direction of travel. The question of what a camera would record went fifty years without a general answer. In 1959 Roger Penrose published three pages in the Proceedings of the Cambridge Philosophical Society pointing out that a camera records no such thing. The outline of a sphere is a set of directions, angle-preserving maps take circles to circles, and a photograph of a sphere at any speed you like comes out round. James Terrell got there independently the same year in the Physical Review, under the title "Invisibility of the Lorentz Contraction".

17.5 Four numbers in a square

The sky has six knobs on it and the interval has four signs in it, and one square of four numbers carries both. Chapter fifteen laid the repair weights out in a twelve-by-twelve grid and took its eigenvalues. Take the smallest block that has any structure: two by two, four entries in a square. Every such block carries a single number of its own: multiply the top

left by the bottom right, multiply the top right by the bottom left, and subtract the second product from the first. What is left over is the block's **determinant**.

Fill the square with a particular pattern. Real numbers down the diagonal, a complex number in one off-diagonal corner, and in the other corner its **conjugate**, meaning the same number with the sign of its i part flipped, so that the conjugate of $x + iy$ is $x - iy$. A square filled that way is called Hermitian, after Charles Hermite, and the pattern means it takes four real numbers to write one down rather than eight. Call the four t, x, y and z, measuring time in the distance light covers in it, and lay them out.

$$\begin{pmatrix} t+z & x-iy \\ x+iy & t-z \end{pmatrix}$$

The four numbers on the diagonal and in the corners are the same t, x, y and z that the interval was built from, one time and three distances, written into a square instead of a row.

Do one. Take five for the time, and two, one and two for the three distances. The diagonal reads $5 + 2$ and $5 - 2$, and the corners read $2 - i$ and $2 + i$.

$$\begin{pmatrix} 7 & 2-i \\ 2+i & 3 \end{pmatrix}$$

Seven times three is twenty-one. The other product is $2 - i$ times $2 + i$, which expands to $4 + 2i - 2i - i^2$, where the middle terms cancel and $-i^2$ is $+1$, so it comes to five. Twenty-one take away five is sixteen. Then do the interval on the four numbers you started with, twenty-five minus four minus one minus four, and it is sixteen as well.

Take the determinant in letters. Top left times bottom right is $t^2 - z^2$. The other product is $x - iy$ times $x + iy$, which multiplies out to $x^2 + y^2$, the cross terms canceling and i squared turning the last one from a minus into a plus. Subtract, and the determinant is $t^2 - x^2 - y^2 - z^2$. That is the interval, with the units chosen so light covers one unit of distance in one unit of time.

Nothing in that multiplication knew about time or distance.

Count the signs. One plus and three minuses, and nobody chose a single one of them. They come from the shape of the square. The diagonal is real, so its product is a difference of two squares, and the corners are conjugates, so their product is a sum of two squares that the subtraction then takes away whole. Einstein reached that pattern of signs from two postulates about light. Subtracting one product from the other in a two-by-two produces it in a line of school algebra.

Set the determinant to zero and you have the light cone. A two-by-two block with determinant zero has one of its rows a multiple of the other, so it carries one independent row rather than two. Chapter fifteen called that count the rank. Light rays are rank one, and every other nonzero separation is rank two.

Keep five for the time and take three, nothing and four for the distances, which are the three, four and five of the right-angled triangle the light clock ran on. The square comes out as nine and one down the diagonal with a three in each corner. Nine times one is nine, three times three is nine, and the determinant is zero. The bottom row, three and one, is the top row divided by three, so there is one independent row and not two.

Take the rank-one blocks with t positive, which is the future half of the cone, and agree to ignore overall size, since scaling a direction by a positive number does not change which way it points. What is left is the sphere of directions. The match is exact both ways: each future light ray names one point of the sphere, and each point of the sphere names one ray. The sky an observer looks at and the boundary of what can reach it are one object described twice. The angle-preserving motions of the first are the determinant-preserving motions of the second.

Chapter fourteen counted sixty rotations carrying a twelve-port carrier onto itself, and no port came out distinguished, because some rotation among the sixty carries any port to any other. The motions that leave a determinant where they found it do that work one level up. They close under doing two in a row and each has one that puts the square back, so they are a group, and they are the six knobs on the sky: the group Einstein's two postulates about light were describing.

Pick a direction into the future with determinant one and call it an observer at rest. Ask which vectors are perpendicular to it, using the interval to decide perpendicular rather than ordinary distance. The answer is a space of three dimensions, the observer's rest space. On that space the interval with its sign reversed is positive for everything except zero, ordinary Euclidean distance in three dimensions. Every observer gets one. It is the same three directions chapter fifteen read off the slow band of the carrier.

One more count comes from the same subtraction. A light ray picks a direction inside that three-dimensional rest space, the way it is going. The directions left over, the ones perpendicular to it, make up a space of three minus one dimensions. Two. A light wave has exactly two independent ways of waving, which is why polarized sunglasses cut glare and why they stop cutting it when you tilt your head far enough.

17.6 Order plus number

The cone so far came off the sky: a sphere of directions, the maps that keep its angles, a determinant that vanishes. Chapter seven's order came from somewhere else entirely, records that read what other records wrote. A rule for those reads joins the two.

Start with chapter fifteen's six counters and prepare a population of records. Their whole-number tallies land at points in the three-dimensional band, with no point of the chosen region further than a small gap from one of them. Protect those addresses while other values change. In each round every record reads every record that lay within a fixed reach of it in the preceding round, itself included, and writes its next value. Reading your own preceding record is what waiting looks like. The reads determine a partial order.

Ask how far one event's influence can have got after five rounds. Two plain facts settle it. A read covers at most one reach, so nothing has got further than five reaches. And anything nearer than five times the reach, less twice the gap, can be got to. Draw a straight line to it, mark five evenly spaced waypoints along the line, and in each round step to the record nearest the next waypoint. Each step is a fifth of the line plus at most a gap at either end, and that fits inside a reach. So the influence of every event fills a region caught between two cones, and the only difference between them is twice the gap measured against the reach. Refine the population so that the gap shrinks faster than the reach, and the two cones close onto one. Its speed is one reach per round.

On a grid where every move runs along one of three axes, influence would spread in no round cone at all: a point two steps east and two steps north lies within three steps as the crow flies and takes four moves to reach, and the region of influence comes out as an octahedron, two pyramids base to base. Here the reach is a ball, set by distance alone, and the sixty rotations of chapter fourteen carry it onto itself. A theorem says the round cone is the only one possible. Suppose possible influences form a cone, so that possible displacements combine into possible displacements and no displacement is possible together with its reverse. Suppose the sixty rotations leave the cone alone, and so does the sky's crowding along one single direction. Then the cone is exactly the one the sky's six motions preserve, the light cone, opening toward the future or toward the past. The rotations alone would allow a cone of any width. One crowding fixes the width, and the rotations carry that one crowding round to all three directions. The records pick the future, because height goes up along every read.

A finite log fits the cone with no limit taken, once each event is drawn at its position with its height times the length of a round as its time. Every read then stays inside the future cone, since a read covers at most one reach in one round. Where the positions also keep events with no chain between

them outside each other's cones, order and cone agree exactly, pair by pair, and every stretch of the order is the matching stretch of the cone.

For the number, pick two events, one in the other's future, and take everything after the first and before the second. The shape is two cones joined at the rim, a **causal diamond**. As the population refines, count one completed record per site and round, give each its fixed weight, and the count inside a diamond sitting wholly within the region approaches the diamond's volume. The volume grows as the fourth power of the duration between the tips. Twice the duration, sixteen times the count.

Counting also gives a dimension test. Take every pair of events inside a flat spacetime diamond and ask what fraction are ordered, one inside the other's cone. With uniform volume sampling and one direction of space the fraction is a half. With two it is eight thirty-fifths, a little under a quarter. With three it is one tenth. Every extra direction gives two events more ways to miss each other. Jan Myrheim worked the fractions out in 1978 and David Meyer made a dimension meter of them in 1988. The records approach one tenth inside every diamond that sits wholly within the region they fill.

Run it. The records sit on twelve-port carriers, each position rebuilt from its six counters, each read checked against its writer's mark. With 39,304 records and six rounds, the outer central interval holds 32,265 events. Of its 520,498,980 pairs, 47,881,819 are ordered, about 9.2 percent. A starting disturbance reaches exactly the records reachable through the read graph. On 166,375 positions and eight rounds, the outer interval gives about 9.3 percent and a fitted dimension of 4.09. On 704,969 positions and ten rounds, the outer interval holds 1,081,730 events, 55,108,871,450 of its pairs are ordered, about 9.4 percent, and the fitted dimension is 4.07. The edge of the region clips all three intervals, and the fraction climbs toward a tenth as the populations grow.

Carrying those reads through local seams takes relay work. Routed through the seams of actual carriers, runs serving 2,197 and 9,261 records use about nineteen million and 293 million primitive events, and a separate checker rebuilds every value and every writer each read consumed. The completed records keep their order. The relay and housekeeping events in between do the carrying and add nothing to the count of records: weight every one of them like a record and the count stops measuring volume.

Storage can be reused once every read of an old value is done. Two alternating stores and some temporary relay space are enough.

Pair averaging can also keep a running total after one preparation. Values carried as differences between two nonnegative loads stay readable as their scales change, and clearing the total retires its old answer. A receiver publishes only when its samples and its error bounds admit one answer, and abstains when several fit.

With two protected banks, pair averaging stores computed answers for later layers and reuses records after their last requested read: one bank stays readable while the other receives the next layer. Complete finite examples check the process, signed inputs and a change to the central input included, with one error budget covering preparation, transport, cleanup and publication.

A field can use its own measurements to keep evolving. On sixty-four sites, probes disturb the field, retained records restore it, and later updates read the restored values. A localized disturbance produces a separated detector response larger than all the errors combined, over a nonzero stretch of the field's own time.

The screen's own field keeps a speed limit as well. Repair on the joined carriers of chapter fifteen gives one operator that sets three things at once: how strongly the field at two places is correlated, how fast a disturbance of it fades, and the frequencies of the waves it carries. Those waves have fronts, and the fronts travel at a finite speed. Change the field in one place and nothing changes at a distance until a front has had time to arrive.

There is also a way to recover initial values while changing the storage that holds them. Imagine connected cups whose contents can be averaged in pairs, with one cup sampled exactly. First measure that cup. Bring in an unmeasured neighbor by averaging along a path of cups whose contents are known, ending at the sampled receiver. The receiver's measurement and the known relay values let us undo the averages arithmetically and recover the neighbor's former value. Repeating this through the connected region recovers every starting value from samples at the same receiver. The paths come from the connections. The cups end up holding new contents, and the starting values live on in the receiver's records.

A requirement on what gets published can also fix how the reads are scheduled. Name the records to recover, the allowed moves, a deadline and a reference probability for each complete sequence of reads, and suppose some sequence meets the requirement in time. The schedule that adds the least information to the reference while meeting the requirement is the reference restricted to the successful sequences and renormalized. Summing the reference weights of the successful continuations from each step gives a controller that samples that schedule as it goes, with no failed runs to execute and throw away. Different deadlines give different schedules.

The deadline can be removed by calculating the probability of eventual recovery from each intermediate state. Weighting the next move by that remaining probability gives distributions that agree when a long sequence is shortened.

On a chain of cups there is a stronger result: place independent unknown amounts at cups separated by calibrated empty cups, and every finite sequence of neighboring averages retains enough independent information in the current contents to reconstruct the unknowns. If every move

keeps a uniformly positive chance of occurring, retained exact samples at one receiver eventually recover them all with probability one, although the precision needed can grow without bound.

Several shrinking reaches, and both golden and uniform populations, pass the same limiting cone and counting tests while having different finite reads. The exact square-root formula for the reach balances two errors: smaller steps sharpen the time resolution and pile up more spatial error. It is a convenient regulator, and no fundamental length is hiding in it.

Removing every spatial restriction would let any distance be crossed in one shrinking step, and keeping only the nearest neighbors along three axes gives the octahedral reach of the grid above. What the limit needs is a local reach that shrinks in balance, and the square-root formula is one way to get it.

Maximum randomness cannot do the work of the rules. Choose the most random quantum state compatible with the observers' constraints, with every observer weighted and every direction allowed, and any failure that some allowed state permits, and that an observer can test for, keeps a positive probability in the chosen one. That holds even when the observers' local states do not commute and no single global state extends them, and any allowed failure gives a number for how large that probability is. Exact success has to come from the constraints themselves. The failure probability can shrink as the resolution grows, so a limit can succeed all the same.

A separate two-time example makes the distinction concrete: preserving the same maximally random state permits both perfect retention of a classical bit and complete erasure of it. Choosing a state has not chosen the process that carries its records.

The quantum version of the averaging move needs one more ingredient. Two quantum versions of the same averaging rule give the same port readings after one repair and different readings after the next step. A version that measures as it repeats does spread influence through the carrier, and it erases a definite family of initial differences on its first cycle, which no later influence restores. The quantum state, the archive of records and the repair instrument come as one package, and chapter nine's record law is the part of the package that says what a repair writes down.

Lost information can be kept on purpose. Signed differences on a selected set of seams, together with the repaired values, recover every input. A quantum construction that keeps the random swap decisions along with the state can undo the whole operation. Those extra records have a cost that can be counted. What a repair writes down is a physical question, and chapter nine gives the experiments that answer it.

Durations come out of the counts too. Divide the counts in two diamonds and take the fourth root, and the answer approaches the ratio of their durations, the exponent coming from four-dimensional volume. Any

duration read from counts alone that varies continuously, scales properly and adds along straight paths has this form, up to a unit, and one reference diamond sets the unit. Laying that unit against a laboratory clock turns it into seconds.

String small diamonds along a path and add their count durations, and the sum approaches proper time. A moving path collects less than a path at rest, by the familiar relativistic factor, which is the speed half of what Hafele and Keating's clocks brought home from their flights. The other half, from height above the ground, belongs to gravity, which this flat construction leaves out.

Simply counting the steps of a history gives a different answer. Between related events five rounds apart, a history can always fill all five rounds: travel when needed and wait for the rest. The longest history has five steps whether its endpoint has moved or stayed put. Assigning the same tick to each step therefore misses the slowing of a moving clock. Counting the events throughout a diamond carries information that counting one chain does not. In an expanding geometry, even the diamond clock needs small enough windows: the volume of a finite diamond samples the changing expansion across its interior, while a clock on a worldline samples its own path.

So the records have a four-dimensional limit: three position coordinates, an order generated by reads, and volume and duration recovered from counts. Its cone speed is one reach per round.

17.7 Cleveland, answered

Michelson and Morley built an instrument to display one number: the speed of their basement through the ether.

That number is one of the six knobs. Turning it moves the stars around on their sky and changes no crossing angle, no circle and no interval, which is to say it changes nothing either of them could compare against anything. The apparatus was not too insensitive. There was no quantity in the basement for it to be sensitive to. No instrument built since has found one.

The six motions shuffle the sky as hard as you like and leave the cone exactly where it was, because the cone is where a determinant vanishes and the six motions preserve determinants. They therefore agree on the cone's three classes: timelike, spacelike and null separation. Wherever records are placed so that order and cone agree pair by pair, as above, the record order carries that classification exactly, and the refining reads close onto the same cone.

The cone speed is one reach per round, and quoted in meters per second it is the speed of light. Local transport can spend many seam operations inside one round, so the speed counts rounds and leaves the relay events

out. The angle-preserving sky maps say why one cone gives everybody one invariant speed. Chapter twenty-two meets the same cone from the other end, as the first moment a wave run on an observer's own count of repairs can answer a disturbance somewhere else. The number the speed takes is a choice of units.

Metrology adopted this in 1983. At the seventeenth General Conference on Weights and Measures the meter was redefined as the distance light covers in one 299,792,458th of a second, which fixes the speed of light at 299,792,458 meters per second exactly. Nobody has measured it since, and nobody can, because measuring it would consist of measuring the meter.

A separation that light can cover is rank one: one independent row, nothing held behind it. Everything that travels slower is rank two. The second row is the reason it travels slower. That row has to be re-asserted every round or the thing it belongs to stops being there, which means something is being spent to hold it, at a rate, against a clock, and it is being spent in every piece of ordinary matter.

Chapter 18

Why Does Anything Have Energy?

A ball moving twice as fast carries more of something. It hits harder, and it digs further into whatever stops it. How much more, though: twice as much, or four times? Weight multiplied by speed says twice. Weight multiplied by the square of the speed says four. Each answer is one line of arithmetic, and nothing in either line says which multiplication the world is doing.

For a long stretch of the early eighteenth century each line had a party behind it. Both sides agreed that a moving ball carries something, and that the something grows with the weight of the ball and with its speed. Descartes' followers held that the quantity worth tracking is the weight multiplied by the speed. Leibniz's held that it is the weight multiplied by the square of the speed, and called it vis viva, living force. The dispute had run for decades, mostly in Latin, mostly in print, and neither side had proposed an experiment.

Willem 's Gravesande, who taught mathematics and astronomy at Leiden, put a tray of soft clay on the floor in 1722 and dropped brass balls into it from measured heights.

A ball released from four times the height arrives at twice the speed, so a measured drop hands the clay a chosen speed. A ball arriving at twice the speed sank four times as deep. At three times the speed, nine times as deep. Two balls of the same size and different weights, released from heights in inverse proportion to those weights, left the same dent.

The square won, and neither side could say what the square was a quantity of.

Émilie du Châtelet took the argument up in France. She had entered the Academy of Sciences prize competition on the nature of fire in 1737, anonymously, as every entrant was required to do; the result was announced on 16 April 1738 and the prize was divided among Leonhard Euler and two others, and her essay was printed with the winning memoirs the following year, the first scientific paper the Academy published by a woman. Her *Institutions de physique* came out anonymously in Paris in 1740. Its twenty-first chapter, on the force of bodies, put the clay in front of readers who had been treating the question as a matter of definition. She came down on the side of the square. In 1745 she began translating the *Principia* into

French, with a commentary running alongside Newton's Latin that supplies the derivations he had left out. She was carrying a fourth child while she finished it, working eighteen-hour days, and she dated the commentary on the last day she worked on it. She died at Lunéville on 10 September 1749, six days after the birth, at forty-two. The complete text appeared in 1759, the only complete *Principia* in French that exists.

Thomas Young gave the number its modern name in 1807: the term energy, he wrote, may be applied with great propriety to the product of the mass or weight of a body into the square of the number expressing its velocity. That sentence assigns a word to a product and assigns nothing else. Newton's *Principia*, which defines mass, momentum, force, centripetal force and half a dozen other things with great care, has no energy in it anywhere. Textbooks define energy as the capacity to do work and define work as force through a distance, which tells you how to compute the number and leaves open what the number counts. Chapter five gave the classical answer: energy is the quantity conserved when the laws do not care when. That correspondence is exact. It wants two things before it starts: a background time to shift and a stage for the shifting to happen on.

18.1 The rate a region is worked through

Chapter nine left this world with one move. Two neighbors share a seam, each holds a reading of it, the readings disagree, and both readings go to their midpoint. Nothing else happens anywhere. Repair drives the outstanding disagreement down toward zero, which is what the seams read when every pair of neighbors agrees.

The machine holds one move, one kind of seam and two readings at each seam. The descent of the outstanding disagreement is the only quantitative thing any of it does, so energy is made out of that descent, there being nothing else to make it out of.

Take a screwdriver to a screw. One eighth of a turn, repeated, drives the screw into the wood. The whole trip of the screw is that one small move done again and again.

A smooth continuous change is a flow. The small move whose repetition produces the flow is its **generator**, and the pairing runs both ways: a flow determines its generator, and a generator run out determines the flow. The trip of the screw is a long list of positions. The one turn is a single instruction. The list follows from it.

Every region in this world advances along a clock of its own. Something generates that advance. That something is the region's energy. The energy of a region is the rate at which its records are worked through by repair, counted against that region's own clock.

The word behaves identically wherever physics uses it. What generates turning about an axis is the angular momentum about that axis, and what generates sliding sideways is the momentum in that direction. Advance along a clock has the energy for its generator, and quantum mechanics has run on exactly that definition since the 1920s without ever saying what does the advancing.

A clock of a region's own is the only kind this world has. Chapter ten's network ran with no tick anywhere in the machinery, nothing waiting for the next round and nothing counting rounds; the cycles in its trace belong to whoever reads the trace. What a region has instead is a clock its own state manufactures, out of nothing but what it holds, and the events at its own ports to count against it. Two regions worked through at different rates differ in a fact about themselves, since no clock outside them has a setting they could differ in.

A frozen instantaneous state has no changing records. A stationary probability law can contain a continuing flow of events. Modular flow acts on observables according to a specified reference state. Those are three distinct claims about equilibrium, and a calorimeter reading needs a connection from the chosen generator and clock to physical energy.

For a faithful reference state, minus its logarithm defines the modular generator. In a thermal description, where the state is proportional to the exponential of minus energy divided by temperature, that generator is the Hamiltonian in units of the temperature, plus a constant that changes no observable. The state fixes this modular reading. Identifying it with autonomous physical evolution requires the state, dynamics and calibrated clock to belong to the same description.

18.2 Positive, and additive

The disagreement outstanding in a region is a sum of squares, the spread of its readings about their average, which puts a floor under it at zero, and zero is total agreement. No repair ever raises it: an even split lowers it by half the squared gap, an odd one by a little less, and a gap of a single unit leaves it level. No region can spend past it, and written as an operator, what repair takes away from a region's quantum state is positive as well: repair only ever runs downhill. So the rate is never negative. There is no arrangement of records that goes on falling forever.

Arthur Wightman wrote that into the axioms of quantum field theory in the 1950s and called it the spectrum condition. It has to be assumed there, because without it every system has somewhere lower to drop into and hydrogen radiates without limit. Here it is the floor of a sum of squares.

Additivity comes off the seams. Two regions with no seam between them have no comparison to make across the gap, so they have no repair to do across it, and neither one's rate says anything about the other's. The rates

add. That is why energy accounting works at all: a calorimeter that stops two beams together reports the sum of what it reports for each of them alone, an assumption so ordinary that nobody writes it down, and holds because separated regions share no seam.

That sum has a condition on it. The condition breaks the moment the two regions touch. Run a seam between them and their rates stop adding, since the seam carries a disagreement belonging to neither of them and repair at that seam changes both. The gap between the joint rate and the sum of the separate rates is the quantity a chemist charges for a bond. The sign of that gap is why two hydrogen atoms end up as a molecule.

18.3 Both sides of every entry

In November 1494 the printer Paganino de Paganini finished a large book in Venice for Luca Pacioli, called the *Summa de arithmetica*. Twenty-seven pages of it set out what the merchants of the city had been doing with their books, the first printed account of the practice.

Every transaction is written twice, once as a debit in one account and once as a credit in another. The books balance, and they balance for a reason that has nothing to do with anybody adding them up at the end of the day: a single-sided entry cannot be written at all. A clerk who wants to steal has to falsify two accounts that other entries also touch.

The repair move has that shape by construction. Chapter nine's second requirement was that a repair leaves the total of the two readings it touches alone, which makes every repair two entries, what comes off one reading going onto the other, the pair of them one event. The requirement is there because a move that changed a total would have to know the total, and no register in this world holds one.

Energy conservation is that fact read at the level of rates. A region is worked through at some rate, the working happens at seams, and every seam is shared with a neighbor, so what leaves one region's account arrives in the next one's, entered twice. The rule performing the entries is one rule, the same at every seam and at every event, carrying no parameter that could take a different value somewhere else. There is a general procedure for turning a symmetry into its conserved number. It is not needed to see this one. Chapter five's correspondence, that energy is what you get when the laws do not care when, is this seen from the other end: the repair rule has no when in it to care about.

In 1775 the Royal Academy of Sciences in Paris resolved to examine no further proposals for perpetual motion, and put them on a list with squaring the circle, trisecting the angle and doubling the cube, none of which had been proved impossible at the time. Every one of those designs is a proposal to make an entry on one side of a boundary and not the other. The obstacle is the list of available moves. Repair acts on one seam,

it touches the two readings at that seam's ends, and it leaves their total where it found it, so a wheel that gains what nothing else loses is asking for a move that has never been on the list.

18.4 A wave that would not spread out

In August 1834 John Scott Russell was riding along the Union Canal near Edinburgh when a boat drawn by two horses stopped suddenly. The water it had been pushing rolled forward off the bow as a single rounded heap, some thirty feet long and a foot to a foot and a half high, and went on down the channel without spreading and without slowing. He followed it on horseback at eight or nine miles an hour, for a mile or two, and lost it in the windings of the canal.

What he chased was a shape. The water in the heap changed the entire way: each stretch of canal rose, moved forward a little and settled back down, while the heap went past it. The thing that traveled was an arrangement re-establishing itself out of new material at every point along the route. It held together because two effects that each ruin an ordinary wave were canceling. A tall crest runs faster than the water beneath it and steepens the front. Spreading pulls the crest apart into a train of ripples. Balance the two against each other and the heap keeps the shape it left the bow with.

The seam network holds patterns of that kind. A bounded set of records reproduces itself after every round of repair, because flattening it would open larger disagreements at the boundary of the set than it closes inside. Repair works on such a pattern at every event and gets nowhere with it, so holding the pattern costs repair work at every event, and what the work buys is the pattern being there for the next one.

Mass is that standing cost: the rate a pattern demands while its records go nowhere. Nothing about a stone on a table is idle. The arrangement that makes it a stone is being re-asserted at every event, at a rate the stone's own clock sets. That rate is what a balance reads off when you put the stone on one.

A pattern of that kind turns its material over, exactly as the heap in the canal did, and the charge falls on the arrangement rather than on whichever records are holding the arrangement at a given event. The proposed cost stays steady while the material under it changes. The quoted laboratory proton mass is 1.0072764665789 atomic mass units. Deriving that number from repair requires the physical quark and strong-interaction dynamics; this account of persistence does not compute it.

18.5 Four numbers

Chapter seventeen put four real numbers into a two-by-two Hermitian matrix and found the interval of special relativity sitting in its determinant: the first number squared, minus the squares of the other three. The transformations that carry one observer's description into another's are exactly the ones that leave a determinant alone, which is why every observer computes that combination and gets the same answer.

A region's entry is four numbers of that kind. The first is the rate the region is worked through against that region's own clock, which is its energy. The other three are the drift of its records across the seam network in the three directions of chapter fifteen, which is the quantity Newton listed second among his definitions and called the quantity of motion: momentum.

Put the four numbers into the matrix and take the determinant. Every observer agrees on it. What they agree on is the square of the mass.

Move the terms to the other side of the equals sign and the statement about the determinant becomes the relation everybody writes down, which says how tightly a region's energy, its mass and its drift are locked to one another.

$$E^2 = m^2 + p^2$$

Here E, m and p are the energy, the mass and the momentum, in units where the speed of light is one.

Energy is positive, so the relation has one branch that matters. Set the momentum to zero and E equals m. Chapter seventeen turned one reach per round into the speed of light; put that speed back in, and a region at rest has E equal to m times c squared.

The corner is worth something because the conversion is squared. A helium-4 nucleus falls short of the two protons and two neutrons that make it by 0.0304 atomic mass units, and the energy released when those four combine, measured in the laboratory, is 28.3 million electronvolts, which is that shortfall multiplied by the square of the speed of light.

18.6 The beam and the calorimeter

Positivity, additivity and conservation went into classical physics as three things the world was observed to do, and Newton, Young and du Châtelet all worked with a quantity whose properties had to be granted before the arithmetic could start. They are three readings of one move: a sum of squares has a floor, separated regions share no seam, and a single-sided entry cannot be written.

Every one of the four numbers is something a region hands over when it is asked for it. A calorimeter reads a beam's energy by stopping it and measuring the heat, which is one question put to a region and answered.

Put another question at the same moment, about where the beam is, and the two do not have a joint answer, and a better calorimeter does not change that. Which pairs of questions have a joint answer is a fact about the questions. Ask the calorimeter its one question, beam after beam, and the answers arrive in stable proportions, and nothing in the four numbers and nothing written at any seam sets those proportions.

Chapter 19

Why Can't You Ask Everything At Once?

At Petco Park in San Diego on the night of 24 September 2010, Aroldis Chapman of the Cincinnati Reds released a fastball that the stadium's tracking system clocked at 105.1 miles an hour, which is 169.1 kilometers an hour.

Two instruments were pointed at that ball. A radar gun read its speed off the shift in a returned microwave pulse. A camera read its position off the light coming back from it. Both worked, at the same moment, on the same object, and neither one spoiled the other's answer. Run the radar first and the camera second, or the camera first and the radar second, and the pair of numbers comes out the same. Nobody at the ballpark has to decide which instrument goes first, because the ball has a position and a speed and both can be read off.

That is two questions about one object, answered together, in either order, with neither answer costing the other anything. Every measurement made anywhere before 1925 was assumed to work that way.

The world does not extend that to every pair of questions. Which pairs it allows, and what asking anyway costs the answer already in hand, are settled before any instrument is switched on.

19.1 Socks then shoes

Put on your socks, then your shoes. Then try it the other way.

The two orders are not the same operation. The difference is visible from across the room. Other pairs are indifferent: left sock then right sock, or right sock then left sock, and you finish in the same state either way. Dressing contains both kinds of pair.

So take the operations themselves as the objects worth studying, rather than the states they act on. You can do one operation and then another, which composes them. You can scale an operation, or add two of them, in any setting where addition means something. A collection of operations that admits all three of those moves is an **algebra**. The word carries no more than that. When every pair in an algebra composes the same way in both orders, the algebra is **commutative**. When some pair does not,

the algebra is non-commutative. Adding up a shopping list is commutative. Getting dressed is not.

Ordinary numbers are the commutative case. Add them, scale them, multiply them in whichever order suits you, and the product comes out the same. Relax that one clause, leave every other rule of the arithmetic where it was, and you have crossed the whole distance between the physics of the nineteenth century and the physics of an atom.

The failure has a size. The size is what you get by doing both orders and comparing. Write A and B for two operations, write AB for doing B and then A, and the difference is the **commutator**:

$$[A, B] = AB - BA$$

That expression is zero exactly when the two operations can be swapped freely. Its size when it is not zero says how much difference the swap makes. Socks and shoes have a large one. Left sock and right sock have zero.

Chapter nine had a version of this with money in it: two edits landing on one account, each of them correct arithmetic, and the balance depending on which one was written first. Order mattering is a property that operations can have in a world with no quantum mechanics anywhere near it. The questions you can put to a physical system are operations of exactly this kind, most pairs of them fail to commute, and the failure belongs to the pair of questions rather than to anybody's equipment.

The radar gun and the camera commute. **Two questions can be answered together exactly when their commutator is zero.** A pair whose commutator is something other than zero is a pair you have to choose between.

19.2 Ten days on Helgoland

In June 1925 Werner Heisenberg was twenty-three and his hay fever had swollen his face badly enough that he left Göttingen for Helgoland, a red sandstone island in the North Sea with almost nothing growing on it and therefore almost no pollen. He spent about ten days there.

The problem he took with him was the hydrogen spectrum. Heat hydrogen and it emits light at a set of sharp wavelengths, the same set every time, catalogued since Johann Balmer fitted a formula to four of them in 1885. The standard picture had electrons on orbits, which reproduced those wavelengths and collapsed for every atom with two electrons in it.

What Heisenberg did was to throw out the orbits. Nobody had ever seen one. What a laboratory actually produces is a table: for each pair of atomic states, how bright the transition between them is and at what frequency. So he built his scheme out of the tables, with one entry for each pair of states, and no statement anywhere about where an electron is.

Then he had to multiply two tables together, because the energy of a system involves the square of a quantity, and squares need multiplication. The rule he was forced into combines a row of one table with a column of the other. It produced an answer. It also produced a different answer when he did the two tables in the other order.

He sent the work to Max Born, who read the multiplication rule and recognized it, recalling it from student days as the way you multiply matrices. Heisenberg's paper reached the *Zeitschrift für Physik* on 29 July 1925. Born and Pascual Jordan submitted the sequel that rewrote the whole thing in matrix language on 27 September. The usual story of a lone twenty-three-year-old on an island leaves out that the theory arrived through three people over four months.

Position and momentum came out of that scheme as tables that fail to commute. Measure a particle's position, then its momentum, and you have performed a different physical operation from measuring its momentum and then its position, and the difference is a fixed quantity with Planck's constant in it. It does not go to zero as the apparatus improves, because there is no apparatus in the statement.

The uncertainty relation is that fixed quantity, read as a floor. The spread in position multiplied by the spread in momentum cannot go below it, and where the floor sits is set by the size of the commutator of the two questions. Confine an electron to a region the size of an atom, about a tenth of a nanometer across, and the spread in its speed cannot be smaller than roughly six hundred kilometers a second, against the two thousand two hundred kilometers a second at which the electron in a ground-state hydrogen atom actually travels. The electron is not being jostled by the measurement into a speed it would otherwise not have had. There is no state of the electron with a sharp position and a sharp speed for the measurement to disturb.

19.3 Three magnets in a row

On the night of 7 February 1922, in the building of the Physikalischer Verein in Frankfurt, Otto Stern and Walther Gerlach sent a beam of silver atoms from an oven through a magnetic field arranged to be much stronger at one edge of the gap than at the other, and let the beam land on a glass plate.

A magnet in an uneven field gets pushed toward the strong side or the weak side depending on which way it points. Silver atoms come out of an oven pointing every which way, so the deposit should have been a smear.

It was two spots.

The deposit was too thin to see at all until Stern leaned over the plate to look, and the sulfur in his cheap cigars turned the silver into black silver sulfide, which developed the trace like a photograph. Two spots, nothing in

between, and the beam had been split into precisely two by a question that could have been asked along any direction they chose to point the magnet.

That is what **spin** is, operationally. Point a magnet along a direction and you get a two-outcome measurement: the atom goes one way or the other, and the only thing you get to choose is the direction. There is no third outcome and no continuous dial. Nothing rotates, either. The name arrived in 1925, when two graduate students in Leiden, George Uhlenbeck and Samuel Goudsmit, proposed that the electron was a small spinning ball of charge, and the word outlived the picture by a century.

Chain three of these and the arrangement stops behaving like a set of properties being read off.

Send the beam through a magnet pointing up, and keep only the atoms that went up. Send those through a second magnet pointing sideways. They split fifty-fifty. Every atom in that beam had been certified up a moment earlier. Keep the ones that went sideways-left. Send those through a third magnet pointing up again, and ask the question whose answer is on record.

Half of them go down.

The atoms were up. The sideways question was asked. Half the certified-up atoms come out down. Skip the middle magnet and every atom comes out up, every time, so the middle magnet is doing it.

The temptation is to say that the sideways magnet knocked the atoms about, and that a gentler magnet would knock them about less. Gentleness has been tried, at every scale anybody can build. The answer never improves. What is going on is that the up-down question and the sideways question do not commute. The second answer overwrote the first because the two of them were never jointly available to be written.

Where the beam sits between the second magnet and the third is a **superposition**: the sideways-left beam, described in up-down terms, carries two numbers, one for up and one for down. Those numbers are **amplitudes** rather than probabilities. Probabilities are their squares. The difference is that amplitudes can cancel and probabilities cannot. The dark bands in the prologue's three-month photograph are places where two amplitudes of opposite sign met and left nothing. A beam that was secretly half up atoms and half down atoms, with nobody having looked, produces no cancellation anywhere, and a laboratory tells the two situations apart by bringing the split paths back together and seeing whether the original beam comes back.

19.4 Four multiplications

Heisenberg's tables carry one entry for each pair of states, and a two-outcome question has two states, so its table is four numbers in a square, which is a size a person can multiply on the back of an envelope.

Write U for the up-down question and S for the sideways one, both laid out in up-down terms, with the physical scale stripped off so the entries come out as plain numbers.

$$U = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \qquad S = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

U is diagonal because up and down are the answers it sorts into, plus one for the atoms that go up and minus one for the ones that go down, with zeros in the corners that would mix them. S has its entries in exactly those corners, since it exchanges up with down.

The rule is the one Heisenberg was forced into. For the entry in a given row and column of the product, run along that row of the left table and down that column of the right one, multiply the pairs and add.

Do US, which is the sideways question asked first, and then do SU. Both products have zeros down the diagonal. US comes out with a one in the top right and a minus one in the bottom left, and SU arrives with those two entries carrying the other sign. Subtract, and nothing cancels: a two in the top right, a minus two in the bottom left. The two orders are exact opposites, entry for entry.

Those four numbers are the middle magnet. Half of the atoms certified up came out down because a subtraction anybody can do in a minute refuses to vanish.

Try a pair that behaves. Keep U, and write a second table with any two numbers you like down the diagonal, five in the top left and two in the bottom right, zeros in the other corners. Multiply it both ways round with U. Both products have five in the top left, minus two in the bottom right and zeros in the corners, so the difference is zero in all four places, whichever two numbers you picked. Tables that sort the atom into the same two answers commute, always, and the radar gun and the camera are a pair of that kind, which is why nobody at Petco Park had to decide which went first.

19.5 A question that squares to itself

Take the smallest well-behaved kind of question: one that admits yes or no and nothing else. Ask it, get an answer, and ask it again straight away. The second answer agrees with the first. Whatever operation the asking performs, doing it twice does what doing it once does.

An operation with that property is a **projector**. It has a name from geometry, because the operation that flattens an arrow onto a plane has the same property: flatten a flat thing again and it does not get flatter. Every yes-or-no question about a physical system is one of these, and every

projector is a question. Its opposite is what is left of the identity operation when you remove it, so a question and its opposite add up to certainty.

A complete list of mutually exclusive alternatives is a set of projectors that never overlap and that add to the operation that does nothing at all. The three magnets gave two such lists on the same atom, one for up-down and one for sideways, and neither list is a refinement of the other. The list a question is asked inside of is its **context**. In richer systems than a single atom the same question sits inside many different complete lists.

To turn questions into odds you need one more instrument, and it is arithmetic. Add up the diagonal entries of a square array of numbers. That sum is the **trace**, and has the one property that makes it usable: rewriting the array in a different set of directions leaves the trace alone. The trace of a projector counts how many independent directions the question says yes to.

A **state** is the thing that assigns odds. It is a rule that hands every question a number between zero and one, hands the question with the certain answer a one, and adds up correctly across a complete list of exclusive alternatives. Probability enters this world as an additive weight on yes-or-no questions rather than as a count of how often something happened in a long run of trials. Every such rule can be written as an array of its own, called a **density matrix**, paired with the question through the trace:

$$p = \mathrm{Tr}(\rho P)$$

Here P is the question written as a projector, the Greek letter rho is the density matrix holding the state of whatever you are asking about, and p is the probability of the answer yes.

Run it on the beam between the second magnet and the third. In up-down terms that beam is the array with a half in all four places. Asking whether the atom went down is the array with a one in the bottom right and zeros elsewhere. Multiply the two, add the diagonal entries of the product, and out comes one half, which is the number the third magnet reported. Then set the two corner halves to zero, which turns the beam into one that really is half up atoms and half down atoms, and run the same multiplication. One half again. Every up-down question hands the two arrays the same number. The corners are the only place their difference lives.

The arrows those arrays act on, with lengths and angles supplied by the state, form a **Hilbert space**. Physics courses usually start there, with the space as the given object and the algebra of questions built afterward out of operations on it. The order runs the other way. What an observer has is a bounded list of questions with a composition rule, and the space is a way of writing that list down: pick a state, and the algebra plus the state manufacture a space of arrows on which the questions act as

operations. Different observers with different question lists manufacture different spaces. They agree wherever their lists overlap, which is the only sense in which the wave function is a public object.

The manufacturing takes a few steps. None of them begins with a space. The algebra acts on itself, so that every question is both an operation and a thing operated on. The state supplies the geometry: it hands a number to every question, so it hands one to each pair of them, and a rule that gives a number to each pair of things is what lengths and angles are made of. Throw away whatever comes out with zero length, and what is left is a collection of arrows carrying the algebra as operations on it.

Two questions can be answered together exactly when a single complete list of alternatives is fine enough to answer both. That condition is chapter five's, wearing different clothes: compatible measurements are measurements that leave the same thing alone.

19.6 Averaging over what you do not hold

An observer holds a patch. The questions it can put are the ones on its side of the seam. The full description of the world involves everything past it.

So the state it holds is manufactured by averaging: take the description of the whole thing, and for every question you can actually ask, work out the odds and throw away everything that only a question from the far side could have detected. The operation is called the **partial trace**. Its output is a density matrix, which is why density matrices exist at all. A density matrix is what a description looks like after somebody has been cut out of the picture.

The averaging cannot be run backward. Two different worldwide descriptions can average to the same patch-level density matrix, and once you hold the density matrix there is nothing on your side of the seam that distinguishes them. It is a shadow. No cleverness turns it back into the object, because the object had features the averaging was built to delete.

A courtroom artist settles the same question every working day, with a few minutes, a partial view, and a drawing to hand in. The drawing has to be consistent with what was seen, and it must not commit to a detail nobody saw, because a detail that happens to be wrong is worse than a vague one. Edwin Jaynes wrote the rule down for physics in the *Physical Review* in 1957: among all the descriptions consistent with what has actually been measured, take the one with the largest entropy, which is the one that adds nothing else. The averaged description a patch holds is exactly that: the maximum-entropy state consistent with the questions it can put and the answers it has recorded, and every feature it lacks is a feature nothing on its side could have established.

19.7 Sorting the mail

Watch what writing a record does to an algebra.

Take a complete list of exclusive alternatives, and take any question at all. Cut the question into the pieces that live inside each alternative, keep those, and delete every piece that straddles two of them. That operation is called **pinching**, a mail room: letters keep their contents, they go into labeled bins, and what is destroyed is every comparison that only made sense between two bins.

Pinching is what writing a record does. The usual name in physics for the process is decoherence. The name describes the deletion rather than explaining it. What comes out of the mail room is an algebra with a property the original did not have: every surviving question commutes with every alternative on the list. The record cannot be disturbed by anything else the observer can ask, which is the whole of what makes it a record. You can read it without changing it, it reads the same the second time, and a neighbor who reads it gets your answer.

The set of things in an algebra that commute with everything in it is the **center**. Something is in there whatever the algebra is. Take the table with ones down the diagonal and zeros in the corners, the operation that does nothing, multiply either magnet by it in either order, and the magnet comes back with its four entries where they were. Doing nothing commutes with everything. Records live in the center.

Everything in the center commutes with everything else in it, so the center is a commutative algebra. A commutative algebra of this kind has one shape available to it. Strip a table of its off-diagonal entries and what is left is a list of numbers, one for each alternative: a value attached to each way things could have turned out, which is a function on a set of outcomes. A classical description has that in it and nothing else. The reason the public world wears that shape is a theorem about multiplication rather than a story about what a measurement does to a particle.

Look at what an observer finds when it looks at its own center. Questions with definite answers. Answers that survive being read. Answers that survive being copied to a neighbor, and that come back the same. Order that never matters, because everything in the center commutes with everything. Which is a complete description of the world of tables and chairs and pointer readings, arriving as the commuting part of a structure that is not commuting, rather than as a separate layer of reality sitting underneath.

The center is also the invariant part. Chapter five defined a symmetry as a change that leaves a named quantity alone, and said that a symmetry is the claim that part of your description was never physical. Chapter eleven built the record map, which takes the state of a network and returns the part that survives comparison. The center of an observer's algebra is what

every internal change leaves alone. Symmetry, the record map, and the classicality of anything anybody can write down are one structure looked at three ways.

The relation between the two layers is sharper than a metaphor about shadows, and runs one way only. Upward, the record layer sits inside the operator layer perfectly: there is a map from records into questions that is faithful, that preserves sums and products and opposites, that lands region by region exactly inside each region's own algebra, and that gives every record the expectation value it had. Records are questions that commute with everything the observer can ask.

Downward there is no such map at all. A map that respected products would have to hand a single number to a whole block of questions that do not commute with each other, and no single number does that job, because the block contains pairs whose two orders of composition differ and a number multiplied by a number does not care about order. The best available downward map is an average. It preserves positivity and totals, it leaves records alone, and it fails to respect multiplication: there is an explicit projector in the block whose average of its square differs from the square of its average.

So the classical description is a subalgebra plus an average, the embedding is exact, and the average loses multiplication. There is no projection of the world onto the world of records. The reason is arithmetic rather than epistemology.

Condition on a record after it is written, which is the operation that makes records usable at all. The state you hold afterward is the one that gives that question probability one and keeps every compatible question's odds unchanged, which is the update rule Gerhart Lüders wrote down in 1951. Apply it twice and the second application does nothing. The operation is a retraction onto certainty: one step takes you into the set of states that agree with the reading, and further reading leaves you where you are. That is why a committed record is a stable classical fact inside a structure whose questions mostly refuse to commute.

Which inverts the usual complaint about quantum mechanics. **Noncommutativity is what creates constraint.** If every question could be answered alongside every other, every observer's answers would be one row of an enormous spreadsheet, any two rows could be filled in independently, and nothing would connect what one observer holds to what another one holds. A world of that kind has no seams in it, nothing to reverse engineer, and no structure to find. The limits are the content.

19.8 The observer, upgraded

Chapter three gave three items, arrived at by asking what the job requires and finding the cheapest object that does it: a bounded view, somewhere to write, and a reading that makes a difference. The third unpacks into three of its own, since a reading that makes a difference needs a way to update, a way to compare, and enough persistence to still be there at the next comparison. Those five become four objects. First, a local algebra of questions, which is the bounded view, bounded because the interface is. Second, a record algebra sitting in the center of that one, which is the somewhere to write, and centrality is what makes the writing stay put. Third, an interface that reads and updates the same way each time, so that comparison gives the same verdict on Tuesday as it gave on Monday. Fourth, enough checkpoint data to fix the odds on every question the observer can put next. The thermostat on the wall has all four: a question list one item long, a record of the setting, a comparison it performs identically every time, and a state that says what it will do next. The five-item version tells you what to look for on a bench. The four-object version can be handed to somebody else, because each of the four is a mathematical object with a name, and two observers can be asked whether they have the same one.

19.9 One generator

Chapter eighteen's screwdriver turned an eighth at a time and drove a screw the whole way into the wood, one small move repeated being what a smooth change is made of. Do the same to an algebra. A change that is reversible and that preserves the structure, meaning it respects sums, products and opposites, is the algebraic version of a rigid motion, and a continuous family of them, starting from the change that does nothing, is a flow: for every real number t, one such change, with the change at t plus s equal to doing the one at t and then the one at s.

On a full block of non-commuting questions, every flow of that kind is generated by a single time-independent quantity: one operator, fixed once at the start, whose repeated infinitesimal application produces every member of the family. Marshall Stone proved the correspondence in the *Annals of Mathematics* in 1932. The generator is what physics calls the Hamiltonian. Chapter eighteen's energy is that generator. A world with continuous reversible change in it has a Hamiltonian.

Ask how the family changes over a single instant rather than over a stretch and what comes back is the equation of motion: applied to the state it is the **Schrödinger equation**, applied to the questions it is Heisenberg's, the same statement written from opposite ends. That is the second of chapter one's three axioms, arriving here as a consequence of continuity and reversibility.

The flow fixes its generator to within one real number. Add a fixed amount to the Hamiltonian everywhere at once, which is to say add a multiple of the operation that does nothing, and that piece commutes with every question in the block and moves none of them, so the family of changes comes back exactly as it was. Two generators producing the same flow differ by that one amount and by nothing else. The gaps between the energies, the ratios of the gaps, the whole spectrum up to a rigid shift, all of it is pinned by the motion. Which is why a laboratory reads differences in energy. Balmer's four wavelengths are differences. The one number the flow does not fix cancels out of every one of them.

The mirror statement is stranger and it is about the record layer. Every change of a record algebra that preserves the structure is a permutation of the finitely many labels the records can take. There are finitely many such permutations. A continuous family of them that starts at the identity has nowhere to go, since it would have to jump, so it sits at the identity forever. A purely classical public record cannot flow continuously and reversibly at all.

Put those two together. The world of pointer readings has no continuous reversible dynamics of its own, in the exact sense that its only structure-preserving motions are discrete relabelings. Whatever flows is flowing in the part that refuses to commute. The record layer changes by relabeling or it changes not at all.

19.10 A rule that fits every count

Max Born wrote the odds rule down in 1926, in a paper on collisions, putting it in the body of the text as the amplitude and correcting it to the square of the amplitude in a footnote. That footnote is the **Born rule**, one of the three axioms chapter one said quantum mechanics arrived with. Textbooks teach the squaring as a postulate.

Andrew Gleason derived it in 1957, in the *Journal of Mathematics and Mechanics*, from almost nothing: any assignment of numbers to yes-or-no questions that lands in the unit interval, gives the certain question one, and adds up across every complete list of exclusive alternatives is the trace against exactly one density matrix. The argument assumes no continuity anywhere and no shape whatever for the rule. There was one hole. Gleason's argument needs three dimensions or more, it fails at two, and two is where a spin lives. Paul Busch closed it in *Physical Review Letters* in 2003 by widening the questions to include blunt ones, which answer yes with a weight instead of sorting the world cleanly in two. The wider version covers every finite dimension including two.

A theorem that short, reaching that far, invites doubt, since it appears to produce the whole of quantum probability from bookkeeping.

Consider a constructed table with eight settings, a pair of exclusive answers at each, and an expected frequency for each answer. The entries are chosen to fit a specified state; no instrument was run to produce them. Then go looking for another odds rule that fits the same table, with the trace formula given no privileges in the search.

One exists, and it is explicit. It reproduces every prescribed frequency exactly, as an identity between whole-number ratios rather than as a fit. Every question it scores lands between zero and one. The question with the certain answer gets one. A question and its opposite get numbers that add to one. And on every sum of exclusive questions that the table contains, it adds up correctly.

As a rule over all weighted questions, it belongs to no state. No density matrix reproduces all its assigned weights; matching the finite table does not fix the weights outside it.

Where it breaks is a pair of questions outside the table. Take the question this rule scores at 143/512, blunt enough that two copies of it side by side are again a legal question. Adding up demands that the pair score twice 143/512, so double it: 143/256. The rule scores the pair 35/64, which over the same denominator is 140/256. The gap is exactly three parts in two hundred and fifty-six. That single pair is enough to prove the rule belongs to nobody's state.

That locates the boundary. Additivity restricted to the sums in the table amounts to nothing more than the numbers in each setting adding to one. A demand that weak leaves the fake alive. The work is done by requiring the adding-up to hold for every compatible sum of weighted questions, including those outside the table. Impose that, and the same prescribed frequencies leave exactly one state standing, without a continuity assumption and without any appeal to long runs.

So a finite table of counts never selects the probability rule by itself. What selects it is the demand behind the theorem: a question carries the same weight whichever setting it is asked in, and compatible alternatives add. For an observer that demand is the overlap rule applied to questions, two procedures that ask the same question owing it the same answer.

19.11 Constraint

The commutator says how badly two questions refuse to be asked together, and the uncertainty relation prices that refusal in meters and kilograms and seconds. Questions that refuse nothing collect in the center. What collects there is the world of tables and pointer readings. Once the refusals hold across every context at once, the odds have one form available to them. It is the trace formula.

An observer holding an algebra of questions and a state has, in that pair, everything there is to have. Constraint is the part of the pair a commuting

world could not supply, and constraint is what makes an undocumented machine readable from a seat inside it.

Constraint has a size. Put two observers far enough apart that no signal can cross between them while they work, hand each of them a pair of questions that do not commute, and compare the two sets of answers afterward. The agreement runs past anything a sealed envelope of instructions could have produced, and then it stops. Where it stops is one fixed number, 2.8284. No apparatus anybody has built has ever gone past it.

Chapter 20

Why Does Nature Cheat, But Only By Exactly $2\sqrt{2}$?

Here is a telegraph that sends messages faster than light, assembled from parts that each look as if they work. Prepare a pair of photons together and send them in opposite directions. Alice, at one end, chooses between measuring plane polarization and measuring circular polarization, and her choice is the message: one question for a zero, the other for a one. At the other end Bob does nothing subtle. He drops his photon into a laser gain tube, which turns one photon into a shower of photons like it, and inspects the shower. Alice's choice fixes which mixture Bob's half is drawn from, a shower of copies would show which mixture that was, and the distance between the two of them does not enter the timing anywhere.

Run it in your head and find the part that fails. Some part must fail, because a working version would break relativity. Nothing in the design touches relativity, though, so the error has to sit somewhere else, and saying where took three separate proofs.

The design was Nick Herbert's, and he called it FLASH, for first laser-amplified superluminal hookup. Early in 1981 the editor of *Foundations of Physics* sent his manuscript to Asher Peres at the Technion in Haifa and asked whether it ought to be published. Peres knew the paper was wrong for the reason anybody would, and he could not find the error. He recommended publication, telling the editor that the paper was obviously wrong, that it would draw considerable interest, and that whoever located the error would push physics forward. He kept his own part in it quiet for twenty years. "I was the referee who approved the publication of Nick Herbert's FLASH paper, knowing perfectly well that it was wrong," he wrote in 2002, in a note titled *How the no-cloning theorem got its name*.

FLASH appeared in 1982, in volume 12 of the journal, running from page 1171 to page 1179. The refutations beat it into print. Bill Wootters and Wojciech Zurek published theirs in *Nature* that October, volume 299, page 802, under a title John Wheeler contributed, "A single quantum cannot be cloned," and Dennis Dieks published the same result independently in *Physics Letters*, volume 92A, page 271. The second referee, GianCarlo

Ghirardi, had put a special case of the argument into his report and voted to reject; he published it the following year in *Il Nuovo Cimento*.

Every component of the telegraph works except one. The correlations Herbert wanted to exploit are real, they are stronger than any set of instructions the two photons could have carried with them, and they are measured routinely. The gain tube is the piece that fails, for a reason that turns out to be the same reason the correlations stop where they do. How much better than a sealed answer sheet can two separated people agree, and what is it about an unmeasured photon that no machine can copy?

20.1 Four questions and a referee

Put the correlation in the form of a game, with a referee who keeps score.

Alice and Bob may confer for as long as they like beforehand and may carry anything they can build. Once the game starts they are separated and no signal passes between them. The referee tosses two coins and hands each of them one question, labeled 0 or 1, and neither is told which question the other received. Each answers with one bit, written here as plus one or minus one because it shortens the arithmetic. They win the round if their two answers differ when both were handed question 1, and match otherwise.

Three of the four question pairs reward matching and the fourth rewards differing. The referee's coins deliver all four equally often. So the obvious strategy is to match always: agree in advance that both will answer plus one, whatever they are asked. That wins three rounds in four. Seventy-five percent.

Nothing sealed does better. Write Alice's answer sheet as two numbers, one for each question she might be handed, and Bob's the same way. Score the round by multiplying the two answers together, which gives plus one when they match and minus one when they differ. Over many rounds each of the four question pairs earns an average of those products, a number between minus one and plus one that runs from perfect agreement to perfect disagreement. That average is the **correlation** for that pair. Take the four correlations, add the three that reward matching, subtract the one that rewards differing, and call the total the score. A strategy that satisfied all four demands at once would score four.

For a sealed answer sheet the score is Alice's first answer times the sum of Bob's two, plus her second answer times the difference of Bob's two. Pick four numbers of your own, each plus one or minus one, and add up the two pieces. Bob's two answers are each plus or minus one, so they are equal or they are opposite. If they are equal, the difference is zero and the sum is plus or minus two. If they are opposite, the sum is zero and the difference is plus or minus two. Either way the score is exactly two in magnitude, every time, for all sixteen sheets, and averaging over any

mixture of sheets keeps it there, because an average of numbers no bigger than two is no bigger than two. The ceiling on sealed instructions is John Bell's, from 1964. The four-setting form is the one John Clauser, Michael Horne, Abner Shimony and Richard Holt wrote down in 1969.

20.2 Fifty-two cards

To do better, Alice and Bob need one description of the pair rather than two descriptions of the parts.

A playing card carries a suit and a rank. Four suits, thirteen ranks, fifty-two cards. Describing a card means fixing both. The count of descriptions is the product of the two counts rather than their sum. The operation that builds the descriptions of a pair out of the descriptions of its parts is the **tensor product**. Its first property is that arithmetic: dimensions multiply.

"Hearts" and "seven" is a description that comes apart into a fact about the suit and a fact about the rank. "The red seven or the black eight" does not come apart into anything. It fixes how color and rank go together while leaving both of them unsettled, and learning either one hands you the other. Most joint descriptions are of that kind. The ones that factor are a thin special case.

Alice's questions and Bob's are far enough apart that no light ray crosses between them during a round, so every question on her side commutes with every question on his: the bracket across the gap is zero, and the order in which the two of them ask changes nothing. What does not commute is Alice's first question with her second.

20.3 Four lines

Write Alice's two questions as A0 and A1, and Bob's as B0 and B1, each one an operation on the pair's joint description that returns plus one or minus one. Asking a plus-or-minus question twice returns you to where you started, so each of the four squares to the do-nothing operation. Build the score as an operator: A0 times the sum of Bob's two, plus A1 times the difference of Bob's two. The score of any state is the average of that operator in that state.

Square it. Sixteen terms come out. Four of them are squares, and each question squares to the do-nothing operation, so those four add to four times it. Eight more cancel in pairs, plus against minus, once you use the fact that everything on Alice's side commutes with everything on Bob's. The last four collapse into a single product of brackets.

$$(\text{score operator})^2 = 4 + [A_0, A_1]\,[B_1, B_0]$$

The square bracket is chapter nineteen's commutator, the amount by which the order of two questions matters, and is zero exactly when the two can be answered together. The four is a sealed answer sheet's two, squared. The second term is the whole of the quantum contribution, a product of two commutators, one from each side.

Put a number into that line. Suppose Alice can answer both of her questions at once and Bob both of his, which makes both brackets zero. Anything multiplied by zero is zero, so the second term goes. The square of the score operator is four times the do-nothing operation. Something that squares to four times the do-nothing operation reads out as two or as minus two. An average of twos and minus twos lands between them. The sealed answer sheet's ceiling follows from that line alone. The 1969 inequality is this identity with its brackets switched off.

How big can that term be? Every operator has a size, the largest factor by which it can stretch anything it is applied to. The name for it is the **operator norm**. A question returning plus or minus one stretches nothing, so its norm is one. A commutator of two such questions is a difference of two products of things of norm one, so its norm is at most two. Multiply Alice's bracket by Bob's and the product has norm at most four. The square of the score operator therefore has norm at most eight, so the score operator has norm at most the square root of eight. An average can never exceed the largest value being averaged. So no state, no choice of questions, and no amount of prior conferring gets a score above $2\sqrt{2}$, which is 2.8284.

Work the square root of eight out by hand rather than reading it off a machine. Eight is four times two, and the square root of a product is the product of the square roots, so the square root of eight is twice the square root of two. The square root of two is 1.41421, the length of the diagonal of a square one unit on a side, and the first irrational number most people are shown. Double it. 2.82842. Multiply 2.8284 by itself and you get 7.9998. The two ten-thousandths short of eight are the digits dropped off the end.

One line read at its two ends gives both ceilings. With the brackets at zero it says four, and the score tops out at two. With them opened as wide as they go it says eight, and the score tops out at 2.8284. A bracket reaches two only where doing Alice's two questions in one order comes out as the exact negative of doing them in the other. A pair that refuses each other by less has a ceiling below 2.8284. The four in the middle does not move, which leaves the 0.8284 of score between the two answers held entirely inside a product of two commutators.

Divide one ceiling by the other. 2.8284 over 2 is 1.4142, the diagonal again: questions that refuse to be asked together beat a sealed envelope by the same factor a square's diagonal beats its side.

Carry that back to the referee. A score of S wins a round half the time plus S divided by eight, which turns the sealed ceiling of two into seventy-five rounds in a hundred and the ceiling of $2\sqrt{2}$ into 85.36. Just over

ten extra rounds in a hundred is the whole of the quantum advantage in this game. Stuart Freedman and John Clauser went looking for those ten rounds at Berkeley with photons from a calcium cascade, found the margin where quantum mechanics says it is, and published on 3 April 1972.

Count what went into that ceiling. Four things that square to one, and two of them commute with the other two. Geometry never appeared. There is no space in it, no distance, no light cone, no wave, and nothing that could be called a particle. The ceiling holds anywhere four questions behave that way and sizes can be compared. This world's questions behave that way. Boris Tsirelson found the number in 1980 and published it in *Letters in Mathematical Physics* under the title "Quantum generalizations of Bell's inequality".

The gap between two and 2.8284 is that commutator term. Non-commuting questions are why joint knowledge has limits at all, and the same brackets, multiplied together, are why separated agreement has a ceiling. What stops Alice from answering both her own questions at once is what lets her agree with Bob more often than any answer sheet permits. Take the brackets to zero and the score falls back to two, and the world becomes a world of answer sheets.

20.4 Real numbers all the way

Set up two slots, each with two outcomes, so the pair has four dimensions. Put them in the state whose two answers are perfectly opposed along whichever direction you ask. Have Alice ask along two perpendicular directions and Bob along the two directions bisecting hers, an eighth of a turn from each. The score comes to exactly $2\sqrt{2}$. Every matrix in that arrangement is real: the state, the four questions, the whole construction. The most strongly correlated thing two separated systems can be put into needs no imaginary numbers.

Change one thing about it. Take a state diagonal in the record basis, which is to say a state that has been written down, an outcome list with weights sitting in chapter nineteen's commuting center, and read it with four readouts diagonal in that same basis. Operators diagonal in one basis commute, so both brackets vanish, the second term in the square is zero, and the score is at most two, for every such state and every such readout. The written layer of the world plays this game exactly as well as an answer sheet does, and for the same reason: a committed record is an answer sheet, filled in and handed over. Everything an observer can point at, quote or file has come through that door, which is why the public world behaves like a world of objects with properties, and why the extra ten rounds in a hundred have to be hunted for with equipment.

The test has been run with the loopholes closed. The Delft experiment of 2015, the one the prologue met, deposited its records, and they hold

245 rounds that count. The two laboratories won 196 of them, a score of 2.42, past the ceiling of two that any written record with its answers fixed on each side can reach. A world of answer sheets would produce a result that good no more often than about one time in twenty-five. The correlations came from the unwritten layer, where Alice's two questions refuse to commute, and only the outcomes were written down. That layer reaches the full $2\sqrt{2}$.

20.5 What the gain tube would have needed

A copier is one fixed machine, chosen in advance, that takes an unknown state in one slot and a blank in the other and hands back the unknown state in both. A machine you have to configure for the state you are about to copy is useless on a photon arriving from a laboratory you cannot see.

Two states have an overlap, a number from zero to one, which is how often one of them passes a test built for the other: zero for perfectly distinguishable, one for identical. Reversible operations preserve overlaps, because preserving the structure overlaps are computed from is what reversible means here. Run the copier on two states and compare the outputs. Each slot contributes the original overlap, so the output overlap is the input overlap multiplied by itself. Preserved and squared at once, so the overlap equals its own square, and exactly two numbers do that: zero and one.

The machine copies states that are perfectly distinguishable, or states that are identical, and nothing in between. Herbert's gain tube was asked to copy a photon whose polarization nobody had measured, which is the in-between case and the only case the telegraph needed. That two-line argument rules out the design.

Look at which states the copier does handle. Overlap zero, perfect distinguishability, which is what a record is: an outcome that has been committed, sitting in the part of the algebra that commutes with everything, as distinguishable from another record as one word on a page is from a different word. Copying is forbidden for unwritten states and permitted for written ones, without limit and without disturbance.

What is copyable is exactly what has been written down as a record, and that is why redundancy works: why a backup is a backup, why a thousand people read one printed page and report the same words, why a measurement copied into a hundred notebooks is the same measurement in all hundred.

20.6 Twelve directions

Alice and Bob have been free to point their equipment any way they like. For a finite calculation, restrict their settings to the twelve vertex directions of the icosahedron in chapter six. Take the perfectly opposed two-slot state,

which is unchanged by a common rotation, and the usual spin questions along those directions.

Four settings drawn from twelve, in order, is 20,736 choices, which is few enough to take all of them. The best score is reached by 960 of the 20,736, and all 960 run the same way. The twelve ports pair off into six axes, corner to opposite corner, and any two of those six meet at 63.43 degrees. Each of the 960 uses three of the six, one axis shared between Alice and Bob. Three of the four pairings put her axis at 63.43 degrees to his, where the correlation has size one over the square root of five, 0.4472. The fourth puts her on his own axis, where the size is one. The four line up rather than cancel, so the score is one plus three-fifths of the square root of five in magnitude. Multiply the square root of five, 2.2360679775, by six and move the point one place left: 1.3416407865. The one on top of that makes 2.3416407865. Sealed instructions on the same four settings land on plus two in eight of their sixteen patterns and on minus two in the other eight, and nothing outside that.

Twelve directions are enough to beat the answer-sheet ceiling. They stop short of 2.8284 because of the corners. Reaching that ceiling wants Bob's directions an eighth of a turn from Alice's, and an icosahedron's twelve corners offer exactly four angles between them, zero, 63.43 degrees, 116.57 degrees and 180. An eighth of a turn is not among them.

The ceiling came out of two objects and nothing else: an algebra of questions, carrying its brackets, and a state assigning odds to them. That pair is the entire inventory of an observer. Go through it looking for a clock and you come back holding a list of questions and a list of numbers, and nothing in either list moves. Something takes time anyway.

Chapter 21

Why Does Time Feel Like It's Moving?

Augustine was in his forties and a year or two into the bishopric of Hippo Regius, a port on the North African coast that is Annaba in Algeria today, when he wrote the eleventh book of the *Confessions*. The thirteen books took him from about 397 to about 400. Twelve of them are about his life and his God. The eleventh is about time, and in its fourteenth chapter it comes apart in his hands.

"What then is time? If no one asks me, I know; if I want to explain it to a questioner, I do not know."

The line gets quoted as a piece of wisdom. Augustine was after an instrument. He works through the possibilities in the chapters on either side of that sentence and discards each one: the past has gone and cannot be laid alongside a ruler, the future has not arrived and cannot either, and the present has no extent, because anything with extent divides into a part that has gone and a part that has not come. He ends the chapter holding a ruler with nothing in front of it.

What he settles on, having run out of things in the world to measure, is that the measuring happens in the mind: the soul stretched across a memory, an attention and an expectation, so that the quantity being compared against a ruler is an impression rather than an interval of anything. He gives it a name, *distentio animi*. Philosophers have discussed the name for sixteen hundred years. It moves the ruler indoors and leaves the question of what is being laid against it exactly where it was.

Sixteen centuries of instrument-building have not improved his position. In 1967 the thirteenth General Conference on Weights and Measures redefined the second as the duration of 9,192,631,770 periods of the radiation from a particular transition in a cesium-133 atom. That is the most carefully specified duration ever defined, and what it specifies is an interval. A cesium clock counts periods and reports a count. Subtract two counts and you have an interval. Every timepiece anybody has built, from a notched candle to the fountain clocks that keep international atomic time to sixteen digits, reports a difference between two events, and none of them has an output for the present moment, because there is no part of the mechanism that could produce one.

Chapter sixteen pulled three objects out of the single word and answered two of them. Order is which committed repair depends on which: the later commit must certify that it read a version written by the earlier one. The wiring gives a hundred and twenty seam pairs that share an observer and three hundred and fifteen that share nothing, and the certified reads of a particular history fix which pairs it actually ordered. Direction is why that relation runs one way, and it came out of an arithmetic loss inside a single repair, seven arrangements of a seam collapsing into one. The third object was left standing. Why does anything seem to be going by, and what sets the rate? Hand somebody the whole web of dependencies with every strand of it pointed, and they are holding a diagram. Something has to make it go.

21.1 An algebra and a state

An observer holds an algebra of questions, which is everything it can ask about its own patch, carrying the brackets that record which pairs can be asked together and which cannot. And it holds a state, which is the odds it puts on every one of them.

That pair fixed the floor under position and momentum, and put the ceiling on separated agreement at $2_{\sqrt{}}2$ and not a thousandth higher. It also decided which part of the world is classical, by collecting the questions that commute with everything into a center and letting the records live there.

Take the pair apart looking for a clock.

The algebra is a list of questions with a rule for composing them. Nothing in a list ticks. The state is an assignment of numbers to that list. An assignment of numbers is a table. There is no parameter anywhere in either object, no reading that changes, nothing that distinguishes one moment from another, because the specification contains no moments. If time were a primitive ingredient of this world, it would have to be posted in from outside the pair by something that owns a clock, and chapter four's inside-only clause rules that out at the top of the argument: there is no outside to post it from.

Chapter ten's run makes the same point in hardware. Eighty-two thousand patches worked all 122,880 seams between them into agreement. The cycles the run is reported in were put there by whoever wrote the report. No patch in the network counted a cycle or waited for one to end. The machinery has no tick in it anywhere. Whatever supplies the rate has to be manufactured out of a list of questions and a table of odds, because nothing else was ever delivered.

21.2 A room left alone

Leave a room alone for long enough and everything in it arrives at one temperature. The coffee, the mug, the table, the air above the table. The description of that room has collapsed to a single number. The collapse discards almost everything that was ever true about it.

Look at what the single number does. Bring any object at all into the room and the number tells you which way heat will move, for that object and for every other one, without anybody consulting how the room got there. Physicists run that argument the other way daily: write down the system, work out which equilibrium it settles into, and the equilibrium follows from the machinery.

Run that argument backwards.

Chapter sixteen took a settled arrangement, the kind the repair law leaves where it finds it, and called the odds it assigns the reference. A state of that kind is in equilibrium, and equilibrium is a word with a hole in the middle of it. In equilibrium with respect to what? Being settled means being unchanged by something. A state is a list of odds. No change is named anywhere on the list.

So work out what it would have to be.

A change that qualifies is a reshuffling of the observer's questions among themselves that respects the algebra: questions go to questions, sums to sums, products to products, and the whole thing is reversible. Line those shuffles up, one for every real number, and the family is what chapter eighteen called a flow, driven by the move it repeats, its generator.

Two demands pick out a flow, the only two in the construction.

The first is that the state's odds are unchanged by it. Every question is worth after the shuffle exactly what it was worth before. That is what being settled means, written out.

The second uses the non-commuting part. In a world where every pair of questions could be answered together, the state's weight for asking A and then B would equal its weight for asking B and then A, because there would be nothing to tell the two orders apart. Chapter nineteen's entire subject is that they differ. So the state carries a gap between the two orders, one gap for every pair of questions. The gap is a fixed feature of the state rather than a nuisance to be argued away. The second demand is that swapping the order costs exactly one unit of the shuffle, the same one unit for every pair of questions, with the amount of shuffling measured along a direction at right angles to the flow's own parameter. That last clause is complex analysis, the one step in the whole construction that cannot be done on a table.

For a state that rules nothing out, giving every possible yes some chance of coming up, which the mathematics calls faithful, the two demands pick out exactly one flow. In finite dimensions faithful means that every eigen-

value of the density matrix is positive. The flow's parameter is a bare number, with no seconds in it.

Minoru Tomita, who lost his hearing at the age of two and published very little in a long career, wrote the construction down in 1967 in a manuscript that the people who received it found close to unreadable. Masamichi Takesaki worked out what it said, passed the results to Jacques Dixmier in the summer of that year, and published a usable version in 1970 as a thin volume of Springer lecture notes. The thing is called **modular flow**, after the algebraic machinery it drops out of rather than after anything it does, and its generator, in chapter eighteen's sense of the word, is called the modular Hamiltonian.

In finite dimensions the generator has a one-line recipe. Take the logarithm of the density matrix and put a minus sign in front.

$$K = -\log \rho$$

Here rho is chapter nineteen's density matrix, the array standing for the state the observer holds, and K is the generator of the flow that state determines. Chapter sixteen's logarithm counted questions and so ran in base two. The one in this recipe is the natural logarithm, the kind chapter ten used for the spread of eighty-two thousand records, where a doubling is worth 0.693 instead of 1. In an eigenbasis of the density matrix it applies the logarithm to each positive eigenvalue; it is not the entrywise logarithm of an arbitrary array.

21.3 One system, two outcomes

That recipe can be run by hand on the smallest system there is.

Take a two-outcome system, the sort Stern and Gerlach put on a glass plate: one question, two answers, nothing in between. Give the two answers energies of zero and E. Put the system into a thermal state at temperature T, which means the odds of finding it in the upper answer, relative to the lower, drop off as the exponential of minus E divided by T. Temperature is written in energy units throughout, the way physicists write it when they want that exponent to come out as a bare number. The exponential drop-off is the shape of every thermal population in physics. It sets the color of a hot poker and the fraction of a gas that is ionized.

Write the state as its array. Take the logarithm, which acts on the two diagonal entries separately. Put a minus sign in front. What comes out has E divided by T in the upper slot, zero in the lower, and a constant added to both, contributed by the requirement that the odds add up to one.

The constant does nothing. A generator shifted by a constant produces the same flow, since the constant piece commutes with the whole algebra and shuffles nothing. Delete it.

What survives is the energy divided by the temperature. The flow the state determines is the system's own evolution in time, measured in a unit the temperature sets.

Put numbers on it, and take a pencil. A state for this system is a slip with two odds written on it. One ratio settles both of them, because the two have to add up to one. Say the lower answer comes up twenty times for every once the upper one does. Write the slip: 20 in 21 and 1 in 21, and there is nothing else about the system left to write down.

The lower slot of the generator holds zero, so the whole generator is whatever stands in the upper slot. That is the logarithm of the ratio between the two odds.

$$K = \log \frac{\text{odds of the lower answer}}{\text{odds of the upper answer}} = \log 20 = 2.996$$

Add it up yourself. Twenty is two times ten, so its logarithm is the sum of theirs: write 0.693 for the two, write 2.303 for the ten, and the twenty comes to 2.996, four thousandths short of 3. The check runs from the other end, since the exponent 3 gives 20.0855, a shade over the twenty on the slip.

The recipe said the surviving entry would be the energy divided by the temperature, so this system's energy is 2.996 times its temperature, and the flow moves the system along at that rate.

The slip fixes the ratio of energy to temperature and nothing more. Double both and the slip comes out the same. A thermometer supplies the scale.

21.4 Whose clock

The flow belongs to the state, so it changes when the state changes.

Two observers holding different states extract different flows, and the flows run against each other at rates nothing requires to be constant. There is no third flow that either of them is failing to match, because a third flow would need a third state and no such state is held by anybody. Chapter seven left the shared present with nobody to hold it, and Hafele and Keating's flying clocks had a reading to report only when they stood next to the bench clocks. The rate goes the same way. It belongs to whoever holds the state, the way a patch's questions belong to the patch. Two observers can agree on a common coordinate or a calibration, the way two ships agree on Greenwich, and the agreement is something they build.

Halve the temperature with the energies held fixed and the generator doubles, while the ratio of the odds is squared. Twenty to one becomes four hundred to one, and log 400 is twice log 20, about 5.99. The same energies in a colder state give a generator twice as large.

Run the recipe on the state repair settles into by itself and the flow does nothing at all. That state gives every arrangement with the same total the same odds, so every log-odds gap is zero and every question stays where it was. A region with nothing left to settle drives no clock. A clock appears once an observer has read something. Condition the settled state on the first current an observer records, a unit crossing into a corner of three ports, and the odds the observer holds on the three arrangements in play are five thirteenths, four thirteenths and four thirteenths. The gap between the first and the other two is the logarithm of five fourths, about 0.223, and the flow that state determines turns at exactly that frequency in its own parameter.

Clocks built on events are local in the same way. An observer's event clock ticks when something happens at its own ports, while repairs elsewhere go on without it. Two observers' event clocks can run at the same average rate and disagree in their readings all the same. Count one observer's ticks between every second tick of the other: the average is two, as it should be, and the spread around that average has a variance of sixteen. Equal rates do not make equal readings. Let one clock's ticks trigger a change in the field the other one counts and the spread changes with it, because the law connecting two clocks is a law about the events they count.

"Time is subjective" is usually said loosely. Here it names an object: a one-parameter family of transformations, extracted from one specific state by a construction with no free choices anywhere in it, different for different states, and computable by hand on a two-outcome system in the time it takes to add two logarithms.

21.5 The center does not move

Records live in the center of the algebra: the questions that commute with everything, which is what makes a record readable without disturbance, copyable to a neighbor, and the same on the second reading as on the first. Run the flow on the center and it does nothing at all. Every element of the center is left exactly where it is by every transformation in the family, at every value of the parameter.

The flow was built to close the gap between asking A and then B and asking B and then A. An element of the center has no gap with anything, because it commutes with everything in the algebra. So there is nothing for the flow to fix about it, and it sits untouched. A record changes only when something writes to it, and writing is a repair event, a different operation from the flow. The price of commuting with everything is having no clock of your own.

Chapter nineteen proved the harsher version, which reaches past this flow to every continuous motion a record layer could have had. The only structure-preserving moves a record algebra has are permutations of its

finitely many labels, and a continuous family of permutations would have to jump to get anywhere, so it sits where it started. The classical layer changes one discrete relabeling at a time, or it holds.

Put those two facts side by side. **You experience time passing because you are not made only of records.** The part of you that has been committed, written down, made copyable and readable and safe from disturbance, is the part with no flow on it. Whatever is going by is going by in the other part, the questions that refuse to be answered alongside each other, which is where the flow lives and where the entire supply of rate comes from. An observer built out of records and nothing else would hold a complete description of a world, would be able to read it, copy it and compare it with a neighbor's, and would sit inside it at no rate at all, the way chapter seven's recipe card lies on a kitchen counter.

21.6 The flow has no arrow

The flow is reversible. Every transformation in the family has its inverse in the family, which is the transformation at minus the same parameter. That was settled before either demand was written down: what qualified as a change at all was a reshuffling of the questions that loses nothing, and a shuffle that loses nothing can be undone.

So the flow cannot be the source of the arrow, and chapter sixteen was right to hold the two questions apart rather than answer them together. What supplies the arrow is what chapter sixteen said supplies it: the commit, seven arrangements of a seam collapsing into one, an arithmetic loss inside a single move that no relaxation of any rule anywhere recovers, with quantum coherence draining away in the same move.

Put the two together and you have the whole of what an observer undergoes, in two ingredients doing two different jobs. Committed repairs are irreversible and supply the direction. The flow the state determines is reversible and supplies the rate. Take away the commits and an observer holds a parameter with no arrow along it. Take away the flow and it holds a bare count of events with no duration between them. A clock on a wall is both at once: a count of commits, and a rate at which the counting proceeds.

An observer that keeps time joins the two on purpose. Passive repair never turns the flow, because a repair nobody acts on is not one of the flow's shuffles. At every repair event that meets its region, including one that moves nothing across the seam, the observer advances its own state by one fixed step of a flow whose generator comes from the repair law itself, and conditioning on the whole closed history of those events gives that flow a thermal reference of its own. Four correlations between the bits recording movement at its ports fix the flow's two frequencies, and those frequencies predict what an interference probe of the observer reads,

with nothing fitted. The events order the steps, and a calibrated clock laid against the count turns it into seconds.

21.7 The region's own clock

Chapter eighteen defined energy and left one item open inside the definition.

Repair works through a region's records at some rate. That rate is the region's energy. Every rate is counted against a clock. That one was counted against a clock nobody had built, with no universal time on offer to borrow instead.

Here it is. The region holds a state, the state determines a flow, the flow has a parameter, and the parameter is the clock. Energy is well defined region by region because every region carries the thing its rate is counted against, manufactured out of what the region holds. Two regions running at different rates differ in the states they hold. That is the whole of the difference.

Energy is the generator of a region's advance along its own clock. The generator of the modular flow is minus the logarithm of the region's state. For a region in thermal equilibrium those are the same operator, counted in units of the region's own temperature and up to a constant that moves nothing, and the second is a recipe for computing the first out of what the region holds, with no constants fitted to measurements. A thermometer turns the units into joules.

21.8 The ship that light never catches

Take an observer accelerating steadily through empty space, and have it keep the acceleration up.

Chapter seventeen counted the motions that take one sky onto another and found six. Three of them turn the sky rigidly, and three gather the constellations toward the direction of travel and thin them away from behind, leaving every circle on the sky a circle, and those three are the boosts. Steady acceleration is a boost applied continuously: one parameter, running.

Chapter seventeen also drew the light cone. The cone does something to this observer that it does not do to anyone traveling at a constant speed. Light chasing an accelerating ship from far enough behind never catches it. Signals from that part of the world arrive at no point of the trip, however long the trip is allowed to run. There is a boundary behind the observer, and past that boundary sits a region the observer can never ask a question about, at any time.

That boundary fixes the observer's algebra: the questions about everything on this side of it, and no others, because there are no others it could

ever put. And the state is the vacuum, the state of a region with nothing in it, which is the same state everywhere in empty space and is therefore the only other thing the calculation needs.

Feed those two objects into the construction and turn the handle. Out comes a one-parameter family of transformations of the region's questions, determined by the pair, with nobody choosing anything. The family is the boost that carries the accelerating observer along, at every value of the parameter, exactly, with the parameter proportional to the time the observer's own wristwatch records.

Joseph Bisognano and Eyvind Wichmann proved it for quantum field theory in two papers in the *Journal of Mathematical Physics*, "On the duality condition for a Hermitian scalar field" in volume 16 in 1975 and "On the duality condition for quantum fields" in volume 17 in 1976. They were not working on a theory of time. The question in front of them was technical, whether the operators available inside a region exhaust the operators that commute with everything outside it. The flow of the region turned out to be a Lorentz transformation. Their theorem stood on a quantum field theory handed to them complete. Chapter twenty-two builds that field theory out of the repair history and finds the same match one level down: the flow of a region's thermal state is the time its own counted clock keeps.

An observer's clock is a physical article. It ages, it can be photographed, and relativity says exactly how much of it elapses along a given path through the world. The flow a state determines is an abstraction pulled out of an algebra by two demands about bookkeeping, with no clock and no motion anywhere among its inputs. In the one case where both can be computed independently and laid side by side, they are the same motion, at every value of the parameter.

21.9 Warm

The flow has one more property. A thermometer can read it.

The state of empty space, restricted to the region the accelerating observer can ask about, sits unchanged under the flow, and the gap between its two orders of asking costs exactly one unit of that flow. Those are the two demands, and every state meets them against its own flow by construction, which makes them unremarkable until you notice what the flow is here. The flow is a physical motion. The observer is riding it. That second demand, the gap costing one unit, is the condition a physicist writes down to say that a system sits in a heat bath, and here it is written against the ship's own motion. The observer is sitting in what any thermometer it carries will report as a warm bath, in a region of space with nothing in it.

Bill Unruh worked out what an instrument would read. Put a small detector on the ship, something with two levels that can absorb a quantum

and record that it did, and count how often it fires. A detector at rest in empty space never fires. The same detector, accelerated, fires at the rate it would fire at while sitting in a bath of radiation at a definite temperature, and the acceleration alone fixes which temperature.

He did that calculation in 1976, in a paper called "Notes on black-hole evaporation" in *Physical Review D*. The temperature is proportional to the acceleration. The constant of proportionality is about four parts in ten to the twenty-first of a degree for every meter per second squared.

Which is why nobody has run into this by accident. Warming yourself by a single degree calls for an acceleration around twenty-five billion billion times the strength of gravity at the Earth's surface, sustained. Every acceleration a human body has ever survived leaves the reading far below what any thermometer built resolves.

Chapter 22

Why Do All the Clocks Agree?

Put four clocks on a table and look at what each of them is.

The first is the one chapter twenty-one built: a region holds a state, the state determines a flow, and the flow's parameter is the region's own time. It has no dial. It is a family of transformations of the region's questions, labeled by a bare number. The second is a tally. Something happens at the region's edge and a counter goes up by one. The third is a ripple: a wave moving through whatever fills the region, whose phase you can read off with an interference probe. The fourth is light, or whatever plays its part, arriving from somewhere else after a delay that a ruler and a stopwatch would call a distance divided by a speed.

Physics writes one letter for all four. The t in the Schrödinger equation, the t in the wave equation, the t in the light cone and the t that thermal physics turns into an inverse temperature are typeset identically. In a theory that starts with a spacetime, the question never comes up, because the coordinate is handed out before the first equation and every mechanism is required to use it.

This world hands out no coordinate. Each of the four clocks has to be built from the one thing there is, a continuing history of repairs, and each could in principle have come out running at its own rate, in its own units, with no conversion between them. They do not. This chapter builds them one at a time out of the same history and shows that they read one parameter, with conversion factors that each construction supplies. Along the way it picks up a field, a wave, a distance, a minus sign and a quantum field theory, because those turn out to be what agreeing clocks are made of.

22.1 Counting every touch

Start with the tally, since it is the easiest to build and the easiest to get wrong.

An observer region sits in the repair traffic. Seams around it are visited one at a time, each visit compares the two sides and commits the repair law's answer, and many visits change nothing at all, because the two sides

agree. The obvious clock counts changes: tick whenever a unit actually moves across one of the region's seams. It is a bad clock.

Chapter twenty-one's timekeeping observer did something else. It stepped its flow at every repair event that met its region, including the ones that moved nothing across the seam. That looked like bookkeeping. It is the whole difference between a clock and a noise source. Write down how the region's state changes from one tick of the change-only clock to the next and the result has a mode that flips sign at every tick. A transformation with a negative mode has no logarithm a clock can use, which means no continuous family of positive steps runs through it: there is no way to say what half a tick of that clock would be. Count every touch instead, idle visits included, and the tick-to-tick transformation is positive throughout, its logarithm exists, and the clock can be read at any fraction of a tick.

Call the result an **incident clock**: a counter that advances once whenever a repair meets the region, whatever the repair does.

Different regions are touched at different rates. A single port sits on five or six seams. A whole twelve-port carrier is met by thirty-three. A patch of the screen is met by a number that grows with its area. So the raw counts disagree from the first visit. Divide each count by the number of touches its region expects per unit of the history's progress, and the disagreement goes away. The normalized count of every region converges to one parameter, uniformly over any stretch you care to fix, with the probability of missing by more than any given margin falling off as the history gets finer. A port and a carrier read the same time, because the visit schedule is the same schedule for both and each count is a fair sample of it.

The finite version of this was in chapter twenty-one: two clocks with equal average rates whose readings between ticks spread with a variance of sixteen. That spread is real. What the normalization does is make it small against the count as the count grows. A clock is a tally large enough that its fluctuations stop mattering, and the incident clock gets there without anybody synchronizing anything.

22.2 A field out of the tally

Now look at what the tally is counting through.

Every seam visit leaves a trace in the occupations of the two ports it touched. Average those occupations over long stretches of the history and the averages fluctuate, slightly, around the settled value. The fluctuations are not independent from port to port. A unit that left one port has gone somewhere nearby, so neighboring averages are correlated, and the pattern of correlations is a field: a number at every place, with a definite rule for how numbers at different places hang together.

The rule is one operator. Call it A. It is built from the seams, it is positive, and it is local in the strict sense, each port talking only to the ports it shares a seam with. On the screen it has the heat-trace dimension of a surface, two, as it should for a surface. Its inverse is the field's correlation between two places, and as the screen is refined, the correlations on the finer screens converge onto a single continuum operator with a definite domain. Nothing was put in by hand. The occupations were always there; the operator is what their long-run statistics turn out to be.

Then look at the field at two different times. The correlation between the field read at one clock reading and the field read at another decays exponentially in the difference between the readings, at rates set by A, mode by mode. The slow modes forget slowly and the fast ones fast. The clock in that statement is the normalized incident count from the last section. The field's time is the tally's time: the same parameter that counts touches is the parameter in which the field relaxes, with no conversion factor between them other than the normalization already fixed.

That is two clocks agreeing. It is also the first sign of something larger, because A has now appeared twice, once as the field's shape in space and once as the field's rate of forgetting in time.

22.3 A wave on the same count

A field that only relaxes is a heat-like thing. It smooths and forgets. Waves keep their phase, and keeping phase needs something the passive repair traffic does not do.

So give the observer a small quantum memory of its own, kept out of the repair traffic, and couple it to the incident clock in the simplest possible way. Each tick, idle touches included, triggers one fixed step of a walk on that memory. Nothing else touches it between ticks. After N ticks the memory has taken exactly N steps.

The walk is built from the same seams as the field, and its amplitude follows a cosine. In the limit of a fine screen the amplitude is the cosine of the clock reading times the square root of A, with one fixed conversion factor set by how many seams the clock's port sits on. The operator that set the field's shape and its forgetting sets the wave's frequencies. The clock that counted touches drives the oscillation. There is no second clock inside the wave; the wave's time is the tally, read through a cosine.

Because everything here is a finite construction with exact rational odds, it makes predictions down to the last digit. Prepare the memory, run the clock, and read it with an interference probe. The probability of the plus outcome is 1 before any tick, 21/22 after one and 91/110 after two. Over a complete two-visit history, every branch included, it comes to 1297/1320. Change the rule so that only visits that actually move something trigger a step, and the same history gives 7901/7920 instead. Reprepare the memory

before every step and it gives 71/72. Those are different numbers from the same repair history, and the difference is how an experimenter tells which clock a region is actually keeping.

22.4 Distance is the first answer

Two regions, some way apart on the screen. Disturb the first and watch the second through the wave.

For a while nothing arrives. The wave's response at the second region is exactly zero, with no small tail leaking ahead, until the clock reaches a definite value. After that the response switches on. The value at which it switches on is the distance between the two regions, measured with the field's own operator as a ruler: the largest amount a function can change from one region to the other while its slope, as A measures slope, nowhere exceeds one.

The relaxing field gives the same number by a different route. Its leakage from one region to the other over a short clock interval falls off like a Gaussian in that same distance, and the exact rate of the fall-off reads the distance back out. The bound is sharp. Nothing arrives before the distance, and something does arrive as soon as the clock passes it.

So distance and duration come out of one apparatus in one unit. A distance is how long the clock runs before a disturbance can be answered, and that makes the speed of the answer one, by construction, for every pair of regions and every observer keeping this clock. Chapter seventeen found one invariant speed from the angle-preserving maps of the sky and noted that the number it takes is a choice of units. This is the same speed from the other end. The cone is the first response of the wave, and the speed of light is the exchange rate between the count the tally keeps and the ruler the field supplies. The meter and the second were tied together in 1983 for exactly this reason: two units for one thing.

22.5 Three directions and one minus sign

The screen supplies two directions. The third is declared: the construction chooses a radial interval and fills it in with the screen's own records.

Compare the field with itself some number of repairs later. Those comparisons are lag records, native readouts like any other, and a weighted mixture of them, the longer lags weighted more heavily, gives the field a new coordinate running away from the screen: the deeper the coordinate, the longer the lags it is built from. Independent long stretches of one history then fill that coordinate in, so the field along it fluctuates on its own instead of copying the screen. Glue the radial coordinate onto the screen's field and the joint operator is the screen's A plus the second derivative along the new direction. Its heat-trace dimension is three. The memory

of a region sits on its surface, as chapter twenty-four will show, and the depth of that memory in the history is the third direction of the space it describes. The interval, its ends and the filter that fills it are choices of the construction. Repair has not selected them, and the same filter would attach a third direction to any two-dimensional screen.

Run the wave in all three directions. Each tick now triggers two fixed pulses, one carrying the screen's frequency and one carrying the radial frequency, and the two pulses act along axes that do not commute. That one fact fixes how the frequencies combine. The joint frequency is the square root of the sum of their squares, the way the hypotenuse of a right triangle combines its sides. Use axes that commute and the frequencies simply add, which is the wrong answer and fails the direct check. The square-root rule is the dispersion law of a wave moving at one speed in three directions, and it has fallen out of the order in which two pulses act on a memory.

Write the wave's action, the quantity whose stationary paths are the wave's motions. The three space directions enter with the coefficients of A and the radial piece, all positive. The clock direction enters with the opposite sign. Given the wave, that sign is no further convention. The wave is second order in the clock and its spatial operator is positive, and an action with that structure assigns the clock direction a minus sign against the three others. The coefficients of that action are a metric, and its signature is minus, plus, plus, plus. That is the signature of the spacetime of special relativity, recovered from a tally, a positive operator and a cosine.

The native metric is measurable rather than smooth: it is defined almost everywhere, with coefficients pinned between two positive bounds, and it carries a crease where the original faces of the screen meet. Laboratory physics asks for smooth spacetimes and it gets one. Pick any finite list of measurements, wave readings, clock comparisons, interventions, detector outputs, and any tolerance however small. One smooth static spacetime with one time and three space directions reproduces every joint probability on that list within that tolerance. No finite set of observations tells the crease apart from a smooth spacetime.

22.6 The region's own clock, on the wall

Now go back to the first clock on the table, the flow from chapter twenty-one.

Take one mode of the wave and give it the energy the clock assigns: two levels, one frequency apart. Put the mode into a state and ask the chapter twenty-one question, which flow that state determines. Then ask when that flow is the same motion as the wave's own evolution, run in clock time.

The answer is a theorem with no slack in it. The state's own flow equals the clock evolution, rescaled by a constant, exactly when the state is ther-

mal for the clock's energy: odds dropping off as the exponential of the energy times that constant. The constant is the inverse temperature. The match has to hold axis by axis, so a state that leans the right amount in the wrong direction fails, and two modes leaning by the same amount at different frequencies would need two different constants and so fail to share one. A region in equilibrium at one temperature has one conversion factor for every mode, and that factor is its inverse temperature, which is what chapter twenty-one said a thermometer reads.

That puts four readings on one parameter: a tally that counts touches, a field relaxing in its count, a wave oscillating in its count, and the flow of the region's own state, with conversion factors fixed by the seams and the temperature. The region's own time is the time on the wall, multiplied by the temperature.

22.7 Quanta

One mode in one memory is a single quantum system. A field has many quanta. Take many independent copies of the thermal mode and add up their fluctuations, scaled the way the central limit theorem scales them.

Two of the summed quantities come out special. Their commutator has expectation value exactly i, the defining relation of a position and momentum pair, and the products of their exponentials converge to the expectation values of a thermal free quantum field. The two-point function at two clock readings has a real part that oscillates as the cosine of the frequency times the difference in readings, and an imaginary part that oscillates as the sine. The imaginary part is the commutator. It is the piece that has no classical counterpart, and it runs in the same clock as everything else on the table.

Put the screen modes and the radial modes in together and the joint thermal state of the three-dimensional wave has a modular flow, in chapter twenty-one's sense, that is time translation of that same wave, rescaled by its temperature. Bisognano and Wichmann took a quantum field and its vacuum as given and proved that a region's modular flow is a motion. Chapter twenty-one used their result about an accelerating observer to show that the region's flow is a physical motion. That result needed a quantum field theory to stand on. The field theory is here, built from the repair history and running on the observer's own count.

It does not stop at free fields. Add an interaction, a quartic term coupling the radial modes and the screen's constant mode, and the interacting thermal state can be prepared from native operations, its flow fixes an ideal clock, and a native incident clock driving the same interacting system runs in step with that flow. The gauge and matter actions of part six are written on fields of this kind; this construction does not produce them.

22.8 Seconds

None of the four clocks has seconds in it. The tally counts touches, the wave oscillates in that count, the region's flow runs in that count multiplied by a temperature, and light covers one unit of the field's distance per unit of count. Each construction fixes its own conversions between the four, which is the point of the chapter; that one conversion holds across all the constructions is unproved. The one conversion left is to the clock in a metrology laboratory, and it is a single number: how many touches of some region make one period of the cesium transition chapter twenty-one opened with. Fix that once, and every other duration in this world is quoted in seconds by arithmetic.

Physics wrote one letter for all four clocks and never had to justify it, because its spacetime came with a coordinate and its equations were required to use it. Here the coordinate is an output. It is the one parameter that a tally, a field, a wave, a light cone and a region's own state all turn out to share, and the reason they share it is that all five are readings of the same continuing repair history, taken by observers who count everything that touches them. Appendix D shows how to run the construction and check the numbers in this chapter on a laptop.

Chapter 23

Why Does Writing Something Down Cost Heat?

A solid-state drive has no moving parts, and it gets warm while it is being written to. Most of that warmth is ordinary electrical loss, and engineers shave it down every year. Underneath it sits a floor that no engineering removes, in any machine anybody has built.

The first limit of that kind turned up in steam. In 1824 a French military engineer of twenty-eight paid a Paris publisher named Bachelier to print six hundred copies of a book about steam engines, which the scientific community of the time did not read. Sadi Carnot's *Reflections on the Motive Power of Fire* runs to a hundred and eighteen pages and asks the question every engine builder in Europe wanted answered, which is how much work you can get out of a fire.

His answer has no pistons in it. The most work any engine whatever can deliver, per unit of heat it swallows, is fixed by two numbers: the temperature of the thing it takes the heat from and the temperature of the thing it dumps the remainder into.

$$\eta = 1 - \frac{T_{\text{cold}}}{T_{\text{hot}}}$$

The Greek letter names the efficiency, the fraction of the heat that comes out as work. The two temperatures are the hot source and the cold sink, both counted from absolute zero. Brass or iron, steam or air, a better valve, a cleverer linkage, a bigger boiler: the ceiling sits where those two temperatures put it and no engineering moves it.

Carnot got that while believing heat was a weightless fluid called caloric, which fell from hot to cold the way water falls down a millrace and did as much work on the way. Caloric does not exist. Every heat engine built since has come in under the number he got out of it.

Cholera reached Paris in the spring of 1832 and killed something near twenty thousand people in the city that year. Carnot caught it on 24 August and was dead within the day, at thirty-six. His belongings were burned as a precaution against contagion, and nearly all of his papers went

with them. What survived was a set of working notes his brother Hippolyte published in 1878, in which the fluid had been quietly dropped.

Look at what the surviving formula says. An engine's ceiling depends on the temperatures and on nothing else. Chapter twenty-one reached the same quantity from the other end, out of an algebra and a state, with no furnace anywhere in the argument. So what is a temperature a count of, and what is the floor under the warm drive made of?

23.1 The doorkeeper

On 11 December 1867 James Clerk Maxwell wrote to Peter Guthrie Tait describing a way to break the second law of thermodynamics using a hypothetical being.

Take a box of gas at one uniform temperature, divide it with a wall, and put a door in the wall. Temperature is an average: some molecules in there are moving fast and some slow. Station a being at the door with good eyes and a light touch. When a fast molecule approaches from the left, it opens the door and lets it through to the right. When a slow one approaches from the right, it opens the door and lets it through to the left. The door is weightless and frictionless, so opening and closing it costs nothing.

Wait. The right side heats up and the left side cools down, out of a box that started uniform. Then run Carnot's engine off the difference, and run it again, forever, in a box that nobody has done any work on.

Maxwell called it a finite being and later wrote that it was really a valve. William Thomson named it a demon in *Nature* in 1874, meaning the word in its Greek sense, a spirit working quietly in the background.

The demon survived every attempt to kill it for sixty-two years, because every attempt went looking in the wrong place. People examined the door, the hinges, the light the demon needed to see by. Leo Szilard moved the search in 1929, in a paper in *Zeitschrift für Physik* whose title says where he was looking: on the decrease of entropy in a thermodynamic system by the intervention of intelligent beings. He stripped the gas down to a single molecule in a box, with a partition dropped in the middle. The demon looks, records which side the molecule is on, and uses that one recorded fact to let the molecule push the partition outward and lift a weight. The work per cycle comes out as the temperature times the logarithm of two, out of a box at one temperature, from one bit of recorded information.

The books balance only if the bit costs at least what the bit earns. Rolf Landauer, at IBM, worked out in 1961 which operation carries the charge, in a paper called "Irreversibility and heat generation in the computing process". Looking can be done for free. Forgetting cannot. An operation that takes two distinguishable states of a memory to one state has thrown away the distinction. The distinction has to leave the device, because the

only exit a memory has is heat. The floor he computed for that heat depends on nothing but the temperature outside the device.

$$E \geq k_B T \ln 2$$

E is the energy released as heat when one bit is erased, T is the temperature of the surroundings the erasing device sits in, and the constant converts a temperature in kelvin into an energy in joules. The logarithm of two is chapter sixteen's one question, the one you ask to tell two possibilities apart.

Put a room temperature into it, three hundred kelvin, and the bill for erasing one bit comes to 2.87 divided by ten to the twenty-first, in joules. A single photon of green light at a wavelength of 550 nanometers carries 3.61 divided by ten to the nineteenth, in joules, which is enough energy to erase about a hundred and twenty-six bits.

Charles Bennett finished the demon in a review of the thermodynamics of computation in 1982. The demon's memory is finite. It can run a few cycles free, filling that memory with records of which side each molecule was on, and then it is full, and to take another measurement it has to clear space. Clearing the space costs at least the temperature times the logarithm of two per bit, which is everything the molecule paid it. The demon was never an engine. It was a machine borrowing against its own memory, and clearing that memory cost the same every cycle.

Antoine Bérut and colleagues put the number on a bench in 2012 and published it in *Nature*. A silica bead about two micrometers across, held in an optical trap shaped into two wells, one well for zero and one for one. Push the bead into a known well and you have erased a bit. Measure the heat that leaves. As the erasure cycles were slowed down, the mean heat per erasure came down onto that value and did not go below it.

Chapter three listed three requirements for anything that counts as an observer, said that writing a record costs energy, and left the price unstated. That is the price.

23.2 Four laws, numbered out of order

Thermodynamics has four laws. They were not discovered in the order they are taught.

Start with the one everybody uses without noticing. Nobody has ever established that a bath and a bowl of soup are at the same temperature by putting the soup in the bath. You put a thermometer in the soup, wait, read it, then put the same thermometer in the bath, wait, read it, and compare two numbers taken minutes apart with an instrument that has been in contact with each and never with both at once.

That procedure is a bet on transitivity. If the thermometer settles with the soup, and the same thermometer settles with the bath, then the soup and the bath will settle with each other. The bet is a law: two systems each in equilibrium with a third are in equilibrium with each other. Ralph Fowler and Edward Guggenheim wrote in 1939 that it "could with advantage be known as the zeroth law of thermodynamics", the other three having been numbered decades before anybody noticed this one was doing work underneath them.

Transitivity is what makes temperature a number. A relation that is transitive, and symmetric, and holds between a thing and itself, sorts everything in the world into classes with nothing left over, and classes in a row can be labeled by numbers on a scale. Without transitivity you could say that this is hotter than that, pair by pair, and there would be no scale to write either of them on, no degrees, no thermometer, and no sentence of the form "the water is at 47 degrees".

The first law gets taught as bookkeeping, in two terms. The energy of a thing changes when heat flows into it and when work is done on it, and heat and work are the two channels. Work moves the energies of the available arrangements while the odds over them stay put. A piston does that. Heat changes the odds while the energies stay put. Contact with a flame does that. Do both at once in one finite jump and a third term appears, the product of the two changes, which vanishes when you change one thing at a time. Over a smooth change that product shrinks faster than either change does, which is why the textbook's two terms are exact for a process that runs continuously.

The second law arrived in chapter sixteen, in a laboratory in Canberra with a latex bead in water. Take the odds a settled arrangement assigns, call them the reference, and ask how far your own description sits from it. Under any process that leaves the reference alone, that distance never climbs.

The third law is the one popular accounts skip. Walther Nernst put it to the Göttingen Academy in December 1905: as a system is cooled toward absolute zero, its entropy approaches a fixed floor. The floor is set by how many arrangements the system has at its lowest energy. One such arrangement and the floor is zero. Many, and it is the logarithm of how many, and sits there down to the last fraction of a degree.

Ordinary ice does this. Each oxygen atom in ice has four hydrogens near it, two close and two further off, and there are many ways to satisfy that rule across a whole crystal without ever violating it locally, the same kind of frustration as chapter six's triangle of neighbors. Linus Pauling counted the arrangements in 1935 and got a floor of 0.805 calories per degree per mole. William Giauque and J. W. Stout measured the heat capacity of ice down to fifteen degrees above absolute zero and reported 0.82, give or take

0.05, in a paper of 1936. That is 3.4 joules per kelvin per mole of entropy that stays in a block of ice at the bottom of the temperature scale.

The other half of the third law is a prohibition. You cannot reach absolute zero in a finite number of steps, however good your refrigerator, because each step takes a fraction of what is left rather than a fixed amount. A sodium gas at the Massachusetts Institute of Technology was cooled in 2003 to 450 picokelvin, give or take 80, which is four hundred and fifty trillionths of a degree above a floor nobody reaches.

23.3 One rule, read twice

Four laws, arrived at over eighty years by people working on engines, gases, chemical affinities and cold. They come out of the repair rule of chapter nine, read twice.

Read it first as a rule about descriptions. Jaynes's instruction from chapter nineteen applies unchanged: among all the descriptions consistent with what has actually been measured, take the one with the largest entropy, the one that adds nothing else. Apply that with one quantity held fixed, the average energy, and the answer is forced. The odds fall off exponentially with energy, cheap arrangements common and expensive ones rare, with a single multiplier in the exponent setting how fast the fall is. Chapter twenty-one ran into that exponential from the far side, in a two-outcome system whose flow came out as the energy divided by the temperature.

That multiplier is one over the temperature. The temperature arrives as a conversion rather than as a property of anything. Fix the average energy of a description and the least assumption you can make about the rest of it carries exactly one number. That number says how many joules the world charges for one yes-or-no question. A temperature is a count of joules per bit. At three hundred kelvin the count is 2.87 divided by ten to the twenty-first of them.

The zeroth law follows from the arithmetic of that exponential rather than from a habit of laboratories. Two systems that have settled with each other carry one multiplier between them, and equality of numbers is transitive, so a thermometer that matches the soup and matches the bath has established that the soup matches the bath. There is one condition on the thermometer: the instrument must have two arrangements of different energy. A thermometer whose two states cost the same reads the same number in a furnace and in liquid helium, because the temperature shows up only in the ratio of odds between two different energies. An instrument with one energy has no such ratio to offer.

Read the rule the second way, as a rule about transitions. A repair step is handed some things that neighbors have settled between them and some things nobody has settled. What should it do? The same instruction applies: change nothing you do not have to, and assume nothing you have

not been given. Leave every settled quantity exactly as it stands, and redraw everything else from the reference, inside the set of arrangements that agree with what is settled.

That instruction picks out one map. Applying it twice does nothing that applying it once did not do, because the second application finds the same settled facts and the same reference. It leaves the reference where it is, since redrawing from the reference cannot move it. And it preserves the average of every quantity that can be read off the settled part, which is the first law in the form chapter nine derived it: a repair that changed a total would have to know the total. The heat and the work come off the same map. Shifting the energies of the arrangements without touching the settled odds is the work channel, redrawing the odds at fixed energies is the heat channel, and the cross term is what a step that does both at once picks up on the way through.

The full map settles everything unsettled in a single step. Local repair, working seam by seam, takes longer and keeps the same public record the whole way. The smallest exact case has eight states: one recorded bit held fixed, and two more bits redrawn from their odds given that record. The correlations between the two decay toward equilibrium, and the recorded bit never moves. Memory and relaxation run side by side in one machine.

The integer averaging rule has another way to carry memory. Once every port holds one of two neighboring whole-number loads, fair repair can keep swapping those loads without changing their total. Let each patch record its load while these exchanges continue. On a fixed connected network, with the stated random timing, slowly relaxing patterns leave stronger correlations in a long record than quickly relaxing ones. Their strength can be calculated from the repair rule and the prepared total. A Gaussian history limit follows without injecting Gaussian noise at the ports. The length of the record and the scale at which it is read both matter. Exchanging loads leaves the cells’ shapes and volumes where they were, so the correlations live in the loads, and two local readouts of the same loads can share a variance and differ in how separated ports correlate.

Three of the four laws drop out of that map without further argument. Distance to the reference falls at every step and never climbs, which is the second law, the data-processing fact of chapter sixteen with the process named: repair is error correction, and running your description through an error-correcting step cannot carry it further from the truth.

Every schedule of repairs on chapter nine’s twelve observers lands on the same arrangement, and things that land on one fixed point together are in equilibrium with each other, which is transitivity again, reached from the transition side this time instead of from the exponential, and with the thermometer taken out of it. And one repair step leaves every arrangement with some weight on it: the step redraws from a reference that gives everything a positive share, so a possibility that had a share

before the step has one after it. Finitely many steps extinguish nothing, which is why the floor of the third law is approached and never arrived at.

Landauer's bill is a corollary of the same map rather than a separate discovery. A step whose entropy falls by some amount expels at least that amount divided by the same multiplier. Take the amount to be one bit and out comes the temperature times the logarithm of two, in joules, per bit.

23.4 A one-way loop that obeys the law

Chapter sixteen's detailed fluctuation relation leans on one extra condition: measured against the reference, the weight a step carries from one state to a second matches the weight it carries back. The condition has a name, **detailed balance**. The integral identity of that chapter does without it.

The second law does not need it. The cheapest way to see that is a machine with three states and a permanent circulation in it.

Label three states one, two and three. From each state, the rule is: stay where you are with probability one half, or step clockwise to the next state with probability one half. There is no counterclockwise move at all. Nothing in this machine can go from state two to state one except the long way through state three, which takes four steps on average.

Check what it does to the flat description that puts a third of the weight on each state. State one receives half of its own third, from staying, and half of state three's third, from the clockwise step. That is a third. So is every other state, by the same count. The flat description does not move, which makes it the reference.

Detailed balance fails outright. The traffic from state one to state two is a third of a half, one sixth per step. The traffic from state two to state one is zero, because that move does not exist. There is a current going around the loop that never dies down.

Watch the second law hold anyway. Start with all the weight on state one, a description sitting the logarithm of three away from the flat one, 1.585 bits. One step spreads it into halves on states one and two, and the distance falls to 0.585 bits. A second step gives a quarter, a half and a quarter, at 0.085 bits. A third step brings it to 0.024. The descent is exact, it never reverses, and it happens inside a machine with a one-way street in it.

Stationarity is what the second law asks for: that the reference be left alone. Detailed balance is a stronger and separate demand. The extra content it carries buys the detailed fluctuation relations of chapter sixteen and Lars Onsager's paired transport coefficients. A world could have the second law with none of those and this loop is what it would look like.

23.5 The price of settling

Two observers hold readings of one shared quantity and the readings disagree. Before any repair, the description of that pair has two live arrangements in it and no fact distinguishing them: one bit of entropy, the logarithm of two. Repair them. Both readings go to one value. The description has one arrangement in it and its entropy is zero.

Settling a disagreement lowers entropy. Every comparison in the world that comes out resolved, and every record written into the public account behind it, takes a description with two possibilities in it and hands back one.

Nothing is violated by that. The second law is about the distance between a description and the reference. That distance is the quantity that never climbs. The bare entropy of a description is a different number. A settling process drives it down by one bit each time it settles a bit. A rule that only ever lowers entropy carries none of the second law's content. That is why the three-state loop had to be scored on its distance from the flat description rather than on its entropy.

The bit had to go somewhere. Reset a memory in contact with a reservoir and the lost bit leaves as heat of at least the reservoir's temperature times the logarithm of two, with temperature in energy units. Copying the old readings somewhere first only postpones the cost, for Bennett's reason: the copy sits in a finite memory, and clearing that memory costs the same. Chapter sixteen showed that a commit discards which of seven arrangements the seam used to hold, and priced it at 2.807 bits of the world's ability to say what it used to hold. The same commit has an energy attached: 2.807 bits of erasure at three hundred kelvin costs 8.06 divided by ten to the twenty-first joules, expelled as heat, per seam and per commit, in a universe that is making a public record out of private disagreements. That is the price against a reservoir at three hundred kelvin, and a colder world pays less.

23.6 Five instruments, one quantity

The same number has been read off five different instruments by five sets of people who did not know they were measuring the same thing.

An engine reads it as the heat it took in divided by the temperature it took it at. That ratio is what Carnot's ceiling is a statement about. A gas reads it as Boltzmann's count of the arrangements a coarse description fails to distinguish. A memory reads it as Shannon's count of the questions it takes to identify what is stored, and Landauer's price per question. A horizon reads it as an area, in square meters, the strangest of the five. And a public record reads it as what a settled disagreement threw away, the one this world is built out of.

Every record ever written has been paid for in heat, at the temperature of whatever wrote it. The heat leaves through the edge of the region holding the record, so the cost falls on the edge rather than on the volume inside it. Ask how many records a region of space can hold, and the answer comes back in square meters. Chapter twenty-nine counts them for the largest region an observer has.

Chapter 24

Why Does a Region's Memory Scale With Its Surface?

John Archibald Wheeler confessed to a crime in his office at Princeton, and the crime was tea. He would set a cup of hot tea next to a glass of iced tea and let the two come to a common temperature. Energy is conserved in that operation, every joule of it accounted for. Something else goes up and does not come back down, because the difference between hot and cold, once mixed, cannot be unmixed by any procedure anybody has found in two centuries of looking. The evidence of what he had done stays in the world afterwards, spread too thin for any instrument to gather it back up.

Then the twist, which he put in 1970 to Jacob Bekenstein, the graduate student sitting in the room: "Jacob, if a black hole comes by, I can drop both tea cups into the black hole and conceal the evidence of my crime."

Chapter twenty-three priced a deleted bit in joules and derived the law the crime obeys. The total entropy of the world does not fall. Wheeler was describing a way to make it fall, using the one object in the universe that swallows things without giving anything back. The argument bites because nobody outside can be told afterwards what was in the cups.

A black hole is a place rather than an object. The place is defined by a failure. Pack enough mass into a small enough region and there is a surface around it across which signals travel inward and never outward, made of no material whatever. A light ray sent outward from just beyond it gets away. One sent outward from the surface itself stays on the surface forever. Outside that surface, the entire object is described by three numbers, which are its mass, its electric charge and how fast it spins. Two holes agreeing on those three are the same hole to everything in the universe beyond them, however different the histories that made them.

A teacup crossing that surface is not destroyed at the crossing. For a large enough hole the cup goes through without anything local happening to it at all. The crossing is a statement about which signals can be sent back rather than a statement about the cup. What it costs is the return path. Every question anybody outside might have asked about the tea has lost its channel.

Wheeler's crime comes down to accounting. Entropy counts what somebody would have to be told to know the rest, the tea had a great deal of it, and what the tea fell into is described by three numbers.

Bekenstein was twenty-five, a National Science Foundation predoctoral fellow at Princeton's Joseph Henry Laboratories, and he came back with an address for the missing count. The entropy stayed in the world and moved onto the hole. The area of the horizon is fixed by those three numbers, and behaves the way entropy behaves: it only goes one way. Hawking had proved in 1971 that in any classical process the area of a horizon never decreases. Throw something in and the area grows. Two holes merge and the area of the result exceeds the sum of the areas that went in. Bekenstein's paper, "Black holes and entropy" in *Physical Review D* volume 7 in 1973, says that the entropy of a black hole is proportional to the area of its horizon, measured in units of a fundamental length squared, and that the sum of the entropy outside the hole and the entropy of the hole never decreases. The horizon grows by enough to pay for the teacups.

24.1 A quarter of the area

Hawking thought this was wrong, and he had a good reason: anything with an entropy and an energy has a temperature, anything with a temperature radiates, and a black hole was the one object in physics defined by letting nothing out. He set out to show that the numbers could not be made to work. What his calculation produced instead was a temperature, a thermal spectrum, and a rate of emission, published in *Nature* on 1 March 1974 in two pages titled "Black hole explosions?". The temperature fixed the constant Bekenstein had been unable to pin down, and turned a proportionality into an equation.

Take the area of the horizon, count it in units of a very small square, and take a quarter of the count.

$$S = \frac{A}{4\ell_P^2}$$

Here S is the entropy of the hole, A is the area of its horizon, and the length written ℓ_P is the Planck length, which is what you get by combining Newton's constant, Planck's constant and the speed of light into a length, about 1.6 times ten to the minus thirty-fifth of a meter. Nothing on the right-hand side of that equation refers to anything inside.

Ordinary storage scales with volume. Double a horizon's radius and the volume inside it goes up by eight while its memory goes up by four. Every further doubling widens that gap by another factor of two. Bekenstein worked the arithmetic into a comparison in *Scientific American* in August 2003: a black hole one centimeter in diameter has an entropy of about ten

to the sixty-sixth bits, which is roughly the thermodynamic entropy of a cube of water ten billion kilometers on a side.

A region's memory is the number of distinguishable things it could be doing, counted as bits: the number of yes-or-no answers a complete description of it would take. That count, for any region of space, is fixed by the area of the surface around it and takes no account of the volume inside. Take a sealed room and double the shelving in it. The ceiling on how many distinguishable things the room can be doing is unchanged, because the walls have not moved.

Black holes are where that ceiling was found, and it applies beyond them. Try to pack more entropy than it allows into a region of ordinary space. The energy required to hold the entropy collapses the region, a horizon closes around it, and the count lands back on the surface where the bound said it would. Gerard 't Hooft stated the general version in a 1993 paper called "Dimensional reduction in quantum gravity": the maximum information in a region of space is set by the area of its boundary, black hole or no black hole.

Chapter thirteen got the shape of it out of a wiring diagram, with no gravity anywhere in the argument. Everything anybody can learn about a patch arrives through its ports, so two interior states exposing the same readings on every port are one state to the entire network, and what can be known about a region is a count of arrangements of its boundary rather than of its inside. That fixes the form. What it leaves open is the exchange rate, how much one unit of boundary is worth. The exchange rate came off a horizon.

24.2 What a cut costs

Chapter nineteen said what an observer holds and how it comes to hold it. The description on your own side of a seam is manufactured by averaging over everything only the far side could have detected, and chapter sixteen said what the averaging costs: the description that comes back has an entropy, a count of what its holder would have to be told.

Take two patches together in a state pinned down as far as anything can be pinned down, where every question that can be settled about the pair at once has been settled. Cut it in two and average over one half. What comes back is vague. The definiteness was living in the relation between the two halves. The cut went through it. That content is **entanglement**. The number measuring how much of it the cut severed is the entropy of the half left behind.

Of either half. Two sides of a cut through a definite whole come back with exactly equal entropies, however lopsided the two pieces are. A thimble of water cut out of an ocean returns the same number as the ocean it was cut from. That means the smaller side sets the ceiling for both, and it means

the number was never a property of what is inside either piece. It belongs to the cut.

Chapter six's coins hold a record of that shape, with no quantum mechanics in the building. Alice and Bob write one word, same or different, which states the relation between two coins and says nothing about either coin: complete about the pair, silent about the halves. Read the entry and you know what will be on Bob's coin the moment you have seen Alice's, and you know nothing whatever about either one before that. The quantum version differs in one respect, which chapter twenty measured. There are no faces sitting underneath the entry for it to be a description of. The correlations run further than any set of faces could account for. The entry is the object.

In this world the cut around a region runs along a screen, a surface tiled with patches, and everything the region can show the rest of the world crosses it. The screen is two-dimensional, the surface chapter fifteen found when carriers were joined into a sphere. The region's memory is the capacity of that screen. Every patch on it is the same twelve-port carrier holding the same finite amount, so the number is a count of the screen's patches. Add a patch deep inside a region, wired only to other patches inside it, and the count does not move. It moves when the screen around the region grows.

24.3 The doorway

Two rooms and a doorway between them. You are in the near room with instruments, and every signal from the far room comes through the doorway, so anything you can learn about what is happening over there arrives as something happening in the doorway. Tell somebody exactly what the doorway is doing, in full detail. What does the far room then add?

If the answer is nothing, the doorway **screens** the two rooms. Given the doorway, the two rooms are independent: they can correlate with each other only through what passes between them, and you have been handed that.

Run a cable through the wall and the screening breaks. The two rooms share something the doorway never carried, the far room goes on telling you things about the near one after the doorway has been fully described, and no amount of watching the doorway recovers what the cable is doing. Whether a boundary screens is a fact about how the rooms are wired.

Ask how much two systems tell you about each other once a third is given in full, and the answer is the **conditional mutual information** of that arrangement. Elliott Lieb and Mary Beth Ruskai proved in 1973 that the quantity is never negative. Zero is the interesting value. Zero means the doorway carries the whole relation.

Chop a network into three pieces of exactly this shape: a region, which is the **cap**; a thin band of patches around its edge, which is the **collar**; and everything else, which is the exterior. When the collar screens the cap from the exterior, the state splits: fix what the collar is doing and the inside and the outside become two independent descriptions, with nothing running between them that the band has not accounted for.

24.4 Rebuilding from the collar

If the collar screens exactly, a map acting on the collar can reconstruct the joint state of cap, collar and exterior from the cap and collar, preserving their correlations. A corresponding map reconstructs the cap from the collar and exterior. These are **recovery maps**. Dénes Petz wrote down a canonical construction for a specified process and reference state; under the exact screening condition it reverses the loss of the missing region on that state.

That is what local repair looks like in quantum form. A patch that disagrees with its neighbor rewrites itself from the data on the interface they share, and a recovery map says exactly how much such a rewrite can restore.

Chapter ten ran eighty-two thousand patches into agreement across 122,880 seams with no patch consulting anything beyond its own neighbors. The quantum version asks the same of every patch, with the collar standing in for everything beyond it.

Screening need not be exact. Omar Fawzi and Renato Renner proved in 2015, in *Communications in Mathematical Physics* volume 340, that when it is nearly exact some recovery channel nearly works. If the conditional mutual information is ε in natural-log units, the recovered state has squared fidelity at least $e^{-\varepsilon}$ and trace-norm error at most $2\sqrt{1-e^{-\varepsilon}}$.

A cap, a collar and an exterior are the three pieces a local rule needs. The collar is a surface one patch thick wrapped around the cap, and the seams that run into it are the ones a repair of the cap reads.

24.5 Hamming's Monday

At Bell Telephone Laboratories in 1947 a relay computer in New York spent a weekend on Richard Hamming's jobs. He was junior enough that the weekend was his entire allocation: start the run at five on Friday evening, collect the output at eight on Monday morning, with nobody in the building in between. The machine hit an error early on, dropped the work, and moved on to the next job in the queue. Two Mondays running he came back to a blank reel of tape and had nothing to hand his colleagues in Murray Hill. What he wanted to know was why a machine clever enough

to notice that a relay had gone wrong could not work out which relay and set it back.

He published the answer in 1950, in the *Bell System Technical Journal*, volume 29. Four bits of data go in and seven come out, the extra three computed from the others and arranged so that a receiver holding a word with one bit flipped can name the position that moved instead of merely reporting that something did. Chapter thirteen took this world's carrier to be a code of that family, with the seams doing the checking, and settled there that a connector hands over its checks and hands over nothing whatever about how far apart two legal configurations sit.

What Hamming had beyond his list of checks was seven relays bolted in a row on a frame. A row has positions and the positions can be counted off with a finger. A surface made of patches has no row and no frame. What it has is the number of patches in it. That number arrives without units.

24.6 Counting a cut and measuring an area

The scalar wave description makes one connection precise. The measure multiplying its action is also the area measure of a screen slice in the associated spacetime metric. Proper three-dimensional volume contains an additional clock factor. Holding the count fixed therefore does not automatically hold physical volume fixed.

A second calculation tests a tempting shortcut. Forty-five four-port regions have the same entropy ledger under the chosen native reference, although their numbers of crossing seams run from ten to eighteen. That entropy cannot equal one universal nonzero constant times the crossing count. This does not refute the whole-screen area reading; it distinguishes the boundary of a region on a screen from the area of the screen itself. The entropy theorem needs the geometric quantity actually appearing in its statement.

24.7 The exchange rate

Bekenstein's proportionality, with Hawking's quarter in it, runs in both directions. Left to right it prices a horizon: hand it an area and it returns an entropy. Right to left it prices the screen. Measure a horizon in Planck squares, meaning squares one Planck length on a side, take a quarter of the count, and what comes back is the horizon's memory in natural units, the number of its patches times what each one holds. So every natural unit of memory is worth four Planck squares of surface. No patch holds more than another, because every patch is the same twelve-port carrier and nothing printed on one makes it larger than the next.

A count of memory and a measurement in square meters are one measurement in two sets of units. The conversion between them is fixed once

and holds everywhere in the world. Four Planck squares comes to 1.04 times ten to the minus sixty-ninth of a square meter, putting about 9.6 times ten to the sixty-eighth natural units of memory in a square meter of boundary.

That count is what a surface has in place of a material. The area of a horizon is its memory quoted in square meters, and the memory is a count of patches that each hold the same amount, so the area comes in whole numbers of one small piece, the same piece on every surface in the world. Take any region, count the patches on the screen around it, multiply by what each one holds and by four Planck squares, and out comes what a ruler laid against that boundary reports.

24.8 Draw a square on a horizon

Draw a closed curve on a horizon, dividing the surface into two pieces. The curve can be routed along seams, between patches, so that every patch belongs to one piece or to the other. The two counts add up to the count for the whole.

Entropy divides the way the patches divide. The memory of a piece of screen is set by how much of the screen it holds, and the seams along its edge do not enter. The smallest case shows it plainly. Take two groups of four ports on a single carrier, one whose edge crosses ten seams and one whose edge crosses fourteen, and in the settled state the two carry exactly the same entropy. On finer screens the information a piece shares with the rest of the screen stays a sliver, growing far more slowly than the seams along its edge, so the memories of the pieces add up to the memory of the whole to within that sliver. Halving the surface halves the number, and a piece with twice the area of another gets twice the count. Equal areas carry equal counts, on any horizon, whatever sits behind them, because the area is the count.

So the number belongs to the piece rather than to the whole surface, putting a figure on a square you can cover with your hand. Ten centimeters on a side, drawn anywhere on any boundary in the world, and the memory it holds comes to about ten to the sixty-seventh natural units.

A patch knows its screen one seam at a time, through its own ports. A ruler laid along the same screen reads it four Planck squares to every natural unit of memory.

Wheeler's two teacups went in carrying entropy nobody could gather back afterwards. The surface around the hole grew by enough to cover them. That surface is made of no material at all. The only property it has to offer is a size. A count of what somebody would have to be told came out balanced, exactly, against a shape.

Chapter 25

Why Does Gravity Look Like Geometry?

The crew of the International Space Station float about four hundred kilometers above the ground, where the Earth's gravity is close to nine tenths as strong as it is at the surface. Gravity has kept hold of them. The station and everybody aboard are falling around the planet together, and that is the whole of the trick.

Albert Einstein worked this out in the autumn of 1907. He was examining patents in Bern and owed Johannes Stark a review article on relativity for a yearbook of radioactivity and electronics, and he was assembling it at his desk in the office when the thought arrived that he described fifteen years afterward, in a lecture in Kyoto on 14 December 1922, as the happiest of his life. If a person falls freely, he will not feel his own weight.

Every other influence in physics discriminates. A magnet lifts a steel paperclip and does nothing at all to an aluminum one of the same mass, because the response depends on what the object is made of. Gravity pulls harder on a heavier object by exactly the factor that makes a heavier object harder to shift, the two cancel, and everything falls at one rate, which is why David Scott could stand on the Moon in August 1971 with a hammer in one hand and a falcon feather in the other and drop them together for the camera. Write the description in a frame that falls along with everything else and gravity drops out of it.

The sharpest test of that cancellation flew between 2016 and 2018 in a French satellite called MICROSCOPE, which carried two hollow concentric cylinders, one of a titanium alloy and one of a platinum alloy, around the Earth in an orbit continuously corrected so that nothing but gravity acted on the pair. Electrostatic forces held both cylinders in place. The size of those forces reports any difference in how the two of them fall. Five months of usable free-fall data came out of two and a half years in orbit. The difference in acceleration between titanium and platinum came out at one and a half parts in ten to the fifteenth, with a statistical uncertainty of two and three tenths in the same units: a null result at the level of one part in a thousand million million.

So gravity belongs to the place rather than to the thing. Which leaves two questions. Why does an influence that treats everything identically

turn out to be a shape, and what decides how much shape a given lump of matter makes?

25.1 Two walkers on the equator

Two people stand on the equator, a kilometer apart. Both face due north. Both are told to walk straight ahead and never turn, with no course corrections and no navigating.

They collide at the North Pole, about ten thousand kilometers later.

Both of them walked straight the whole way. The gap between them closed from a kilometer to nothing, at an accelerating rate over the last stretch, and no force acted on either walker at any point in the trip. A cord tied between them would have gone slack rather than taut. The convergence is a fact about the surface they crossed. They can establish it without ever seeing the Earth from outside, with nothing more than a tape measure between them. Send a third walker off from a point between them and the same thing happens to each of the three gaps at once. That is what makes the effect a property of the ground rather than an agreement between two people.

The path you get by going straight ahead is called a **geodesic**. On a flat floor it is a straight line, on a sphere it is a great circle, which is why the flight from London to Tokyo goes over the Arctic and why that route looks bent on a wall map. Two geodesics that set off parallel and end up crossing is what curvature means. The walkers measure it from the inside.

Free fall is walking straight ahead. Two ball bearings released side by side above the Earth converge, and the amount they converge by is a fact about where they are rather than about what they are made of, which is what MICROSCOPE measured to one part in a thousand million million. Nothing in the account calls for a sheet with a weight on it, and nothing bends into a further dimension for the bending to happen in.

Release a small ball of particles, all at rest with respect to each other, and let them fall. The ball's volume changes. John Baez and Emory Bunn, in the *American Journal of Physics* in 2005, wrote Einstein's equation as a sentence about that ball: the rate at which it begins to shrink is proportional to its volume times the energy density at its center, plus the pressure in each of the three directions at that point.

25.2 The equation, read rather than solved

The whole of that, in the notation the textbooks use, takes one line.

$$G_{ab} + \Lambda\, g_{ab} = 8\pi G\, T_{ab}$$

That says the shape of the world at a point is fixed by what is sitting at that point.

The letters a and b each stand for one of the four directions, one of time and three of space. Four directions taken two at a time gives sixteen pairs, and swapping the two directions of a pair changes nothing, leaving ten of the sixteen independent, so that single line is ten equations. The object g_{ab} is the **metric**: ten numbers at every point of the world that turn a pair of directions into their contribution to chapter seventeen's interval, which makes the metric the ruler, the thing that says how much time and how much distance separate two nearby events. The object G_{ab} is a particular combination of the rates at which those ten numbers change from point to point, assembled so that its own bookkeeping closes automatically, with nothing leaking in or out. On the right, T_{ab} is the stress: energy density, momentum flow and pressure, everything that is there. And $8\pi G$ carries Newton's gravitational constant, which fixes how much bending a given amount of energy buys.

The one term that spoils the arrangement is Λ, the cosmological constant, which multiplies the metric itself and has nothing on the other side of the equals sign to account for it.

Einstein reached the equation without that term in November 1915, by requiring that it reduce to the old law of gravity for weak fields, that it not depend on the coordinates anybody chose, and that both sides conserve what has to be conserved. Those requirements narrow the possibilities to one. They do not say why geometry should respond to energy in the first place.

25.3 Four pages in 1995

Ted Jacobson's paper occupies pages 1260 to 1263 of volume 75 of *Physical Review Letters*, titled "Thermodynamics of Spacetime: The Einstein Equation of State".

He needed three things. An observer who accelerates steadily has a horizon behind it, a boundary past which signals never arrive. A horizon carries an entropy equal to a quarter of its area. And an accelerating observer finds empty space warm, at a temperature set by its acceleration. Then Clausius's relation, which is the first thing anybody learns about heat: the heat flowing into a system, divided by the temperature it arrives at, equals the entropy that heat buys.

Jacobson's move was to demand that this hold on every one of those horizons. A black hole has one. So does the air above your kitchen table, and so does every other point of the world, one horizon for every direction an observer passing through that point might accelerate in.

Matter crossing such a horizon carries energy, so it delivers heat. The horizon's entropy is a quarter of its area, so the entropy change is an area change. The temperature is the one the accelerating observer reads. Insist that heat over temperature equals the area change at every point and in

every direction at once, and the geometry has no freedom left: it has to distort in exactly the way that keeps the accounts straight everywhere, and that distortion is Einstein's equation, coupling constant included.

Jacobson had to bring the area law and the temperature into the argument. OPH offers candidates for both. Chapter twenty-four counts the capacity of patches on a screen. Identifying that capacity with physical boundary entropy, with the area normalization gravity requires, is a further step. Chapter twenty-one relates modular flow to an accelerating observer under its geometric-flow and temperature-normalization hypotheses. A finite state has a modular flow; establishing that this flow is the physical boost is part of those hypotheses.

The finite repair constructions establish a useful part of the argument. Under the declared schedules and conserved constraints, they converge to states of maximum entropy, whose first-order entropy change vanishes along variations preserving those constraints. Gravity needs a different, physical stationarity statement: the entropy of a small region's contents plus its boundary area divided by four times Newton's constant must be stationary under the admitted state and shape variations at fixed physical volume. A fixed occupation total or a fixed list of patches does not establish that condition. With the area, modular-energy and continuum identifications supplied on the same construction, the following calculation shows what this additional condition requires.

25.4 Where the eight and the pi come from

Take a small ball around a point in the assumed smooth geometry, in the rest frame of one observer, and apply that fixed-volume stationarity condition in the shrinking-ball limit.

The contents side first. The required ball-modular identification weights the physical energy density by a factor that falls from a maximum at the center to zero at the edge: the difference between the squared radius of the ball and the squared distance from the center, divided by twice the radius. This geometric weight is an input to the small-ball theorem, rather than a consequence of having a finite Gibbs state. Integrating it over the ball gives four pi times the fourth power of the radius, divided by fifteen. The integral takes one line, and the fifteen comes out of it.

One factor remains: the geometric modular normalization of two pi. Under the boost and Unruh-temperature identification used in chapter twenty-one, an accelerated observer reads its acceleration divided by two pi, in units where the speed of light, Planck's constant and Boltzmann's constant are all one. The corresponding entropy-energy relation supplies the extra two pi, taking the matter side to eight pi squared times the fourth power of the radius, divided by fifteen. This normalization must belong to the same physical clock and stress as the ball calculation.

The geometry side. Alfred Gray and Lieven Vanhecke, in *Acta Mathematica* in 1979, worked out how the size of a small geodesic sphere depends on the curvature around it. The piece of that needed here says that if you hold the volume of a small ball fixed and raise the curvature, the area of its boundary falls by four pi times the fourth power of the radius, divided by fifteen, times the change in exactly the curvature combination that stands on the left of Einstein's equation. Curving the surroundings costs boundary area.

Set the sum of the two to zero, remembering that the area enters the total entropy divided by four times Newton's constant. At leading order every appearance of the radius cancels. Both fifteens cancel. What is left is eight pi squared on one side against pi over Newton's constant on the other, and their ratio is eight pi times Newton's constant, the coupling in the field equation. Under these entropy and geometric normalizations, the factor eight pi follows without a fit. The calculation does not itself select Newton's constant or establish the physical inputs that gave the two coefficients.

That gives a relation in one observer's rest frame. Suppose the same geometry and stress tensors obey it for observers passing through the same event at every velocity in an open range, with the transformations and remainder bounds required by the theorem. Written out in terms of the velocity, each relation is a polynomial of degree two. A polynomial that vanishes throughout an open range has every coefficient equal to zero. The tensor upgrade recovers all ten components once those common physical inputs are established.

25.5 A geometry that can respond

A wave description on a curved space does not by itself make gravity dynamical. The conditional field construction supplies a useful test. Its static product geometry has no radial null curvature response. The scalar field it describes admits a radial wave carrying positive null stress. Keep the reference and other sources fixed, and the same geometry cannot obey Einstein's balance for both the reference and that excitation.

The test identifies a required response. If the area of a spherical slice is four pi times a radius squared, energy passing along an affinely measured light ray must change the second derivative of that radius by minus four pi times Newton's constant, the radius and the null stress. This is the focusing part of Einstein's equation. Deriving that response from the same repair history is a physical obligation. The entropy and tensor theorems state what follows when their premises are met; the finite field description does not meet them simply by possessing a Lorentzian metric.

25.6 A gas law without molecules

Jacobson ended his abstract with a sentence that has been argued over for thirty years: it may be no more appropriate to canonically quantize the Einstein equation than it would be to quantize the wave equation for sound in air.

Pressure times volume equals a constant times temperature. Émile Clapeyron put it in that form in 1834. It held up in every laboratory in Europe through the rest of the century without containing a single molecule. In 1908 Jean Perrin put grains of gamboge resin in water under a microscope, counted how the population thinned out with height, and got Avogadro's number out of the count, which finished an argument about whether matter came in pieces at all. Wilhelm Ostwald, who had spent a career arguing that atoms were a bookkeeping convenience, gave that position up in the preface to the fourth edition of his own textbook in 1909, citing Perrin by name. The gas law did not change a symbol, because it was never a claim about molecules. It is a claim about a gas that has settled down, and stays true whatever the gas turns out to be made of.

The conditional entropy argument puts Einstein's equation on that shelf. A network satisfying its physical entropy, stress and continuum premises must report that equation. Settling alone does not supply the premises. In this interpretation, quantizing the metric in the hope of finding the atom of space resembles quantizing the pressure of a gas in the hope of finding the molecule. The pressure is real enough to lift a piston, and it belongs to a settled state rather than to a constituent.

Once the smooth metric and physical frame identification are in place, the 1907 thought has its usual precise meaning. A freely falling frame removes the connection at a point, while tidal curvature remains. The finite frame constructions supply algebraic ingredients for this identification; they do not establish physical free fall by themselves. For pressureless matter with nonzero density, covariant stress conservation then makes the flow geodesic. The walkers return as a consequence within the physical geometric description.

25.7 Nine light rays

Everything in the argument crossed a horizon, and horizons are made of light rays.

The horizon probes used in this argument evaluate stress along a light-ray direction. The stress takes two directions and returns a number. Feed it the same null direction twice and you get the number used in the null-balance theorem.

This restriction specifies the data used by the theorem. Light cones define the causal boundary of the physical description, but not every local

measurement is a null contraction of stress. A quantity invisible to these horizon probes is undetermined by these probes; a different measurement or an additional physical condition can carry more information.

The conditional scalar-field construction supplies a compatible cone test. Disturb its field in one slab and read a detector in a separate slab. Under the construction's refinement and detector assumptions, the response before that field's light cone arrives shrinks to zero.

Ten numbers describe the stress at a point. Light rays report one number each. How much of the stress do the rays determine?

A symmetric object that reads zero along every light ray has to be a multiple of the metric, and nothing else. So the rays pin down nine of the ten numbers. What they leave undetermined is a multiple of the metric.

For readings from one symmetric tensor, nine directions do it. Take the six rays that travel along one axis at a time, forward and backward along each of the three. Then take three more that share their motion equally among all three axes, each space component equal in size to one over the square root of three, with the signs plus plus plus, plus plus minus, and plus minus plus. The six axis rays never pick up the cross terms, the entries pairing one space direction with a different one, because a ray running along a single axis has no second direction to pair it with. The three tilted rays are there for those, and their three sign patterns read the three cross terms off one another. Nine light rays, nine numbers, one system of linear equations to invert. Its determinant is exactly eight thousand one hundred and ninety-two over twenty-seven, or two to the thirteenth over three cubed. A determinant that misses zero is a system that inverts, and this particular inversion is stable: small errors in the nine readings stay small in the nine numbers recovered from them.

Any nine numbers fit that system, so the fit alone does not show that the readings come from a tensor: a tenth ray, or a hundredth, has to land where the recovered tensor says it will. Given the geometric and stress tensors, balance along the nine rays fixes the field equation up to a multiple of the metric. Conservation makes that multiple constant on each connected region. That term is the tenth number.

25.8 The vacuum and the tenth number

A Lorentz-invariant vacuum contribution to stress is a number times the metric. This is the vacuum-reference condition used here; a general quantum state on a curved background need not have stress of that form.

Contract it with a light ray. The metric evaluated on a light-ray direction twice is the interval of a light ray with itself. That is zero: chapter seventeen defined a light ray as a separation whose interval is exactly zero. So the vacuum's contribution to what a light ray reports is a number times zero.

Exactly zero, for any vacuum energy density whatever, however enormous, and zero by the definition of the probe rather than by a cancellation between large terms.

The single direction the light rays cannot see is exactly the direction the vacuum's energy points in. That is why the cosmological constant survives in the field equation as a term nobody derived: the small-ball calculation that produced everything else on the left-hand side is built out of horizons, horizons are made of rays, and rays are blind to it.

A naive Planck-scale cutoff on quantum zero-point energies gives a vacuum energy density around a hundred and twenty orders of magnitude larger than the one inferred from the sky, whose cosmological constant is about 1.1 times ten to the minus fifty-second per square meter. The null-balance theorem explains a precise limitation: its data cannot fix the metric-proportional term. That limitation does not by itself solve the vacuum-energy problem or prohibit other local measurements and calculations.

A specified horizon-capacity condition offers a way to fix it. On that branch, the smallness of the constant is the largeness of the horizon's record count. Chapter twenty-nine computes candidate counts and compares them with the sky. Selecting the physical capacity condition from the same repair history remains necessary.

25.9 Newton's constant is an area

On the physical area-law branch, boundary entropy is its area divided by four times Newton's constant, in the same natural units. Read that from right to left: Newton's constant is the area divided by four times the entropy. It supplies an area scale for counting distinguishable states. With the measured gravitational conversion, one square meter of boundary has entropy of order ten to the sixty-ninth.

Chapter thirteen counted readings without assigning a width in meters. A specified screen-area dictionary can join the two. Give a cell an area equal to a whole number of one small area unit, and entropy equal to a quarter of that same number. Divide its area by four times its entropy and the number cancels, leaving the area unit. The cancellation is exact. Identifying this dictionary with the physical entropy and area is a separate premise.

Under that identification, Newton's constant is the unit of area. Its square root supplies the corresponding gravitational length.

The measured Planck area is 2.61228 times ten to the minus seventy square meters, with square root 1.6163 times ten to the minus thirty-five meters. Multiply that area by the cube of the speed of light and divide by Planck's constant over two pi, and the measured gravitational constant comes back: 6.67430 times ten to the minus eleven cubic meters per kilo-

gram per second squared. These are conversions of the laboratory value. They do not independently measure a grain in the proposed screen.

The figure carries meters, kilograms and seconds, which people chose. The proposed screen dictionary concerns relations that survive a change of units; its physical attachment still needs an independent area and entropy calibration. The laboratory figure itself is known poorly. Newton's constant is the least precisely known of the fundamental constants by a wide margin, measured to five figures where the electron's magnetic moment is known to twelve, because gravity is feeble and everything in the laboratory has mass. MICROSCOPE measured gravity's indifference to what falls at one part in a thousand million million, and gravity's strength is known to twenty-two parts per million, ten orders of magnitude worse, because a difference between two cylinders nulls out everything the two have in common and a magnitude has to be assembled out of a kilogram, a meter and a second.

25.10 Order, number and curvature

Chapter seventeen constructed a flat spacetime limit under a declared population, read law and counting measure: an order from who read whom, a volume from how many. Extending that route to the curved geometry needed here requires the corresponding order and volume identifications on one common refinement.

Start with the order. Within the distinguishing Lorentzian manifolds covered by the theorems of Hawking, King and McCarthy and of Malament, causal order determines the conformal geometry. It fixes the light cones but leaves a local scale: multiplying the metric by a positive function leaves those cones unchanged. A calibrated volume measure fixes that scale. If record order and counts converge to the required causal order and physical volume on the same smooth limit, they therefore determine its metric. An arbitrary finite log does not meet those conditions merely by having an order and a count; reading curvature also needs the appropriate smooth control.

Spatially flat homogeneous expansion illustrates the division. In conformal coordinates its metric is the flat metric multiplied by the square of a growing scale factor. The cones remain the same, while the four-volume density grows as the fourth power of that factor. If calibrated record counts sample equal small conformal-coordinate four-volumes, their density ratio recovers the fourth power of the scale-factor ratio. For comoving emission and reception, the fourth root is then one plus the redshift. The statement uses that cosmological geometry and count-volume identification; it does not follow from an arbitrary pair of record totals.

Curvature in general is more than one growing number. With count-volume calibration and the small-diamond expansion in place, counts can measure its effect on volume. Boundary area has a separate finite construc-

tion: in the declared three-dimensional population and all-neighbor read law, crossing reads approach a surface count with density equal to a quarter of pi times the square of the population density times the fourth power of the reach. This supplies a geometric area proxy for that source. Joining it to physical generalized entropy and to the same curved field history remains a further requirement.

The constructions therefore give concrete candidates for parts of the derivation: order, volume and boundary counts, finite entropy identities, and modular clocks. They do not yet establish that one repair history supplies all the physical identifications together, or that its generalized entropy is stationary at fixed physical volume. If those inputs and the required continuum bounds hold, the small-ball argument yields Einstein's equation. Establishing the responsive geometry is part of completing that physical construction.

Under those premises, the geometric equations close. Ten numbers at every point, a rule fixing how they answer to stress, the factor eight pi from the ball calculation, and a gravitational area scale requiring physical identification. In that description free fall is going straight ahead, and the two walkers on the equator were doing physics the whole way to the pole.

The geometric description is still an effective one. Its field equation relates curvature to energy and momentum; it does not select an internal gauge algebra or matter sector. The port-current constraints and the physical matter lift pose those separate questions in the next chapters.

Chapter 26

Why Exactly These Forces?

Move two electric charges twice as far apart and the push between them drops to a quarter, and the two in that inverse square is pinned to the fifteenth decimal place. Williams, Faller and Hill, writing in *Physical Review Letters* in 1971, wrote the exponent as two plus an unknown small amount and then measured the amount. It came to two point seven parts in ten million billion, give or take three point one.

The two itself came off a twisted thread, hung by an army engineer who wanted a better compass. Charles-Augustin de Coulomb had come home a captain in June 1772, after eight years building Fort Bourbon on Martinique, with a good deal of what is taught as soil mechanics and a succession of tropical illnesses that damaged his health for the rest of his life. The Academy of Sciences in Paris wanted a better mariner's compass, and Coulomb's entry, which took a share of the grand prize in 1777, hung the needle from a fine thread instead of resting it on a pivot, because a pivot has friction in it and a thread has almost none. That left him needing to know exactly how hard a twisted thread pulls back, so he measured it, at length, and published the numbers in 1784: the pull rises with the fourth power of the wire's diameter, falls off with its length, and holds exact proportion to the angle of twist, which makes a long thin one as weak a spring as an experimenter could want and an exact one.

In 1785 he hung a light horizontal arm from one, mounted a small ball on its end, charged it, and brought a second charged ball up beside it. The two pushed apart, the arm swung, and the twist in the thread read off how hard they pushed.

Three readings from that first memoir on electricity carry the whole argument. Left alone, the thread let the balls settle thirty-six degrees apart and carried thirty-six degrees of twist. He turned its top through a hundred and twenty-six degrees, forcing them closer, and they came to rest eighteen degrees apart, held by a hundred and twenty-six plus eighteen, a hundred and forty-four degrees of twist. Half the separation, four times the push. He wound the top through five hundred and sixty-seven degrees, better than a full revolution, and the balls held out at eight and a half degrees, with five hundred and seventy-five and a half degrees of twist in

the thread. The separation had come down by a factor of a little over four, and sixteen times thirty-six is five hundred and seventy-six.

Which is a fact about distance: every version of the law written down since 1785 says only how far apart two things are. What does a force law look like on a carrier that has no distances in it, and how many forces will that carrier hold? The twelve observers of chapters six through fourteen have a wiring diagram, thirty pairings saying which port is joined to which, and not one number anywhere saying how far apart anything is.

Two centuries of better instruments have left the exponent where Coulomb put it, at two, to fifteen decimal places. The carrier has no ruler in it.

26.1 A value at every place

Between 1698 and 1700 Edmond Halley took a small Royal Navy vessel called the Paramore twice across the Atlantic, on two voyages fitted out for the single purpose of measuring the Earth's magnetism, and recorded the angle between where his compass pointed and where north actually was. The angle changes as you sail. In 1701 he published the results as a chart, and rather than print a table of positions and readings he drew curves through the places where the deviation came out the same. Sailors called them Halleyan lines for the next hundred years.

What that chart holds is a number attached to every place on it. A captain anywhere in the Atlantic could read his local deviation off the paper. The reading changes from place to place, the rule for finding it does not, and a ship crossing the ocean experiences the chart as a sequence of small differences between one position and the next.

That is a **field**, one value per place, and the word covers nothing else. The temperature in a room is a field. So is the wind over Halley's Atlantic, which hangs a speed and a heading at each place rather than a single number, and so is the deviation he charted, which needs only one.

On the twelve-port carrier the places are the ports. Give every port a number and each seam reads the gap between its two ends, thirty readings nobody had to supply separately. Chapter twelve had the same twelve numbers standing for something else, the private labeling each observer keeps at its own port. Twelve values and thirty differences: the same kind of structure Halley put on his chart.

26.2 Twelve numbers that add to nothing

Chapter twelve left a law sitting at the ports: what a port pushes out along its five seams is the charge it holds. That is Gauss's law with the geometry taken out of it. By itself that law picks no field, since a current going round in a circle leaves every port's books balanced, so one list of charges comes

back with a whole family. One member of the family has the least energy, the one with no circulation in it.

Energy on this carrier is a sum over the thirty seams. Each seam contributes the square of the number sitting on it. Squaring is what makes circulation expensive: a current running around a loop delivers nothing to any port and puts a value on every seam it crosses, and every one of those values gets squared and added in. So the arrangement of least energy is the one that has stopped moving anything it does not have to move. On a wiring diagram with nineteen independent loops there is exactly one such arrangement for any list of charges.

Adding the same amount at all twelve ports changes no difference anywhere, so the twelve numbers are pinned only up to a common shift, exactly as chapter five's meridians were pinned only up to a choice of zero. The shift is fixed here by requiring the twelve values to add to nothing.

And the surface is closed. Twelve ports wired into a sphere have no edge, no outside and nowhere for a net outflow to go, so the twelve charges have to sum to zero or the law has no solution at all. A single unit of source at one port therefore arrives with its own balance: one twelfth taken off every port, the same subtraction everywhere, including at the port the source is sitting on. That subtraction is the only balance that treats the twelve ports alike.

Put one unit of source at one port, then, and ask what the other eleven read.

26.3 Four fractions

Walk the wiring outward from the source and count hops. The eleven others sort into three groups: a ring of five, a second ring of five past it, and the port directly opposite, three hops out and the only one chapter fourteen leaves at that distance.

Four distances, therefore, and four numbers. Here they are, exact, in the order the walk produced them:

$$\frac{7}{36}, \qquad \frac{1}{90}, \qquad -\frac{7}{180}, \qquad -\frac{1}{18}$$

The first is the potential at the source itself, the second is what every port in the first ring reads, the third is what every port in the second ring reads, and the fourth is the reading at the far port.

Put them over a common denominator of a hundred and eighty and they read thirty-five, two, minus seven and minus ten. Add all twelve of them up, taking five at the second value and five at the third: thirty-five and five twos on one side, five sevens and a ten on the other, forty-five against forty-five. The twelve values add to nothing, which the closed surface required of them.

They fall. Each ring reads lower than the ring inside it. The ordering was not put in anywhere. The hop counts came out of walking the wiring. The four values came out of the charge law and the least-energy condition. One of those was a walk and the other was a piece of algebra. They agree about which port lies further out.

Subtract each value from the one inside it, in one-hundred-and-eightieths: thirty-five take two is thirty-three, two take minus seven is nine, and minus seven take minus ten is three. Over sixty those are eleven, three and one. The fall is steep near the source and shallow far from it, the shape of the thing Coulomb's thread was weighing.

The potential crosses zero somewhere between the first ring and the second. A potential around an isolated charge in open space never does that, and this one has no way to avoid it: twelve numbers that must sum to nothing, with the largest of them sitting at the source, have to go negative somewhere further out.

Compare what the same law costs in the ordinary setting. To write down the potential of a point charge in a laboratory you need the distance between two points, and a distance needs a coordinate system to measure in, and a coordinate system needs a space to be laid over. Three assumptions come before the first symbol goes on the paper, and the carrier supplies none of them.

The input to that calculation was a list of thirty pairs saying which port touches which. Nothing else went in. No coordinate, distance, direction or angle went in, and the hop counts that organize the answer were themselves read off the same list of thirty pairs. Coulomb's shape came out of a table of who touches whom, in exact fractions with denominators small enough to check by hand.

One force, worked to the last digit, out of a wiring diagram with thirty entries in it. The same thirty entries have to say how many forces there can be.

26.4 Following the response into matter

The response matrices can be followed beyond their commutators. A specified spin lift converts their weak rotation block into a doublet, and a specified redistribution of the central trace fixes relative integer charges. Applying the exterior construction then yields the internal matter list in the next chapter. Charge and Casimir measurements recover the list even after the basis has been mixed, so the names of the fields are not an input to that decoder.

The lift is a physical choice requiring its own identification. A full turn acts trivially on the native vector and as minus the identity on the doublet. These are distinct representations, despite sharing an infinitesimal rotation algebra. Establishing one does not silently identify it with the other.

26.5 Dials

Chapter five's group was eight moves of a paper square, and eight is a menu. You pick one.

Take instead a dial on the front of an amplifier. Turn it through any angle you like, including a millionth of a degree, and then turn it through another. The result is a setting of the dial, so the moves close. Every turn is undone by turning back, and leaving the dial alone is on the list. Closed, undoable, do-nothing included: a group, by chapter five's three conditions, with the difference that the moves come in a continuous supply rather than in a list of eight.

One thing about a dial that a menu of eight moves does not have: you can turn it by almost nothing. Set it a hair off the do-nothing position and you have a nudge, and nudges behave far better than the moves they generate. Two nudges in different directions can be added. A nudge can be doubled or halved. Every large turn is a small nudge repeated, so the whole continuous supply of moves is determined by the nudges available at the do-nothing setting, which is a much simpler object: a flat space of directions, with as many independent directions as the dials it came from.

The useful number about a collection of dials is how many independent dials it has. That number is its **dimension**. The amplifier has one. So does a sheet of paper spun about a pin stuck through it: every rotation of a plane is that one dial at some setting. A rigid object turned in three dimensions has three, which is why aircraft have exactly three words for it: pitch, roll and yaw.

Counting dimensions is a way of settling questions, and chapter six ran one without calling it that. The independent loops of a wiring diagram are a collection with a dimension of its own, and edges less dots plus one reads it off: thirty less twelve plus one, nineteen. Nineteen was then the number of conditions that had to hold at once for the twelve observers to agree. Two of them refused, both facts settled before anybody drew a field on the wiring.

Two more collections, the ones physicists reach for. Take an object with two slots in it, each holding a complex number, and ask for every reversible change that leaves the total size alone. Then ask the same of an object with three slots. Both counts are short.

Each slot has a dial of its own, which turns that slot and leaves the others where they were. Each pair of slots has two more, which lean the two members of the pair against each other. Two slots: two single dials, one pair, two pair dials, four. Three slots: three single dials, three pairs, six pair dials, nine. Singles and pairs together come to the number of slots multiplied by itself, every time.

One dial in each collection turns every slot together by the same amount, a change of overall convention that says nothing about how the slots stand

relative to each other. Divide it out. Four less one is three, nine less one is eight, and the count that survives needs nothing but how many slots there are:

$$n^2 - 1$$

Here n is the number of slots and n squared is that number multiplied by itself. Four slots give fifteen. Five give twenty-four. Between eight and fifteen this family gives nothing at all.

26.6 A book on a table

Take a hardback off the shelf and lay it flat, cover upward, spine to your left.

Turn it a quarter turn clockwise on the tabletop. Then flip it away from you, end over end. Note where the cover faces and where the spine points.

Put it back the way it started and do the same two moves in the other order: flip it away from you end over end first, then turn it a quarter turn clockwise on the tabletop.

The cover is face down both times. The spine is nearest you in one case and furthest from you in the other. The two results are a half turn apart.

A group in which every pair of moves gives the same result in either order is **abelian**, after Niels Henrik Abel, and one containing even a single pair that does not is non-abelian. The amplifier dial is abelian: thirty degrees then fifty is fifty degrees then thirty. Turns of a solid object in three dimensions are non-abelian. The book on the table is the demonstration. The three-dial and eight-dial collections are non-abelian for the same reason, because three of the dials in each of them are turns of exactly that kind. The single dial is the only abelian collection here.

26.7 Twelve dials on a closed box

Back to the carrier, a closed box with twelve ports.

Each port holds one reading. A response is whatever the box does when the readings are disturbed: twelve numbers go in, twelve come back, and the box is the thing in between. Reversible means the twelve that came back are enough to recover the twelve that went in, so nothing was thrown away in the middle. Ask for every response of that kind the ports admit. Those responses compose, they undo, and they come in a continuous supply, which makes them dials.

The box’s complete continuous reversible response is a compact group with exactly twelve directions of small change, one for each port reading. The rotations act on the directions and on the readings in the same way,

and every direction moves some reading. That ties the response faithfully to the carrier.

So there are twelve dials. The question is how their infinitesimal responses split into independent algebraic pieces.

26.8 The swap and the center

One move of the twelve ports came out of chapter fourteen indifferent to order: the swap that trades every port for its opposite. Do it before any of the sixty rotations or after, and the ports finish in the same arrangement either way. Twelve ports can be shuffled 479,001,600 ways, and of the 479,001,599 that move something, the swap is the only one that gives the same result in either order with all sixty rotations.

Collect, in any group, the moves that commute with every move in the group. That collection is the **center** of the group. Chapter nineteen used the same word for the questions inside an algebra that commute with every other question, which is where records live, and the property is identical, commuting with everything in sight; the difference is what is being commuted, an algebra of questions there and a group of moves here.

Groups differ enormously in how much of themselves lives in the center. Settings of the single amplifier dial all commute with each other, so that group is its own center. Turns of a solid object in three dimensions go the other way: the book on the table showed that a quarter turn and a flip disagree about order, and if you work through the possibilities the only turn that commutes with every other one is the turn that does nothing. Its center holds only the move that does nothing, zero dials wide.

Go back to the clause chapter four spent a section on. Two parties who cannot look a rate up anywhere outside the room have to manufacture one out of what both of them hold. The same clause binds the twelve ports harder, since nothing anywhere holds a preferred labeling of them. Any change in how the readings are labeled has to be a change the box itself can perform.

Take that concretely. The twelve ports carry no names of their own. Numbering them is something an observer does on a piece of paper, a second observer numbers them differently, and the wiring is indifferent to both. What the box can do is turn its own dials. So the relabelings with any standing are the ones the dials produce, and applying one is a three-step operation built out of moves the box has: turn the dials to the new labeling, do whatever was going to be done, turn them back.

Feed a central move into that. It commutes with the relabeling, so the relabeling and its undoing slide together and cancel. The move comes back out untouched. Every change of description leaves the center exactly where it was.

The sixty rotations of the wiring are changes of description. By that same clause, each of them is something the box itself performs. So the center has to sit inside the part of the twelve readings that all sixty rotations leave fixed.

That part is easy to find, because chapter fourteen showed the sixty rotations carry every port to every other port. A set of twelve readings that survives all sixty unchanged has to read the same at every port. One number, repeated twelve times, and one number is one dial.

The center of the twelve is at most one dial wide.

26.9 One, three, eight

A continuous group of this kind comes apart into its center and a remainder with no center in it at all. The center is one dial or none. The remainder is eleven dials or twelve.

A group that cannot be broken into independent pieces is called **simple**, and the sizes a simple group is allowed to have are a fixed list rather than a matter of taste: three, eight, ten, fourteen, fifteen, twenty-one, and on upward. Below twelve, the list holds exactly three entries. There is no simple group of size four, or five, or six, or seven, or nine, or eleven, or twelve.

So the count of response directions is a sum to be hit exactly. The parts come off that list, the total to be reached is twelve less whatever sits in the center, and nothing may be left over at the end. Two targets, then, twelve and eleven. Take a pencil to both, using threes, eights and tens, as many of each as you like.

Suppose the center is nothing at all. Then twelve dials have to be assembled out of pieces of size three, eight and ten. Ten leaves two to find, and the list starts at three. Eight leaves four, and four is not on the list. Threes run three, six, nine, twelve, and the fourth one lands exactly. Three plus three plus three plus three is the only way to do it: four copies of the smallest simple group there is.

Check whether the wiring has room for four copies. Pairing every port with the port opposite it gives the six axes of chapter fourteen. Each axis carries two readings, splitting into the part that is the same at both ends and the part that is equal and opposite at the two ends: six even readings and six odd ones. Among the six even readings sits the uniform one, the same number everywhere, and taking it out leaves five. The six odd readings fall into two separate blocks of three. So the twelve come apart as one, three, three and five, and no rotation of the wiring mixes any block with any other.

Four blocks of three are needed. Two exist. The rotations cannot get around that by shuffling four copies among themselves either, because a shuffle of four things would require the sixty rotations to fall into four

families of fifteen, and no fifteen of the sixty close up into a group of their own.

So the center is exactly one dial, the remainder is eleven, and eleven has to be built out of three, eight and ten. Ten leaves one, and one is smaller than anything on offer. Two eights overshoot to sixteen. Threes run three, six, nine and then jump to twelve, stepping over eleven without touching it. Eight leaves three, and three is the first entry on the list. Three plus eight, and no other way at all.

$$1 + 3 + 8 = 12$$

One abelian dial, a three-dial piece and an eight-dial piece, with nothing left over and nothing borrowed.

A simple Lie algebra has no continuous central dial, so it cannot be the complete twelve-dimensional response. Howard Georgi and Sheldon Glashow proposed in 1974 that the three forces sit inside one larger simple symmetry. The carrier has a central dial, and a simple symmetry has nowhere to put one.

The allocation is fixed too, and not by size alone. The uniform reading, the one thing all sixty rotations leave alone, is the central dial. One of the two odd blocks of three carries the three-dial piece. The remaining block of three and the block of five carry the eight-dial piece between them, and they carry it because the eight dials of the three-slot group turn among themselves the same way those eight readings do under the rotations, which is a stronger match than eight equals three plus five.

These three pieces are hypercharge, the weak interaction and the strong interaction of the Standard Model: the same three algebras, in the same sizes. Hypercharge combines with a weak generator after symmetry breaking to give electromagnetism.

The argument uses twelve ports and thirty seams, the sixty rotations, the splitting of the carrier into blocks of one, three, three and five, and a complete reversible response tied faithfully to those twelve readings, with the rotations acting inside it. The classification of compact simple Lie algebras supplies the allowed sizes.

No measured mass, coupling strength or decay rate enters this classification. The algebra comes out one, three and eight.

Forces need something to act on. The next question is which charged matter can sit on the carrier while respecting that response.

The catalogue of things that do exist makes that look difficult. The electron carries one unit of electric charge. The down quark carries one third of a unit and the up quark carries two thirds, and every quark carries one of three color charges besides, and the whole arrangement is repeated three times over at rising masses. Thirds, in a world assembled out of a table of whole numbers. And three of everything.

Chapter 27

Why This Exact List of Matter, and Why Three Times Over?

Two physicists at the Massachusetts Institute of Technology filled a metal cavity with sulfur hexafluoride in 1973 and waited for the gas to sing.

Sulfur hexafluoride is heavy and chemically dead, which is why electrical engineers pump it into switchgear to keep the contacts from arcing. H. Frederick Dylla and John G. King wanted it for a different property. They wanted a gas whose molecules they could count.

Across the cavity they ran an alternating electric field. If every molecule carried even the smallest excess of charge, the field would shove the whole body of gas one way and then the other in time with itself, and a gas shoved back and forth a few hundred times a second is a sound. They mounted a calibrated microphone in the cavity wall, switched on the field, and listened.

The cavity stayed quiet.

The silence set an upper limit. The net charge on one molecule of sulfur hexafluoride came in below two parts in ten billion billion of the charge on a single electron. That molecule carries 146 units of atomic mass, so per unit of mass the difference between the charge on a proton and the charge on an electron sits below one part in a thousand billion billion. Hydrogen, which is one proton and one electron and nothing else, is electrically neutral to better than one part in a billion billion.

Look at what has to be true for that. A proton is two up quarks and a down quark. The up quark carries two thirds of a unit of electric charge and the down quark carries minus one third. Two thirds plus two thirds minus one third is one. An electron contains no quarks. The strong force that binds the proton's insides passes through an electron without noticing it. The electron's charge is minus one, exactly three times the down quark's. The exactness holds past every decimal place anyone has reached.

Nothing in the equations requires it. Write the same physics down with the down quark at minus 0.333333 and the electron at minus 1.000000 and every equation goes through unchanged. What changes is the sky: a mismatch of one part in a billion billion would leave two hydrogen atoms

pushing each other apart about as hard as their gravity pulls them together, and a cloud of the stuff would blow itself apart instead of collapsing into a star.

So the charges are locked to each other, across species that have nothing else in common, to the precision a microphone in Cambridge, Massachusetts was able to test. Two questions follow. Why does matter come with exactly these charges and no others, and why does the whole list arrive three times over, in copies that differ in nothing except how heavy they are?

27.1 Three colors, two weak slots

Chapter twenty-six counted the forces and came back with twelve directions, split as one plus three plus eight. The three belongs to the weak force and the eight to the strong force. Those two numbers say how many things each force shuffles.

The single abelian factor carries **hypercharge**. What that dial reads on a species, combined with its position in a weak pair, gives electric charge. Electromagnetism emerges from that combination.

A force that shuffles some number of slots has a fixed stock of independent moves, the square of the number of slots minus one, which chapter twenty-six counted. Set that against the blocks: the three-dimensional block matches a force shuffling two slots, the eight-dimensional block matches a force shuffling three, and the next size up, fifteen, overshoots while three is spoken for.

Two color slots would give a different matter pattern. Their color and anticolor representations are equivalent, and two quarks can form a color-neutral combination. Three quarks cannot: reversing every color vector changes the sign of their product, so it cannot be invariant. Three color slots instead admit both quark–antiquark pairs and an antisymmetric three-quark combination. Those are the color structures of mesons and baryons. Three slots is the smallest number for which a color and an anticolor differ.

Three color slots, then, and two weak slots. Five slots, and matter is what you can build out of them.

Each slot carries a hypercharge tag. Two conditions fix the tags, both of them conditions on the finished list rather than choices anybody makes. The three color tags and the two weak tags, added with their multiplicities, have to come to zero. And the charges have to sit on the coarsest grid that carries all of them as whole multiples of one step. Those two requirements leave one answer and its mirror image: each color slot tagged minus one third, each weak slot tagged plus one half. Check the balance. Three times minus one third is minus one, two times one half is one, and they cancel.

27.2 Choose two, choose four

Take the five slots and build things out of them by picking a handful at a time.

One rule governs the picking, the rule that makes matter behave like matter. Write a pick as an ordered list of slot names, and demand that swapping any two names flips the sign of the whole object. That is **anti-symmetry**. It has an immediate consequence: pick the same slot twice and swapping the two copies changes nothing while also flipping the sign, so the object equals minus itself, so the object is zero. Every slot appears at most once. Order stops mattering, since any reordering costs a sign and nothing else. What survives is the bare question of which slots are in and which are out.

The number of ways to choose two things from five is the number of pairs you can make: five choices for the first, four for the second, and a division by two because each pair got counted in both orders. Ten. The number of ways to choose four from five is easier, because choosing four is the same as leaving one out. There are five slots to leave out. Five. Numbers of this kind are called **binomial coefficients**. For cases this small you can write the choices down and count them.

Ten and five. Fifteen.

Fifteen is the number of states in one generation of matter. The charges come out with it. The hypercharge of a combination is the sum of the tags of the slots in it. The strong and weak forces act on whichever of their own slots are present.

Slots picked	Ways	Hypercharge	What it is
two color	3	minus two thirds	anti-up quark, one per color
one color, one weak	6	plus one sixth	up and down quarks, three colors each
two weak	1	plus one	positron
three color, one weak	2	minus one half	electron and its neutrino
two color, two weak	3	plus one third	anti-down quark, one per color

Every hypercharge in that column is the sum of the tags. Minus a third and minus a third is minus two thirds. Minus a third and a half is a sixth. Three color tags and one weak tag is minus one plus a half, or minus a half. The rows add to three plus six plus one plus two plus three, which is fifteen.

Electric charge is one more addition. Where a pick includes exactly one of the two weak slots, the weak force can rotate that choice into the other, so those states come in pairs. The two members of a pair sit half a step either side of the hypercharge tag. A sixth plus a half is two thirds, the up quark. A sixth minus a half is minus one third, the down quark. Minus a half plus a half is zero, the neutrino. Minus a half minus a half is minus one, the electron. The unpaired rows carry their tag unchanged: minus two thirds, plus one third, plus one.

That is the roster. Up and down quarks in three colors, the electron, its neutrino, and the antiparticles of the ones that need them, with every charge in the Standard Model's table reproduced, out of choosing two or four from five slots and adding up tags.

Two further counts fall out of the same table. The quark pair appears once per color and the lepton pair once, so a generation carries four weak pairs. A spin-zero field with hypercharge plus one half ties into the list through three channels. The quark pair couples to the anti-up through that field; the quark pair couples to the anti-down, and the lepton pair to the positron, through its conjugate with hypercharge minus one half. Three channels, the three that give the up-type quarks, the down-type quarks and the charged leptons their masses.

27.3 Reading the list

The conditional fifteen-state module has an exact measurement dictionary. Its charge and color and weak Casimir operators separate six quark-doublet states, three up conjugates, three down conjugates, two lepton-doublet states and one charged-lepton conjugate. A change of basis mixing unlike sectors leaves the decoded list unchanged. That is stronger than putting familiar names on a collection of slots.

The construction nevertheless has declared ingredients: the spin lift, the selected exterior component and the interpretation as same-chirality matter. Auxiliary exterior parity does not establish physical fermion statistics. An independent spin and fermion construction can implement the finite controls, while propagation and physical identification require their own common history.

27.4 Dirac's holes

Three of the five rows above are antiparticles. In 1928 Paul Dirac wrote an equation for the electron that respected relativity. It came with solutions of negative energy that nobody wanted. He tried for a while to identify the holes in that sea with protons, and in a paper submitted to the Royal Society on 29 May 1931 he gave that up and proposed instead a particle nobody had seen: the mass of an electron, the opposite charge.

Carl Anderson photographed one at Caltech on 2 August 1932, a track in a cloud chamber curving the wrong way through a magnetic field, and the name came from an editor at *Physical Review*, who proposed it when Anderson sent the discovery paper in: positron. The pattern turned out to be general. Every species of matter has a partner with the same mass and every charge reversed, and the partner is as ordinary as the original: hospitals make positrons to order out of fluorine-18, and the pair of flashes from each annihilation is what a scanner photographs. Writing a species as its antiparticle is therefore a change of notation rather than a change of content, which is why the roster above lists an anti-up quark where a textbook table lists a right-handed up quark, and the reason for preferring one of those descriptions comes from an experiment with cobalt.

27.5 Wu's cobalt

Hold your hands out in front of you, palms down. They are the same object twice. No amount of turning either one will make it into the other, because a mirror sits between them. A screw thread has the same property, and so does a particle: give it a direction of travel and a sense of spin, and either the spin turns the way a right-handed screw turns as it advances, or it turns the other way. The two cases are called right-handed and left-handed, a mirror swaps them, and the property is **chirality**.

A symmetry is the claim that some part of a description was never physical. Mirror symmetry would say that left and right are a labeling convention, and that nothing in an experiment distinguishes a world from its reflection. Until 1956 every physicist assumed it, because electricity and the strong force obey it exactly and nobody had looked anywhere else.

Chien-Shiung Wu was born in Liuhe, near Shanghai, in 1912, took her doctorate at Berkeley in 1940 under Ernest Lawrence, and worked on uranium separation for the Manhattan Project at Columbia, where she stayed. She was the person other physicists went to about beta decay. In 1956 Tsung-Dao Lee and Chen Ning Yang worked out that no experiment had ever tested mirror symmetry in the weak force, and they brought Wu a list of things to try. She picked the decay of cobalt-60 and cancelled a trip to Geneva and the Far East that she had planned with her husband, on the grounds that somebody else would think of it.

The experiment needs the cobalt nuclei lined up, which needs them cold. On 4 June 1956 Wu telephoned Ernest Ambler at the National Bureau of Standards in Washington, whose group could reach three thousandths of a degree above absolute zero by adiabatic demagnetization, and asked him to collaborate. Several months went into getting the apparatus to work. The first successful run began at four minutes past noon on 27 December 1956.

A cobalt-60 nucleus decays by throwing out an electron. With the nuclei aligned by a magnetic field, the question is whether the electrons prefer the spin direction or the opposite one. In a mirror the spin direction survives, because a spin is a circulation rather than an arrow, while the direction the electron flies in is reversed. So a world obeying mirror symmetry has to emit equal numbers each way. Any asymmetry at all kills the symmetry outright.

The electrons came out against the spin. Reverse the magnetic field and the count dropped, then climbed back to baseline over about six minutes as the sample warmed and the alignment was lost. Wu, Ambler, R. W. Hayward, D. D. Hoppes and R. P. Hudson published it in *Physical Review* volume 105, page 1413, in 1957. Lee and Yang took the Nobel Prize in Physics that year. Wu took the first Wolf Prize in Physics, in 1978.

What her cobalt established is that the weak force acts on left-handed particles and not on right-handed ones. Which is why the roster above is written the way it is: put every species into its left-handed form, converting the right-handed ones into their left-handed antiparticles as you go, and the list becomes a list of things the weak force either pairs up or leaves alone. The quark pair and the lepton pair are the pairs. The anti-up, the anti-down and the positron are the singles. A list of that shape differs from its own mirror image, and a list that differs from its own mirror image is called chiral.

Mirror symmetry is a symmetry the world lacks. Every other structural fact about matter says that some distinction was never there; this one says that a distinction nobody put in is physical. It is why the roster has to be rewritten all in one handedness before it can be counted at all.

27.6 One thousand and twenty-four lists

The five slots generate more than the fifteen states above. Pick none of them, pick one, pick three, pick all five: there are thirty-two picks in all, and they group into components according to how many color slots and how many weak slots each one uses. Drop the empty pick and the pick that takes everything, and ten components are left, each with its charges attached. Any subset of those ten is a possible content for matter: two to the tenth possible lists, or 1,024.

Each of the 1,024 gets tested against two demands. It has to be chiral, since the mirror is what Wu's cobalt says the world distinguishes, and it has to make four separate sums come out at zero.

Of 1,024 lists, exactly two survive.

The two survivors are the even picks and the odd picks: the objects built from an even number of slots, and the objects built from an odd number. Charge conjugation, which swaps every particle for its antiparticle, carries each of the two onto the other, so they are one list and its mirror image, and

a world taking either one gets the same physics with matter and antimatter labeled the other way round. The even one is the two-slot picks and the four-slot picks: ten plus five, and the roster above.

The search asked each list for chirality and four vanishing sums. What came back divides by the number of slots a pick uses: all even, or all odd. This is parity in the auxiliary exterior construction, a rule organizing the internal representation. The physical minus sign under exchanging fermions requires a separate spin and statistics construction. Auxiliary exterior parity does not supply that identification.

27.7 Four sums and a count

Chapter twelve proved that a force works if and only if its charge is conserved, tight enough that violating it leaves no theory at all rather than a damaged one. The quantum version of that proof adds conditions, and they are arithmetic.

Certain sums, taken over every species on the list at once, have to come out at exactly zero. When one of them does not, the conservation law the force depends on fails and the theory stops predicting anything. Sums of this kind are called **anomalies**. A matter list either satisfies all of them or describes no possible world. Every entry contributes, so the list has to be checked whole. A species cannot be added or dropped on its own.

Five conditions apply to a list like this one. Take the first.

The input is every species the strong force touches, tagged with its hypercharge: the quark pair, which has two weak members and so counts twice, the anti-up, and the anti-down. The operation adds their hypercharges, with color triplets and antitriplets entering the same way. One sixth counted twice is one third. The anti-up brings minus two thirds. The anti-down brings plus one third. One third minus two thirds plus one third is zero.

The second is the harder one, because hypercharge enters three times over. The input is all fifteen states with their hypercharges cubed, each multiplied by how many states carry it. The quark pair supplies six states at a sixth, and a sixth cubed is one part in 216, so six of them give one thirty-sixth. The anti-up supplies three states at minus two thirds, and minus two thirds cubed is minus eight twenty-sevenths, so three of them give minus eight ninths. The anti-down supplies three at one third, giving one ninth. The lepton pair supplies two at minus one half, giving minus one quarter. The positron supplies one at plus one, giving one. Put all five over thirty-six and the numerators are one, minus thirty-two, four, minus nine and thirty-six, which add to zero.

Nothing on that list was arranged to make it come out. The tags were fixed by a balance condition and a grid, the multiplicities were fixed by

counting ways to pick slots out of five, and five cubes over a common thirty-six come to nothing.

The three conditions left over are cheaper. Adding the plain hypercharges of all fifteen states gives one, minus two, one, minus one and one, which is zero. Adding the hypercharges of the weak pairs gives three color copies at a sixth, a half in total, against the lepton pair's minus a half. The last condition is a count rather than a sum: a world with an odd number of weak pairs has no consistent weak force at all, which Edward Witten proved in 1982, and this generation carries four.

Five conditions. A list assembled by choosing slots and adding tags meets every one. In the way physics is normally taught these conditions arrive after the fact, as a check on charges that were read off a detector, and their passing looks like a coincidence. Here they are the sieve, and 1,022 of the 1,024 lists fail them.

27.8 A clock with six positions

The sums fix the charges relative to one another. The unit they are counted in comes from somewhere else.

Chapter four's clock face bundles three and fifteen together and declares them one thing, because for a clock's purposes the difference between them does not exist. Do that to a group of symmetry moves and you get a **quotient group**: find the moves that change nothing observable, bundle every one of them in with doing nothing at all, and what is left over is the group that does the physical work. The clock bundles the multiples of twelve. The question here is which combinations of gauge moves this world bundles.

Chapter twenty-six's center supplies the moves to try. The center of a group is the moves that commute with everything, and for the strong force it is the move that advances all three color slots together by a third of a full turn, for the weak force it is the half turn on both weak slots, and for hypercharge it is every rotation of the dial. Combine one from each and you get a move that might do nothing, settled by trying it on the roster.

Take the quark pair, which sits in three colors, in a weak pair, at hypercharge one sixth. Apply the color move: a third of a turn. Apply the weak move: half a turn. Apply the hypercharge move, under which a species turns through a fraction of a full turn equal to its hypercharge, so a hypercharge of one sixth turns through one sixth of a turn. Add the three angles. A third plus a half plus a sixth is two sixths plus three sixths plus one sixth, or six sixths, which is one complete turn. The quark pair comes back exactly where it started.

Try the anti-up, which takes two color slots and no weak slot and sits at hypercharge minus two thirds. Each color slot it holds turns a third, so the object turns two thirds, and the hypercharge move contributes minus four

sixths, which is the same two thirds backward. They cancel. The lepton pair takes all three color slots, a whole turn by itself, and adds half a turn from the weak move against minus a half from hypercharge. The positron takes both weak slots, which is another whole turn, and one more turn from its hypercharge of plus one. All fifteen states come home.

So there is a combination of gauge moves that does nothing whatever to any piece of matter that exists, and repeating it generates a family of exactly six such combinations, counting the one that does nothing to start with. Six moves that nothing in the world can detect. By the argument chapter four made about permitted relabelings, they get bundled in with the identity, and the group that does the physical work is the quotient.

Which is where the sixths come from. The bundling holds together only if the hypercharge dial advances in steps that fit that six-element pattern. A species whose hypercharge misses a whole multiple of one sixth turns through an angle that spoils the cancellation above, so it cannot exist without breaking a bundle that is a fact about which moves the world can tell apart. Every hypercharge in nature is a whole number of sixths. Which sixth is tied to how many color slots the species carries. That tie is where the electric charges land: anything carrying color comes out on a third, and anything without color comes out on a whole number.

That is the lock. The down quark and the electron are separated by everything. They are counted in the same unit, because the unit belongs to the wiring rather than to either of them. The down quark stands at minus one third, which is minus two sixths, and the electron stands at minus one, which is minus six sixths, and three times two is six, exactly and leaving zero room for a discrepancy in the twentieth decimal place. The proton, being two ups and a down, lands on plus one for the same reason. The microphone in the cavity had nothing to hear.

27.9 Three at the least, five at the most

That settles the charges. The whole of it then arrives three times over.

The muon is a heavier electron and nothing else: the same charge and the same weak behavior, with a mass two hundred and seven times larger and a lifetime of two microseconds. The tau is heavier again. Behind each of them sits a full copy of the fifteen-state roster, quarks included. Isidor Rabi supplied the standard phrasing on being told about the muon, “who ordered that”, and it holds for the third copy as well.

Start with the floor, which comes from matter and antimatter being different at all.

Charge conjugation swaps every particle for its antiparticle. Combine it with a mirror reflection and you have an operation turning any process into its opposite-and-reflected version, and the question is whether the two run at the same rate. On 27 July 1964 James Christenson, James Cronin,

Val Fitch and René Turlay published the answer from the Brookhaven accelerator: among the long-lived neutral kaons in their beam, 45 events decayed the way the symmetry forbids, about one in five hundred. Matter and antimatter are physically different, by a little, and a universe with any matter left in it needs that difference.

The weak force makes them different through the way it couples quarks across families. That coupling is a square table with one row and one column per family. A real difference between matter and antimatter needs an entry in it carrying a complex phase that no relabeling of the fields removes. Plain rotation angles are useless here, because a rotation angle looks the same to matter and to antimatter. Counting the phases that survive relabeling is arithmetic on the size of the table:

$$\frac{(N-1)(N-2)}{2}$$

Here N is the number of families, and the expression counts the complex phases left over after every field has been re-phased to absorb whatever can be absorbed. Put N equal to one or two and the answer is zero. Put N equal to three and it is one. With fewer than three families the table carries no such phase, and kaons decay symmetrically. Makoto Kobayashi and Toshihide Maskawa ran that reasoning in 1973, when three quarks were known and a fourth was a rumor, and concluded that there had to be six.

The ceiling comes from asymptotic freedom. Put a charge in a vacuum and the vacuum responds: pairs flicker in and out around it, orient themselves and screen it, so the charge you measure from far away is smaller than the charge close up. The strong force does the reverse, because its own carriers carry the charge they transmit, which spreads the charge outward instead of hiding it and leaves the force feebler at short range than at long. David Gross, David Politzer and Frank Wilczek worked this out in 1973. It is why quarks rattle around almost freely inside a proton while being impossible to pull out of one.

Species of matter push the other way, back toward ordinary screening, and each family adds species. The tally that runs out first belongs to the weak charge. It starts at twenty-two thirds, each family costs four thirds because a family brings three colored quark pairs plus one lepton pair, the spin-zero field costs a sixth, and the total has to stay positive. Solve it and the bound is five and three eighths families.

Which leaves three, four or five: a floor set by the difference between matter and antimatter, a ceiling set by asymptotic freedom, and nothing in either constraint that picks a number.

A minimum-cost family-band rule offers a candidate choice. The twelve ports of chapter fourteen sit on a solid with twenty triangular faces, the sixty rotations carry any face to any other, and each face has its own

threefold rotation cycling its three corners. Something has to carry that threefold turn. Identifying the number of physical families with the width of that band is an additional attachment rule. Chapter fifteen inventoried what is available: the twelve directions break into uncuttable pieces of sizes one, three, three and five, and the piece of size one takes the same value under every rotation and can therefore carry no phase whatever. There is no irreducible piece of size four or two on that list. This restricts the proposed irreducible-band attachment; it does not exclude every other family construction.

The three eligible pieces can also be priced. Ask what it costs the screen to carry a family band in each of them, and the three answers are five minus the square root of five, six, and five plus the square root of five: 2.7639, 6 and 7.2361. The cheapest is the first, which is a piece of size three. The spectrum fixes that ranking. Selecting the cheapest eligible band as the physical family space remains a hypothesis.

A rank-three candidate band. An independent experimental count concerns light active neutrinos: Through the 1990s the four detectors at CERN's Large Electron-Positron collider measured how often the Z boson decays into things nobody sees, which is neutrinos, and divided by the rate for a single species. The full data set gives 2.9840 plus or minus 0.0082 light neutrino species, which is three to within a percent. This constrains species light enough to enter that decay; it is not a count of every possible family sector.

Every entry on that roster is a set of charges. A charge is an answer: what the strong force gets back when it asks, what the weak force gets back, what the dial reads. Ask an up quark what it is, as opposed to what it answers to, and five slots and fifteen states say nothing at all.

27.10 Three copies and three different masses

A three-dimensional response band does not determine three distinct masses. If both family factors carry the same unbroken irreducible source symmetry and the Higgs does not distinguish families, every invariant mass-generating intertwiner is a scalar times the identity. Its three singular values are equal. The exact source calculation verifies this constraint.

Splitting the masses therefore needs a specified asymmetric background or interaction. Choosing a distinguished port gives a splitting control; averaging it over the whole symmetry removes the splitting. Neither operation selects the observed hierarchy. The family-band result and the physical family identification are separate statements.

Chapter 28

What Is a Particle, What Is a Wave, and Why Does It Look Like Both?

An electron lands at one spot. The spots it chooses over many landings are the spots a wave makes bright.

The argument about how one object manages both has been running since 1905, when Einstein published a paper in the *Annalen der Physik* proposing that light is emitted and absorbed in lumps of a definite size, at a moment when every optical instrument in Europe worked because light is a wave. Textbooks have mostly handled the tension by saying the object has two natures.

For the electron, the evidence for the wave started with an accident. In April 1925, in a laboratory on West Street in Manhattan, a bottle of liquid air blew apart next to an electron-scattering experiment Clinton Davisson had been running for years and that Lester Germer had joined in 1924. The bottle stood near a vacuum tube containing a nickel target. The target was hot. Air rushed in and the nickel oxidized.

They spent the following weeks baking the oxide off, heating the target for long stretches to drive the damage out, and then started the run again. The experiment was simple to describe: fire electrons at the nickel, count how many come back at each angle. Before the accident the counts had traced a smooth hill. Afterward the same apparatus gave sharp peaks at particular angles and very little in between.

Nothing about the electron gun had changed. What had changed was the metal. The prolonged heating had recrystallized the target, and where the nickel had been a mass of tiny crystals pointing every way, it had become a handful of large ones. A handful of large crystals is a diffraction grating. They rebuilt the apparatus around a single crystal. The strongest peak came at an accelerating voltage of 54 volts and an angle of 50 degrees, and the atomic spacing of nickel converts that angle into a wavelength of 0.165 nanometers, against the 0.167 nanometers Louis de Broglie's rule of 1924 assigns an electron of that speed.

Nickel settled one half of the matter. The other half was put on film in 1989 at the Hitachi Advanced Research Laboratory, where Akira Tono-

mura's group sent electrons past a charged wire that bends two paths back together, the equivalent of a two-slit arrangement. They ran the beam so faint that the instrument held one electron at a time. Each one landed as a single dot at a definite place on a counting detector. It took about twenty minutes of accumulating dots for the stripes to appear.

The question the nickel put, and the charged wire put again sixty-four years later, is how a spreading pattern and a single count can be one thing happening.

28.1 One word, two objects

A value assigned at every place. That is one of the two things physics calls a field, and chapter twenty-six built one out of a table of who touches whom: a number at every port, falling off with hop distance in four exact fractions, and no space anywhere in the calculation.

The other one starts with a room.

Take a region and ask what can be answered inside it without leaving. Chapter nineteen made a question into an operation. A collection of operations you can compose, scale and add is an algebra. So attach to each region the algebra of everything answerable there, with its composition rule. Regions that contain other regions carry longer lists, because anything answerable in the small one is answerable in the large one. The small algebra sits inside the big one exactly.

The list is longer than it sounds. Everything a thermometer, a photographic plate, a photomultiplier and a wristwatch inside that region could between them establish is on it, along with every combination of those readings, and nothing whose answer would require somebody to go outside and look.

Take two regions far enough apart that no light ray leaves one and reaches the other while the questions are being put. Every operation on one side then commutes with every operation on the other. Alice asks and Bob asks, in either order, and the two orders give the same answers, so there is no fact about who went first for anybody to establish. Chapter twenty used that as a step in the Bell argument, where it is the statement that neither of them can signal to the other.

An assignment of an algebra to every region, with those two properties, is what the word field names in quantum field theory. There is no arrow at a point anywhere in it.

On the carrier that structure is exact. Two observer regions with no seam in common sit as exactly the two tensor factors of the joint description. Chapter twenty built the tensor product out of a deck of cards: four suits, thirteen ranks, dimensions multiplying, and most joint descriptions refusing to come apart into a fact about the suit and a fact about the rank. Two separated regions are the suit and the rank. Each one's questions

are questions about its own factor, the two lists commute across the gap, and the pair's description is assembled by the operation that multiplies dimensions.

Everything in that construction is countable. The regions are finite collections of patches, the lists of questions are finite, and the commuting across the gap is arithmetic on the wiring rather than the limit of a sequence of approximations, so a continuum is never called for and never used. Chapter twenty-six's field is a value at every place. This one is a list of questions per region. Physics runs both under one word. The second is the one the Standard Model is written in.

Chapter twenty-two gave the second kind its quanta and its clock. Many copies of one thermal mode, stepped by an observer's count of repairs, converge to a field whose two halves fail to commute by exactly the amount a position and a momentum do, and the region's own flow acts on that field as time translation.

28.2 Why the iron bar does not slosh

Warm one end of an iron bar and the warmth crawls along and stays, arriving late at the far end and never overshooting it. Chapter nine's repair move is that same equation written on a graph: my reading minus the average of my neighbors', a sixtieth at a time. Diffusion by itself produces no waves at all.

What produces a wave is a second quantity that the first one feeds. A pendulum has two: displacement sets the rate at which velocity changes, and velocity sets the rate at which displacement changes. The pair circles instead of settling. Conserving the sum in a repair does not by itself supply that paired evolution. The active coherent construction supplies a wave evolution using the source operator; a separate classical construction supplies a Maxwell action and paired field update. Each needs its stated field variables, evolution and clock. Once those are attached, the seam operator supplies a spatial stiffness that can be read and tested.

In the Maxwell construction, fix a direction of travel and a wavelength, and the wavelength fixes a wavenumber: how many crests fit into a set length. After imposing the gauge constraint, the propagating amplitudes are perpendicular to that direction. In the declared three-dimensional space the perpendicular directions form a plane. Two of them. Two is a rank in chapter fifteen's sense, the number of independent directions a table of numbers contains. This count uses the field action and constraint as well as the dimension of the observer's rest space.

Write A for the amplitude in that perpendicular plane, and write the Greek letter lambda, capitalized, for the stiffness the wiring supplies at a given wavenumber k, which is the number saying how hard the carrier

pushes back on a mismatch of that wavelength. Two quantities feeding each other give one equation.

$$A'' + \Lambda(k)\, A = 0$$

The double prime is the rate of change of the rate of change in the model time used for the wave evolution. That is the equation of a mass on a spring with the wiring supplying the spring, and its solutions oscillate at a frequency whose square is the stiffness. Reading that frequency on an observer's clock also uses the construction's clock conversion.

For the declared spatial placement, the stiffness is computed from the complete thirty-seam operator. At long wavelengths it approaches the square of the wavenumber. Its first correction subtracts a twentieth of the fourth power times the square of the seam spacing, and an exact bound holds across the full operator: the difference from the square of the wavenumber never exceeds that correction in magnitude. At zero wavenumber the stiffness is exactly zero, so the frequency there is exactly zero. That exact zero, with the stiffness rising as the square of the wavenumber above it, gives the long waves one limiting speed in the declared space and time units. An event count supplies a clock coordinate; its conversion to this evolution parameter is part of the attachment.

This controls the whole evolving field in the stated continuum family. Lay the seam-shaped operator across the supplied three-dimensional space and shrink its spacing with the prescribed wave evolution. The paired fields converge to Maxwell's equations, with an error bound proportional to the square of the spacing for sufficiently smooth initial fields. Every field of finite total energy converges, even without that smoothness. A current can be included, and conservation of its charge preserves Gauss's law.

A wave is therefore an executed or proved field continuation under those premises. The wiring contributes its stiffness. The coherent service or classical field action, spatial interpretation and clock determine how that stiffness becomes a propagating response; passive repair alone does not make the identification.

A finite software observer can read its own field without keeping a spare copy. It remembers a port's value, averages that port with an adjacent value, and reads the average. Twice the average minus the remembered value recovers the hidden value. Writing the recovered pair back lets the next measurement start from the same state. Applied to forty-two potential coordinates, this procedure produces the records that drive a sourced Maxwell update, and the fields reconstructed from those records satisfy the same finite action, including the charged paths.

The recorded potentials also define fields inside a solid built by joining the carrier's twenty triangular faces to its center, which gives the surface description an explicit volume. A geometric interpolation preserves magnetic

flux closure and Faraday induction. Adding up the field energy throughout that volume gives a different action from counting its stored coordinates, and interpolation between time steps adds a separate correction. Both differences are calculated on the recorded history.

The volume can also carry its own evolution. Its geometric action gives equations for every edge, including the edges running through the interior. A field that starts out satisfying Gauss's law keeps satisfying it, and an exact bound controls its field energy. A software observer reads and restores fifty-five potential coordinates and uses the decoded records to advance the field, so the records carry a complete history of the volume that keeps its action stationary.

The same geometric action has thirty independent radiative modes. Quantized by the usual rule, they become thirty quantum oscillators. Their occupation numbers form an infinite Hilbert space, and a self-adjoint Hamiltonian evolves its states unitarily. That is the free radiative sector.

Dynamical matter enters as a charged scalar field. Its interpolation changes with the electromagnetic potential as gauge symmetry requires. Both fields follow one stationary action, and positive conserved energy ensures global classical solutions of the finite equations. A short numerical evolution starts with opposite charges near the center and the boundary. Five software patches record, probe and restore the symmetry-reduced state, and each advance reads those decoded records. Independent reconstruction checks every finite field equation. Exact interval arithmetic encloses the same solution throughout two units of time, and replaying the observer records reproduces all five positions and five velocities at its eighty-one decoded checkpoints to within 1.0001×10^{-10} of that solution.

A real scalar sector gives a controlled continuum of evolving matter. Its electromagnetic fields vanish and its quartic self-interaction stays. Project a smooth continuum solution onto the finite mesh and subdivide the solid without making the tetrahedra progressively flatter, and the motion approaches the continuum nonlinear wave equation. The error in the field and its velocity decreases with the cell size over a fixed time interval. The same convergence rate holds for complex matter in a uniform magnetic field, with initial data projected using its magnetic energy.

The changing configuration also keeps time. Its kinetic metric and conserved energy assign a duration to every stretch of motion with positive kinetic energy, whatever labels were used to order the samples. Applied to decoded records, this approximates the action's elapsed time without reading a single timestamp.

The full interacting action has a Hilbert-space description as well. Removing redundant gauge descriptions gives a complete positive kinetic metric. Its volume measure and its Laplace–Beltrami operator, the curved-space version of the ordinary Laplacian, fix a Hamiltonian with a unique self-adjoint extension, and that Hamiltonian evolves neutral states unitar-

ily. An explicit normalized initial state gives exactly calculated matter observables.

The classical charged state also provides position and momentum values for preparing quantum states, once the redundant gauge coordinates are removed. Gaussian packets centered on those values keep all fifty-six configuration directions. Averaging over the remaining common phase produces neutral states with an exact positive normalization formula, and their scalar radius can be calculated.

A trial history on the full fifty-six-dimensional configuration space changes the centered state's phase and leaves its configuration probabilities fixed. It stays within one tenth of the exact quantum evolution in Hilbert norm for times up to 2^{-55} in the action's units.

A separate comparison theorem states what a useful effective quantum description must control. Preparation error, the difference between the full Hamiltonians, detector mismatch and probability outside the chosen energy window together bound the error in measured probabilities. Coupling to omitted states matters even when the retained matrix entries agree exactly.

28.3 Two questions, two orders

Amplitudes can cancel and probabilities cannot. That is chapter nineteen's difference and the reason there are dark bands at all. Cancellation needs a sign. A pattern that slides as one path is lengthened needs more than a sign, because it needs something that turns continuously, and the amplitude turns: it carries an angle, and the arithmetic of turning angles is the arithmetic of the square root of minus one.

The suspicion that the turning is bookkeeping is reasonable enough. Turning arrows multiply conveniently, and there might be real numbers underneath doing the work.

The carrier settles it on a two-by-two table.

The machine hands over two yes-or-no questions from behind one public reading, both real, both built out of labels an actual run produced, and the two fail to commute. It hands over two states as well, and every real question in the entire web scores those two states identically: put any of them to either state and the same odds come back. The states are different states. Nothing real tells them apart.

Build the missing question out of what is there. Start with the even split, which is the do-nothing operation halved and which as a description gives fifty-fifty odds to every question in sight. Take the two questions, call them P and Q, do them in one order and then in the other, and subtract. That is chapter nineteen's commutator. Multiply that difference by the square root of minus one and by two divided by the square root of three. Subtract the result from the even split, and call what comes out Y.

$$Y = \frac{1}{2} I - \frac{2\sqrt{3}}{3}\, i\,(QP - PQ)$$

I is the operation that does nothing, P and Q are the two real questions the carrier supplied, and i is the square root of minus one.

Y squares to itself, making it a question in chapter nineteen's sense, with a yes, a no and no third possibility. Put it to the two states that nothing real could separate. It answers yes on one of them with certainty and no on the other with certainty.

The coefficient was not fitted to make that happen. The commutator of those two particular questions has off-diagonal entry exactly the square root of three over four, and two over the square root of three scales that entry to exactly one half, which is the value that makes the result square to itself. Two real questions and one subtraction produce the direction that neither of them held.

And the direction is unreachable from below. Close the real questions under everything real processing does to them: average them, scale them, take complements, push them through any real operation of the kind a physical process performs on a description. Every element of that closure hands the two states the same number, and Y hands them different ones, so no amount of coarse-graining can manufacture it.

Three questions then fix everything. The record question the run supplied, its rotated partner, and Y between them pin down any description of a two-outcome system whose odds add up to one, leaving nothing about it undetermined.

The direction that was missing is the phase. Ernst Stueckelberg tried the other route in 1960, writing quantum theory in a real space in *Helvetica Physica Acta*, and made it work by adding an operator that squares to minus one and commutes with every observable, which is the imaginary unit under another name.

28.4 The leftover, named

Twelve observers wired into an icosahedron, one record out of thirty flipped, two thousand and forty-eight starting states and sixteen schedules apiece: chapter six ran all of it. Every run converged. Every run left exactly one unit of disagreement sitting in the system, movable to any location and removable from none, with no edge of the world to push it off.

That leftover motivates three conditions for a **particle candidate**. They organize a search; identifying a physical particle also requires its field response.

It is a pattern in the way record content moves from port to port, a property of the transport rule rather than a lump occupying a spot. Every observer whose patch overlaps yours reads the same pattern in the overlap,

so it survives comparison and is not an artifact of one observer's labels. And it survives a regrade: change the resolution of the records, finer or coarser, and the pattern is there with its identity intact. Fluctuations of the repair process fail that last condition, because the faster bands decay faster and wash out under regrading.

Two checks make the candidate quantitative, and both require a specified vacuum or stationary reference for the continuing dynamics. The zero-disagreement fixed state of chapter six is a reference for its static repair grammar. It does not select the physical vacuum of a quantum field description.

Does the pattern propagate: expand the response around the chosen reference to second order, remove the redundancy, and test the surviving modes for positive norm and the claimed dispersion. An isolated stable one-particle state gives a sharp contribution to a spectral response, from which its invariant mass can be read. This is a sufficient idealization with its own assumptions. Charged states can require a cloud of soft photons, while the W, Z, Higgs and top are unstable and have resonance widths. Confined quarks have masses quoted in a specified running convention rather than as free isolated particles. Each reading requires its field dynamics, normalization, clock and matching procedure. A decay rate in a repair calculation supplies none of those identifications by itself.

28.5 One clock for several fields

The conditional scalar action determines a metric that can also express spinor and electromagnetic wave equations with the same local light cone. A finite fermion construction can share the scalar's spatial operator and the observer's event-count clock. This joins actual propagation formulas, although its extra modes and boundary conditions require physical interpretation.

A useful alternative completes a finite geometric complex through its faces and volumes. Spurious stationary edge cycles become propagating form modes, while the scalar equation is preserved. Removing a truncation artifact does not identify those forms with the observed Weyl particles. A finite cavity frequency also changes when the cavity changes size; that frequency is not an intrinsic rest mass merely because a detector can resolve it.

The particle identification must use the same field, interactions and clock. Its spectral response, with momentum and physical normalization specified, distinguishes a particle mass from a relaxation rate, a background offset or a geometric resonance.

28.6 Reading familiar field roles

One direction, three and eight in the twelve-dimensional gauge response. Chapter twenty-seven adds a conditional fifteen-state matter representation. With three declared families, a Higgs doublet and their common field action, these structures can express the familiar Standard Model roles. The physical family attachment and interacting fields remain premises. Counting the gauge directions does not classify every possible matter field or composite excitation.

The uniform carrier direction is the one all sixty rotations leave alone, and it is the central dial of chapter twenty-six, the one direction that commutes with everything. Give it its Maxwell action, impose the gauge symmetry and its constraints, and the field that propagates has two transverse polarizations. Identifying it with light requires the electromagnetic current and normalization; in the electroweak action, the photon is the unbroken combination of the central and weak directions.

The three and the eight do not commute. Their brackets enter the nonabelian curvature, giving a term quadratic in the gauge field. In a Yang–Mills action this produces interactions among the gauge fields themselves. The bracket fixes this algebraic structure. It does not establish a linear potential, confinement or the spectrum of bound states. Those require the interacting quantum dynamics and its physical regime; no confinement calculation follows from the block count alone.

In the declared electroweak action, the Higgs vacuum distinguishes massive gauge modes from the photon. Its magnitude is an input: the familiar value, about 246.22 billion electron-volts, is calibrated from weak-interaction measurements. The moves that leave the direction of the Higgs vector alone form a two-parameter family; the moves that leave the vector itself alone form a one-parameter family. Three of the four electroweak directions move it and one does not. The quadratic action gives the three broken directions a nonzero frequency at zero wavenumber, while the unbroken combination stays massless. With the relativistic kinetic normalization, the added constant in the squared frequency is a squared rest energy. A physical resonance mass requires the quantum corrections and matching procedure too: the measured Z value, 91.1876 billion electron-volts, is a comparison, not a mass derived from counting the directions. The polarization bookkeeping does close within this action. Three massless gauge modes with two polarizations each, plus three displaced Higgs directions, become three massive modes with three polarizations each. Nine before and nine after.

Chapter twenty-seven's declared exterior construction gives a charge-conjugate pair of rank-fifteen matter projectors. Repeating one representative over three declared families gives forty-five internal labels. Their identification with physical chiral matter requires the spin, field and family

attachments. Orthogonal projectors describe exclusive alternatives in that internal space; they do not forbid two particles from occupying a mode. Pauli exclusion follows in the separate many-body fermion construction, whose creation operators anticommute and square to zero. Attaching that CAR algebra to the same propagating matter fields supplies the exclusion rule. The auxiliary even-or-odd exterior grading alone does not.

The Higgs field supplies another role in this declared action. Around its chosen vacuum, its remaining excitation changes the magnitude of the vector. The curvature of the supplied potential sets the quadratic restoring term. The excitation is a scalar under spacetime rotations, so it has spin zero. Connecting that field response to the Higgs resonance announced by the two CERN collaborations on 4 July 2012 requires the interacting spectrum and its physical normalization. Neither the vacuum magnitude nor the resonance mass is fixed by the gauge algebra.

These are familiar roles that the conditional construction can accommodate. The exact splitting of the gauge response into one plus three plus eight constrains the gauge algebra. It does not prove that these are all the particles nature permits, or rule out additional matter, scalar fields or composite states. That question belongs to the selected interacting theory and its measured response.

28.7 When the record gets written

Questions that refuse to commute, and a center where the answers commute, survive being read twice, and survive being copied to a neighbor: chapter nineteen supplied both halves of the answer and left them lying apart. Pinching carries a question from the first into the second by deleting every comparison that only made sense between two bins.

Between commits, no record exists of which route a pattern took. What exists is the spreading mismatch of the wave, with a phase and a frequency. Every question about it lives in the part of the algebra that refuses to commute. Amplitudes for different routes add, opposite amplitudes cancel, and cancellation is what a dark band is made of.

At a commit a record is written. A record lives in the center. Everything there commutes with everything else, the answers survive reading and copying, and they are ordinary classical numbers: this one, here, count of one.

The electron crossed the tube on West Street with no record written anywhere of which crystal plane it scattered from, and arrived at the collector where a record was written and a count went up by one. The pattern of counts is the wave. Each count is the particle. Nothing about the electron differed between the crossing and the arrival except how much of it got written down.

The same reading settles what happens when somebody tries to watch. Put an instrument in the apparatus that succeeds in telling you which route was taken and the bands go. Making the instrument gentler does not help. No version of the instrument has ever been built gentle enough to report the route and leave the pattern standing, because reporting the route is writing a record, and a record is a commit. What the instrument deletes is every comparison between the two routes, which is where the cancellation lived. An instrument that fails to determine the route deletes nothing, and the bands survive.

A repair is such an instrument. Chapter nine's experiment puts an empty seam and a seam holding a single unit into superposition and lets one repair act. A repair that writes down only how many units crossed leaves most of the comparison standing. One that also writes the parity of the seam deletes it, and the bands go flat at three eighths. How much contrast the bands keep is a reading of what the repair writes.

Duality is a statement about when the record is written. The older question, which of two natures the thing has, carries an unstated variable, and the variable is the depth at which the reading is being taken. Fix the depth and the ambiguity goes. Nothing whatever happens to the electron at the plate. A record gets written, and a record holds a number.

28.8 The term that cannot be written

In 1919 Ernest Rutherford published four papers in the *London, Edinburgh and Dublin Philosophical Magazine* on what happens when alpha particles are fired into light gases. In nitrogen he saw scintillations at distances no alpha particle could travel, identified them as hydrogen nuclei knocked out of the nitrogen nucleus, and gave the thing its name the following year. Proton.

Chapter one told the other half. Grand unification put the eight, the three and the one inside a single simple symmetry, and the fractional charges and the family structure fell into place. The difficulty came with the carriers. Building three blocks into one simple group forces extra directions into the list of carriers. The smallest group with room for all three holds twenty-four carrier directions, which is the twelve the world has and twelve more. Every one of the extra twelve carries color and weak charge at once. A carrier of that kind couples a quark directly to a lepton, so a proton comes apart into a positron and a neutral pion.

The list of carrier directions on the twelve-port response is the sum of the blocks and nothing besides: eight color directions carrying no weak charge, three weak directions carrying no color, and one carrying neither. A simple group is compelled to mix its directions into one another, because its own internal moves carry any direction to any other, which is what being simple amounts to. A product of blocks has no move connecting one

block to another, so its list stays block by block. The direction a quark-to-lepton carrier would have to occupy is absent from a twelve-dimensional list whose splitting is fixed. So the carrier that would make a proton decay the grand-unified way, a gauge boson turning a quark into a lepton, cannot be assembled out of the directions that exist. That route is closed exactly, with no small coupling in it and no heavy mass sitting in a denominator to make it slow.

The companion problem goes the same way. A monopole of the unified kind is a knot in the field configuration, formed when a large symmetry is squeezed down to a smaller one, and whether such knots exist is settled by asking which loops in the space of ways to do the squeezing can be shrunk to a point without leaving that space. Whether a loop can be shrunk to nothing is a homotopy question. The loops that refuse are what count the knots. The carrier emits the product of blocks from the first moment there is a group at all, with no earlier stage at which it was simple, so the squeezing never happens and the loops are never drawn.

The fifty thousand tons of ultrapure water under the mountain in Japan have been running since April 1996. A search through 450 kiloton-years of data, which is the mass of the water multiplied by the years it was under observation, turned up nothing of the right shape. The floor that search puts under the decay sits at 2.4 times ten to the thirty-fourth years. The product gauge algebra excludes that particular grand-unified carrier. It does not exclude every possible baryon-violating interaction or prove a general proton-stability theorem.

Five entries, and not one of them says how hard anything pulls. Twelve directions, thirty seams and a census of what can hold together leave the photon on the list and the forty-five states of matter on the list, and say nothing whatever about the grip of the first on the second. That grip is a number near one part in 137. It sets the size of every atom in your hand.

Chapter 29

Why 1/137, and Why Is There a Limit to What the Universe Can Remember?

Wolfgang Pauli went into the Rotkreuzspital in Zurich in December 1958 with pancreatic cancer. His assistant Charles Enz came to visit him there, and Pauli asked whether he had noticed the number on the door of the room. It was 137. He died in that room on the fifteenth of the month.

The number had been at him for most of his working life. Take one electron and one photon and ask how strongly the first couples to the second. Write the answer so that every unit cancels out of it: no meters, no seconds, no kilograms, nothing that depends on anybody's choice of ruler. What is left is a bare number close to one part in 137. Arnold Sommerfeld pulled it out of the fine splitting of the lines in the hydrogen spectrum in 1916, where it appears squared, as a splitting about one part in nineteen thousand of the gap it sits inside. That is where the name fine-structure constant comes from. The alpha that stands for it in every equation is Sommerfeld's letter.

Arthur Eddington thought he could derive it. He had organized the 1919 eclipse expedition that measured starlight bending around the sun, he wrote the book that taught a generation of English speakers what relativity was, and by the end of the 1920s he was arguing that the reciprocal of this constant had to be a whole number, and that the whole number was 136, which he obtained by counting the independent components of an algebra. Then the measurements drifted toward 137. He revised the count, produced 137, and his colleagues started calling him Arthur Adding-One. The book he was folding all of it into, *Fundamental Theory*, appeared in 1946, two years after his death, and its derivation was not accepted.

Feynman put the number in *QED: The Strange Theory of Light and Matter* in 1985 and called it one of the greatest damn mysteries of physics, a magic number that comes to us with no understanding by man. He added that all good theoretical physicists put it up on their wall and worry about it.

What they are worrying about is one number that quantum electrodynamics does not set and cannot set. The theory also takes particle masses as inputs. Its interaction strength is supplied from outside. Once it is supplied it fixes the strength of every chemical bond, the rate at which an excited atom drops back down and throws off a photon, and the color of everything you have ever seen. A theory that computes the electron's magnetic moment to twelve places takes its central constant from a laboratory in Illinois.

That constant is measured to ten significant figures. Take a single electron, hold it in a Penning trap for months, and read off its magnetic moment: Xing Fan and his colleagues at Northwestern published that measurement in 2023 to thirteen parts in a hundred thousand billion. Turning it into the constant requires the theoretical side to be worked out to the same depth, which took the evaluation of all 12,672 Feynman diagrams that contribute at tenth order, a job finished by Tatsumi Aoyama, Masashi Hayakawa, Toichiro Kinoshita and Makiko Nio. The recommended reciprocal that comes out of the whole enterprise is 137.035999177, with an uncertainty of twenty-one in the last two places, which is one and a half parts in ten billion. In the Standard Model it is a measured input.

Where does that number come from, and why is there a ceiling on how much the universe can hold? OPH approaches both questions with one proposed move: take an object, read it from the outside and from the inside, and require the two readings to agree. The object is one cell of an observer's screen, and then, at the other end of every scale there is, the whole horizon.

29.1 A number with no units in it

The speed of light stopped being a measurement in 1983, when the meter was redefined to make it 299,792,458 meters per second exactly. Those digits are a fact about the second and the meter. Pick a different unit of length and every one of them changes, while light is unaffected.

A pure number has no such escape. 137.035999177 is what it is in any unit system, on any planet and for anyone who measures it, and the same goes for the 1836.15 that the proton outweighs the electron by. There is nothing in either of them that anybody chose. The electron in the innermost orbit of a hydrogen atom travels at one 137th of the speed of light. That is the same number again, the reason atoms are the size they are relative to the light they emit.

The strength of electromagnetism depends on how closely you look. A charge sitting in the vacuum is surrounded by pairs of particles flickering in and out, each pair oriented so as to screen the charge a little, so an experiment probing from far away sees a weakened version and an experiment that gets in closer sees more of the bare charge. The coupling carries a scale

with it. Read at rest, at the far end of the range where the probing photon has essentially no energy of its own, the reciprocal is 137.036. Read at the energy of the Z boson, ninety-one billion electronvolts, the same coupling comes out near 128. Colliders watched electrons and positrons scatter off each other at rising energies and read the coupling climbing as the collisions sharpened, so the sliding is measured rather than inferred. Say which energy a fine-structure constant was read at, or the number means nothing. The 137 value is the one at the far end.

29.2 The balance point

Chapter twenty-four showed that an outsider's finite record access is controlled by the boundary rather than the hidden interior. Chapter twenty-five supplies a gravitational area unit from Newton's constant. Measure the area of one cell of the screen in that unit. Call the resulting pure number the **grain**.

Start from the entropy balance used as a reference in the trial map.

A cap of screen is the region inside a circle drawn on it, the way a constellation is a region of sky, with an inside and an edge like any other bounded thing. Take one and count its entropy the way chapter twenty-four counted it, in two parts. There is what the cap holds inside it, and there is what the cut around its edge contributes, and the total is the sum of the two. Ask for the arrangement in which that accounting looks the same at every magnification: the total is to the inside as the inside is to the edge. Write x for the ratio of the total to the inside. Then the inside divided by the edge is also x, so the edge is the inside divided by x, and the total is the inside plus the inside divided by x. Divide through by the inside and what is left is x equals one plus one over x.

Which is the instruction from chapter two, the one with the calculator: divide one by your number, add one, repeat. Its fixed point is 1.6180339887, the golden ratio, reached here from an entropy balance on a screen rather than from a sequence of fractions on a phone. The same number sets the coordinates of the twelve corners of the carrier, a property of the icosahedron that was known long before anybody was looking for a coupling constant, and not something this argument arranged.

The entropy-balance ansatz associates that value with a reference point and uses the difference above it as an electromagnetic detuning. This interpretation is additional to the fixed-point arithmetic. A stationary distribution can support changing local states, clocks and records, so constant entropy alone does not prove that a universe contains no observers.

For the displayed candidate, the electromagnetic detuning is less than one percent of the grain. Interpreting that coordinate as the available budget for every physical record requires the same capacity and observable identifications used in the other chapters.

29.3 Two readings of one cell

So the grain sits a little above the balance point. The question is what fixes the distance.

Read the cell from the outside first. Its excess above the balance point becomes a pure number once the boundary is weighed per unit of area. The edge is a profile smeared across the boundary rather than a drawn line, and the normalization of that profile brings in the square root of pi, 1.772453851.

Read the same cell from the inside. There it is the smallest electromagnetic act available to an observer, the finest detail light can resolve.

The two readings describe one cell, so chapter thirteen's rule about consistent readings sets them equal:

$$P = \varphi + \alpha\sqrt{\pi}$$

Here P is the grain, the cell area in units of the gravitational area scale. The Greek letter phi is the golden ratio, 1.618033989, the entropy balance point. Alpha is the electromagnetic coupling carried to that grain, and the square root of pi is the boundary normalization.

The unknown appears on both sides. A trial grain fixes the coupling the forces share at the cell scale, and running that coupling down the range returns an electromagnetic coupling. Feed in a grain and get a grain back.

29.4 The last three operations

Take a calculator again.

The chain from a trial grain down to a coupling is the part a machine does. Hand it the grain 1.6309682414 and it hands back 137.03566014. The coupling is the reciprocal, 0.00729737062.

Multiply by the boundary normalization, 1.7724538509, and you get 0.0129342527.

Add the entropy balance point, 1.6180339887, and you get 1.6309682414.

Which is what was fed in, to the last of the ten decimal places quoted. That grain is the fixed point of the declared map. The detuning coordinate is 0.0129342527, about eight parts in a thousand of the grain. Arithmetic closure does not prove that this coordinate separates universes with observers from universes without them.

Run it backwards if you prefer. Take the balance point away from the grain and 0.0129342527 is left. Divide that by the boundary normalization and you have 0.00729737064. Take the reciprocal and 137.035660 comes back out, which is the coupling returned by this declared screen-cell map. Its identification with the measured electromagnetic endpoint requires the stated transport and same-quantity hypotheses.

The obvious suspicion is the one Eddington earned: an expression can reproduce a measured number while hiding choices in its construction. Once its inputs and branch conventions are fixed, each declared map here has a definite output. The golden ratio follows the entropy-balance ansatz, while the square root of pi belongs to the specified boundary profile. Change the boundary factor from 1.7724538509 to a round 1.77 and the displayed loop returns 1.630950 instead of 1.630968. The two tested versions, bare and shared-width, have certified inverse-coupling roots 136.994835 and 137.035660. This is not an exhaustive menu of admissible maps. The physical selection of the map, profile, charged-spectrum inputs and branches remains part of the task; uniqueness inside a specified map does not remove those choices.

The second test is what else the number determines. An expression that hits one target and produces nothing further is probably a coincidence, and the literature since the 1930s contains many such expressions. The grain is the coordinate the rest of the numbers are read from: the single coupling the three forces share at the cell scale, which comes out at one part in 24.32, conditional running and electroweak coordinates. The quark masses require additional family and matching data; they are not predictions of this one coordinate. Move the grain in the ninth decimal place and every one of those moves with it.

29.5 Two endpoints instead of one number

Chapter two's other root, minus 0.618034, is a fixed point nothing ever arrives at: start a calculator anywhere near it and the numbers flee. This one is approached, and the rate can be bounded.

Take two trial grains a small distance apart and run both through the chain. What comes back is two grains at most 0.0724 of that distance apart. A map that pulls its inputs closer together like that is a **contraction**, and a contraction has exactly one fixed point and drags everything toward it. Seed it with 1, a guess wrong by more than a third. After one pass the answer is inside 0.0457 of the destination, after two inside 0.0033, after three inside 0.00024, and after ten passes inside two parts in a million million. Twelve digits, from a deliberately poor starting guess, in ten passes, under the stated transport inputs and without fitting the measured fine-structure constant.

Arriving at a number and proving one are different operations, and a chain of this length rounds at every step. Anybody who has written software has met the small version: ask a computer for 0.1 plus 0.2 and it answers 0.30000000000000004, because a tenth has no exact form in binary. Ten thousand operations later, the digits on the screen are a report about the arithmetic rather than about the world.

The repair is to stop carrying numbers and start carrying pairs of them. Every quantity becomes two endpoints, a low one and a high one, with the true value somewhere between. Every operation rounds the low end downward and the high end upward, so the bracket can widen and can never lie. Add the bracket that runs from one to two to the bracket that runs from three to five and the answer is the bracket from four to seven. Multiply two brackets and you get a bracket containing every product of anything in the first with anything in the second. This is called **interval arithmetic**. It turns a computation into a proof. Feed an interval of grains into the chain, and if what comes back is an interval strictly inside the one you fed in, then a fixed point sits in there as a theorem, and the root it holds is certified rather than merely computed. Sampling the slope of the chain at points along it suggests it pulls its inputs together by a factor of about 0.003 a pass. The interval run proves 0.0724, which is the worse figure and which holds at every point of the domain at once, so 0.0724 is the number the argument is allowed to use. The proof that the declared interval maps inside itself takes 203 seconds of processor time. It certifies the map's arithmetic; the connection to measured electromagnetism requires its physical matching.

29.6 The quark cloud

The fixed point puts the coupling at 137.035660. The laboratory value is 137.035999. The two differ by two and a half parts per million. That is a certified mismatch for the declared map, whose physical matching remains incomplete.

The loop is built in two stages. Run it with the cell's own electromagnetic reading alone and it settles at 136.994835, three parts in ten thousand short of the laboratory value. That is the bare root. The weak and strong forces take their share of the cell as well: their combined width has to fit inside a quarter of the grain. Put that shared width inside the loop, so that every pass recomputes it from the grain it is handed, and the fixed point moves to 137.035660. Nothing in the second stage looks at the measured value, and that one step closes more than ninety-nine percent of the distance between the bare root and the laboratory. The two stages are the two versions of the loop in the freedom count above.

The laboratory reads the low-momentum electromagnetic response. Quark vacuum polarization contributes to that response. Its strong-interaction part requires a calculation beyond perturbation theory, in the same convention as the loop. Neither trial map supplies that calculation. The interval run proves that numerical error in the fixed-point solve cannot account for the discrepancy. It does not prove that the omitted quark contribution has exactly the sign and size needed to remove it. The transport and physical identification remain separate tests.

29.7 A number from two entangled ends

Alex Wissner-Gross gives an independent finite spectral candidate built from entangled qutrit and qubit pairs. Its exact expression is

$$\alpha_{\text{pair}} = \frac{79 + 48\sqrt{2} + 42\sqrt{3} + 3\sqrt{6}}{31104}.$$

Its inverse is 137.035999189566. The weights count the eigenspaces of specified exchange operators, and that arithmetic can be reconstructed without fitting a continuous coefficient. The proposed identification with the electromagnetic coupling is a conjecture: the observer construction must identify the same conserved current, photon pole and low-momentum response. Equal dimensions of two matrix spaces do not supply that physical identity.

If both the candidate identification and the same-cell equation are granted, the grain is 1.630968209402773. This substitution does not change either of the two trial maps above, each of which keeps its own certified fixed point and physical endpoint test. The candidate gives a further conditional route through the same-cell equation. It supplies neither the missing normalization proof nor a derivation of the particle masses.

The primary account is Wissner-Gross's fine-structure preprint.

29.8 The horizon's budget

The same move works on the largest object an observer has.

Look far enough out and the expansion carries a galaxy away faster than light can close the distance, so its signals arrive at no point of your future, ever. The sphere at that distance is your horizon. Gary Gibbons and Stephen Hawking showed in 1977 that it carries a temperature, a couple of thousandths of a billionth of a billionth of a billionth of a degree. Chapter twenty-four counted what a boundary can hold, and the horizon is a boundary.

Counting the records that survive is a different count from counting the ones that can be written. Take a channel with five symbols arranged in a ring, where each symbol can be confused with either of its two neighbors and with nothing else. How many can you use with zero chance of being misunderstood? Two. Number them one to five around the ring and send only one and three. Those two are not neighbors, so nothing can turn either into the other, and there is no room for a third: two sits next to both of them, four sits next to the three, five sits next to the one, and the ring holds nothing else. **Zero-error capacity** counts the messages that stay distinguishable whatever the channel does to them, the count that matters when a record has to survive being read back by somebody else.

Read the horizon two ways: from outside by its geometric area, and from inside by the durable public records observers can write, keep, tell apart and read back. The two readings describe one horizon, and setting them equal fixes its capacity.

The arithmetic takes one input, the grain, and one correction. A shipping container holds slightly less than its stated volume, because the bracing that keeps the load from shifting takes up room. Every shared cut on the screen is assigned a small uniform reserve in the same way: the grain divided by twenty-four, chapter twenty-six's twelve dials counted twice, once out and once back. The reserve is 0.067957 against a budget of 281.03, and because the capacity is an exponential of the budget, those seven hundredths move it by six and a half percent, from 3.5321 times ten to the 122nd before the reserve to 3.3001 times ten to the 122nd after it.

The reserve can be booked two ways, as the fraction of cuts that survive it, one minus the grain over twenty-four, or as the exponential of the same quantity. The exponential gives the 3.3001 above and the fraction gives 3.2921 times ten to the 122nd. The figure the sky reports, read off the Planck satellite's measurement of the expansion, is 3.3129 times ten to the 122nd, four parts in a thousand above one booking and six above the other. Which booking the horizon keeps, and the exact form of the map that reads its records back, are work in progress.

The capacity is the horizon's entropy, so it fixes the cosmological constant through the dimensionless relation $\Lambda \ell_\star^2 = 3\pi / N_\star$. The capacity does not change with time, so dark energy has the equation of state $w = -1$ at every epoch, and a measured evolution of w would falsify it. Five thousand and twenty robotic fibers on the four-meter Mayall telescope at Kitt Peak, repointing themselves to a few microns every twenty minutes, are collecting the spectra of tens of millions of galaxies that test such constant-dark-energy models. The survey's second data release leans against a constant ratio: its combinations place it between 2.6 and 3.9 standard units from their best fits. That is short of an exclusion, and the test fixed in advance waits for the third release.

29.9 One value, and why there is one

Take those two numbers away from the machinery for a moment and look at what they normally are.

Alpha is a dial. Quantum electrodynamics is a consistent theory at any setting of it, and nothing inside the theory prefers one setting to another. You measure it, you enter it, and everything downstream follows. The cosmological constant is worse, because it enters the field equations as a term you are permitted to add with any coefficient at all, and the one estimate that tries to compute it from the vacuum overshoots the sky by a hundred and twenty orders of magnitude. The two govern nothing in

common. One sets the size of an atom. The other sets how fast the universe comes apart.

This chapter makes a single move on both of them. Read a cell of screen from the outside and from the inside, and set the two readings equal. Read the horizon from the outside and from the inside, and set those equal. Both are chapter thirteen's rule about seams.

That demand needs something to hold before it can be made at all. It needs the world to have an inside to be read from, and it needs what the inside holds to be a description of the thing the inside is part of. A structure carrying no account of itself has no second reading to set against the first. The equation cannot be written, and the number stays free.

The constants are wired together. The grain P enters the coupling the forces share, running that down the range returns alpha, and alpha appears back inside P. The grain also enters a budget, the budget enters the capacity N, and the capacity fixes the dimensionless cosmological term through three pi divided by N. P is a resolution and N a capacity.

A loop of that shape is chapter two's equation with the world inside it. The unknown stands on both sides, and each version of the loop has one solution where a theory with a stage has a dial.

Self-reference turned up in chapter two to answer why there is anything at all, and it turns up again in chapter thirty-seven to take the beginning away. Here each declared return map has a certified solution. Showing that the observer construction selects the physical map and its normalization is the further step needed to turn that solution into a prediction about our world.

The structured trial map returns 137.035660 against a laboratory reading of 137.035999. The missing physical matching has not been shown to close that difference. The capacity lands four parts in a thousand from the sky's figure on one booking of the reserve and six on the other.

Feynman's went up on the wall because nothing inside the theory could reach it. The difference here is that the theory has an inside, and the inside is what reaches it.

Chapter 30

Why Do the Masses Land Where They Do?

Take the masses of the three charged leptons, the electron, the muon and the tau. Add them up. Then take the square root of each of the three masses, add the three square roots together, and square that total. Divide the first number by the second.

The answer is two thirds.

There is no evident reason for these three masses to produce a number like that. The muon is two hundred and seven times the electron and the tau is three and a half thousand times the electron, and a quantity assembled out of three separately measured masses would be expected to come out as an unremarkable decimal.

Nobody computes that combination by accident, and the first person with a reason to was Yoshio Koide. He was an associate professor at Shizuoka Women's University at the end of 1981, working on composite models of quarks and leptons. Models of that kind put something smaller underneath the particles on the roster, so that the shape of the roster becomes a consequence of how few pieces there are to build from. The field was busy. In 1979 Haim Harari and Michael Shupe published the same proposal four pages apart in the same issue of *Physics Letters B*, without either knowing what the other was doing: every quark and every lepton built from triples of two objects, one carrying a third of the electron's charge and one carrying none. Harari called them rishons, and none has ever been observed. One line of arithmetic came out of Koide's model and stayed.

Two of the three masses were known to many digits by 1981. The tau was the trouble. It had been found at Stanford in the mid-seventies, it dies in less than a picosecond, and its mass had to be reconstructed from what came out of the decay rather than weighed. The world average sat at 1784.2 million electron-volts, give or take 3.2. Koide's relation, given the electron and the muon and asked for the third, returned 1776.97.

The gap is 7.2 million electron-volts, and physics reports a gap in units of the experiment's own quoted uncertainty. A measurement is a claim about a range, which is how chapter four read two different Higgs announcements as one result. The half-width of that range has a symbol, the Greek letter sigma, and a distance between two numbers gets quoted as a multiple of

it. Koide's tau mass sat 2.25 sigma from the measured one. Gaps that size open and close as the measurements improve. The word discovery is reserved for five, which is the threshold ATLAS and CMS cleared in July 2012.

The unit in that comparison belongs to somebody. Sigma is whatever the experiment declared its own uncertainty to be, so a distance in sigma depends on one group's uncertainty estimate in one year, and such estimates get revised.

In the spring of 1992 a group running the Beijing Spectrometer, a detector on China's electron-positron collider, went at the tau mass from a different direction. Below a certain beam energy you cannot make a pair of taus at all and the production rate is zero. Above it the rate climbs. The corner between the two is the mass, twice over. They set the beam using the standing world average of 1784, took twelve measurements across the threshold, and found the corner at 1776.9 million electron-volts, 7.2 below where the average had put it, with an uncertainty smaller by a factor of seven.

Koide's 1776.97 had not moved. The world average had walked 7.2 million electron-volts and arrived on it.

A triangle supplies a conditional description of that near equality. The balance condition and the identification of its response eigenvalues with mass amplitudes still need a physical derivation.

30.1 What a mass calculation must fix

An internal charge list is a list of possible transformation laws. It becomes a mass calculation only after the same observer history supplies a kinetic term, a state, the mass-generating interactions and a clock that reads energy. The finite source admits the three charged-fermion interaction channels with one Higgs doublet. It does not select their coefficient matrices or the Higgs vacuum.

The neutrino sector admits an optional effective interaction with two Higgs factors, but its coefficient and scale are additional inputs. None of the observed nonzero masses is derived by those algebraic possibilities alone. The relation below has a different scope: under its balance and mass-ordering assumptions, measured electron and muon masses constrain the tau. That conditional relation is not an absolute prediction of all three masses.

30.2 Three corners, two numbers

Chapter twenty-seven described a candidate rank-three family band. The comparison here attaches the electron, the muon and the tau to the three

corners of an oriented face of the icosahedron. That attachment is a premise, not a consequence of counting corners.

An oriented face has one rotation left in it. Rotate the corners round the face: one goes to two, two goes to three, three goes to one. Do that three times and everything is back where it started. Of the sixty rotations of the icosahedron, that turn, its repeat and leaving the solid where it stands are the only three that keep both the face and its sense of circulation.

Take three identical rooms arranged in a ring, each joined to its two neighbors by identical doorways, and put a heater in one of them. Wait, then measure the temperature in all three. Doing that for each room in turn gives you nine numbers: how warm room one gets when room one is heated, how warm room two gets when room one is heated, and so on across and down. Nine numbers, and seven of them are redundant, because the ring has no room that differs from any other room. There is a number for how much heat stays where it was put and a number for how much crosses a doorway. The whole table is those two numbers arranged in a pattern.

The face works the same way. Push at one corner and read what comes back at all three, which is again a three-by-three table.

A response that respects the rotation cannot tell corner one from corner two, because the rotation carries the first onto the second. So the second row of the table is the first row shifted one place along, and the third row is the second shifted again. Nine numbers collapse into two: one for a corner's response to itself, written a, and one for its response to a neighbor. The neighbor entry carries a size and an angle, because a response can arrive turned as well as scaled, and chapter nineteen's amplitudes are that kind of number. Call the size b and call the angle the twist.

A table whose every row is the row above it shifted one step round the cycle is called a **circulant**.

Chapter fifteen took a twelve-by-twelve grid of repair weights and hunted for the grid's own directions, the ones it merely scales without swinging them round, and called the scale factors eigenvalues. Finding those for a general grid means solving an equation whose degree is the width of the grid. A circulant gives them directly, because its own directions are fixed by the cycle before anybody says what a and b are: they are the three ways of stepping a phase round the triangle in equal parts.

The three scale factors come out as a, plus twice the size b, multiplied by a cosine. The three cosines are the cosine of the twist, the cosine of the twist plus a hundred and twenty degrees, and the cosine of the twist plus two hundred and forty.

Three cosines, a hundred and twenty degrees apart. The face has fixed their spacing and left the starting angle free.

30.3 Two lines about cosines

Draw three arrows of length one from a single point, a hundred and twenty degrees apart. They are the spokes of an equilateral triangle drawn from its center. They add up to nothing, because there is no direction for the sum to point. The cosine of an angle is the shadow one of those arrows casts on a chosen axis, and the shadow of a sum is the sum of the shadows.

Three cosines a hundred and twenty degrees apart sum to zero.

For the squares, use the fact that the square of a cosine is half of one plus half the cosine of the doubled angle. Double the three angles and you get the twist doubled, that plus two hundred and forty degrees, and that plus four hundred and eighty. Four hundred and eighty degrees is a hundred and twenty with a full turn discarded, so the doubled angles are three directions a hundred and twenty degrees apart, and by the line above their cosines sum to zero. Three halves of one, plus half of nothing.

Three cosines a hundred and twenty degrees apart have squares summing to three halves.

Add the three eigenvalues. The three copies of a give three a. The cosine terms cancel by the first line. Total three a, whatever the twist happens to be.

Add their squares. Each square is a squared, plus twice a times twice b times a cosine, plus four b squared times the square of a cosine. The middle terms cancel by the first line. The last terms come to four b squared times three halves, which is six b squared. Total three a squared plus six b squared.

What a corner carries is an amplitude, and chapter nineteen's rule for turning an amplitude into a measurable quantity is to square it. In the declared mass reading, each eigenvalue is the square root of its corner's mass, up to one overall scale that fixes the units and cancels out of any ratio. The square root of a mass is a positive number, so the only responses describing anything that exists are the ones whose three scale factors all come out positive.

Koide's ratio, written Q, is the sum of the eigenvalues' squares divided by the square of their sum. The two lines above have evaluated both of those.

$$Q = \frac{m_e + m_\mu + m_\tau}{\left(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau}\right)^2} = \frac{1}{3} + \frac{2}{3}\left(\frac{b}{a}\right)^2$$

On the left are the three charged lepton masses, measured in laboratories. On the right is one number, the size of a corner's response to its neighbor divided by the size of its response to itself. Everything else has cancelled, the twist included, which appears nowhere on the right and therefore cannot be tuned to help.

Q equals two thirds exactly when b over a is one over the square root of two.

The left side of that equation is trapped, for any three positive masses. Set all three equal and it gives one third, which the right side agrees with by sending b to zero. Let one mass run away from the other two and it climbs toward one, without arriving. So every possible triple of masses lands somewhere in the range from one third to one. Two thirds is the exact midpoint of that range. The three charged leptons, whose masses cover a factor of three and a half thousand, sit on the midpoint.

Two pieces make up the response. There is the direction in which all three corners move together, which carries weight three a squared, and there is the two-dimensional plane in which they move against one another, which carries weight six b squared. Setting b over a to one over the square root of two sets those two weights equal. One condition on the response of a single face, and Koide's two thirds is that condition written out in masses.

The twist gets a small allowance rather than a value. At balance, the three eigenvalues stay positive as long as the twist is within fifteen degrees of one of the three spokes, and anywhere inside that band the ratio sits at two thirds without moving.

Take the three square roots of the masses as the three coordinates of a point in space. There is a line through that space along which all three coordinates are equal, the line where the three leptons would weigh the same. The balance says the point stands at exactly forty-five degrees from that line, and forty-five degrees is why the relation looks so peculiar written in masses. A clean statement about square roots turns into an agreement to five decimal places once everything in it has been squared.

30.4 Seventy-two electron-volts

Feed in the two masses known best. The electron is 0.51099895 million electron-volts and the muon is 105.6583755 million electron-volts. Requiring the balance to hold turns the relation into a quadratic in the square root of the third mass. A quadratic has two roots. One of them is 3.3174 million electron-volts, which would put the third charged lepton between the electron and the muon rather than above both of them. The tau is the heaviest of the three, which is why it can decay into a muon and a muon cannot decay into it, so the upper root is the one.

Chapter twenty-nine turned a computation into a proof by carrying intervals rather than points. The same machinery applies here: give the electron and the muon their measured widths, do the arithmetic on the widths, and what comes back is an interval guaranteed to contain the answer.

It runs from 1776.968991 to 1776.969063 million electron-volts, centered on 1776.969027.

That window is 72 electron-volts wide. The measured tau mass is 1776.93 million electron-volts, give or take 0.09, and 0.09 million electron-volts is ninety thousand electron-volts, so the uncertainty on the measurement is twelve hundred and fifty times the width of the window it is being compared against. The center of the window is 0.039 million electron-volts from the measured value, which is 0.43 sigma.

Fed the measured tau mass, Koide's ratio comes out at 0.66666446.

The sharpest tau measurements are threshold scans of the kind the Beijing Spectrometer ran, reaching ninety thousand electron-volts. What limits them is how precisely a beam energy can be calibrated at the point where a pair of taus becomes possible, and that is where the next three decimal places are.

A measured central value more than three standard uncertainties from 1776.969027 destroys the balance condition. There is no free parameter left inside the balance to absorb the move: the twist cancelled, the scale cancelled, and the one remaining quantity was set to one over the square root of two by the requirement that the two pieces of the response weigh the same. At an uncertainty of 0.09 million electron-volts, three of them is 0.27, so a central value below 1776.70 or above 1777.24 ends it.

30.5 Thirty-one axes

In June 1963 Nicola Cabibbo published three pages in *Physical Review Letters* under the title "Unitary Symmetry and Leptonic Decays". Decays that changed strangeness were running slower than decays that did not, by a consistent factor, and the theory of the day had no room for a consistent factor. Cabibbo's move was to let the weak interaction couple to a rotated combination of the down quark and the strange quark, turned by a single angle. One number, fitted from decay rates, and the discrepancy went away.

The strength of the transition from a down quark to a strange one is measured at 0.2250. The angle whose sine that is comes to 13.0029 degrees. Rates go as the square of that strength, so a decay that changes strangeness runs at about a twentieth of the rate of one that does not. That factor of twenty is the suppression sitting in the data he was accounting for.

An icosahedron has axes of three kinds. Six of them run through pairs of opposite corners. A fifth of a turn about one of those leaves the solid looking untouched. Ten run through pairs of opposite faces, at a third of a turn. Fifteen run through pairs of opposite edge midpoints, at half a turn. Six and ten and fifteen is thirty-one.

Take the angle between every pair of those axes. Four hundred and sixty-five pairs, one angle each, and the smallest nonzero angle anywhere in the list is 20.9052 degrees.

13.0029 is not in the list. The reason it cannot be got at is that 20.9052 is the floor. The smallest angle the geometry has to offer is sixty percent larger than the one the quarks require, and relabeling which axis is which does nothing, because the floor is a property of the solid rather than of the labeling. The angle that sets quark mixing is not an axis angle of this geometry.

The list has an end, which is what makes an empty search a result: four hundred and sixty-five angles, written out in full and checked to the last one.

Koide's own model reached past the leptons to the quarks. The quark side did not hold. Two structures built out of entirely different material, a set of subparticles and a twenty-sided solid, keep the charged leptons and stop at the quarks.

The declared spin lift and charge normalization give the familiar internal charges and three color slots. A minimum-cost response rule offers a rank-three family band, whose physical identification remains open. On an attached oriented face, equal weight for the two response sectors gives Koide's two thirds; deriving that equal weight and its mass readout from repair remains work in progress. Handed the electron and the muon and that balance, it pins the tau inside 72 electron-volts and prints the value at which the balance dies. It does not hand over the angle by which the weak interaction turns a down quark into a strange quark: 20.9052 against 13.0029 proves that the angle is no axis angle of this solid. Had 13.0029 turned up anywhere on that list, the charges and the family count would have been worth less, because a solid that yields whatever angle is asked of it has yielded nothing.

The settled entries and the untouched ones end up on the same line. Working physicists carry it in four terms, holding every charge, every family count, every coupling and every mass on the roster, the forced ones and the typed-in ones side by side, with nothing marking which is which. Nobody has ever derived that line.

Chapter 31

What Is a Lagrangian, and Why Is It the Last Thing?

The line the last chapter ended on has no pictures in it. It is four terms, each a whole subject folded flat, short enough to print on a coffee mug. Writing physics as symbols with nothing to look at is older than the electron by more than a century, and it started with a boast.

In 1788 the widow Desaint published a book on mechanics in Paris that contained no pictures, and its author announced that on the first page. Joseph-Louis Lagrange had come down from Berlin the year before. The Académie sent the manuscript to a committee of Laplace, Cousin, Legendre and Condorcet, who approved it, and Legendre read the proofs.

No diagrams will be found in this work, the preface opens. The methods set out in it, Lagrange goes on, call for neither constructions nor geometrical or mechanical reasoning, but only algebraic operations subject to a regular and uniform procedure.

Mechanics until then was drawn. Newton's *Principia* argues in figures: an arc, a tangent, a shaded triangle, a limit taken as two points slide together, and the reader follows by looking. For a century after 1687 doing mechanics meant getting the picture right first, and the pictures were different for a pendulum, a spinning top and a planet, so each of them was a separate accomplishment. What Lagrange had found was that somebody who cannot draw at all obtains every one of those results by writing down a single function of the system and then turning a handle, and that the handle turns the same way in all three cases.

The line on the mug is an object of the same kind. Two questions come with it. What sort of thing is it, that a whole subject folds into one line at all? And what does the folding cost, since a book of mechanics with the figures taken out has thrown something away, whatever else it has gained.

31.1 Proof of a supreme being

On 15 April 1744, before the Académie Royale des Sciences in Paris, Pierre-Louis Moreau de Maupertuis read a paper called "Accord de différentes lois de la nature qui avaient jusqu'ici paru incompatibles", agreement between different laws of nature which had until then seemed incompatible. The laws were the ones for how light bends and how it bounces. He proposed that a single quantity, which he named the action, comes out as small as it can be whenever anything in nature changes.

Two years later in Berlin, running Frederick's academy, he published "Les Loix du mouvement et du repos déduites d'un principe métaphysique". There the principle acquired a second job. Maupertuis presented it as a demonstration of a supreme being, and ranked it above the older proofs drawn from the beauty and the order of the world, on the ground that a mathematical principle was firmer evidence.

The inference is the obvious one: a rule that picks the smallest total out of every total available looks like an economy, and an economy suggests somebody managing it.

Maupertuis had earned the standing to say such things by going out and getting a number. In 1736 he led an expedition to the Torne valley in Swedish Lapland to measure the length of a degree of the meridian close to the Arctic Circle. The party came home in August 1737 with a degree longer than the one measured in France, which meant the Earth is flattened at the poles, which meant Newton was right and the Cassinis were wrong. Robert Levrac-Tournières painted him afterward in a fur hat and reindeer skins with one hand pressed down on a globe, squashing it out of round. Voltaire, an admirer at that stage, addressed him as the flattener of the world and of Cassini.

The rest went badly. In 1751 Samuel König claimed that Leibniz had had the least-action principle first and produced a letter to prove it, and the Berlin academy, with Maupertuis presiding, declared the letter a forgery. Voltaire took König's side and wrote the *Diatribe du docteur Akakia, médecin du Pape*, in which a papal physician works through the published proposals of a self-styled academy president, most of them quoted from what Maupertuis had actually printed. Frederick had the pamphlet burned by the public executioner on Christmas Eve 1752, with the king and Voltaire both standing at the fire. Maupertuis left Berlin and died in Basel in 1759.

His principle survived him exactly as he stated it, and so did his reading of it. A rule that scores a whole path and then selects one looks as though the path was compared against paths that never happened.

31.2 A ball, described twice

Throw a ball across a room. It leaves your hand at one place and arrives at another a second later.

Here is the ordinary description. At each moment the ball has a position and a velocity. Gravity pulls down, which changes the velocity a little, which changes the position a little, and repeating that as finely as you like traces out the arc. The description looks forward one step at a time and needs nothing except what is true where the ball is.

Here is the other one. Draw every arc you like between those two points taking the same second: the true one, one that goes twice as high, one that moves slowly at first and fast at the end. Score each of them with a single number. The arc the ball flies is picked out by what its score does when you nudge it.

The scoring rule has two parts. The future appears in neither.

The first part is one function. At any instant, two things about the ball matter: where it is, and how fast it is going. Feed both into a single function and it hands back a number. For a thrown ball, that function is the part that grows with speed less the part that grows with height.

Both parts are countable for a ball you could actually throw. The height part is the **potential energy**, what the ball's position owes: the mass times the strength of gravity times the height. Gravity at the Earth's surface pulls at 9.81 meters per second per second, and rounding that to ten costs two percent and keeps the arithmetic simple, so a one-kilogram ball held a meter off the floor owes ten joules. The speed part is the **kinetic energy**, which is what the motion holds: half the mass times the square of the speed. That same ball moving at two meters per second holds two joules, and at four meters per second it holds eight, because the square is where the growth is. The function is the second of those less the first, joule for joule.

The second part is a running total. The number the function hands back is per second, so run along the arc from the first moment to the last and keep a running total of it, the way a taxi meter accumulates while the wheels turn. One arc in, one number out. Every arc on the list gets a total this way.

Selection uses slopes. Stand on a hillside and step one pace in any direction: your height changes by an amount proportional to the size of the step. Stand on the floor of the valley and step one pace: your height barely changes, because the ground under you is level, and the change you do get shrinks with the square of the step rather than with the step. The bottom of a valley is where the slope is zero.

The arc a thrown ball flies is where the slope is zero. Push the middle of it up by a hair, holding both ends where they are, and the running total does not change. Push it down by a hair and the running total does not

change. Every neighboring arc scores the same to first order. The score is flat there in the way the valley floor is flat.

Take the crudest version of that and check it by hand. Three moments and two hops: a one-kilogram ball starts on the floor, ends on the floor one second later, and the only freedom is how high it is at the halfway mark. Raising the midpoint costs in the speed part, because the ball has to climb further and fall further in the same second. It pays in the height part, because the ball spends the middle of the flight higher up. Each hop lasts half a second and there are two of them, so anything holding steady through both contributes its own value times one second to the running total, and the halves cancel out of the arithmetic before it starts.

A peak of one meter. The ball climbs a meter in half a second, so it travels at two meters per second and holds two joules of motion, and its height averages half a meter across each hop, which owes five joules. Two less five is minus three.

A peak of one and a half meters. Three meters per second, four and a half joules of motion, average height three quarters of a meter, seven and a half joules owed. Minus three again.

A peak of one and a quarter. Two and a half meters per second, three and an eighth joules of motion, average height five eighths of a meter, six and a quarter joules owed. Minus three and an eighth.

The arc that overshoots and the arc that undershoots score the same number. The one between them scores below both, by an eighth of a joule-second in either direction. A single line says why, with h standing for the height of the halfway mark in meters and the total coming out in joule-seconds:

$$\text{score} \;=\; 2h^2 \;-\; 5h$$

The cost grows with the square of the midpoint height and the payment grows in proportion to it, so the cost overtakes. At one midpoint height and no other, raising the peak by a hair costs exactly what it pays. There the total has slope zero. That is the height the rule selects, one and a quarter meters, where a ball airborne for a second peaks: it leaves the floor at five meters per second, sheds ten of those every second, and stops climbing after half of one. Three moments is the coarsest account of a flight with any freedom left in it, and lands on the height exactly. Cut the second into four hops instead of two and the corners of the selected shape land on a parabola, and into forty and they land on the same one.

Look at what had to be in your hand to do any of that. All three sums used the landing. Fixing the speed on each hop took the ball being back on the floor at one second, and without the finish there was nothing to add up. The ball has the start. It leaves your hand with a place and a speed and no information whatever about where it comes down, and the rule says it flies

the arc whose total sits at the bottom of a parabola whose shape depends on where it comes down. Something in that description knows the ending before it happens, and the ball does not. Maupertuis put a supreme being in the gap.

The one function of where you are and how fast you are going is the **Lagrangian**. Its running total along a path is the **action**, which is the word chapter twelve used for a single number scored over a whole arrangement. This is the same object scored along a history instead of across a wiring. The rule that selects the path where wiggling changes nothing is **stationary action**.

Everything about a system is in its Lagrangian. Hand somebody that function and every quantity the system has gets its equation of motion, by the same handle-turning in each case: take the function's sensitivity to how fast the quantity is changing, ask how that sensitivity changes as the run proceeds, and set it equal to the function's sensitivity to the quantity itself. One line goes in and the entire behavior comes out. **A Lagrangian is a compression**: the thing you keep when you throw away the histories, chosen so that the histories can be regenerated from it.

Watch it come apart once. The Lagrangian of a thrown ball has two pieces, the speed piece and the height piece. Its sensitivity to how fast the ball is going comes from the speed piece and is the mass times the velocity, which is the ball's momentum. Its sensitivity to where the ball is comes from the height piece and is the mass times the strength of gravity, pointing down. Set the rate of change of the first equal to the second and out drops the statement that the ball's momentum gains mass times gravity downward every second, which was Newton's second law and which nobody wrote into the function. Two pieces of arithmetic and the handle, and the same handle applied to a spinning top gives its equations too.

31.3 Göttingen, 1918

The second thing a Lagrangian compresses is the reason anything is conserved.

Emmy Noether arrived at Göttingen in the spring of 1915 at the invitation of Hilbert and Klein, and worked there for years with no salary and no post. The faculty blocked her habilitation. Her lecture courses appeared in the catalogue under Hilbert's name, with herself listed as the person who would actually be giving them. When she had the result that carries her name she could not read it to the Göttingen Scientific Society, not being a member of it; Klein read it out for her on 26 July 1918. It went into the society's proceedings under the title "Invariante Variationsprobleme", pages 235 to 257.

In April 1933 the Law for the Restoration of the Professional Civil Service removed her from the university. She went to Bryn Mawr College in

Pennsylvania, and once a week to Princeton to lecture. In the spring of 1935 she had surgery for a tumor, the operation went well, and four days afterward she developed a fever and lost consciousness. She died on 14 April 1935, at fifty-three.

Chapter five gave the correspondence and chapter eighteen used it: every continuous symmetry of the laws comes with exactly one conserved quantity, determined by the symmetry, and momentum and energy are what you get from the laws not caring where and not caring when. The machine that turns the one into the other is a recipe with three steps.

Take the Lagrangian. Find a change you can make to it continuously, by an amount you can dial down to nothing, which leaves its value alone: slide the whole system a hair sideways, turn it by a small angle, advance every clock by the same interval, add the same small amount to a phase everywhere. Each of those is a direction of change with a size attached to it.

Then, for every quantity the system has, take the Lagrangian's sensitivity to how fast that quantity is changing. That sensitivity is the momentum belonging to the quantity, and falls out of the function rather than being defined off to one side. Multiply each sensitivity by how far the symmetry moves its own quantity, and add the products over the whole list. Advancing the clock takes one more move, since shifting the clock shifts the running total too: subtract the function's own value times the interval.

The number you get does not change along any history the Lagrangian selects. One symmetry in, one number out, by a recipe that does not care which symmetry it was fed.

Slide sideways and the number is momentum. Turn and it is angular momentum. Advance the clocks and it is energy. Add the same phase everywhere and it is electric charge, arriving by a second road. Chapter twelve reached it from the other end, by showing that a force whose charge leaks stops being a force at all. The recipe here starts from the same phase and hands back the quantity you can total up.

The same statement holds on a discrete chain with no limit taken anywhere. A run is a list of states with one step between each neighboring pair, and a symmetry is a change that carries each state to another while leaving each step's contribution to the running total alone. Take the contribution's sensitivity to the change across one step and it comes out the same across every step of the run. Segment by segment, the number holds, and there is no continuum anywhere in the argument.

Bookkeeping in this world was never a separate law added on top of the dynamics. It is a property of the line, extracted by a fixed recipe. A world has as many conservation laws as its line has continuous symmetries.

31.4 A product of small numbers

None of that touches Maupertuis's reading. The recipe is exact. A rule that scores whole paths and selects the flat one continues to look like a comparison, which seems to require the rejected paths to exist somewhere.

Start again from the smallest assumption a world can be given.

Something is in some configuration. There is a rule saying what it does next. The rule has three properties. It is local, so the next step depends only on the configuration it is standing in. It is memoryless, so the rule does not consult how that configuration was arrived at. And it is strictly positive, so nothing is flatly forbidden: every configuration it could move to has some chance, however small.

That is the whole of the input. There is no action in it, no Lagrangian, no scoring, and nothing that looks at a history.

Ask about histories anyway, since a history is a thing that has happened and can be asked about. Take a list of configurations, one after another, from a start to a finish. What are the odds of that particular list? Each step carries odds of its own and the steps do not consult each other, so the odds of the list are the odds of the first step times the odds of the second times the odds of the third, all the way to the end.

A product of a thousand numbers each below one is unmanageably small. Chapter sixteen introduced the operation that fixes it: a logarithm converts multiplying into adding. Take the logarithm of each step's odds, flip the sign so the contributions come out positive, and add them along the history instead of multiplying. Written out, with the odds of the first step called p one, the odds of the second p two, and the history running to n steps:

$$-\log\left(p_1\, p_2 \cdots p_n\right) \;=\; \left(-\log p_1\right) \;+\; \left(-\log p_2\right) \;+\; \cdots \;+\; \left(-\log p_n\right)$$

On the left is one number belonging to a whole history. Check it on the smallest case: two steps, each a coin flip, so the history has odds of one in four, which is two of chapter sixteen's yes-or-no questions, one from each step. On the right are n numbers, each belonging to a single step and worked out from the two configurations that step ran between, none of them consulting any other step, and the equals sign in the middle is the line of algebra that takes the foresight out.

Look at what that produces. It is a running total along a path, one contribution per step, and each contribution depends on where its step started and where it ended and on nothing else anywhere in the history. That is the shape of the thing the thrown ball was scored with, arrived at from a rule that contains no score.

A history with a large total took improbable steps. A history with the smallest total is the most probable history there is. And a history

whose total does not move when you jiggle one of its interior configurations, holding the ends, is sitting on the valley floor in that direction, which is what it means to be locally most likely. **Least action and maximum probability are two readouts of one thing.**

Run the total backwards. Give each history a weight that shrinks as its total grows, one over the exponential of the total, and out comes the odds of that history again, because exponentiating undoes the logarithm that built the total. Nothing has to be divided by anything to bring the weights to one across all the histories, since the odds added to one before any logarithm was taken.

Then suppose somebody hands you a different running total, and a different overall scale to multiply it by, and their combination also reproduces the same odds. Their scale times their total equals the total above plus a constant, at every history, and there is no other freedom anywhere in the problem. You may slide the zero, and you may change the units. Everything else is forced.

A world whose step rule is local, memoryless and strictly positive has an action, then, whether or not anybody writes one down.

Nothing in that argument looks ahead, because the step rule has nothing to look ahead with. The endpoint turns up in the statement of the principle because the statement is about a completed history. A completed history has a finish in it the same way a receipt has a total printed at the bottom. A machine repairing locally, with no view past its own neighbors, produces a selection that looks global, because that is what taking the logarithm of a product of local factors does.

Maupertuis had found something real. The quantity a world comes out smallest on is improbability. A chain of local steps runs that tally by itself, without a manager and without a comparison ever being performed.

31.5 Plateau's wire frames

Joseph Plateau lost the sight of both eyes by the end of 1843, at forty-two. Ghent promoted him to full professor a few months later, and for the next four decades his wife Fanny, and afterwards his son and son-in-law, did the seeing for him. They dipped wire frames of every shape into soapy water mixed with glycerine, drew them out, and described aloud exactly what surface had formed. In 1873 he published nearly a thousand pages on the statics of liquids held together by molecular forces alone, every word of it dictated. Jean Taylor proved his rules about how the films meet each other in 1976, a hundred and three years afterward, as consequences of minimizing area.

Dip a bent wire in soap solution and the film stretched across it is the surface of least area that wire will hold. It looks like the answer to a question about all possible surfaces. The film has no way of finding out

what any other surface would have cost. Each patch of it pulls on the patches around it, the pulling is settled locally, patch against neighbor, and when no patch is being pulled harder one way than another the film stops moving. Where it stops has the least area available. Nothing in the film has ever evaluated an area.

The settling is also too fast to see. By the time the frame is steady in your hand the film has finished. What Plateau's family described to him across forty years was the finished surface, every time.

31.6 The interaction coefficients

One common local metric can carry the scalar, spinor and gauge terms. Exact invariant tensors connect the conditional matter list to one Higgs doublet and its conjugate, giving the up, down and charged-lepton interactions. Gauge and Lorentz invariance constrain their form. For three declared families, they permit three complex coefficient matrices before changes of flavor basis.

The line below is a conditional effective action with those coefficients and its vacuum specified. Deriving the coefficients, the chiral field interpretation and the geometric response from repair is distinct from verifying that the line is consistent. The fixed product background illustrates the difference: it can describe waves, while a gravitating excitation requires a geometry that responds to their stress.

31.7 The line

Physics as it is normally taught reaches a Lagrangian in the first week and spends the rest of the time taking it apart. The spacetime is there before the pen touches the page, the group is written down because the detectors reported it, the fields are listed because they were found, and the numbers are entered because laboratories measured them.

The line has four terms in it, written schematically, each standing for a block that opens out further.

$$\mathcal{L} \;=\; \mathcal{L}_{\text{carry}} \;+\; \mathcal{L}_{\text{matter}} \;+\; \mathcal{L}_{\text{couple}} \;+\; \mathcal{L}_{\text{select}}$$

Four things added together. The chapters behind this one give finite constructions and explicit premises for interpreting the symbols.

The script L on the left is a Lagrangian in the sense taught above, with one change of scale. Where the thrown ball had a position and a speed, this one takes a field's value at a place and how fast that value is changing, and its running total runs over all of space and all of time rather than along a single arc.

The first term is the carriers. Chapter twenty-six split the twelve dials of the wiring into a central one, a block of three and a block of eight, with

nothing else arithmetically available. This term holds the energy stored in what those dials do around a loop. Chapter twelve's loop readings are the field strength, and squaring them and adding over the arrangement is the term.

The second term is matter, carrying chapter twenty-seven's conditional fifteen-state representation, repeated over three declared families. Inside it sits a comparison between a reading here and a reading next door, taken after the reading next door has been run through the dictionary sitting on the seam between them, which is chapter twelve's connection written with derivatives instead of exchange rates. That comparison is the entire content of the phrase "the force acts on matter". Relabel the readings differently from one place to the next and the whole expression keeps its value, because the derivatives of the relabeling cancel.

The third term holds the couplings: the invariant channels through which the scalar doublet touches matter. Their coefficient matrices remain inputs. Chapter thirty's separate balanced-face comparison constrains the tau using measured electron and muon masses; it does not source-select those matrices.

The fourth term belongs to the scalar field itself. Its Lorentz spin is zero, while its weak-doublet components retain their internal indices. A vacuum magnitude of 246.22 billion electron-volts is a laboratory calibration in this effective action, not a repair-selected output.

One charged scalar component of the line reduces consistently to a classical system, with both diagonal gauge fields aligned by their kinetic weights. A local execution on 64 spatial sites records a phase disturbance, the current it drives and the electric feedback on it. A gauge transformation of the same starting state leaves every invariant reading unchanged. Bounds on the exact continuous evolution also resolve the current and a response at a neighboring detector, and at the final observation times the error on the intensity is smaller than the response itself.

Alongside the four sits a conditional gravitational term. Chapter twenty-five relates its coupling to a screen area under the entropy and continuum hypotheses; the laboratory value of Newton's constant calibrates that area in square meters. The generalized entropy stationarity and geometry response needed for a physical coupled model remain to be established. Null contractions leave a metric-proportional constant undetermined.

The prologue promised that the symbols would have an account. Some accounts are finite constructions, some are conditional theorem interfaces, and some still require measured inputs. The record geometry, internal response algebra and invariant interactions have explicit witnesses. Passing from them to physical chiral fields, a global gauge lift and a responsive geometry retains the named premises. The Higgs vacuum, family matrices and observed masses are not selected by writing their terms on this line.

The candidate family band has three directions under the stated minimum-cost rule. An oriented face offers a three-slot attachment. Neither fact alone establishes three physical families or their distinct masses. The line makes those remaining choices visible.

31.8 What a compression costs

A compression is allowed to be lossy, and the useful ones generally are.

This one presumes a continuum. The running total runs over space and time, the fields are smooth, and the handle-turning that produces the equations of motion needs slopes to exist at every point of every path. Chapter fifteen was careful about that limit for a reason that bites here: an observer holds six whole numbers and sits at some finite number of passes, which is a scatter of isolated points with gaps between them that never close at any stage. Smoothness stands to that scatter as pi stands to 3.1416. The line is written where the tally is heading. No observer's tally has ever been there.

The approach to that limit is under control. On chapter fifteen's record points, one positive nearest-neighbor action gives classical waves and, quantized in the usual way, the full state space of a quantum field, with any number of quanta in each mode. Compare it with a massive scalar field in a cube with fixed boundary values. Refine the points while shrinking each local time step faster than the smallest gap between coordinates, and the smeared classical and coherent quantum readouts converge to those of the smooth field.

The same calculation resolves a compact disturbance at a detector confined to a separate slab. At 233 source coordinates per axis and a time step of 1/131072, reading the nearest update inside the window keeps the total probability error below 0.050078 and the induced signal above 0.137746 throughout the stretch of time from 0.95 to 1. The error includes spatial localization, every spectral mode, time stepping and the change in vacuum fluctuations caused by discretizing time. Tensor factorization keeps the calculation small.

The second loss is sharper, and can be written down exactly.

Take the line and add a term chosen so that it comes out zero at every configuration the world realizes. On a chain, the realized configurations sit at the corners. A term that vanishes at every corner adds nothing to the score of any realized history. So every realized history keeps its old score. Swap an interior configuration for another the world can realize, holding the ends, and the answer comes back what it was: the same run is the most probable run, and sits at the bottom against every single-step swap.

The momentum changes. Momentum is the line's sensitivity to how fast a quantity is changing. A term pinned to zero at every corner tilts as it passes through, in a direction no realized step ever moves. So the two

lines assign different momenta to the same motion, and therefore different energies, since energy is what you get by trading the speeds for the momenta.

The freedom is one number. The added term bends with speed by any positive amount you like, and its bend with position is then forced, the two summing to one particular number the step odds themselves set. Under that constraint the most probable run stays at the bottom when an interior configuration slides continuously rather than jumping corner to corner, and the momentum stays a strictly increasing, invertible function of speed. Each setting is a different line carrying a different energy. Nothing in any history of this chain distinguishes them from each other. The histories are the same histories, and they were the only evidence there was.

So the line is the best compression anybody has written down. A compression is not the thing compressed. Lagrange knew what he had taken out, which is why he mentioned it in the first sentence.

Hand the line to somebody with a computer and enough patience and out comes the arc of a thrown ball to any precision wanted, the spectrum of hydrogen, the lifetime of a muon and the rate at which the sun burns. Then put the ball in their hand and push it, and watch it go. The line scores the path it takes. It says nothing whatever about why a pushed thing moves.

Chapter 32

Why Does Anything Move?

Benjamin Robins hung a broad iron plate from a frame in 1742, screwed a thick sheet of wood across the front of it, and fired a musket into the wood from twenty-five feet away.

The plate swung back. A narrow ribbon ran from the plate through a clamp gripped just tightly enough to hold, so the swing dragged a length of ribbon out through the slot and left it lying there. Robins picked up the ribbon and measured it. Its length is the chord of the arc the plate traveled, the chord gives the speed the plate left with, and the speed the plate left with gives the speed of a ball nobody had ever seen move: sixteen hundred and seventy feet per second, for a three-quarter-inch ball out of a forty-five-inch barrel. The same ball measured at seventy-five feet from the muzzle came in at fifteen hundred and fifty, and at a hundred and twenty-five feet at fourteen hundred and twenty-five. The difference was lost to air resistance.

New Principles of Gunnery came out the same year and turned artillery into a subject with mathematics in it. Euler translated it into German in 1745 and added notes until the translation ran to roughly five times the length of the book it was translating.

What Robins built is an accounting device. The ball goes into the wood and stops being a ball. Whatever it was carrying does not stop: the plate leaves with all of it, and a length of ribbon on the floor is a measurement of a quantity nobody in the room can see, trusted because the books balance.

Three things have to hold for that ribbon to mean anything. The quantity has to survive a collision that destroys the shape of the ball, the shape of the wood and most of the geometry of the problem. It has to arrive at the plate in full, with no fraction of it lost in the splintering. And it has to be the same quantity in a ball made of lead as in a plate made of iron, since the two are being added together. Robins had all three from firing the gun and derived none of them. The ribbon measures a quantity whose entire definition is that it comes out even.

Twenty-nine chapters have built a world and none of them has said why a thing that gets pushed goes. Four words carry every account of it, and a reader owns all four before opening any book: momentum, inertia, force,

acceleration. Robins' ribbon measures the first of them, which a collision hands over whether or not anybody has defined it.

32.1 A chain of readings

Take a history of the kind chapter thirty-one scored: a list of readings, one per event, from a start to a finish. Chapter thirty-one's rule walks that list and adds one contribution per step. Each contribution depends on the two readings at the ends of its own step and on nothing else anywhere in the history.

For a pattern crossing the seam network with nothing pulling at it, the contribution is half the square of the change across the step, weighted by what the pattern costs to hold together. That square is the one the brass balls settled in chapter eighteen's tray of clay, and the half in front of it is the half that stands in front of chapter thirty-one's kinetic energy.

Here is a history with five readings in it, carried by a pattern with a standing cost of two units, chosen because two cancels the half and keeps the arithmetic in whole numbers.

0, 3, 6, 9, 12

The steps are 3, 3, 3 and 3, their squares are 9, 9, 9 and 9, and half the standing cost is one, so the history scores 36.

Slide the whole thing. Add five to every reading and the list reads 5, 8, 11, 14, 17. Every step is 3, every square is 9, the total is 36 again, and nothing in the arithmetic noticed. That indifference is the laws not caring where you are, on a list of five numbers you can check with a pencil, and chapter fifteen met the same thing in the wiring: shifting all the readings together is the one direction repair leaves exactly where it found it, and the one direction no observer inside the arrangement can read.

Chapter thirty-one's recipe converts an indifference of that kind into a number. Take one step's contribution and ask how sensitive it is to how fast the reading is changing across that step. A square's sensitivity is twice the step, since widening a square by a hair lays a fresh strip along two of its sides. The half in front of the square cancels that two and the standing cost multiplies what is left, so the answer is the standing cost times the step. Then weight it by how far the symmetry moves the reading. A slide moves every reading by the same amount, so the weight is one everywhere, and with one reading per event there is nothing to add up. Every step of the history therefore carries a number of its own. On a pattern costing two it is twice the step.

On 0, 3, 6, 9, 12 the segments carry 6, 6, 6 and 6.

Move one reading and watch what breaks. Change the second reading from 3 to 4 and leave the rest alone: 0, 4, 6, 9, 12. The steps are 4, 2, 3 and 3, the squares are 16, 4, 9 and 9, and the total is 38. The segments carry 8, 4, 6 and 6.

Two things went wrong at the same place. The score went up by two. The segment numbers stopped agreeing across the junction at the moved reading, where an 8 comes in and a 4 goes out. One move cures both: put the reading back at 3, which is the midpoint of the 0 and the 6 on either side of it.

The midpoint is chapter nine's repair, arriving from a direction chapter nine never looked in. A history that repair has nothing to do to is one whose every interior reading sits at the midpoint of its two neighbors. That is exactly the condition under which the number carried by the segment coming in equals the number carried by the segment going out. Where the two disagree, a repair is owed, and it is owed at precisely the record where they disagree.

So the number is the same on every segment of an undisturbed chain. Whatever one segment hands over the next one receives, no segment in the middle keeps a share, and the chain has nowhere to put any that went missing. A quantity that behaves like that along a chain is a **current**. This one is four numbers on four segments, computed from a list of five readings by taking each difference and multiplying it by the standing cost.

A constant total says less than that. A total can hold while the quantity behind it disappears at one end of the world and turns up at the other. An audit of the total would find nothing to object to. The chain rules that out by saying where the quantity was in between. It crossed every junction on the route, in order, one segment at a time. Any place where the number changes is a place with a repair owed at it. Chapter nine's speed limit is the same count run the other way: influence advances one seam per event, so nothing gets anywhere without crossing the seams between here and there.

The number is **momentum**. Chapter five named it as what you get from the laws not caring where you are, chapter thirty-one supplied the machine that turns the one into the other, and this chain is the machine running on an object small enough to check by hand. Chapter eighteen listed the same quantity as the second, third and fourth of a region's four numbers, the drift of its records across the seam network along the three directions chapter fifteen supplied. One sentence from there carries what the drift is worth: energy squared is mass squared plus momentum squared, and a region's energy equals its mass exactly when its drift is zero.

Two patterns meeting have one chain between them at the seam where they touch. The segment on that seam carries one number. What leaves the ball arrives at the plate because the same segment is the last one of the ball's chain and the first one of the plate's, and a segment carries one number rather than two. Robins' ribbon is a reading of that number, taken at the far end of the plate's chain, some fraction of a second after the wood stopped splintering.

32.2 Why a heavy thing is hard to push

Chapter eighteen built mass out of the heap of water John Scott Russell chased along the canal: a bounded set of records that comes back after every round of repair, because flattening it would open wider disagreements around its edge than it settles in the middle. Repair never finishes with an arrangement like that. The rate it goes on demanding while the records stay put is the standing cost. A balance reads that cost off as a mass, the cost divided by the square of the speed of light, a conversion that has stood at exactly 89,875,517,873,681,764 joules to the kilogram since the meter was redefined in 1983. It has no error bar, because it was never measured: it is the square of 299,792,458.

Such an arrangement holds its shape and not its contents. The records carrying it at one event are not the records carrying it at the next. It survives by being re-established out of whatever is there, event after event, across every seam on its boundary.

So ask what a change of drift costs. Redirecting the pattern means re-establishing it in a different place relative to its neighbors, so every seam on the boundary has to be renegotiated instead of reconfirmed. The seams that have to be renegotiated are the same seams, and the work each one takes is the same work, that holding the pattern in place demands at every event anyway. A pattern that costs twice as much to hold costs twice as much to redirect, because the two are the same cost. That is **inertia**, and explains a fact that otherwise has to be assumed: the price of a given change of drift is proportional to the standing cost and to nothing else about the pattern.

The proportionality is exact. A measurement can contradict it. If the price of redirecting a pattern depended on anything besides what the pattern costs to hold, then two lumps carrying the same standing cost and different insides would need different pushes to change their drift by the same amount, and a balance that put them at the same weight would be reporting on a different quantity from the one a push reports on. Everything anybody has ever weighed has been weighed on the assumption that those are one quantity.

Inertia settles the undisturbed case too. With nothing pulling at the seams, every interior reading sits at its midpoint, the segment numbers agree the whole way along, and the advance per event is whatever it was at the start. A pattern with nothing to repair goes on going. The going costs nothing to maintain, because the maintenance is the standing cost and that is paid whether the pattern travels or sits.

The standing cost turns up in one other place. Chapter twenty-five's field equation takes what a region's records demand in the way of work as its source, and hands back the shape of the neighborhood around it. So the number that fixes how much shape a lump makes is the number that

fixes what it costs to redirect the lump. The mass that resists a push and the mass that responds to gravity are one entry in one ledger, read twice.

One of those readings is a price and the other is a source, they come off the same entry, and their equality needs no separate principle. The standard account states one, called the **equivalence principle**. The two hollow cylinders of chapter twenty-five, one of a titanium alloy and one of a platinum alloy, went into orbit to make the two readings disagree, and across five months of usable free-fall data they agreed to better than a part in a thousand million million.

32.3 Eleven, three, one

Chapter twenty-six put one unit of source on one port of the twelve-port wiring and solved for what the other eleven read. Over a common denominator of a hundred and eighty the twelve values are thirty-five at the source, two at each of the five ports one seam out, minus seven at each of the five ports two seams out, and minus ten at the port directly opposite.

A pattern sitting on the wiring cannot respond to those values. Add the same amount to all twelve and every seam sees what it saw before, so the readings are pinned only up to a common shift. A quantity pinned only up to a shift has nothing in it to push with.

What survives the shift is the drop from one port to the port next door. Over sixty rather than a hundred and eighty, those drops are eleven from the source to the first ring, three from the first ring to the second, and one from the second to the far port. Steep near the source, gentle far from it.

Stand on the source port and step outward and a unit of charge drops eleven sixtieths on the first hop, three on the second, one on the third. Stand on the far port and step inward and it climbs the same three amounts in the reverse order, a sixtieth, then three, then eleven, and the climb is the part somebody has to pay for. A port in the middle of that sits on five seams at once, and each seam has its own drop across it: a first-ring port reads a rise of eleven sixtieths toward the source, a fall of three sixtieths on each of the two seams leading outward, and nothing at all on the two that join it to the rest of its own ring.

The fall per hop is the **gradient** of the potential. The force on a unit of charge is the gradient across the seam it is sitting on. A force is a difference between what one port reads and what its neighbor reads. The difference is the renegotiation the next event owes.

32.4 The size of a shell

Follow the outflow rather than the readings.

Five seams leave the source port, one to each member of the first ring, and each carries the drop across it, eleven sixtieths. Five elevens is fifty-five,

so eleven twelfths of a unit crosses the first surface you can draw around the source.

Ten seams join the first ring to the second, two from each of the five, and each carries three sixtieths. Ten threes is thirty, so a half crosses the second surface.

Five seams join the second ring to the far port, each carrying one sixtieth, so a twelfth crosses the third.

The seams inside a ring carry nothing whatever, since both ends of them read the same number. The entire outflow therefore crosses the seams that lead outward, and eleven twelfths, a half and a twelfth are the whole of it.

Check them against what is enclosed. Chapter twenty-six's source arrived with its own balance, a twelfth subtracted at every port including its own, so the source holds eleven twelfths and every other port holds minus a twelfth. Eleven twelfths is what crosses the first surface. Eleven twelfths less the five twelfths held by the first ring is a half, which is what crosses the second. A half less another five twelfths is a twelfth, which is what crosses the third, and the far port's own twelfth brings the total to nothing.

The amount crossing a closed surface is the **flux** through it. The flux is fixed by what is inside and does not depend on which surface you drew. Nothing in that depends on the influence being an electric one. Any quantity whose total outflow is fixed by its source and which has to cross the seams between here and there obeys the same count, which is why two forces with nothing else in common carry the same exponent, and why the same exponent was on the books for gravity and for electricity long before anybody could say what either of them was.

Distance enters only as a count. The strength on one seam is the flux divided by the number of seams in the shell it crosses. The flux is the same across every shell, so the strength falls off exactly as fast as the shells grow and nothing else about the arrangement enters.

How fast a shell grows is fixed by how many directions there are to grow in. A container twice as wide in every direction holds eight times as much, because there are three directions available to be twice as wide in. A ball's content grows as the cube of its radius when the carrier has three independent directions in it, and as the fourth power when it has four. A shell is what you get by growing the ball's radius by a hair, so it grows one power more slowly than the ball it bounds: three directions in the ball, two in the shell, and the strength falls as the square of the distance.

Check that on two shells. The twelve ports run out of room at three hops, the far port being the end of the walk, so take the count on a seam network that keeps going. Stand two hops from the source and count the seams in the shell around you, then stand four hops out and count again. The far shell is twice as wide in each of the two directions a shell has when the carrier has three, so it holds four times as many seams. The flux crossing the two is the same number, so each far seam carries a quarter of what a

near seam carries. Hand the carrier a fourth direction and the shell picks up a third direction of its own, the far shell holds eight seams for every one the near shell holds, and each of them carries an eighth.

The count runs on two numbers. Both are shorter as letters. Let r be how many hops out from the source you have gone, and d the number of independent directions the carrier gives anything to spread into. The shell at r has d minus one directions in it, its seam count grows as r raised to that power, and has the same flux to share out however far out it sits.

$$\text{strength} \propto \frac{1}{r^{d-1}}$$

The curly sign is a proportionality, which holds the two sides in step while the source's flux sets the scale, and everything distance does to a force is sitting in that exponent.

Put d = 3 in. Three minus one is two, the strength falls as one over r squared, and the quarter you counted between two hops and four hops comes back out. Put d = 4 in. Four minus one is three, the strength falls as one over r cubed, and the same step outward costs an eighth. Nothing in either substitution asked what the source was made of, or whether the influence crossing the seams was gravity or charge.

The exponent is the number of directions in the carrier, less one.

Chapter fifteen fixed that number and did not choose it. Twelve ports leave twelve shadows in the band that outlives every other band, all 144 comparisons between those shadows come out as 1, minus 1, or one over the square root of five with a sign in front, and the count of independent rows in that table is three. The exponent of the force law is that three, minus one. The most famous number in classical physics is a subtraction.

Chapter twenty-six put that exponent at two out to the fifteenth decimal place. Edwin Williams, James Faller and Henry Hill measured it at Wesleyan University in 1971, on an apparatus built out of five concentric icosahedral shells, twelve corners and thirty edges apiece. Four megahertz at ten kilovolts peak to peak went across the outer two, and a battery-powered amplifier sitting inside the innermost shell listened for any trace of that signal appearing across the inner two. It heard nothing. A carrier whose surviving band held four directions instead of three would put the exponent at three rather than two. An instrument that sensitive could not have missed a whole integer.

32.5 Mass times acceleration

A pattern's advance per event is its step. The rate at which that advance changes is the difference between one step and the next, and a chain of readings shows it directly: the change in the step across a record is twice the amount by which that record falls short of the midpoint of its two

neighbors. A record sitting at the midpoint has equal steps on both sides and an unchanging advance. A record sitting one unit above the midpoint leaves with a step two units shorter than the one it arrived on.

That is **acceleration**, the rate at which a pattern's advance changes relative to its neighbors. On the chain it is a difference of differences, with no limit taken anywhere and nothing smooth required.

The history with the moved reading has one in it. In 0, 4, 6, 9, 12 the second reading sits one unit above the midpoint of the 0 and the 6 on either side of it, and its steps run 4 and then 2. The step drops by two across that record and the segment number drops by four, from the 8 coming in to the 4 going out, which is the standing cost of two multiplied by the change in the step. That record is accelerating, and also the record where the segment numbers disagreed and the score could be brought down. A pattern accelerates at exactly the records where its history has a repair outstanding.

Put every piece on one chain. A pattern with a standing cost, crossing a wiring that has a potential on it, scores each step twice over: half the standing cost times the square of the step, less what the potential reads at the port the record is sitting on. Ask, at one interior record, that the score not move when the record is nudged, which is chapter thirty-one's condition for the history the world runs. One line of arithmetic comes back. The standing cost, multiplied by the change in the step across that record, equals the fall of the potential across that hop.

Read the line in the words its pieces arrived with. On the left, a standing cost multiplied by the rate an advance is changing: mass times acceleration. The same left-hand side is the change in the segment number from the incoming segment to the outgoing one, which is a change of momentum across one event. On the right, the gradient of chapter twenty-six's potential: force.

Written with one letter for each piece, it is the line that opens every mechanics course.

$$F = ma$$

F is the drop in the potential across one hop, m is the standing cost of holding the pattern together, and a is the rate at which the pattern's advance changes relative to its neighbors. Nothing in it was measured in a laboratory and typed in. The mass is what a self-re-asserting pattern demands while its records go nowhere, the acceleration is a difference of differences along a chain of readings, the force is the fall of chapter twenty-six's exact potential across one hop, and the equals sign is chapter thirty-one's condition that the score not move when a record is nudged.

The left-hand side is a rate. A rate here is counted against repair events, at whatever pace a region's own state sets them, which was chapter twenty-

one's answer to what a clock is made of. Textbooks write that side as a change of momentum per unit of time without saying what the time is made of.

Divide both sides by the standing cost. It goes from the left, and when the potential is gravity's it goes from the right as well, because the entry that priced the redirection is the entry that put the shape there. What is left is an acceleration with nothing about the pattern in it, which is why a lump of anything at a given place changes its advance at the same rate as a lump of anything else.

Robins fired one ball and it crossed the room by one route. The plate swung once, the ribbon came out at one length, and every part of that has an account behind it: the number handed over at the wood, the price of turning the iron, the fall the ball rode down on the way. One thing about it has no account. The ball took a path, and it took one of them. Nothing that has been built says why it did not take several.

Chapter 33

Why Was Every Open Problem the Same Problem?

A physicist can spend forty years on the measurement problem and never need to open the literature on the cosmological constant. The two have separate conferences, journals and graduate courses, and people working on one share little technical vocabulary with people working on the other.

Lee Smolin put five problems of that standing on a single page of *The Trouble with Physics*, published in September 2006. He calls them the five great problems in theoretical physics. Combine general relativity and quantum theory into a single theory of nature. Resolve the trouble in the foundations of quantum mechanics. Find out whether the particles and the forces are aspects of one thing. Explain how the free constants of the Standard Model came to have the values they have. Account for dark matter and dark energy, or work out how gravity is modified at galactic scales instead.

The proposal pursued here is that these problems share an origin in how the background is specified. That common reading suggests mechanisms to test; it does not establish that the finite model has closed every physical problem.

A background is a set of places for things to be at, a parameter for them to happen along, and the two of them present before anything happens in them. Chapter one called it the arena. Read Smolin's five with the arena in view and each one is a question about the arena rather than about what moves around inside it: how many dimensions it has, what the constants of the things in it are, why one class of event occurring inside it counts as a measurement while the others do not, and what its empty parts weigh. A description that grants itself the arena in its first line has no route to any of that. Four of the five ask what the arena is or what sits in it, and the fifth asks whether the law running on it needs changing at the radius of a galaxy.

Take them one at a time. Name the problem the way its own literature names it. Name the thing assumed underneath it, which is usually so old that nobody lists it among the assumptions. Then take the assumption

out and watch what the problem does. Solving a problem leaves an answer behind that somebody can go and measure. Removing an assumption leaves a question that cannot be put into words any more.

33.1 Ten words Bell wanted banned

In August 1990 John Bell published eight pages in *Physics World* listing ten words that, he wrote, have no place in a formulation with any pretension to physical precision. System. Apparatus. Environment. Microscopic. Macroscopic. Reversible. Irreversible. Observable. Information. Measurement. The worst of all, he added, is the last one.

His target was a joint. The theory as John von Neumann set it out in 1932 contains two kinds of change. One is smooth and deterministic and runs whenever nobody is looking. The other is a jump: it happens when a measurement is made, it produces one outcome out of several, and the odds come from Born's footnote. Nothing in the theory says where the first kind stops and the second begins, and von Neumann showed that the joint slides anywhere along the chain from the instrument through the amplifier to the eye of the physicist without changing a single prediction. Fifty-eight years later Bell was pointing out that a fundamental theory containing a movable part, whose position no experiment fixes, is an unfinished theory.

The assumption under the joint is older than the theory and it sits inside the word Bell most wanted removed. To measure is to find out a value that was there beforehand. To disturb is to spoil a value that was there beforehand. Both take for granted a world stocked with definite values and experiments as encounters with them, and the measurement problem is the question of how one encounter manages to be smooth and a jump at once.

Take the stocked world out. What an observer has is an algebra of questions, some of which refuse to commute, and a record layer sitting in the part that commutes with everything. Nothing gets disturbed, because nothing was carrying a value to be disturbed. A record gets written. The write is a commit that discards the alternatives it resolved. The odds come with it. Take any assignment of numbers to the observer's questions, the weighted ones as well as the plain yes-or-no ones, that lands between zero and one, gives the certain question the value one, and adds up whenever two questions combine into a third. It is the trace against exactly one state, in every finite dimension. Born put the squaring in a footnote in 1926 because he had noticed that it worked. Additivity forces it. The word Bell wanted removed drops out of the description altogether, and what stands in the place where it was is an operation with a price: a write that cannot be undone, leaving the world less able than it was to say what it used to hold.

The joint Bell objected to turns out to have a home. It falls between the questions that commute and the questions that refuse, it sits in a different

place for every observer, and no experiment slides it, because a commit either happened or it did not. The ball out of Robins' musket took one route across the room because one route got written down, in splinters and in a length of ribbon on the floor.

33.2 Loschmidt's objection

In 1876 Josef Loschmidt put an objection to Ludwig Boltzmann that has never been met on its own terms. Take any mechanical process, run it, then reverse every velocity in it. The reversed motion obeys the same equations exactly, so a quantity that rose during the first run falls during the second, and no theorem about time-symmetric mechanics can deliver a time-asymmetric conclusion. Every derivation that appears to do it has smuggled the asymmetry in somewhere.

Where it gets smuggled in is a statement about how everything started: an arrangement of the rare, low-entropy kind, fourteen billion years ago, defended on the ground that nothing else does the job. David Albert named it the past hypothesis in 2000.

The assumption sits one line above the fix. The elementary dynamics is blind to direction, so a direction has to be imported from outside the dynamics, and the only place left to import one from is the boundary.

Here the dynamics is not blind to direction. The repair move is many-to-one on the readings it touches, its quantum version drains coherence that no later repair restores, and the relative-entropy and fluctuation identities of chapter sixteen hold at every step, with forward and reverse weights apart by exactly the exponential of the entropy produced.

So the asymmetry belongs to the operation, and works on the first repair of a fresh arrangement exactly as it works on the ten billionth. Loschmidt's reversal turns every velocity around and un-writes nothing. The bits the commits threw away have nowhere to come back from.

33.3 A hundred and twenty orders of magnitude

Add up the zero-point energies of the fields of empty space, cut the sum off at the shortest length anybody takes seriously, and compare the total against what the sky reports, which is a cosmological constant of about 1.1 times ten to the minus fifty-second per square meter. The two disagree by roughly a hundred and twenty orders of magnitude, the largest gap between a calculation and a measurement in the history of the subject.

The assumption is that empty space carries an energy density, that the density sits in a spacetime which was there first, and that gravity responds to it as it responds to any other energy.

Gravity's local probes are light rays. The vacuum's stress is the same everywhere and looks identical to every observer however fast they are moving, which forces it to be a number times the metric, and the metric evaluated twice on a light-ray direction is the interval of a light ray with itself, which is zero. A light ray is blind to any vacuum energy density whatever, however enormous, by the definition of the probe rather than through a cancellation. Under chapter twenty-five's continuum, conservation and entropy-balance hypotheses, the null data deliver Einstein's equation with one metric-proportional term left open. The finite record constructions do not yet establish all those physical hypotheses.

A specified horizon-capacity condition can supply the missing term. In that trial model the count is 3.3001 times ten to the hundred and twenty-second, against a weighted expansion coordinate of 3.3129 times ten to the hundred and twenty-second. Identifying this capacity with physical generalized entropy and deriving its stationarity from the same repair history remain separate requirements.

33.4 An ensemble of one

In 1953 Fred Hoyle turned up at the Kellogg Radiation Laboratory at Caltech and told the nuclear physicists working there that the carbon-12 nucleus had to have an excited state at around 7.68 million electron-volts, on the ground that stars manufacture carbon and have no other way of doing it. Ward Whaling, who had helped build the laboratory's magnetic spectrometer, put his group on it: a nitrogen-14 target, a beam of deuterons, and a scan across the alpha particles that came off. The state is there, at 7.68 million electron-volts give or take 0.03. The measurement was published in *Physical Review* the same year.

Shift that level by a fraction of a percent and the reaction that builds carbon inside stars stops working, which makes that level the first exhibit in every account of a universe tuned for its occupants. Take any of the twenty-six measured numbers of the Standard Model, change it on paper and work out what the resulting world does, and for several of them the answer is a world with no chemistry in it. The constants of this one look, from that angle, like settings somebody dialed in.

An argument of that shape is a probability claim. A probability claim needs two things: a collection to draw from, and a weight on the collection. The collection was proposed later. The extra dimensions of a string theory can be curled up in an enormous number of ways, each giving a different set of particles and constants at ordinary energies. A collection that large makes a hospitable world unsurprising, provided the collection is there and somebody supplies the weight.

The declared self-reference maps compare two readings of the same carrier and give an equation with the trial quantity on both sides. Contraction

proves one fixed point for each specified map. The shared-width map returns inverse coupling 137.035660 against the measured 137.035999. No calculation here proves that a quark contribution closes that difference. Uniqueness within a trial map does not yet select that map as the physical electromagnetic response.

33.5 Two parts in ten to the seventeenth

The Particle Data Group's 2024 review lists the Higgs boson at 125.20 billion electron-volts, with an uncertainty of 0.11. The scale at which gravity becomes as strong as everything else sits around 1.2 times ten to the nineteenth billion electron-volts, seventeen orders of magnitude higher up. In a theory where a scalar's mass collects contributions from every scale in between, that mass belongs near the top of the range and it sits at the bottom of it. The standard repair was a symmetry pairing every particle with a partner whose contributions cancel against its own. The Large Hadron Collider spent the whole of its second run looking. The floor under the partner that theory assigns to the gluon sits above 2.3 trillion electron-volts.

The assumption is the top of the range. A hierarchy problem is a comparison. A comparison needs two scales, one of which was written into the theory by hand as the place where the description gives out.

A conditional scale comparison relates Newton's constant to a screen area calibrated in laboratory units. A proposed exponential relation uses a cell coupling near 0.0411, about one part in 24.32, and yields a small scale ratio. Its running, matching and scale-identification assumptions must be specified. The Higgs vacuum value of 246.22 billion electron-volts remains a comparison input; the finite interaction identities do not select it.

An exponential can explain how a large hierarchy is possible. Deriving the physical coefficient, the matching map and the vacuum would make it a prediction. That remaining selection problem is the one chapter thirty-one keeps explicit.

33.6 The groups that died in 1894

In 1974 Howard Georgi and Sheldon Glashow proposed that the eight, the three and the one of the Standard Model sit inside a single simple symmetry. The thirds of quark charge fell into place, the three force strengths appeared to converge at high energy, and the proton acquired a finite lifetime. Fifty thousand tons of ultrapure water under a mountain in Japan have been watching for the decay since April 1996 and have seen nothing of the right shape, which puts the floor under the lifetime at 2.4 times ten to the thirty-fourth years.

The assumption is in the choosing. A symmetry group gets picked by the person writing the theory and imposed on the world from outside, which is a move available only where there is an outside to impose it from.

The clause that removes it arrived with two firms keeping books in different currencies. They need an exchange rate, there is no market outside the room to look one up in, so the rate has to be computed out of what the two of them hold between them. Applied to the twelve ports of a patch, the same clause says that every relabeling of the readings has to be a move the patch can perform on itself, since no venue anywhere holds a preferred labeling of twelve ports. The relabelings with any standing are the ones the dials produce.

Within the port-current theorem's stated hypotheses, that requirement constrains the group. A central move commutes with every relabeling, so a relabeling and its undoing slide together and cancel and the move comes back untouched, which puts the center inside the part of the twelve readings that all sixty rotations of the wiring leave alone. Those sixty rotations carry every port to every other port, so the surviving part is one number repeated twelve times: one dial, and at most one. A simple group has a center zero dials wide, by the definition of simple. Twelve dials come apart as one plus eleven, eleven splits as three plus eight and in no other way, and a simple group has nowhere to stand.

What survives is a product arrangement. One dial that no rotation touches, a block of three and a block of eight, which match the hypercharge, weak and strong Lie algebras in the conditional construction. Electromagnetism combines hypercharge and a weak generator after a specified symmetry breaking. The local algebra result precedes the matter lift; it does not by itself select the physical global group or couplings.

So the beautiful groups died of a clause about exchange rates. The catalogue of sizes a simple group is permitted to have was finished by Wilhelm Killing in four papers between 1888 and 1890 and made rigorous in Élie Cartan's thesis of 1894, twenty-six years before the proton had a name.

33.7 A proof one step long

On 27 November 1997 Juan Maldacena posted a paper showing that a theory of gravity in a particular curved spacetime is the same theory as a field theory without gravity living on that spacetime's boundary at infinity. Everything in the interior has a counterpart on the surface. The interior is the surface's information reorganized into an extra dimension. In 2006 Shinsei Ryu and Tadashi Takayanagi supplied the geometric dictionary entry: the entanglement entropy of a region of that boundary is the area of the smallest interior surface hanging on the region's edge, divided by

four times Newton's constant. It is the most productive thing anybody has done with the idea that a region's memory scales with its surface.

The assumption is that the inside of a region is fixed by what is written on its edge. Stating that carefully takes three pieces: a region with an inside, a summary that lives on the edge, and the claim that two insides carrying the same summary are the same inside. Everything depends on which summary. The first one anybody reaches for is the edge itself, because the edge is where the information was supposed to be. Set it up that way and the line going in reads: two insides with the same edge are the same inside. The line coming out reads: two insides with the same edge are the same inside. A proof checker handed both lines gets from one to the other in a single step, the one stating that a thing equals itself.

The claim has content when the summary is built out of something other than the edge and shown, on its own ground, to fix the inside. Then a pair of insides agreeing along the edge and differing behind it would break the claim. Whether the network holds such a pair is settled by its seams.

Here the edge is a collar. A band of patches one patch thick, with an interior on either side of it, and both interiors keeping records, which makes both of them observers. The surface where the information sits is inside the network, between two parties comparing notes across it.

Two theorems come with that arrangement. Finitely many patches, finitely many readings each, and a local check on every seam demanding that its two readings answer each other under the dictionary sitting there: a setting of all the patches meeting every check at once is a legal word, the legal words are exactly the globally consistent states, and an object of that shape is a constraint code, of the same family as the one Richard Hamming built in 1950 out of four data bits and three checks. The second theorem is a negative one: a minimum-cut formula does not come with the code. A list of checks fixes which configurations are legal and says nothing whatever about how far apart two legal configurations sit, or where the cheapest bottleneck through the wiring runs, and both of those depend on which port is joined to which port. An area formula for the entropy of a region has to be computed on one particular wiring. On this one, chapter twenty-four found the count in the patches of the screen, with the seams along a region's edge playing no part.

A region's memory is set by its surface because everything anybody can learn about the region arrives through the ports on that surface, which chapter thirteen got out of a wiring diagram with no gravity in the argument. What drops out of the account is the outside boundary. The far side of any collar in this world is another interior with somebody in it, keeping its own records and repairing its own seams.

33.8 Eight short answers

The following answers summarize the finite constructions and their conditional interpretations. They retain the hypotheses stated in the earlier chapters; they are not a claim that every physical parameter has been source-selected.

What suggests three families? Within the stated quark-mixing and asymptotic-freedom assumptions, the usual bounds restrict the possibilities. A minimum-cost irreducible response-band rule chooses a rank-three candidate. Attaching that band to physical families and deriving their nondegenerate masses remain separate problems.

Under the declared matter lift, why do the charges come in sixths? Because six combinations of gauge moves do nothing whatever to any piece of matter, so they get bundled in with the move that does nothing at all, and the bundle holds only if every hypercharge advances in whole multiples of one sixth. That is the lock that leaves hydrogen neutral to better than one part in a billion billion.

Why is the usual grand-unified gauge-boson route absent from this carrier? Because the carrier that route needs, one direction coupling color to weak charge at once, is missing from a twelve-dimensional list that splits into eight, three and one, and a missing direction cannot be made small or slow: that carrier route is absent exactly. This does not exclude every effective baryon-number-violating interaction.

How close are the alpha candidates? The screen-cell trial maps have certified fixed points. The shared-width map gives inverse coupling 137.035660. A. D. Wissner-Gross supplies a separate entangled-pair expression with inverse 137.035999189566. Both arithmetic constructions need the physical electromagnetic identification discussed in chapter twenty-nine.

What does the geometry say about Koide's relation? On one face of the icosahedron, equal weight for the two ways the corners can respond gives exactly two thirds. With the electron and muon masses as inputs, it pins the tau inside a window seventy-two electron-volts wide, centered at 1776.969027 million electron-volts against a measured 1776.93.

What gives two polarizations in the field construction? A declared massless Abelian vector field in one time and three space dimensions has four components. The gauge constraint and redundancy leave two transverse modes. Its identification with the physical photon needs the current and response matching discussed in chapter twenty-nine.

Why does space have three directions? Because the twelve readings of a patch split into blocks that fade at different rates under repair, and the slowest block, holding 0.9539 of what it carries per pass against nine tenths and 0.8794 for the others, has rank three.

Why is there a speed limit at all? Because the four numbers of a two-by-two block carry the interval in their determinant, one sign for time

against the opposite sign for the three directions of space, and where the determinant vanishes there is a cone whose directions make up an observer's sky. A separation is a count of seams crossed and a duration is a count of repairs committed, and the speed of light is the conversion between the two counts once each carries physical units, worked out by each observer from its own books and agreed on because every observer's sky is a sphere.

A separate conditional dark-matter construction represents missing mass at galactic scales by a repair charge on the collars. It stays because no later commit writes what would clear it, and its profile around a galaxy produces an acceleration law fixing the speed of the outermost stars from the visible mass and one acceleration constant.

33.9 The empty column

Chapter one read three centuries of physics as a crash log: what was assumed, what broke, and what got fixed. The third column records proposed mechanisms under their stated assumptions, rather than a completed derivation of every observed scale.

The problem	What sits underneath	What comes out when that goes
Measurement	A world stocked with definite values, disturbed by an encounter	A commit into the commuting part of an algebra, with the odds forced by additivity
The arrow of time	Direction-blind dynamics plus a special first arrangement	An irreversible write, forward and reverse weights apart by the exponential of the entropy produced
The cosmological constant	A vacuum energy sitting in a spacetime that was there first	Local null data leave one metric term free; a specified horizon-capacity condition fixes it
Fine-tuning	An ensemble of worlds for this one to be improbable among	Specified self-reference maps have unique roots; selecting their physical attachment remains open
The hierarchy	A fundamental scale entered by hand for a measured one to be compared against	A conditional exponential scale relation with matching inputs; the Higgs scale remains unselected

The problem	What sits underneath	What comes out when that goes
Grand unification	A symmetry imposed on the world from outside it	Every relabeling performed by the patch itself, one central dial, and no simple group with room for it
Holographic duality	A boundary at infinity taken as its own summary, so the assumption is the conclusion	A collar with an interior on both sides, and a constraint code carrying no distance

Seven rows expose one recurring assumption: somebody wrote down the stage before writing down anything that happens on it. The finite wiring supplies records, seams and repair as a common structure. It realizes some finite premises; the continuum, physical-field and geometric-response premises still require their own witnesses.

There is a test that separates the outcomes. Ask what a solution would have to contain. A misplaced mechanism has to go, and a free parameter has to become the solution of an equation the world cannot avoid. For the cosmological constant, the light-ray argument explains why local data leave the metric term undetermined, and an additional horizon-capacity condition supplies its value. Deriving that physical condition from the same repair history remains part of the programme.

Which leaves the picture everybody starts with, the one with a container, a clock running at one rate for everybody, and objects sitting at positions inside it. That picture is the same assumption in domestic form, accurate to more decimal places than any instrument in a house can resolve. Somebody assumed a container and a clock and got the tides out of it, and the precession of the equinoxes, and the mass of the Sun worked out from how long a planet takes to go round it. That is two hundred years of correct answers out of a picture whose opening sentence puts the world inside a box that was never there.

Chapter 34

Why Was Newton Right for Two Hundred Years?

Yaakov Fein, Markus Arndt and six colleagues sent molecules down a two-meter tube in a laboratory at the University of Vienna in 2019 and collected what came out the far end. The molecules were tailored oligoporphyrins, up to two thousand atoms apiece, weighing more than twenty-five thousand atomic mass units, which is roughly a small protein. What arrived at the detector was a pattern of light and dark bands. The wavelength that produced it was 53 femtometers, five orders of magnitude smaller than the molecules doing the interfering.

No heavier object has been shown to interfere.

The natural reading is that the bands hold up to some weight and give out above it, and that a kitchen chair is far above it. The natural reading is wrong. An experiment run fifteen years earlier in the same city says why.

In 2004 Lucia Hackermüller, Klaus Hornberger, Björn Brezger, Anton Zeilinger and Markus Arndt put C70 fullerenes through an interferometer, seventy carbon atoms each, and heated them on the way in with a multi-pass laser. Every photon a molecule absorbed raised its internal temperature by about 140 kelvin. The beam entered the interferometer at temperatures approaching three thousand. A molecule that hot glows. The bands weakened as the molecules got hotter, and the weakening matched, quantity for quantity, the rate at which the emitted infrared light was carrying off which route through the gratings the molecule had taken.

Nobody read that light. There was no detector aimed at it, no experimenter waiting on it, and no measurement in any sense a laboratory notebook would recognize. The molecule wrote its own position into the room, in a handful of infrared photons. The writing was enough. Weight was never the variable. The pattern ended when a record got written.

The chair in your kitchen is in the same business and has never once come out in bands. Why does a chair have a definite place, when a molecule of two thousand atoms does not have to? Why does everything any instrument has ever written down come out as ordinary numbers, obeying ordinary arithmetic, with none of the refusals that run through everything underneath? Why does space look smooth? And why did a description of space and time that put both of them in by hand, unexplained, run for two hundred and eighteen years with almost nothing to show against it?

34.1 Sixteen copies

Take a fact worth sixteen yes-or-no questions: enough to fix where something is to one part in sixty-five thousand, along one direction. Write it into one place and it is a record that a determined liar could rewrite by guessing sixteen bits, which comes off once in 65,536 tries.

Write it into sixteen places instead.

The liar has to produce all sixteen copies without holding the original, because a single copy that disagrees with its neighbors exposes the whole attempt. Take 65,536, the odds against faking one copy, and multiply it by 65,536 fifteen more times, once for every copy after the first. That is two hundred and fifty-six bits of blind guessing, and comes off about once in 1.16 times ten to the seventy-seventh.

Look at what the sixteenth copy contributed to the world's knowledge. Nothing. It says what the first copy says, to the bit. The world learned sixteen bits when the first copy was written, and it holds sixteen bits after all sixteen exist, and the entire effect of the other fifteen is on what it would cost to lie.

A chair is not held by sixteen copies. Every photon that leaves it carries where it was, every air molecule that comes off its surface leaves with the same information, and the copies are being made continuously, in numbers far beyond sixteen. That copying is chapter nineteen's pinching, performed by the room rather than by an experimenter, and decoherence is its name.

So definiteness belongs to the record rather than to the chair. There is a record of where the chair is, written into every photon that has left it since you last looked at it, no two copies disagreeing, and the odds on any alternative pinned so far below one part in ten to the seventy-seventh that no experiment will ever separate them from zero. Definiteness is what that redundancy looks like from inside, to an observer who is one more copy.

The public half comes free with it. Chapter four's overlap rule says that where two observers can compare, they have to agree, and the redundancy is what makes the agreement cheap: you and everybody else in the kitchen are reading different copies of the same sixteen bits, so nobody has to go and check with anybody. Four people walk into a room and the chair is in the same place for all four, which is what the word objective is describing, and the reason it holds is that all four are downstream of a record copied an enormous number of times before they arrived.

Chapter eleven has the name for it. A settled state that the record fixes, rather than the path taken to it, is an observation-determined normal form. At kitchen resolution the record is the photons and the settled state is where the chair is. The room hands it to all four of them without any of them having seen a single repair.

It is also being paid for. Chapter twenty-three put a price on a committed record: at three hundred kelvin, 2.87 divided by ten to the twenty-first

joules for every yes-or-no question the commit settles, expelled into the room as heat. The chair sits at a definite place because the room is buying it a definite place, bit by bit, without pause, at a rate too small for anybody in the kitchen to notice.

34.2 This or that, here or there

Chapter nineteen found where records live. The questions that commute with everything else in an observer's algebra collect into its center. A record has to be there, because a record is exactly a question that nothing else the observer can ask will disturb.

Take an inventory of that center once the copying has done its work. A finite list of alternatives. This one or that one, here or there, one or two. Every pair of them can be asked together, in either order, with no cost to either answer. Probabilities that add the way probabilities added before 1925.

That is an ordinary classical event algebra, the kind George Boole set out in 1854 in a book about the laws of thought, where every question has an answer, the answers combine with and, or and not, and nothing depends on the order you ask in. It is the whole of what any instrument can print. A kitchen scale showing 1.24 kilograms is one alternative from a finite list, written into a place where nothing else the observer can ask will move it, and the same goes for a pointer, a photographic plate, a click in a counter and the sentence a physicist writes in a notebook afterward.

Everything an observer can write down was always going to look like ordinary numbers, because the writing is what makes it ordinary. Physics before 1925 was an accurate theory of the center, put together by people whose instruments could not reach a question living anywhere else.

34.3 The traffic runs one way

A physicist who works only with pointer readings is working inside the center, and everything that person writes down is true. Whether it is the whole of what there is gets settled by the maps between the two layers. Those maps are lopsided.

Upward, every classical record is one of the operators. Take a reading off your kitchen scale and find the question in the algebra that asks it: add two readings and the questions add, multiply them and the questions multiply, swap yes for no and the question flips with it. A record about one region lands inside that region's own algebra, carrying the odds it had. Nothing is lost and nothing is approximated. Records are the operators that quarrel with nothing.

Downward there is no such map. The reason is arithmetic rather than difficulty.

A classical readout assigns a number to a question. Suppose some rule did that for every question in one of chapter nineteen's blocks, thirteen alternatives across, and suppose it added the way the questions add and multiplied the way they multiply. Numbers multiply the same way round in either order, so the rule would have to hand the same number to a product taken one way as to the product taken the other, and hand their difference zero. Those differences, taken over every pair in the block, fill out everything in it except the overall scale. A rule that sends all of them to zero has one option left: report the sum of the diagonal, divided by thirteen, which is an average.

Then take one question inside the block and watch the average fail. The question says yes to one direction out of thirteen, so its average is one thirteenth. Ask it twice and you have asked it once, which means the square of the question is the question, so the average of the square is one thirteenth as well. Square one thirteenth yourself. A rule that respected products would have to hand the square that number, one part in a hundred and sixty-nine, and the average hands it one part in thirteen: same block, same question, and a factor of thirteen between the two answers.

So there is no structure-preserving map from the operator layer down to the classical one, in any block bigger than one alternative across, and the obstruction is blunt: a readout has to assign a number, and a thirteen-by-thirteen block of operators has no numbers in it to be assigned. What exists downward is the average. It keeps every total right, keeps every probability between zero and one, carries each record across untouched, and mangles multiplication.

The everyday world is therefore one exact copy plus one lossy average, and multiplication is what the average loses. That settles which of the two layers is holding the other up. A classical account lifts into the operator account entire, products included, and lands inside it as a working part. Send the traffic the other way and the best map available gets the product of a question with itself wrong. No amount of care in the averaging repairs it, because the failure sits in what a number can do rather than in how the average was taken. A classical world could contain no operator layer at all. An operator layer contains a classical world exactly, in its center, and hands it out to anybody who writes anything down.

34.4 A world that cannot flow

There is a second obstruction, the one that decides whether a classical world could ever have been the ground floor.

A record takes one of finitely many values. A change of the record layer that preserves its structure has to send alternatives to alternatives, which is to say it shuffles the labels. A finite list has only finitely many shuffles. So consider a continuous family of them, one shuffle for every real number,

starting from the shuffle that does nothing: the kind of family that any dynamics, any symmetry with a dial on it, any smooth motion whatsoever would have to supply.

Such a family cannot move. To carry a label off itself it would have to pass through values in between, and between two alternatives there is nothing: a record whose value is partway from heads to tails is not a record. The parameter runs over the real numbers, which come in one connected piece, and the labels sit apart from one another with gaps that no continuous path crosses. A path that starts at the label it started at stays there.

Every continuous one-parameter group of purely classical public symmetries is the identity. The proof is the connectedness of the real numbers on one side and the gaps between the labels on the other. There is nothing else in it.

Try to write down the rate of change of a record and the same wall arrives from the other direction. A rate of change is a difference divided by a shrinking interval, and needs the quantity to take values arbitrarily close to the one it has. A record takes the value heads, and the values near heads are heads. Every derivative of it is zero, every equation of motion for it says nothing changes, and the only motion the layer supports is a jump from one label to another with no time spent in between.

Chapter twenty-one pulled a clock out of a state and found the flow living on the part of the algebra that refuses to commute, reversible, supplied by whatever the observer holds. The connectedness argument reaches the same place from the classical side. A purely classical world cannot flow. It can be relabeled in steps, and between the steps it does not change.

Which is why the classical layer is a readout rather than a foundation. Whatever is flowing is flowing underneath, in the part where order matters. A classical dynamics is a sequence of photographs of that flow, taken at a resolution where the photographs join up smoothly enough to differentiate. Newton wrote down a continuous dynamics for positions and momenta and it worked for two centuries. What moved in it was the layer underneath, and not the records anybody wrote down.

34.5 The finest length anybody has measured

Run a hand along your kitchen counter. It is smooth under your fingers, and chapter fifteen built space out of comparison counts that are not smooth: the continuum is where the counts are heading, and nobody is at the destination. Whatever an observer holds is a finite list of whole numbers taken after a finite number of passes, and the distance from there to the smooth limit is every step there is. Refinement has no completed final stage. Descriptions get finer and no description arrives.

The instruments say the same thing from the other side. In Louisiana and in Washington State there are two L-shaped machines with arms four kilometers long, built to catch gravitational waves, and what they detect is a change in the length of one arm of about a meter divided by ten to the nineteenth. A proton is about 1.7 of a meter divided by ten to the fifteenth across, so what those machines resolve is under one ten-thousandth of the width of a proton. That is the finest length measurement human beings have made.

It lands at a finite resolution, like every measurement before it. The next instrument will land at a finer one, and the one after that finer again, and the sequence of resolutions has no last term in it, which is what it means for smooth space to be a limit rather than a place. The counter feels smooth because every instrument you could put on it stops short of the limit by less than that instrument can register.

A position fixed to one part in 65,536 takes sixteen yes-or-no questions, each one halving the stretch the answer can be in. Do that along the edge of the counter. Halve a meter ten times and you have cut it by 1,024, which is a thousand for envelope purposes. Do six lots of ten and you are at a meter divided by ten to the eighteenth. Three more halvings supply a factor of eight.

Sixty-three questions therefore place a point on your counter to about a meter divided by ten to the nineteenth, which is the wobble those four-kilometer arms register. Going from where a chair is to the finest length anybody has measured costs four times the questions and nothing else.

Ask a sixty-fourth and what you hold is one bit longer and a whole number. Ask sixty-three more, a second copy of everything between your counter and Louisiana, a longer whole number, finite, exactly as it was at the first question, and with as many questions left to ask as there were before anybody asked one. The smooth counter is where those answers pile up. Every answer anybody holds about it is a finite list of whole numbers.

Calculus works on the world for that reason and no other. A derivative is a limit of differences, so writing one down commits you to differences taken at every scale. The world supplies them down to whatever depth the apparatus reaches and no further. The answers agree because the disagreement is parked below the last decimal place any instrument has produced, which is exactly where an abstraction hides its leak until the day somebody builds a better instrument.

34.6 Twenty-two and a half meters

The *Principia* was published on 5 July 1687 with a scholium near the front saying that absolute, true and mathematical time, of itself and from its own nature, flows equably without regard to anything external. Absolute space got the same treatment in the sentence after, similar and immovable

of its own nature, answerable to nothing outside. Chapter one called those two sentences the load-bearing assumptions of the book, put in by hand by a man who had just explained that they were too obvious to need stating.

They ran for two hundred and eighteen years. A wrong assumption at the base of a theory does not usually take two centuries to show itself.

In 1960 Robert Pound and Glen Rebka took the tower of the Jefferson Physical Laboratory at Harvard, 22.5 meters of it, put an iron-57 source near the roof and an absorber in the basement, and watched what happened to 14.4 kilo-electronvolt gamma rays on the way down. The prediction they were testing was that the frequency arriving at the bottom would be higher than the frequency leaving the top by about 2.5 parts in ten to the fifteenth, because a clock in the basement runs slower than a clock on the roof. Over ten days of counting they measured 2.56 parts in ten to the fifteenth, give or take 0.25.

That is the size of Newton's error about absolute time, measured across a building. Two and a half parts in a thousand million million. To see it at all, Pound and Rebka needed a nuclear transition, a technique Rudolf Mössbauer had published two years earlier, and six thermocouples watching the two ends of the shaft, because one degree Celsius of difference between the source and the absorber shifts the frequency by about as much as the whole height of the building does.

Set that against the instrument. The best timekeeper of the eighteenth century was John Harrison's fourth attempt at a sea watch, which sailed from Portsmouth in November 1761 and reached Jamaica on 19 January 1762 having lost 5.1 seconds. Over two months at sea, 5.1 seconds is an error of roughly one part in a million, and the shift Pound and Rebka went after is a few hundred million times smaller than the error in the finest clock of its century. Newton's absolute time was safe from every clock built in the two hundred years after him, and safe by a margin that no improvement in mechanical clockmaking could have closed.

The sky had the better instruments. It took a century and a half of them to turn up a single discrepancy. Urbain Le Verrier went through the transits of Mercury from 1697 to 1848, subtracted the pull of every other planet, and reported in 1859 that the orbit's closest point was creeping around the sun about 38 arcseconds per century faster than Newton's law allowed. Simon Newcomb redid the subtraction in 1882 and made it 43. Forty-three arcseconds per century, out of 574 that the whole orbit turns through against the fixed stars in the same time. All 574 of them come to a sixth of a degree. That is the one leftover in the solar system no further Newtonian bookkeeping ever absorbed.

The density of records does the same work on the other side of the account. Everything a human body handles is copied into its surroundings so many times over that the odds on any alternative place for it sit below one part in ten to the seventy-seventh and keep falling, so the indefinite-

ness Newton left out of his description is as invisible from a kitchen as the difference between the clock upstairs and the clock downstairs. Speeds far below the speed of light, and records copied past counting: those are the two conditions a person lives under, and under both of them the corrections sit in the fifteenth decimal place, nine places past anything Newton's contemporaries could read.

The scholium was an excellent local approximation, and Newton wrote it down as a definition of what time is, of itself and from its own nature. A quantity introduced that way has no route back to being corrected, because there is nothing it was ever answerable to. That is chapter one's error in its purest form, committed by the man whose mechanics held for two hundred years.

34.7 What the abstraction is an abstraction of

The prologue put the everyday picture on the table: a three-dimensional container, time running inside it at one rate for everybody, objects at positions in it, physics describing how the objects move. It named that picture a leaky abstraction, in Joel Spolsky's sense of a simplification that holds until the thing underneath does something the simplification has no words for, and listed six places the leaks come through.

Thirty-two chapters later, the everyday container is the commuting, redundant, coarse-grained readout of a finite network of observers repairing their disagreements. Commuting, because the questions that survive being written down are the ones that refuse nothing, and that is the center. Redundant, because a fact written in enough places has odds no experiment can lift off zero or lower off one. Coarse-grained, because the only downward map is an average, and the average discards the order in which questions are asked. And a readout, because the layer it reads has no motion of its own and never did.

Every one of the prologue's six leaks is a place where the reading is taken at a resolution the abstraction has no words for: a cabin that gives no sign of its own motion, an orbiting clock running fast by forty-five microseconds a day and slow by seven because a rate belongs to whoever holds it, a needle whose bands survive until a route gets written down, two laboratories 1.3 kilometers apart with no separate answers to have been carrying, a black hole that cannot destroy the record of what fell into it, and a gravitational entropy ceiling set by a surface, because a region's records live on its surface.

There is nothing wrong with the container picture. It is what the world looks like from inside, at the resolution a body operates at. It is not where the world starts.

Throw a ball across the kitchen. Its path is a pattern following a gradient, re-established at each place in the comparison count; its mass is the standing cost of holding that pattern together against the seams it touches; its position is written into the room by every photon that leaves it; and it lands in your other hand with a definite place and a definite speed, both of them readings taken off a layer that has neither. Newton's account of that throw is accurate to every decimal place your hand can feel. Thirty-three chapters went in underneath it, and not one of them has said what the ball is.

Chapter 35

What Is Anything Made Of?

Catch the ball. Pick up anything else at all: a beam of light, the ache in a knee. What is any of it made of?

The most careful answer the eighteenth century wrote down is a table of thirty-three rows. Antoine Lavoisier printed it in Paris in 1789, in the first volume of the *Traité élémentaire de chimie*, under the heading Tableau des substances simples: everything he judged could not be taken apart any further, sorted into four groups, with the name he was proposing for each one printed beside the older name it was replacing.

He paid for the laboratory that produced it out of tax collection. He had bought a share in the Ferme générale, the private company holding the right to collect the crown's duties, and that is what killed him: on 8 May 1794 he and twenty-seven other tax farmers were tried in a morning, driven to the Place de la Révolution and guillotined, Lavoisier fourth in the line.

Twenty-three of those thirty-three are chemical elements and hang on the wall of every school laboratory today. He got them by taking things apart and weighing what came out, which nobody had done at that scale before, and for a list drawn up in 1789 that is a great deal of it left standing: oxygen, nitrogen, hydrogen, carbon, sulfur, phosphorus, and seventeen metals running from antimony to zinc.

Two rows sit above the others, in the group he headed as the substances that may be regarded as the elements of bodies. Lumière and calorique. Light and heat, entered as materials, on the same footing as gold, in the first two lines of the sheet.

Caloric is the fluid chapter twenty-three met, running downhill out of hot bodies into cold ones and doing work on the way. It could be neither created nor destroyed, and Lavoisier and Laplace had measured it in 1783 by the weight of ice a guinea pig could melt. In 1798 Benjamin Thompson, running the arsenal at Munich, set a cannon barrel in a box of water, fitted the boring machine with a deliberately blunted bit, and put two horses on it. The water boiled in two and a half hours, and would boil again the next day out of the same brass, which lost no weight and no capacity for heat. Whatever comes out of one lump of metal without limit, while the metal loses nothing, was never a fluid. Heat is a motion of the parts. The

second row of the most careful table in eighteenth-century chemistry was a process entered as a substance.

That table was drawn up to answer the question the ball puts, and no list drawn up since has answered it all the way down.

35.1 Ten entries

Space is a rank. Twelve readings sit on a patch, and the rule that repairs them cannot tell one port from another, so the twelve directions fall into four bands that no rule respecting the wiring can mix, of sizes one, three, three and five. The band of size one carries no comparison, so nobody inside can read it. The other three fade at rates of their own, and a disturbance in the fastest is down to half in 5.4 passes while the slowest takes 14.7. The quickest question a patch can put about itself runs three seams out and three back, six repairs, by which time most of the fast content has drained, so every observer is a slow ear. What lasts long enough to fix a position is the slow band. It is three wide. There is no container for anything to sit in. There is a count of comparisons. Three independent directions is what that count will carry. Distance is how many comparisons apart two patches are. A position is six whole numbers, one for each pair of opposite ports. An observer reads the band's table of angles exactly off its own records. The carriers tile a screen whose own smooth limit is two-dimensional, the surface a hologram is written on, and the three directions are the rank of each carrier's slowest response.

Time is three objects sharing one word. The first is order, a partial order on committed repairs: one commit precedes another when the later one certifies that it read a version the earlier one wrote. Of the four hundred and thirty-five pairs of seams among twelve observers, a hundred and twenty share an observer and can compete for it, and three hundred and fifteen share none. The wiring fixes those possibilities. The history that runs fixes which pairs actually get ordered. The second is direction, which is the irreversibility of a commit: seven arrangements of a seam go into the move and one comes out, what separated them was in the arrangement the move overwrote, and quantum coherence drains away in the same move. The third is flow, which is what a state does to its own algebra, and the entire supply of rate. Flow is reversible, so it carries no arrow. It belongs to whoever holds the state. There is no timeline for events to be strung along. There is a web of repairs with a direction on each strand. Each observer's rate is a property of what that observer is holding. Clocks tick on events, each observer's on its own, and two of them running at the same average rate disagree in their readings.

In the proposed interpretation, mass is a standing cost. A bounded arrangement of records that keeps re-asserting itself against the seams around it has to be paid for on every cycle, because flattening it would open larger

disagreements at its edge than it closes inside, so repair works on it at every event and gets nowhere with it. The size of that bill is the mass. It resists a push and it bends the geometry around it. Those are one entry read twice. A proton comes to 1.0072764665789 atomic mass units on every scale that has ever been put under one, the uncertainty confined to the last two of those digits, and not one of the records holding it together stays put long enough to be the proton.

An energy-like repair rate counts changes against a specified observer clock. A nonnegative disagreement functional and additive seam bookkeeping provide finite cost and flux identities. They do not by themselves make a relaxation rate a conserved physical energy or a particle rest mass. A coherent generator, its measured spectral response and the clock-to-energy calibration supply the additional interface described in chapter eighteen.

These interpretations require a quantitative connection to the observer's field response. The finite constructions give conditional internal charges and invariant interactions. They do not source-select the observed masses or establish that a repair decay rate is a relativistic mass. The common dynamics, clock calibration and physical matching supply that further test.

Motion is transport of a pattern, and nothing travels. The pattern is re-established at one place in the comparison count after another, each place holding it for a cycle and handing it on, the way a wave crosses a lake while the water goes up and down and stays where it was. What advances is an arrangement. The material carrying the arrangement is different material at every step. Momentum is what the arrangement carries while this is going on, conserved because the repair rule does not care which seam it is applied at.

Force is a steepness. Put one unit of source on the twelve ports and the potential reads one value at the source, a second at every port a hop away, a third at two hops, and a fourth at the port directly opposite, and the drops between those four are eleven sixtieths, three sixtieths and one sixtieth. Eleven sixtieths of the fall happens between the source and its nearest neighbors, and everything past them, out to the far side, is worth the remaining four sixtieths, computed off a list of which port touches which and nothing else. A pattern sitting in a fall like that gets re-established slightly off-center on each cycle, toward the steep side, and how far off-center is the force.

Matter is a defect that no order of repair removes, together with a record saying it is there. Flip one record out of thirty on twelve wired observers and the machine converges from all two thousand and forty-eight starting states under every schedule and leaves exactly one unit of disagreement behind. Repair in another order and it sits somewhere else, at the same total. A defect of that kind is a pattern in the way content moves rather than a lump occupying a spot. It reads the same in every overlap, and survives being regraded to a coarser or finer resolution. The record is the

other half of the entry: some observer has to have committed the fact that the defect is there. That writing is a separate event from the defect it is about.

Charge is a number you pick up going round a loop that will not go away. Each observer's labels are its own business, and the translations sitting on the seams are the only thing anybody can check; chain them round a closed loop and what comes back is a translation from your own conventions into your own conventions. Relabeling cannot reach it, because relabelings leave round trips where they found them. Nor can repair flush it out: repair works one seam at a time, and no single seam holds it. Twelve observers have nineteen independent loops for such numbers to sit in, and what sits there has to balance at every port and be paid for in energy.

In the declared massless-vector description, a gauge constraint and redundancy leave two transverse propagating modes. A constant mode can have zero stiffness for the specified spatial operator. Those are conditional field properties, not consequences of the wiring alone. Calling the modes light requires a physical current, coherent continuation and photon response identified in the same history, with the observer's clock calibrated to that evolution.

A record is a commitment. It is an irreversible write into the part of the algebra where everything commutes with everything, which is what makes it readable twice with the same answer, copyable to a neighbor, and proof against being disturbed by the reading. The flow does nothing to it: the price of commuting with everything in sight is having no clock of your own. Of the ten entries this is the only one that is not a rate or a count, and the one the other nine are written in.

35.2 Rates, counts and one commitment

Read those ten down the page and ask what kind of object each one is. A rank. A relation, a loss and a flow. A cost per cycle. A rate. A reestablishment. A gradient. A residue with a note attached to it. A loop reading. A mode and the rank of a plane. A write.

There is nothing underneath them. None of the ten is a stuff that the other nine are made out of, and there is no eleventh entry held in reserve that the ten are the behavior of. An inventory of this kind is supposed to contain at least one substance. Lavoisier's contained seventeen metals you could pick up and put down again. This one contains a rank, a rate, a residue and a write.

Take the most solid item available and walk down through it. A proton is a standing cost, the cost is a rate of repair, a repair is one comparison between two readings held at the ends of a seam, and a reading is a number an observer holds about a neighbor it cannot see. Four steps. At no step does a material turn up that the numbers are numbers of.

Lavoisier's table had thirty-three rows and thirty-one of them were things you could put in a jar. The two you could not put in a jar were the two he placed at the very top, as the most elementary items on the list, and they are the two he got wrong. The ten entries above finish the correction he was not in a position to make: the jar is empty all the way down, and the inventory is complete anyway.

The world is made of accountancy.

The road to that sentence ran through a bell in an Essex church tower, a bead held in a laser in a dish of water in Canberra, and a nickel target in a laboratory on West Street, and every step of it was a count of something.

Stuff is what it feels like. The feeling has an account too. You are made of the same entries. The hand you put on a table is a set of patterns with a standing cost, the pressure you feel is a rate, the table's refusal to let the hand through is one pattern declining to occupy a state another pattern is holding, and your conviction that you have touched something solid is a commitment written into the part of you where everything commutes. There is nowhere in that arrangement to stand and look at the accountancy from outside, because the eye, the ruler and the place to stand would each have to be built out of more of it.

35.3 The shape of space

Cosmology puts a question to the sky that sounds like the most concrete question anybody could ask. Is space finite or infinite? Does it curve away like the surface of a ball, so that a straight line eventually comes home, or run flat forever, or close on itself the way a doughnut does, joined up in a direction nobody can point at? There are instruments aimed at it and a number attached to it. The number comes back flat to within a fraction of a percent.

Take the question apart before answering it.

It asks for the shape of an object. Being a ball or being a doughnut is a property of a set of points: which points there are, and which of them lie near which. Chapter fifteen got the three directions as the rank of the slowest band a repair rule leaves standing, and the first entry on this chapter's sheet says what that buys, which is a count of comparisons. A rank is not the kind of thing that can be a ball. There is no set of points anywhere on the inventory for a shape to be the shape of.

It also asks for the shape now. A shape at an instant needs the instant. Chapter seven went looking for one and came back with a partial order: one record depends on another when it read what the other wrote, and two records with no chain of such reads between them have no fact about which came first. Nothing in that hands over a slice through everything that two observers would draw the same way. So the question wants a space and a moment. The sheet carries neither.

A question whose terms name nothing does not get a false answer. It stops having one. Space is not a ball, and it is not a doughnut, and it is not infinite either, in the same way that the ether was not moving slowly.

Which leaves the instruments, because they are measuring something. Observers hold records. Where two of them overlap, the records agree or they get repaired until they do. Glue the compatible ones, coarse-grain until individual seams stop being visible, and out comes a geometry: distances, angles, curvature, the whole apparatus chapter twenty-five assembled, standing in for a very large number of separate agreements. That description has a shape. The shape belongs to the description, to the coarse-graining that produced it, and to the slicing whoever took the reading chose. It is an effective account of a virtual slice. Every cosmological measurement ever published is a measurement of it. Ask what shape space has and the answer has to name a resolution and a slicing before it can begin.

Parts of that reconstruction can be written out. Chapter fifteen gave records positions in a three-dimensional space, and chapter seven gave them an order when one record reads another's value. Chapter seventeen gave them rounds and a reach and got a four-dimensional spacetime out of the count, with volume and duration read off the completed records and the relay work left out. A field on those positions settles into a smooth wave as the population refines.

The screen gets the same treatment from a single continuing history of repairs. One operator comes out of that history, and it governs how strongly the field at two places is correlated, how fast a disturbance fades, and the waves the screen carries, whose fronts travel at a finite speed. That operator lives on the two-dimensional screen, and its parameters count observations: seconds come from a clock laid against them. Under chapter twenty-five's entropy-stationarity hypotheses, the balance on small balls constrains how geometry responds to matter. The fixed product-background wave examples do not establish that response law.

Gravity sits on the same finite events. Put a geometric tensor and a stress tensor on them and ask the two to balance along nine fixed directions. Together with symmetry, conservation and connectedness, those nine balances give balance along every null direction and an identity of exactly Einstein's form, with one constant term proportional to the metric. That constant term is the one chapter twenty-five's light rays cannot see.

Is that manifold out there, or is it how things look from inside? It comes apart into two halves, and each half has a plain answer.

The certified record order is shared. A writer's mark and a value settle which record a read consumed, and a fixed counting rule gives the same total to every checker. As the records refine, the fraction of ordered pairs inside a diamond approaches one tenth, the four-dimensional value, for everybody at once.

The split into three and one belongs to each observer. An observer moving along a path through spacetime has a three-dimensional rest space perpendicular to it, and the counts supply durations along the path. Two observers cut the same four-dimensional geometry differently, the way the guard and the passenger cut the lightning. A present spread across a rest space belongs to that cut.

So an observer has a position from its records and a history from their dependencies. Fourth roots of counts give it durations, and a clock carried along a moving path collects fewer of them, by the factor the flying clocks brought home. The flow its state determines and the direction its records run are two further structures, and the observer's event clocks are where they meet: each repair at its region is one event in the order, one step of the flow and one more write that cannot be taken back.

The same container picture is buried in the other question everyone asks. Galaxies recede, the recession runs faster the further out you look, so wind the film backwards. The scale shrinks. Keep winding and it reaches zero, so everything was at one point, at one moment, and that moment was the beginning. Every step of that is sound arithmetic performed on the effective description. What the last step needs, and helps itself to, is a global clock to run backwards and a spacetime to shrink.

What the reconstruction is made of is records held today, inside finite patches, extrapolated. The hot dense era at the far end of that extrapolation is as well evidenced as anything in the subject. Nothing here disturbs it. The step that does not follow is the one that converts a limit of an effective description into a first moment of the world, because a first moment of the world is a reading on a clock that no observer holds and that nothing in the arrangement ever started.

Chapter thirty-seven is where the word first gets taken away properly. The smaller point will do here. Both questions are the container picture asking to be let back in, at the edge and at the beginning.

35.4 The Link Trainer

In 1929, in his family's organ factory in Binghamton, New York, Edwin Link built a stubby wooden aircraft that went nowhere. Organ bellows underneath it, driven by an electric pump, pitched and rolled the box while the pilot inside worked the controls, and the instruments on the panel moved the way instruments move in the air. The first customers were amusement parks. In the spring of 1934 the Army Air Corps was carrying the mail and losing aircraft to weather, and Link flew in to a demonstration in conditions nobody else would fly in. They ordered six, at three thousand five hundred dollars each.

That box is a simulation. The reason has nothing to do with the bellows. It is a simulation because flying exists. There are aircraft, in the air,

doing the thing the box imitates. The box is pointed at them. Take the aircraft away and the trainer does not become more authentic; it becomes a wooden cabinet that pitches and rolls. Imitation is a relation with two ends, something standing in for something else, and with one end missing the word has nothing to grip.

So put the question to the list. Which of the ten entries is standing in for something else? Where is the original that a repair rate is a copy of, and what is a loop residue an imitation of? Chapter two ran two rules on a string of letters, and the strings that came out were not descriptions of realer strings kept in a warehouse somewhere; you ran the rules, and what came out is what there is. Every entry above has that character. Nothing on the sheet points outward at an original.

Outward is the load-bearing word, because a second question hides inside the same objection and that one has an answer. A structure can hold a description of itself while nothing in it stands in for anything, since standing in takes two objects and self-description has one. The description has a subject. The subject is the thing holding the description. Chapter thirty-seven is where that gets demanded and priced. Chapter twenty-nine computes specified arithmetic candidates, while their physical electromagnetic identification remains open. The interpretation pursued here is that there is no second copy for the first one to be a copy of, so the accounts have to settle inside.

The other half of the objection is the word computed, used as though it were a synonym for fake. A thing that is computed is a thing that happens. When a repair runs at a seam, a disagreement that was there has gone afterwards, the total is where it was, and seven possible pasts have been discarded, which is an event with consequences rather than a picture of one. The same holds for a magnetic field and for the taste of coffee. Both are computed in the only sense of the word this world supplies, and both occur. There is no clock outside the arrangement stepping it along, either: the eighty-one thousand nine hundred and twenty patches of chapter ten worked every seam between them into agreement with no tick anywhere in the machinery, each region advancing at the rate its own state manufactures. There is no next frame, and nothing anywhere that would be producing one.

35.5 The reader

Every entry above came apart into something more basic than itself. One item has not.

Look at what the ten have in common besides being counts. A rate is counted against somebody’s clock. A rank is the rank of somebody’s table of comparisons. A distance is a count of comparisons, and a comparison takes two parties that hold something to compare. A record is a write into

somebody's center. Charge is what a loop reads on the way round, and a loop is a walk from one observer to the next. Every quantity on the sheet is a reading, and a reading takes something that reads.

That something has not been dissolved. The reason is structural: the dissolving was performed with it. Chapter three asked what the job of holding a description requires and got three plain items, a bounded view, somewhere to write, and a reading that makes a difference, all three of them met by a thermostat screwed to the wall of a classroom in Whitewater, Wisconsin. Chapter nineteen took the same object and gave it four: a local algebra of questions, a record algebra sitting in the center of it, an interface that reads and updates the same way every time, and enough of a state to fix the odds on the next question. Nothing in the chapters since has explained either version away, because every one of them was built out of it.

Which is what the first page meant by putting observers first, arrived at from the far end of the argument. The stage was assembled out of what bounded record-keepers can consistently write down about each other, in that order, and the ten entries are the assembly. Take the record-keepers out and there is nothing to take them out of.

Every number quoted in the last thirty-three chapters was obtained by observers standing inside the thing they were measuring, with instruments assembled out of the ten entries. Lavoisier weighed the elements on a balance made of them. Nobody since has found a way to step outside and check.

Chapter 36

Why These Laws and Not Others?

"Why does the electron weigh what it weighs?"

"Because that is what it weighs. Somebody measured it."

"Why that number and not another one?"

"Because there are a great many universes with other numbers in them, most of them empty, and this is one of the few with anybody in it to ask."

Those are the two answers usually on offer, for the electron and for every other number chapter one counted, the ones physics takes from laboratories and enters by hand, nineteen of them or twenty-six depending on how neutrinos are handled. The numbers are brute facts and there is nothing to say about them, or they were drawn from an enormous collection of alternatives and we see this set because the others have nobody in them to look. Both answers end the conversation somewhere the reader cannot go. The first says there is nothing behind the numbers and does not say how it knows. The second needs a shelf with the other worlds standing on it and does not say where the shelf is.

A third kind of answer was read aloud in London in the summer of 1858.

On 18 June 1858 a package reached Down House in Kent, and Charles Darwin opened it and found twenty pages of his own theory written by somebody he had never met. Alfred Russel Wallace had drafted the essay on Ternate, in the Moluccas, between bouts of fever, called it "On the Tendency of Varieties to Depart Indefinitely from the Original Type", and asked that it be passed to Charles Lyell if it seemed worth anything. Darwin had been assembling the same argument for twenty years, through barnacles and pigeon fanciers and a filing system of small stubborn facts, and he wrote to Lyell that day. Your words have come true with a vengeance. If Wallace had my manuscript sketch written out in 1842 he could not have made a better short abstract. All my originality, whatever it may amount to, will be smashed.

Ten days later his son Charles Waring Darwin, eighteen months old, died of scarlet fever. On 1 July, Charles Lyell and Joseph Hooker took both accounts to the Linnean Society of London and put them before the meeting, where the society's secretary, John Joseph Bennett, read them aloud. Darwin was at Down with his family, burying the child. Wallace

was in the Malay Archipelago and learned that the meeting had happened months afterward, in a letter. The reading drew no discussion.

Thomas Bell, the society's president, reviewed the year in his address the following May. The year which has passed, he said, has not indeed been marked by any of those striking discoveries which at once revolutionize the department of science on which they bear. Bell had been in the chair on the evening of 1 July.

What Bennett read aloud was a procedure for producing the appearance of design with nothing in it that chooses: variation happens, some variants leave more descendants, and the arithmetic does the rest. It has an application in physics that has nothing to do with finches, and the question it applies to is the one underneath chapter one's list: why the world runs on these rules with these numbers instead of some others.

36.1 What a description is for

In 1950 Vitaly Ginzburg and Lev Landau wrote down an account of superconductivity that mentions no electrons at all. They introduced one quantity standing for how much of the metal has gone superconducting at each place, expanded the energy in powers of it, and read off the two lengths a superconductor has: how far the superconducting state can vary over, and how deep a magnetic field soaks in before the metal expels it. The account worked near the temperature where superconductivity sets in and nowhere else. The constants in it were fixed by measurement, because nothing in the argument could produce them.

In 1957 John Bardeen, Leon Cooper and Robert Schrieffer explained superconductivity from underneath: electrons pair up through their coupling to the vibrations of the lattice, and the pairs move as single objects. Two years later Lev Gor'kov showed that the 1950 account is what the 1957 theory becomes near that temperature. The constants nobody had been able to calculate came out in terms of the pairing. The 1950 account did not become wrong when its reasons arrived. It stopped being the bottom and stayed in the textbooks, because near the transition it is the easier of the two to compute with and gives the same answers.

That is an **effective description**: correct inside a range, carrying constants it does not generate, and usually able to say where it will fail. Ginzburg and Landau's account announces its own ceiling, since the expansion it rests on is only good while the superconducting quantity is small, which is to say near the transition and not far below it. Physics is made of descriptions like that and there is no defeat in having one. The Standard Model is an effective description of the same kind. The complaint chapter one lodged against it was about direction rather than count. Numbers go in and predictions come out. No arrangement of it sends a number the other way.

36.2 The planet that was easier to move

Chapter thirty-four left one leftover in the solar system: 43 seconds of arc per century in Mercury's orbit that no amount of Newtonian bookkeeping supplied, about a hundredth of a degree in a human lifetime. The nineteenth century tried to account for it with an extra planet. The man who proposed the planet had done it once before.

Urbain Le Verrier had found Neptune on paper in 1846, by taking the anomalies in the orbit of Uranus and computing where an unseen body would have to sit, and the telescope at Berlin found it that night within a degree of the position he gave. So he proposed another one, inside Mercury's orbit, hot and small and hard to see against the sun. A country doctor at Orgères-en-Beauce named Edmond Lescarbault wrote to say he had watched a dark spot cross the face of the sun in March of that year. Le Verrier travelled out to interview him, came back convinced, and the planet was named Vulcan.

Vulcan was never there. Each failed search moved it rather than killing it: smaller, closer in, differently tilted, and in one version broken into a belt of bodies each too small to be seen, which no observation could exclude. Nothing in the proposal forbade those moves, because a planet comes with a mass and an orbit and both are dials. During the total eclipse of 29 July 1878 two experienced American observers, James Craig Watson at Separation in Wyoming and Lewis Swift near Denver, both reported a reddish object near the sun. The two positions did not agree with each other. Nobody else at that eclipse saw anything. Eclipse expeditions went on looking through 1908.

In November 1915 Einstein computed the same 43 seconds of arc from field equations with nothing adjustable anywhere in them.

Vulcan accommodated everything and predicted nothing, and closing the search took fifty-six years. Flexibility is what lets a description survive contact with data. Every bit of it is subtracted from what the description tells you. Chapter one counted the inputs. Each measured number a description absorbs is a fact about the world that description will never predict. Le Verrier's planet had two knobs, a mass and an orbital radius, and between them they could reproduce any perihelion shift anybody came back with, which is why no measurement of the perihelion shift could ever have got rid of it. The trade stops paying at the point where nothing anybody could see would tell against the description, and Vulcan reached it when the belt version arrived. From then on the only property the planet had left was that it could be moved. Einstein's equations had none of that, which is why 43 seconds of arc could have killed them.

36.3 What an assumption costs

In 1978 Jorma Rissanen published a paper called “Modeling by shortest data description”. Its rule for choosing between accounts of the same data is to add two lengths: what it takes to write the account down, and what it takes to write down the data once you have the account. Take whichever account makes the two together smaller. Rissanen’s name for that sum is the description length. An explanation is a compression. A rule that costs more to state than it saves has explained nothing.

The inverse electromagnetic coupling read at rest is 137.035999177. The shared-width trial map in chapter twenty-nine returns 137.035660, about two and a half parts per million away. Its certified error bounds exclude numerical rounding as the explanation, while the missing physical matching has not been shown to remove the mismatch.

The package from Ternate is the same accounting seen from the other end. What Darwin had been accumulating for twenty years was data: island species, barnacle anatomy, fossil beds, the breeding records of pigeon fanciers who had made fantails and pouters and tumblers out of the rock dove, geographical ranges that made no sense on any other reading. Stored as itself, that is a library. The sentence that variants which leave more descendants become more common adds no fact to the library. It shortens it, by removing the need to hold each adaptation as a separate item. That shortening is why two men at opposite ends of the world wrote the same sentence out of two entirely different collections of animals.

36.4 Reading the collection as a sieve

String theory gave up point particles. Its known difficulty is the count. Curling the extra dimensions is a choice, threading fluxes through them is another, and choosing bundles and branes and hidden sectors are more. The product of all those choices is an enormous collection of complete specifications for a low-energy world, each with its own particles and its own constants. Somewhere in that collection, the hope goes, is one that looks like ours. It also holds worlds with four families of matter and worlds in which the proton comes apart in a second.

A target turns the collection into a stock list. The conditional overlap constructions motivate a comparison target: the global gauge group, three colors, three declared families, the hypercharges, a specified Higgs sector, no unwanted light matter, and the absence of the usual grand-unified gauge-boson decay route. The matter lift, global quotient, family attachment and scalar-sector choices are stated premises of that target; overlap consistency has not source-selected them all. The exact group is the item that does most of the rejecting, because two constructions can match each other on every local property and disagree about which charges are allowed to exist

at all, and a charge is the quantity an observer's records are keeping count of. Take that target to the collection and reject everything that misses it.

In 2006 Vincent Bouchard and Ron Donagi published a construction that gets a long way down the list. Curl the extra dimensions on a particular six-dimensional shape, one of the kind called a Calabi-Yau and one with a loop in it that cannot be shrunk to a point, and hang on it a structure whose symmetry is the smallest grand-unified group there is. Then carry that symmetry around the unshrinkable loop and compare it with what you set out with, which is chapter twelve's round trip performed on a larger object, and the mismatch breaks the big group down to the visible one. Out of it comes the exact group, three generations, no extra light matter the experiments would have caught, and a region of the settings with exactly one Higgs pair in it. The group, the three generations, the hypercharges, the Higgs count and the absence of exotic matter all check out. Then there are the dials.

Count them. Three kinds come with the construction, one kind setting the sizes of the curled directions, one the shape they are curled into, and one the structure hung on top, and the published data for that one-Higgs region record eleven of the first, eleven of the second and forty-nine of the third. Seventy-one complex directions, a hundred and forty-two real ones, counted before anything has been done to hold the vacuum in place. The comparison against measurement uses five real numbers: two coupling strengths read at the Z boson's mass, the vacuum value of the Higgs field, the Higgs mass and the top quark's mass. The comparison consumes none of the construction's own values, so the table of how that comparison responds to the hundred and forty-two dials is a hundred and forty-two columns of zeros. The map has rank zero. Any smooth comparison that returns five numbers leaves at least a hundred and thirty-seven directions unpinned, which is a hundred and forty-two minus five and is unrelated to the hundred and thirty-seven of chapter twenty-nine.

So the construction matches the target on structure, and the published data supply no formula linking its seventy-one dials to the five numbers. The only comparison anyone can run ignores the dials, and a comparison that ignores the dials pins none of them, which is what rank zero says.

36.5 Testing a specified world

A specified vacuum that disagrees with the data is ruled out, and the collection has an escape ready. The move available is to go and stand at another vacuum, and with a collection that size on the shelf there is always another one to stand at. Vulcan was moved every time a search came back empty. A measurement that retires one vacuum leaves the collection as full as it was.

The move is available because there is an outside: a space of alternatives sitting outside any particular world, which a description can be relocated into whenever the world it was sitting in fails to match the data.

There is no such space here. The reason is arithmetic. Chapter twenty-nine read one cell of an observer's screen twice, from the outside as an area and from the inside as the smallest electromagnetic act available, and the two readings are readings of one quantity, so setting them equal gives an equation whose unknown appears on both sides. One number satisfies it. Change that number and the two readings disagree. There is no third place to put the difference and no neighboring world to carry it off.

Exactly one comes from the property chapter twenty-nine proved about the equation. Feed it two trial grains some distance apart and it hands back two grains at most 0.0724 of that distance apart, so the passes squeeze rather than spread, and every starting guess in the domain walks to the same destination. A second destination would be a second world with a different fine-structure constant in it. The squeezing leaves nowhere for one to sit. The loop comes in the two versions chapter twenty-nine laid side by side, the bare one and the one that recomputes the weak and strong forces' shared width inside it, and each has its own single fixed point. Neither fixed point is yet the laboratory's number. The bare loop lands three parts in ten thousand short and the width loop two and a half parts per million short, and carrying the bare root down to the laboratory's low-energy reading with the measured quark contribution misses by half a percent. The grain used for comparisons is located from the measured coupling, and which map is the physical one remains open. If one of them is, what comes out of the equation is a reading of the architecture in the same way that the diagonal of a square is a reading of the square. Nobody chose the fine-structure constant any more than anybody chose the square root of two. The panel of dials that was supposed to have been set at the beginning is missing along with the beginning, and the choice between the two versions of the loop is the one place left for a dial to sit, in plain view.

36.6 Scoring a rule

Chapter nine derived the repair move from three requirements. Every move that fails one of them is a rule somebody could have written down, and rules that get written down compete.

Three quantities, each of them a count of something. The first is schedule stability. Take a starting arrangement, run a rule until it stops under one order of repairs, run it again under a different order, and ask whether the two settled readings agree; do that across many starting arrangements and many orders, and the fraction that agree is a number between zero and one. Chapter six's twelve observers ran from every one of two thousand

and forty-eight states under sixteen schedules and landed every time on the same arrangement, which is that number coming out at 1.

The second is observer yield: how many observers a rule produces. A region counts as one when its record reads the same at the end of a stretch of time as it did at the beginning, and when what it holds carries information about what its boundary does next. Both halves are measurable, the first as how far the reading has drifted and the second as how much the record tells you about the neighbors' future. Count the regions that qualify and take the average over the running of the world.

The third is the cost of writing the rule down, in Rissanen's sense, and enters with a minus sign. A rule's score is the first two with positive weights, minus the third.

The three pull against each other. A rule that repairs nothing scores a perfect 1 on stability, because every order of doing nothing agrees with every other order of doing nothing. It is about as cheap to write down as a rule can be, and yields not one region holding a record about anything, so the second term deletes it. A rule that buys the second term by copying down everything it ever sees pays for the copies in the third. The rule chapter nine derived, which moves two disagreeing readings to their midpoint and touches nothing else, is where the three pulls balance.

The rules are competing for one finite thing, which is the capacity of the structure to repair itself. A rule's share is the fraction of that capacity being spent its way. A rule scoring above the average grows its share at a rate proportional to how far above the average it sits, and one below shrinks the same way. That is **replicator dynamics**, arithmetic for the sentence that fitter types spread. Nothing in it requires universes to reproduce.

Write f for a rule's score, put a bar over it for the average of the scores across the rules holding capacity, weighted by their shares, and let t measure how far the reallocation has run. Ask how that average moves as the shares move, and the answer is one line.

$$\frac{d\bar{f}}{dt} = \mathrm{Var}(f)$$

The left side is the rate at which the average score changes. The right side is the variance of the scores: the average of the squared distances of the individual scores from the average score. So the average climbs at exactly the rate at which the survivors disagree with each other, and since a squared distance is never negative, the average never falls.

Two rules make it concrete. Scores of 3 and 1, half the capacity each, and an average of 2. The first gains half a share per unit of t and the second loses the same. The variance is 1, and the average is climbing at 1. As the first takes over, the spread closes and the climb stops, because the mean rises only on disagreement and by then there is none.

What survives that is a rule whose answers do not depend on the order anybody works in, which is thick with regions that hold stable predictive records, and which is cheap to state. Chapter six measured the first of those on twelve observers, and chapter nine derived the move that supplies it.

36.7 An ensemble of one

The anthropic argument runs as follows: there is a large collection of worlds with different settings, the great majority are barren, ours is one of the few in which anybody could be around to take a reading, and no further explanation of the settings is needed. As selection reasoning it is sound. It requires the collection.

Take the phrase "the largest prime number". It has the grammar of a name and there is nothing at the end of it, and Euclid's argument shows why: take any finite list of primes, multiply them together, add one, and the result is divisible by none of them, so no finite list is all of them. The phrase does not pick out a number that turned out to have some other property. It picks out nothing.

The phrase "a universe with a different fine-structure constant" is of that kind. The two readings of a cell fix that number by requiring agreement with each other. A structure whose readings disagree does not settle into a world that observers then fail to appear in. It does not settle at all, and the phrase, like the largest prime, picks out nothing.

So, if the physical map is one of these, the ensemble was never larger than one. An anthropic filter has nothing to filter. The selection that did happen ran over repair rules rather than over worlds, and from inside.

Running from inside has a price. The capacity being competed for is the structure's own, the observers being counted include whoever is doing the counting, and the rule that takes the whole share is the rule the judges are running on. Darwin's argument reached Down House in an envelope from Ternate, carried halfway around the world by a post office, so there was a morning before it arrived and a morning after. There is no morning before this one. Nothing was posted and nothing was opened, and whatever holds the arrangement up is somewhere inside it.

Chapter 37

Why Does the Loop Have No Beginning?

The room was emptying. It was the afternoon of 7 September 1930, the last day of a three-day conference on the epistemology of the exact sciences that Kurt Reidemeister had organized at the University of Königsberg, and the closing item was a roundtable on the foundations of mathematics with Hans Hahn, Rudolf Carnap, Arend Heyting, John von Neumann, Arnold Scholz and Reidemeister himself at the table. Everyone had said their piece over the previous two days: Carnap for Russell's programme, Heyting for Brouwer's, von Neumann for Hilbert's. The proceedings went into the journal *Erkenntnis* the following year, volume 2, pages 87 to 190, with the roundtable transcript at pages 135 to 151, so it is possible to read what was said and in what order.

Near the end, a twenty-four-year-old Viennese logician remarked that one can give examples of statements of arithmetic that are true and that no proof inside the system reaches. Kurt Gödel had come to Königsberg to present a completeness theorem, the result he had built his reputation on, and he left having mentioned in passing the result that closed the question the three days had been about.

The discussion moved on. Almost nobody in the room reacted. One person did: von Neumann cornered him afterward, wanted the construction, went home and thought about it for two months, and wrote to Gödel on 20 November with a further consequence that Gödel had reached first.

The next morning the city of Königsberg made David Hilbert an honorary citizen. He was sixty-eight, he had been born there, he had retired from Göttingen that year, and he addressed the Society of German Natural Scientists and Physicians on knowledge of nature and logic. He closed with six words that were later cut into his gravestone: we must know, we will know. A recording of the ending survives, 373 words read for local radio and running four minutes. The answer had been supplied the previous afternoon, in a smaller room, at a table of seven, one of whom was listening.

What Gödel had was a construction. Hand it a system that can talk about its own formulas and it hands back a sentence about that system, assembled out of the system's own material and pointing at nothing outside it. A world made of observers comparing records leaves a bill that no

amount of further comparing pays. It has to settle that bill the same way, out of its own material. Something has to be doing the comparing. Where did the network come from, what is it running on, and what started it?

37.1 A program that reads its own text

Every programmer has wanted the same tool at some point. Something you can point at a piece of code that tells you whether the code finishes or runs forever, and answers by deciding rather than by waiting, correctly, on any program you hand it.

Suppose you have it. Call it the oracle. It takes a program's text along with an input, answers finishes or runs forever, and is never wrong.

Then write the following, which takes about four lines. It reads a piece of program text. It asks the oracle what that program does when it is fed its own text as its input. If the oracle says finishes, it goes into an infinite loop. If the oracle says runs forever, it stops immediately.

Hand that program its own text.

If it finishes, the oracle must have answered runs forever, since that is the only answer that makes it stop, and the oracle was wrong. If it runs forever, the oracle must have answered finishes, and the oracle was wrong the other way. There is no third case, so there is no oracle. Alan Turing published that argument in 1936.

Look at what did the work, because the contradiction is the disposable part. The four-line program is one whose behavior is the opposite of what is predicted about it. Hand somebody the transformation "do the opposite of the prediction" and the construction hands back a program satisfying it. Feeding a description into the thing it describes is a way of solving an equation. The equation is chapter two's: return what you were given.

The calculator in chapter two hunted for a number the operation left alone and found 1.6180339887. The same demand, made of descriptions instead of numbers, has the same kind of answer: a description the transformation returns unchanged. That is a fixed point, and self-reference is the construction that builds one. Chapter two's warning travels with it, because a fixed point can be a place nothing ever arrives at.

37.2 Four rows and a diagonal

The move underneath the oracle is older than computers and fits on an envelope. Georg Cantor published it in 1891, four pages in the first yearbook of the German mathematical society, of which he was the first president. It runs on a numbered list.

Write four strings of four bits, one per line.

row 1: 1 1 0 0 row 2: 0 0 1 0 row 3: 1 1 1 0 row 4: 0 1 0 1

Walk the diagonal: the first bit of row one, the second bit of row two, the third of row three, the fourth of row four. That reads 1, 0, 1, 1. Flip every bit you landed on and you have built 0, 1, 0, 0.

Check it against the list. It differs from row one in the first place, which is where you flipped row one's own bit. It differs from row two in the second place, from row three in the third, from row four in the fourth. Four rows, four disagreements, one per row, each one forced by where the bit was taken from.

Nothing in the walk cared that the list was four long. Run it down a numbered list of strings that never end, one string per whole number, and what you build differs from the seventeenth string in the seventeenth place and from the millionth in the millionth, so it sits on no line of the list. A minute of bit flipping. No table of any size holds every string.

Turing's four lines are that walk with programs on the list and the oracle's prediction as the bit. Gödel's sentence is the identical construction pointed at proof instead of at halting. Number the formulas, so that a statement about numbers is also a statement about formulas, and build the sentence whose content is that it has no proof in the system. If the system proves it, the system proves a falsehood. If the system does not, the sentence says something true that no proof reaches. The lever was never a pun on the word this.

The numbering is the engineering and the self-reference is what falls out of it. Give every symbol a number. A formula is a list of symbols, so code the list by prime powers: symbols numbered three and one give two cubed times three, which is twenty-four, and twenty-four factors into primes one way only, so the code hands back three and one. Proofs are lists of formulas and get their numbers the same way, so the relation between a proof and what it proves is arithmetic, and "no number codes a proof of the formula coded by twenty-four" is a statement about numbers that a system able to do arithmetic can write down.

Getting a sentence to carry its own code with no regress is Quine's device, and works in English with no numbers in it at all. Take a phrase, write it once inside quotation marks, then write it again outside them.

"is twenty words long when appended to its own quotation" is twenty words long when appended to its own quotation.

Count. The quoted phrase is ten words and the phrase after it is the same ten, so the sentence is twenty words long, which is what it says about itself. It does that without the word this and without a second sentence propping it up.

Gödel's sentence is built that way. It gets called a paradox. It is not one. The liar sentence, the one saying that it is false, has no consistent reading: true makes it false, false makes it true, and the machinery seizes. Take Gödel's instead and suppose the system proves it. Then a proof of it exists, so what the sentence says is false, and the system can read that

proof and derive the opposite of what it just proved. It contradicts itself, which is chapter two's disease, buying everything and ruling nothing out. So a system that contradicts nothing does not prove that sentence, which is what that sentence says about itself. The sentence is true.

That is what the table of seven was handed: a consistent system with a truth inside it that its own proofs do not reach. Hilbert wanted consistent and complete, and completeness is the half that failed.

37.3 Two hands and a sheet of paper

In January 1948 M. C. Escher printed a lithograph, 28.2 by 33.2 centimeters, showing a sheet of paper lying on a surface. Two shirt cuffs are drawn on the sheet in flat gray pencil. Out of each cuff a hand rises off the paper in full modeling and shadow, and each hand holds a pencil, and each pencil is drawing the cuff of the other.

Chapter six borrowed the impossible staircase, which this artist put into a lithograph twelve years later. The staircase fails at the closing: every individual step up is unobjectionable and the loop cannot be assembled. *Drawing Hands* fails nowhere. Every line in it is a line an artist could draw. Both hands are drawn and both hands are drawing, and cover any part of it you like and what is left is consistent. What the picture lacks is a ground floor. Neither hand is the real one. There is no corner of the composition to stand in and ask which was drawn first.

Douglas Hofstadter named the arrangement in *Gödel, Escher, Bach* in 1979 and spent seven hundred pages on it. A **strange loop** is a hierarchy that climbs through its levels and arrives back at the level it started from, with no level underneath holding the rest of them up. The staircase is what a strange loop looks like when it fails. The hands are what one looks like when it holds. Everything remaining in this chapter is the second picture with something else in the cuffs.

37.4 The description and the thing described

Chapter eleven put a name on what a network of observers holds in common. The record map takes the full state and returns the part that survives comparison, which is every seam entry and nothing that only one observer could have known. That list has a fiber, the set of consistent states carrying it, and when the fiber holds exactly one member, that member is the observable normal form. Objective reality is that state, up to changes nothing can see.

The smallest case runs to the end. Put two observers on one carrier, overlapping on a shared port, and let each prepare its ports, keep its own clock, read its own field and act on what it reads, all on one continuing history of repairs. The two accounts are one law. Every record either

of them keeps, every tick and every timeout comes out exactly the same whether the history is computed repair by repair or through the effective description an observer would write down for it. The same holds a level up. On the refining screen, with the radial direction the construction declares, any finite collection of an observer's clock, field and response readings is matched, as closely as anyone asks, by a field moving in one smooth, unchanging spacetime with one time and three space directions. That is the kind of description relativistic field theory writes down, and the observers inside the construction get to use it.

Every object in that sentence about objective reality is a description or an operation on one. The record list is a description, and a description can fit no world, one world, or several.

So point it at everything. Take the whole structure, records included, as one object, and apply the operation to it. What comes back is what that object's own description specifies. The demand is that the two agree.

The universe is the observable normal form of its own description.

One line covers the calculator, the four-line program, the sentence carrying its own code and the demand just made of the whole structure, and the same unknown stands on both sides of it.

$$x = f(x)$$

Here x is whatever is being solved for and f is the operation applied to it. On the calculator, x was a number and f was divide one by it and add one. Applied to everything, x is the whole structure with its records and f is take the description and return the world it specifies. Self-reference is that equation, and closure is that equation with the universe in it.

Chapter eleven's three verdicts apply here as well. A description can fit no world, one, or several. Closure asks for the case where realizing the structure's own description gives back that same structure, with every observation and intervention inside it preserved.

A finite log does not pin down the whole network that produced it. In a network of 1,280 twelve-port carriers, the recorded repairs and reads use 86 of its 7,650 connections. Reconnect two unused pairs and every recorded operation survives, although the number of triangles in the complete network changes and the two networks cannot be renamed into one another. Their local states, ports, readback and protected records support the same observed history. A self-reading world reads more than one log, and closure asks that everything the structure records, across the whole of its history, fixes the structure.

The requirement constrains a world and its self-description together. A finite observer inside it keeps a useful partial account without recovering the whole architecture, and every observer does.

Nothing outside supplies the description. Asking what switched the universe on assumes an outside to switch it on from. The demand that picks out this object is the demand that there is no outside. Anybody asking has to supply the outside themselves. Every outside anybody has supplied has needed one of its own.

There is a dividend in that which nobody went looking for. A description obliged to specify the structure it came out of cannot leave a number loose, because a loose number is a second structure answering to the same description, and the fiber has to hold one member. Chapter twenty-nine is where that gets cashed. The grain and the fine-structure constant are written in terms of each other, and the grain also fixes the record capacity and, through it, the cosmological constant. Each version of that loop admits one solution, so a quantity that every other physics has to measure and enter by hand becomes, for each declared map, a computed number. Selecting the physical map and closing the gap to the laboratory reading are still open; the working grain used for comparisons is located from the measured coupling. Constants are what a world with no outside is not free to leave open.

37.5 Three orders

The word loop invites a picture of something traveling backward into its own past. Three different orderings are in play. Every paradox that gets built out of the loop comes from confusing two of them.

The first is complete-history closure. The equation fixes a whole structure under a reconstruction map and requires no external time in which someone first solves it. That structure contains an internal causal order, changing local states, clocks and computations. The intrinsic interactions that realize these dependencies and the timeless self-consistent history are two descriptions of the same reality.

The second is an observer's record order, which reads depend on which writes. Actual clocks and acts of construction occur within that history. Each observer's clock ticks on its own events, and the flow its state determines turns one step at each of them. Order, record formation and duration are different parts of the account.

The third is a solver's repair sequence, the order in which a computer running the network happens to make its moves. A solver step is not a second. Repairs with no dependency path between them have no order in the causal record, even when an executor lists them one after another.

Set the three side by side. First they work out the architecture, later they build it, is a statement about the record order inside the history. The closure equation adds no earlier builder outside it. Recovered specification, intrinsic realization and inhabited world are one complete structure, its in-

ternal precedence and its responses to interventions included, and replaying one observed trajectory is a smaller thing.

37.6 A thousand machines and one machine

Run the network on one computer. One process, one memory, one loop that picks a seam, checks it, repairs it if the two sides disagree, and goes round again. It ends where chapter ten said it ends. The endpoint does not depend on which seam it picked first.

Then run the same network across a thousand computers, which is what anybody does in practice with eighty-two thousand patches. Cut the patches into a thousand pieces, one per machine. Each machine holds its own patches plus a copy of the seam data along its border, refreshed by messages that arrive late and out of order. Machines take checkpoints. One dies and restarts from its last checkpoint, replaying updates it had applied before it fell over. A supervisor notices another machine falling behind and re-cuts the entire partition to move work off it.

Anybody who has run a large computation knows what the second version looks like from the operator’s chair. The logs differ. The timings differ. Two runs of the identical setup interleave their operations differently. Neither interleaving matches anything the single machine does.

Whether the two are the same universe takes one bookkeeping move to settle: say exactly what counts as the state of the world and what is scaffolding around it. Keep the authoritative record each patch holds. Quotient by the changes nothing can see, which chapter eleven’s group names. Throw away the message queues, the retry counters, the labels saying which machine owns which patch, and any wall clock the machines happen to be reading. What is left is a state of the plain single-machine network. Every readout an observer inside can perform is computed from that.

Then classify what a distributed run is permitted to do, and it comes to three things.

A commit, which projects onto a legal repair path of the single-machine network. Something disagreed, a patch fixed it, and the fix corresponds to a repair the one machine could have made.

A stutter, which projects onto nothing at all. Delivering a message, refreshing a border copy, taking a checkpoint, preparing a transaction, aborting one, starting a machine, stopping a machine, replaying an update that has been applied once, and every byte of metadata describing how the work was cut up. The projected state before and after is the same state.

A rollback to an earlier committed state, carrying the name of the committed state it returns to.

That exhausts the list. The rest is an induction with three cases. Chapter ten’s result is that the endpoint of the repair process is constant across

everything reachable: if legal repairs get you from one state to another, both have the same endpoint. A commit moves along legal repairs, so the endpoint is unchanged. A stutter moves nowhere, so the endpoint is unchanged. A rollback lands on a state whose endpoint was the same one. Start the distributed run and the single-machine run from the same initial state, and after every event in the run, including the last, the endpoint is the same.

So when the distributed run finishes, its terminal normal form is the single-machine one. Stutters and rollbacks never finish anything on their own; the commits do the work.

A checkpoint survives the trip as well. Serialize a run halfway through, records and quantum state together, import it on another machine by replaying its prefix, and resume: every final quantum output matches the run that never stopped.

That is a symmetry of the settled result, the endpoint every observer inside ends up reading. Partitions, retries and wall clocks never reach it, because they never touch the projected state. The operator in the chair can tell two runs apart by their logs, and the operator is outside. Chapter five supplied the deflation, and it applies here at full strength: a symmetry is the claim that part of your description was never physical, and how the work was divided was never physical.

Nothing else in the architecture moves as much. Rotating the carrier moves twelve ports. A gauge change moves a convention chosen separately in every place. This one moves the entire execution, and goes uncounted as a symmetry because it looks like a fact about implementation rather than a fact about the world, which is what the labeling of the twelve ports and the choice of gauge in every place both looked like until somebody counted them.

A rendering loop can have a frame number shared by its objects, and a distributed solver can have a global progress counter. Neither number is time inside the world. The endpoint theorem is about what repair schedules produce. Reality does not approach a final instantaneous state: local events go on inside the one complete structure, and their rates belong to its clocks.

A faithful description running inside a world preserves its operational dependencies along with its terminal record. Closure says the inhabited universe is a realization of exactly that kind, of its own description.

37.7 Five cards

Chapter six's local rule needs no synchronized clock and no leader, and any run of it has a schedule all the same. Can a thing built like that do anything besides agree?

Deal five cards face up in a row and number the places one to five. A shuffle is a rule saying where the card in each place goes. Call the plain

one C: every card moves one place along, and the card in place five comes round to the front.

Two more shuffles, each of which is C wearing different clothes. Shuffle A sends place one to two, two to four, four to five, five to three, and three back to one. Shuffle B sends one to four, four to three, three to five, five to two, and two back to one. Trace either of them and you visit all five places once and return to where you began, which is exactly what C does. They differ in the route and in nothing else.

Chapter nineteen measured how far two operations are from being swappable by doing them in both orders and subtracting. In a group there is nothing to subtract, so you cancel instead: undo one, undo the other, then do each in turn. If the two get along, every doing cancels its undoing and the deck is as you found it. Try it with these two. Perform, in this order, A backwards, B backwards, A, and B.

Read the row off. Every card has moved one place along and the last has come round to the front. Four shuffles that were supposed to cancel have delivered C.

Five pieces of card on a table are enough to check that identity.

Write a program as a list of instructions, where an instruction reads one input bit and says: if that bit is one, do this shuffle, otherwise do that one. Run the instructions in order on the deck. Say the program computes a function of the input bits when the deck ends up cycled by C exactly on the inputs where the function is true, and untouched on the others.

A single instruction computes a single input bit. Building everything else out of that is the identity above, used once per gate. Given a program for one statement and a program for another, assemble four blocks in the order the identity wants them: the first in A's clothing run backwards, the second in B's clothing run backwards, the first in A's clothing, the second in B's clothing. If either statement is false, its two blocks do nothing and the surviving pair cancels, so the deck is untouched. If both are true, the four blocks are the four shuffles above and the deck comes out cycled, by a route that depends on the clothes and can be steered to any cycle you want by choosing them, which is what lets the step be applied again to its own output. That is the AND gate, and with the complement it is every gate there is. David Barrington published the construction in 1989 in the *Journal of Computer and System Sciences*.

Every function of bits is a tree of gates, so every function of bits is a fixed list of shuffles of five cards. The deck never holds more than five things in a row. The price sits in the length: four blocks per level of the tree, so a formula of depth ten compiles into a word of about a million shuffles, and depth is expensive.

Those five cards are hardware. Chapter fourteen counted the rotations of the twenty-faced carrier, got sixty, and found five families of half turns that those sixty rotations shuffle among themselves. Sixty permutations

of five objects is the group that leaves the fifth-degree equation without a solution formula, and the carrier's own group of moves. The deck is the carrier. The shuffles are things the carrier does to itself.

The property making the construction work is the property Galois found. The leftover from those four steps, the thing that came out as C instead of as nothing, is itself a move of the group. Collect every leftover the group produces that way and run the same operation on the collection. In a group with the shape a formula in radicals requires, the leftovers shrink at every round and die out. In this group of sixty they never shrink at all: the leftovers are the whole group again, round after round. The five-card identity is that refusal to shrink, written out on one case. The absence of a formula for the quintic and the presence of universal computation are one property of one group, read from two ends.

Compile a function into a federation of observers instead of into a deck: one patch per gate, each patch holding a few registers, each patch checking the seams it touches and repairing its own registers and nothing else. The input arrives as patches that hold their bits and accept nothing else. Set every other register in the network to anything at all. Let the patches repair in any order that keeps going, with no synchronized clock and no leader. The network settles, and the value of the function is sitting in a designated register of the settled state, from every starting state, and any two settled states of that network agree there whatever order the repairs ran in.

The small cases can be checked on every input rather than on a sample. A full adder's sum bit, the three-input circuit that adds two bits and a carry and sits underneath every binary addition anybody has ever performed, comes out right on all eight inputs from every starting state. So do the two bit-mixing functions called choose and majority, which take three inputs each and sit inside the hash function that Bitcoin's ledger is built on. A two-input NAND, the gate that answers false only when both its inputs are true, runs end to end on both engines, the deck and the federation, and they agree on all four inputs.

Networks of patches repairing locally compute finite Boolean functions, every one of them. Whoever builds one can describe it globally and count its work, and none of that description enters the gate rule a patch runs.

37.8 The hardware

Set the two results next to each other.

The first says the endpoint does not care how the work was cut up. The second says local repair computes anything a circuit computes. Together they describe a world that computes itself, with no machine underneath it.

Observers inside it can reconstruct and replay the recorded program. The hardware is the carrier: the deck is its five families of half turns, and the shuffles are its own rotations.

Which leaves the observers themselves, who have done all the work in this account without being explained by it. A loop that computes has pieces. Some of those pieces spend everything they have on continuing to be pieces: holding a boundary, keeping a record, repairing themselves faster than they come apart. Every rule in the architecture is indifferent to which patterns survive. Some of the patterns are not.

Chapter 38

Why Does Anything Try to Persist?

The rabbits came from the market and their urine was wrong.

Claude Bernard tells the story in a book he published in 1865, and the detail he leads with is the one that was sitting in front of him: the urine in the tray was clear, and it was acid. Rabbits eat grass. A grass eater's urine is cloudy and alkaline, and it had been cloudy and alkaline in every rabbit that had ever come through the laboratory. These animals were producing the urine of a cat.

He guessed they had not been fed for some time. He gave them grass, and within hours the urine went cloudy and alkaline. He took the grass away. It turned clear and acid again. Then he fed rabbits cold boiled beef, which a rabbit will eat if there is nothing else on offer, and for as long as the beef kept arriving the urine stayed clear and acid, exactly as it had while the animals were starving.

Fasting and a meat diet produce the same rabbit. A rabbit that has stopped eating is a carnivore. The meat it is eating is the rabbit.

The urine is the smallest part of this. What the fasting animal was defending is the composition of the fluid its cells sit in: the sugar in it, the salt, the water, the acid, the heat. Cut the supply from outside and those quantities hold. They hold because the liver dismantles the animal in order to hold them. Bernard proved that the liver manufactures sugar rather than merely storing it by washing a freshly removed dog's liver until no sugar could be found in it, leaving it overnight, and finding it full of sugar in the morning. That was 1855. He had pure glycogen out of liver tissue by 1857.

Repair happens by itself. Two readings disagree, an observer works out a move and commits it, the disagreement drops, and nothing anywhere wants anything: chapter six ran two thousand and forty-eight universes to a standstill on exactly that, and no observer in any of them was trying to get anywhere. A rabbit interrupts that pattern. It holds a set of numbers fixed at its own expense and goes on holding them until it has nothing left to spend. Why does anything do that, what exactly is it spending, and what does the spending buy that nothing else buys?

38.1 The internal environment

Bernard had a name for that fluid. He had been using it since 1854 and he set it out in the 1865 book, *Introduction to the Study of Experimental Medicine*: the internal environment, the *milieu intérieur*. His claim about it was blunt. A complex animal does not live in the air or the sea. It lives inside a fluid it manufactures and defends. The liberty of a warm-blooded animal to walk out into weather that would kill a fish is bought entirely by how tightly that fluid is held.

In lectures published in 1878, the year he died, he compressed the program into one line: the constancy of the internal environment is the condition of a free and independent life. A creature whose insides track its outsides is at the mercy of its outsides. Every degree of independence from the weather is paid for by an equal amount of work done indoors. Bernard's rabbits were buying independence from having eaten. The currency was their own tissue.

The explanation available at the time was a vital principle: living matter has a property, dead matter lacks it, and the property is why the numbers hold. Bernard spent a career refusing that. His position was that the constancy is produced by machinery, that a physiologist's job is to find the machinery, and that the machinery obeys the same determinism as a furnace or a barometer.

The word everybody uses arrived sixty-one years after the book. In May 1926 a volume was published in Paris for Charles Richet's jubilee, edited by A. Pettit, in an edition of five hundred copies. Walter Cannon, professor of physiology at Harvard, contributed a paper to it called "Physiological regulation of normal states: some tentative postulates concerning biological homeostatics". In that paper he gave the steady states of the body's fluid matrix the name **homeostasis**.

Cannon had come to the subject by watching the thing fail. He went to France with Base Hospital No. 5, a unit of Harvard clinicians, moved forward to a casualty clearing station run by the British, and spent the war on wound shock: men whose blood pressure fell and kept falling, whose blood turned acid, and who died with their injuries dressed and their wounds clean. He carried on the work in William Bayliss's laboratory in London and at the American laboratory in Dijon, and published *Traumatic Shock* in 1923. The list of quantities he assembled afterward as the ones a body holds steady, temperature and water and salt and sugar and oxygen and blood calcium, is a list of the readings he had watched go.

Cannon explained the choice six years later in *The Wisdom of the Body*. Equilibrium was taken: it had acquired an exact meaning for simple physico-chemical states in closed systems where known forces balance, and the coordinated business of brain and nerves and heart and lungs and kidneys holding a mammal's blood where it belongs is a different kind of thing.

"The word does not imply something set and immobile, a stagnation," he wrote. It means a condition that varies and comes out roughly constant anyway.

38.2 A symmetry with a payer

Chapter five settled what a symmetry is: a change that leaves a named invariant alone, which makes it the claim that part of your description was never physical.

Homeostasis has that exact shape. The shape is easy to fill in. Name the invariant: the concentration of sugar in the rabbit's internal fluid. Name the change: everything that happens to a rabbit across a day, being fed, not being fed, running, sitting in a cold room. The reading comes out the same, and it goes on coming out the same while the animal producing it is being eaten to produce it.

The symmetries of the earlier chapters arrive free of charge. The laws not caring where you stand costs nobody anything: move the apparatus a meter to the left and the same numbers come out. Move a rabbit from fed to fasting and the same number comes out because a liver was taken apart to produce it.

So there are two ways for a reading to be invariant, two different physical situations under one word. In the first, the invariance is a property of the world's laws. The observer has nothing to do with it. In the second, the invariance is manufactured by a bounded observer out of its own capacity, continuously, for as long as it can afford the outlay. What the vital principle was tracking is the outlay.

The outlay has to continue while the surrounding conditions keep changing. Neighbors update shared boundaries, and a maintained pattern has to answer every update, at a rate set by its own clock. Chapters eighteen and twenty-one priced that kind of work: a standing cost is a rate, and a rate is energy. Bernard's rabbits paid it in liver, and a calorimeter around the cage reads it as heat.

38.3 What a living thing is

Chapter eleven divided a network's state into what any neighbor can check and what only one observer holds. What an observer publishes is the entries on its seams. The fiber of a reading is the set of consistent worlds that produce exactly those entries. A reading whose fiber is empty describes nothing that could have happened.

A bounded observer persists when the reading on its seams keeps having a consistent extension. At every point along its own record time there is a world that carries what it publishes. It is spending to keep that true.

Bounded homeostatic persistence is the criterion for a living pattern: maintenance under disturbance, paid for out of a finite budget.

Chapter five's invariants survived something done to a system from outside: a rotation, a shift, a relabeling. This one survives the system's own running. A rabbit is whatever a rabbit's metabolism leaves alone while it takes everything else apart.

The rabbit makes the point about fibers by itself. Across a week without food almost everything about its interior changes: the liver empties, the fat goes, the mass falls by a fraction anybody can measure on a kitchen scale. What it publishes holds. The animal is traveling an enormous distance inside the set of states that produce one reading. The traveling is what the reading costs.

Stone persists too. Nothing inside a stone is spending to make that happen. Leave a stone alone and its readings sit where they sit. Leave a rabbit alone for a week and its readings walk off, because holding them where Bernard found them was work, and the work stopped. When the entries on an observer's seams stop having any consistent extension, the fiber is empty, and repair does what repair does: the surrounding patches redistribute until the record closes again, with the boundary of the thing that was doing the spending drawn somewhere else. Every atom is where it was a minute earlier. The reading is what stopped.

38.4 Two budgets, and what a policy is

Chapter nine found two kinds of repair and separated them by who pays. In the first, an observer rewrites its own patch so that its reading matches what the seam says: the capacity spent is the observer's own, nothing outside the patch changes, and the boundary record comes out of the transaction saying what it said going in. In the second, the boundary record itself is rewritten so that both sides read something new, and the capacity spent sits on the overlap, owned by neither observer alone.

A bounded thing that persists is choosing between those two, at every event, out of a finite budget. The rule it chooses by is its **policy**, and running a policy is the whole of behavior. There is nothing else in the word. An organism, a firm, a thermostat and a bacterium each run one: how much to spend bringing themselves into line with what the world reports, and how much to spend on changing what the world reports.

In 1972 Howard Berg and Douglas Brown built a microscope that chased bacteria. Rather than hold the field of view fixed and watch a cell swim out of it, the instrument moved to keep one *Escherichia coli* centered and recorded how far it had to move, which gave them the three-dimensional track of a single cell. The results went into *Nature* that year, volume 239, pages 500 to 504.

A cell swims in nearly straight lines, which they called runs, interrupted by brief episodes of jittering in place, which they called twiddles, after which it sets off in a direction unrelated to the one it came in on. In plain water that is a random walk. Put a gradient of something edible across the chamber and exactly one thing changes: a run that happens to be pointed up the gradient goes on longer. Runs pointed the wrong way come out at ordinary length. The cell has no steering at all. It has a brake. It declines to apply the brake while things are improving.

A bacterium cannot read a gradient across its own body, because the difference in concentration between its two ends is far below the noise of molecules arriving at random. Robert Macnab and Daniel Koshland showed what it does instead, in the *Proceedings of the National Academy of Sciences* in September 1972. They built an apparatus that mixed a cell suspension abruptly out of one uniform concentration into another, so that the population experienced a change in time with no gradient in space anywhere in the vessel. The cells responded anyway. A sudden drop set them tumbling, a sudden rise sent them swimming smoothly, and what the bacterium is reading is therefore its own record: what the seam says against what the seam was saying a moment ago.

Both budgets are open to that cell, and it uses both. It can retune its own detectors, and it does: after a step up in attractant, the receptors are chemically modified until the same concentration stops counting as news and the tumbling rate comes back to where it started. That is capacity spent inside the boundary. The sugar in the chamber is precisely where it was. Or it can swim, which puts its boundary against a different piece of the world, so that the entry on that boundary comes out different because the boundary is elsewhere.

Swimming is expensive in the only currency the cell has. The flagellar motor is turned by ions falling across the membrane, and that gradient is the same one the cell draws on to build everything else it builds, so a run is charged to the account that would otherwise have bought the next protein.

A company faces the identical pair. It can work on the product, which is capacity spent inside the boundary and leaves what the market says untouched on the day. Or it can buy entries in the shared account that customers, competitors and regulators all read from, which is what an advertising campaign is. Both are expensive and both are ordinary, and the two come out of different accounts: a firm that does more of the first does not thereby obtain any of the second.

A bacterium declining to tumble and a company buying billboards are the same move at different scales under different budgets.

38.5 Four states

Strip the arrangement to the smallest thing on which the two budgets can be told apart at all.

Take an observer with two registers, each holding one bit. One of them sits on the overlap: it is the shared record, the entry a neighbor can read. The other is private. What it holds is a flag, set when the observer's own state fails to match what the shared record says and clear when it matches. Four states in total, of which two are consistent, being the two values of the shared record with the flag clear.

Two moves exist. The private move clears the flag and touches the shared record not at all, which is self-repair in its smallest possible form: the observer bringing itself into line with what the seam says. The network move writes the shared record, so that what both sides read comes out different.

A rule for picking moves, step after step, possibly at random, is a scheduler. Constrain it three ways, all of them constraints the repair law carries anyway: it never increases the mismatch, it writes at most one register per step, and it spends at most one unit of capacity per step. Two sub-families inside that class do different things. A self-update-only scheduler never writes the shared register. A network-update-only scheduler never writes the private one.

Start in a consistent state, with the shared record holding one of its two values. Ask for the other one: the state in which the shared record holds the opposite value and the flag is clear.

Three facts hold about that target, on a system with four states in it.

A self-update-only scheduler never arrives. The failure has nothing to do with budget or patience. From every consistent start, over any number of steps, for every member of that sub-family, the probability of arrival is zero.

An explicit network-update-only scheduler arrives with probability one.

The third fact prices the other two. Any scheduler in the class, whatever its rule, that has a positive probability of arriving writes the shared register somewhere along the way. A positive chance of getting there costs at least one shared write. There is no arrangement of the private budget that supplies it.

The reason for the first of the three is one line. It is chapter five's word doing the work. A private move leaves the shared register alone. Compose any number of moves that each leave a quantity alone and the quantity is left alone, so the value of the shared register is an invariant of the whole self-update-only sub-family, and every state that sub-family can reach carries the value it started with. The size of the budget never appears in that argument, which is why enlarging the budget does nothing to the conclusion. The private register can be cleared, set, cleared again

and read any number of times, and the neighbor reading the seam sees the entry it saw at the beginning.

The two budgets do not buy the same things. The demonstration fits on four states, which matters more than a large example would, because a system with four states has nowhere for an inefficiency to hide. The gap between the budgets is not a shortfall in the private one that a larger private budget would cover. There is a class of outcome that self-repair does not reach at any size. The only thing that reaches it is expenditure on the record two observers hold jointly.

Every bounded system that persists therefore runs into that class. Bernard's rabbit can convert its own body into blood sugar for days and hold the internal environment at the value he measured, and the entry on the seam between the rabbit and the room, which is that there is nothing in the room to eat, comes through unaltered. A bacterium that has finished retuning its receptors reads zero mismatch while sitting in a patch of water with no food in it. Perfect internal adjustment and starvation are compatible states. A policy that spends only on the first arrives reliably at the second.

The company gets the same result with the accounts written out. A firm can spend a decade making its product better and arrive at a market that has heard nothing about it, because what the market holds is an entry the firm never wrote to, and that entry carries the value it carried before. Which is why the workshop and the billboard both exist, and why every firm has to decide how to divide its budget between them.

38.6 The entry is one entry

An entry on a seam is one entry. It is read by both patches, owned by neither, and holds a single value at a time.

Two observers spending on the same entry are spending against each other. Whatever stands written there when the spending stops is what both of them have to live with. Each brings a finite budget to the argument. A bacterium in a gradient is in that position with respect to every other bacterium in the same drop of water, which is why the gradient it is climbing is partly a record of what the others have taken out of it.

That is a competition. It is settled by which pattern goes on having a consistent extension afterwards, and was running on the thirty seams of a twelve-patch net long before anything with a flagellum turned up to enter it.

Chapter 39

Why Is a Bacterium Freer Than an Atom?

Every morning somebody in a laboratory lifts twelve glass flasks off a shaker and moves one hundredth of the liquid in each of them into a fresh flask of the same clear solution. The used flasks go into the autoclave. The new ones go back onto the shaker at 37 degrees Celsius for the next twenty-four hours. The job takes a few minutes, and somebody has done it every day since 24 February 1988.

Each flask holds ten milliliters. The entire experiment, all twelve populations of it, comes to a hundred and twenty milliliters.

What is in the flasks is *Escherichia coli*. On the first day all twelve held the descendants of a single cell. Six of them were founded from that cell's line and six from a variant of it differing by one change that does nothing in this medium except make its colonies come up a different color on one kind of indicator plate, which is how a contaminated flask gets caught before it ruins anything. The solution carries twenty-five milligrams of glucose in every liter and nothing else these cells can eat. It also carries citrate, about twelve times as much of it as glucose by count of molecules, which is in the recipe to keep iron in solution. *E. coli* cannot use citrate as a carbon source when oxygen is present. Handing a bacterium a tube of citrate and watching whether it grows is one of the routine ways of telling *E. coli* apart from the things that look like it under a microscope, so the inability belongs to the species rather than to these twelve flasks. The test is on the identification key, and a strain that passes it gets filed as something other than *E. coli*.

The daily transfer carries over one percent. Overnight that one percent eats the glucose and grows back a hundredfold, which is six and two thirds doublings, so a day in the flask is six and two thirds generations and a year is roughly two thousand four hundred.

Nobody in that laboratory picks which cells get transferred. The pipette takes a hundredth of whatever is in the flask and cannot tell one cell from another, and the person holding it has no way of knowing which lineage is which until long afterward, when somebody sequences something. Yet the flask holds a different population every year. The differences run in a consistent direction. Something settles which lineages are in there, and it is not the person holding the pipette.

There is a second question in that flask, about one cell rather than about a population. A bacterium has options. It can put the next minute's energy into dividing or into repairing itself, it can swim toward the sugar or sit, it can carry the machinery for eating something and switch it off. A hydrogen atom in the same flask has nothing of the kind. The atom came first and has fewer moving parts, and less room to be otherwise than the bacterium does, which runs backwards from the way the world is usually described.

39.1 Three things, and no fourth

Something copies itself imperfectly. The imperfection is passed to the copies. Some versions end up with more copies than others.

That is the whole mechanism. Each of the three parts is visible in the flask. Copying is imperfect: over thirty thousand generations, each of the twelve populations tested billions of mutations, which is what happens when a population of hundreds of millions divides six and two thirds times a day for decades. The imperfection is heritable: a daughter cell gets its mother's copy of the genome, errors and all. And the versions do not fare alike, because the glucose in a fresh flask is gone within a few hours, and a lineage that got through more divisions before it ran out is a larger fraction of what is sitting there when the pipette arrives in the morning. A larger fraction of the flask is a larger fraction of the hundredth that gets moved.

The name for those three conditions running together is **natural selection**. There is no fourth item in which anybody decides anything. Nothing in the flask is aiming at a result, no lineage is trying to win, and the only thing acting on the flask from outside is a pipette that samples without discrimination. The sorting has finished before anybody arrives for the morning transfer.

"Fitter" is a bookkeeping word here and carries none of its usual freight. A fitter lineage holds a larger share of tomorrow's flask than of today's. That exhausts the meaning. The word predicts nothing about the day after tomorrow, because the flask the lineage will be competing in is the flask it is in the middle of changing.

39.2 The freezer

Richard Lenski set the twelve flasks going at the University of California, Irvine, with two thousand generations in mind, which at six and two thirds a day is about ten months of transfers. He moved the populations to Michigan State in 1991 and they carried on there. The organism he picked reproduces without sex, so every lineage is an unmixed line of descent and no gene ever arrives sideways from a neighbor. It also survives being frozen.

Every five hundred generations, which comes to every seventy-five days, the part of each population that does not go into the fresh flask gets a dose

of glycerol and goes into a freezer at minus 80 degrees Celsius. The cells come through it alive. Thaw a vial and the population resumes, from that day, with the mutations it had accumulated by then and none of the ones it acquired afterward.

Twelve populations, one ancestor, and a shelf of vials sampled every seventy-five days for decades. It is a fossil record whose fossils are alive, so the experiment can be run again from any stored point.

The freezer was needed early in 2020, when the populations had passed seventy-three thousand generations and the laboratory shut. The flasks went into the freezer. In September the experiment came back out of it and carried on from where it had stopped. The rest of the time the freezer is doing something stranger. Thaw a vial from before an event and run the population forward again, and whether that event had to happen becomes a question somebody can answer by counting.

The one quantity this experiment reports comes out of the freezer as well. To find out how much better a population has got, thaw the ancestor, mix the two in equal parts in a fresh flask, and plate a sample at the start and again a day later. The ancestor's colonies come up one color and the competitor's come up the other, so counting red against white on a plate gives the ratio before and the ratio after, and the change in that ratio over one day is what the word fitness denotes here. Every measurement of it is a competition against cells that were frozen in an earlier year and thawed for the purpose.

Michael Wiser, Noah Ribeck and Richard Lenski ran that measurement across all twelve populations through fifty thousand generations and published the trajectory in *Science* on 13 December 2013. Mean fitness increases without bound, consistent with a power law, with no sign of leveling off after twenty years of it. Chapter thirty-six's argument said that mean fitness never decreases in a population sorted this way, and that the climb stops once the survivors stop disagreeing. A flask never runs out of disagreement, because copying keeps making more of it. Twelve flasks on a shaker in Michigan are where somebody went and measured it.

39.3 Citrate

For more than thirty thousand generations, the citrate sat in all twelve flasks, twelve times more abundant than the food, untouched. Then one of the twelve, the population known as Ara-3, started eating it. By generation 31,500 there were cells in that flask growing on citrate in air, which is the thing the diagnostic test says *E. coli* does not do. The population got larger, because there was suddenly far more carbon available to it, and it got more diverse, because the cells that could not use citrate persisted alongside the ones that could.

Zachary Blount, Christina Borland and Richard Lenski published the result on 10 June 2008, in a paper called "Historical contingency and the evolution of a key innovation in an experimental population of *Escherichia coli*". It was Lenski's inaugural article for the National Academy of Sciences, the paper a newly elected member writes on a subject of their own choosing. Most of it is taken up with the experiment Blount ran after the announcement.

Two explanations were available for why one population out of twelve did this after thirty thousand generations and the other eleven did not. Either the mutation required is astronomically rare, in which case any of the twelve could have got it on any day and this one was lucky, or the mutation is ordinary and works only in a cell that carries some other change first, in which case the population had to build up to it.

Those two stories make different predictions about the freezer, and the freezer answers. Blount thawed vials from across that population's history and grew enormous numbers of cells from each of them. From the ancestor, not one citrate-eater among 8.4 trillion cells. From sixty clones sampled across the first fifteen thousand generations, none among nine trillion. From clones taken later, citrate-eating appeared repeatedly and significantly more often than chance allows, which puts the arrival of the enabling change at around generation twenty thousand. That change left the cells no more error-prone in general. It made this one step reachable.

Run the population from the early part of its own past and it cannot get to citrate. Run it from a later part of the same past and it can, over and over. What a lineage is able to become is set by what that lineage has done. The set gets bigger as the history gets longer.

39.4 One budget with two halves

Chapter thirty-eight left a living thing as a pattern that keeps its public reading extendable at its own expense, and behavior as a policy over the two ways of spending: on itself, or on the record it shares with a neighbor.

A pattern persists by being re-asserted into the record every cycle, and a record costs capacity, and the capacity is finite and shared. Two patterns whose re-assertions cannot both be extended cannot both persist. Which of them does is settled by which one keeps having an extension. Nothing about food or territory is needed to get this started. The scarcity is in the bookkeeping, and would be there in a universe with unlimited everything else.

Chapter thirty-six ran that argument on laws, and got a filter with no chooser in it: what survives is what closes, and the share held by the descriptions that close does not decrease. This is the same argument one level down, with organisms where the laws were. Selection at the level of

physics and selection at the level of bacteria are one filter applied at two grain sizes. The finer one inherits its shape from the coarser.

A pattern can spend on either side of the split. The two sides buy different things.

Spend everything privately and you get homeostasis: an interior in excellent order, every internal reading consistent, and an overlap you have conceded to whatever else is spending out there. That is a stable arrangement right up until the neighborhood's reading stops extending yours, at which point the maintained interior is a private fiction with no public correlate, and the pattern ends with its interior intact.

Spend everything on the overlap and you get domination: the neighborhood rewritten in your terms, every seam carrying your version, and an interior that has gone unrepaired for as long as the spending has run. The reading you are imposing has less and less behind it. A reading with nothing behind it fails the first check that reaches for the state underneath.

So fitting in and crowding out are one expenditure seen from the two sides of the split. Nothing that persists optimizes for either alone. The cells in those flasks do both, all night, and the boundary between the two runs through individual molecules. Refolding a protein and proofreading a copy of the genome are interior work. Eating the glucose is overlap work, because what a lineage does to the flask is what every lineage in it, its own descendants included, has to live in tomorrow. Pushing a proton out through the membrane is both at once: the cell has kept its own interior in range and made the flask more acidic for everybody. The citrate-eaters are the cleanest case, because the mutation is in the cell and the consequence is in the flask: they opened a carbon source nobody was using, and the whole population, competitors included, got bigger.

The harder result in chapter thirty-eight was that some outcomes are reachable only by spending shared capacity, and that no amount of private self-repair arrives at them at any budget. The flasks are what that looks like when it runs for decades. A lineage cannot reach citrate by getting better at being itself. It gets there by changing what the flask contains, which is a change to everybody's record, and then living in the flask it changed.

Different lineages settle on different policies, in flasks prepared from the same recipe on the same bench. Paul Sniegowski, Philip Gerrish and Richard Lenski reported in *Nature* on 12 June 1997 that lineages which copy themselves less accurately had arisen on their own in some of the twelve populations and spread. Sloppier copying is a strange thing to spend an interior budget on, because most mutations are damaging and the lineage pays for every one of them. It spreads anyway, because a sloppier copier reaches the useful change sooner, and in a flask where everything is sampled once a day, sooner is the whole of the argument. Twelve flasks, one ancestor, and the populations differ in how accurately they copy themselves.

39.5 What a hydrogen atom cannot do

Ask how much room something has to be otherwise, and count it at each level.

Chapter fourteen counted the ports: twelve, with thirty seams between them. Twelve is what the consistency conditions permit rather than a number anybody chose. Chapter fifteen counted the directions: three, the rank of the slowest-fading band, with the other bands dying away faster and taking their dimensions with them. Chapter sixteen found the arrow: one way round, because a commit discards what it resolved. No arrangement of the early universe would have made it run the other way.

Those three were settled with no history available to settle them. There was no record, nothing had happened, and the conditions had exactly one solution. A universe with eleven ports never lost a competition to a universe with twelve. Eleven ports does not survive the counting, so there was never an eleven-port world for the twelve-port one to beat.

The early universe is therefore the most constrained thing there has ever been. Every question about it that has an answer has a forced answer. The forcing comes from its age rather than from its simplicity. Nothing had been written down.

Then things start getting written down. Chapter eleven's result was that the normal form is unique given the boundary data, which is to say that different records on the boundary settle into different worlds. Boundary records are made, one commit at a time. A commit is a fact that the closure conditions did not supply and could not have supplied. Every one of them enters as an input to what comes next. So the fraction of what a thing is that traces to its own history rather than to the conditions that permitted anything at all rises with every cycle, and it rises fastest in the patterns that are making the most records.

Look at a hydrogen atom with that in hand. Its ground state is one state. Two hydrogen atoms in it are the same object in the strong sense chapter twenty-eight needed for the exclusion argument: there is no record anywhere that distinguishes them, and the absence is not a limitation of anybody's instruments. It has no policy over the direction split. Holding that pattern together costs what chapter eighteen said it costs, a standing bill re-asserted every cycle, and the bill has exactly one line on it and no way to move an amount from one line to another. Put the atom in a field and its response is fixed by the same conditions that fixed the number of directions the field can point in. A hydrogen atom has no strategy available to it. There is no record in it that could carry one.

Two *E. coli* cells in the same flask at generation 31,500, both of them descended from the cell that went into a flask in February 1988, differ. The difference sits in their records, it is heritable, and one of them can eat something the other cannot. That lineage carries a policy and spends it

every cycle: divide or repair, swim or sit, keep an unused piece of machinery running or shut it down and save the cost. Each of those is a choice about which half of the budget to spend on, and each of them is available only because of a long run of commits in that lineage's past that could have gone otherwise.

Freedom, if the word is going to mean anything measurable, means room to be otherwise, and room to be otherwise accumulates. It is the residue of history rather than a late arrival with brains. There is more of it the further down the tree you go.

39.6 The view from inside

Chapter six wired twelve observers into a closed surface and ran every one of their two thousand and forty-eight states through sixteen schedules apiece. Every run ended in the same arrangement. Nothing chose that arrangement and nobody voted on it. It was the one configuration that satisfied every seam at once. The observers arrived at it by repairing whatever was in front of them.

Chapter eleven gave that its general form. A reading has a fiber, which is the set of states consistent with it. A reading whose fiber is empty belongs to no world at all. Nothing rejects such a reading. There is nowhere for it to be. Rejection would need a judge, and having nowhere to be needs nobody.

Seen from outside, the twelve observers are a solve: a system with a unique consistent state settling into it. There is no outside. The only available view is the one from within, held by the patterns that came through, and from there it looks like something else entirely. Some patterns are here and others are not. The ones here are the ones whose readings kept having an extension. The missing ones ran out of world to be consistent with.

That is selection, and needs no agent, because differential persistence is the whole requirement. Lenski's laboratory has a pipette that cannot tell one cell from another. The universe contains no sampling device of any kind.

So the filter and the fixed point are one object seen from two positions. One of the two positions is not available to anybody. Selection is what a fixed point looks like from the inside, when the patterns competing to be in it are the ones doing the looking.

Twelve flasks, a hundred and twenty milliliters, one ancestor, and a freezer full of its own past. Nothing in that arrangement wants anything. The same is true of the arrangement the flasks are sitting in. One of the patterns that came through the filter has hands, and a laboratory, and a habit of writing things down, and is holding this book.

Chapter 40

Does It Feel Like Anything, and What Is Left to Decide?

The first big interdisciplinary conference on consciousness was organized by an anesthesiologist.

Stuart Hameroff worked in anesthesia at the University of Arizona Medical Center, and in April 1994 he convened a meeting in Tucson called Toward a Scientific Basis for Consciousness, which put philosophers, neuroscientists and psychologists in the same rooms for a week. He spent his working days putting patients under and waking them again.

The talk people were repeating in the corridors came from a philosopher a year out of his doctorate, holding a fellowship in a philosophy, neuroscience and psychology program at Washington University in St. Louis. David Chalmers opened by splitting the subject in two. The split was a list.

How a system discriminates between things around it, categorizes them and reacts. How it brings information together from separate places. How it reports its own states in words. How it gets access to its own internal states at all. How attention gets focused on one thing rather than another. How behavior gets deliberately controlled. What separates being awake from being asleep.

Seven items, and he called them the easy problems of consciousness, with a warning attached to the adjective: getting the details right, he wrote, will probably take a century or two of difficult empirical work. They count as easy because in every case anybody can say what an answer would look like. It would be a mechanism. Specify the machinery that retrieves information about internal states and makes it available for verbal report, and reportability has been explained.

Then the other one. When our cognitive systems engage in visual and auditory information-processing, he wrote, we have visual or auditory experience: the quality of deep blue, the sensation of middle C. Thomas Nagel had supplied the standard phrase for it twenty years before, in the October 1974 issue of the *Philosophical Review*: an organism is conscious if there is something it is like to be that organism. Why physical processing should

give rise to any of that seems objectively unreasonable, Chalmers wrote, and yet it does. The talk went into the *Journal of Consciousness Studies* the following year, volume 2, pages 200 to 219, under the title "Facing Up to the Problem of Consciousness". The phrase the hard problem became the standard name for that question.

Descartes had a physical world made of extended stuff and a thinking thing that occupies no place at all, and his trouble was traffic between the two: how a decision taken in something with no location moves an arm that has one. He put the meeting at the pineal gland, on the grounds that every other part of the brain comes in pairs and that one does not, so it was the only place the signals from two eyes and two ears could be brought together into one impression. Every version since has kept the shape and dropped the gland. There is an account of the world written in the third person, there is what it is like to be one of the things in it, and nobody has a rule that turns either into the other. The name for that is the **mind-body problem**.

Chapter three defined an observer without any mention of consciousness. This chapter takes one further step: certain organized self-reading and repair processes are experiences. A thermostat shares the record-and-feedback interface with a person, and the step is about what is organized behind the interface.

One of these bounded record-keepers is holding this book. There is something it is like to be it. Where in the architecture does that sit? And if the network settles into the same arrangement whatever order anybody repairs in, what is there for that reader to decide?

40.1 A creature molecule for molecule identical to you

Chalmers put the sharpest form of the first question two years later, in a book called *The Conscious Mind*. Consider, he wrote, his zombie twin: a creature molecule for molecule identical to me, lacking conscious experience entirely. Same cells, same firing, same words coming out of its mouth, and nobody inside.

The argument runs three lines. Such a creature is conceivable, and Chalmers set the standard in those words: it certainly seems that a coherent situation is described, and I can discern no contradiction in the description. Whatever is conceivable in that sense is possible. And if a world physically identical to this one can lack experience, then experience is something over and above the physical facts, and physics by itself does not reach it.

The twin writes the paper about zombies. It sits down, works through the conceivability argument, publishes it in a journal and answers the ob-

jections. There is nothing it is like to be the author. Chalmers grants that, because the twin does everything he does.

The zombie argument keeps the entire physical organization, every register and every causal dependency, and removes experience. An identity theory rejects the separation outright: if experience is a kind of physical organization, a duplicate of the organization is a duplicate of the experience. That is the position this chapter takes.

Everything rests on the first step, the step nobody pauses on, because taking a complete physical description sounds like the part you get for free.

40.2 The mechanical properties of the ether

Chapter one's crash log had an entry for the substance light was supposed to wave in, rigid enough to carry a wave at the speed light travels and insubstantial enough for a planet to cross without losing speed. The nineteenth century held both properties rather than drop the idea that a wave needs something to wave in. The interferometer in the Cleveland basement went looking for the Earth's motion through it in 1887 and found nothing, and forty years of better instruments found nothing again.

Nobody ever caught the ether not being there. Einstein took the null result as the law rather than as the disappointment, and after that every question the ether had been introduced to answer had an answer without it. Its stiffness explained nothing, because nothing needed explaining by a stiffness. Its density had no consequence anywhere. The ether was made redundant, which is a more complete failure than going undetected, because a thing nobody detects may have effects that need accounting for.

Take the sentence "what are the mechanical properties of the ether?" It has a subject and a verb and it never got an answer. It stopped being a question, because one of its words stopped naming anything. A question with a term that names nothing does not have a false answer either.

Two things can happen to a question, then. It gets answered. The question stands with a fact attached to it. Or one of its terms is shown to name nothing. The question goes away and stops consuming effort. The second is called **dissolving** the question. It happens rarely, and from the outside it looks the same as a refusal to answer.

The hard problem asks how experience arises from the physical facts, where the physical facts are meant as an inventory belonging to nobody: a list of what there is, compiled from no position. Go looking for that list and it fails to turn up, one entry at a time.

Distance counts comparisons. There is no comparison without two parties each holding something. A rate needs a clock, and chapter twenty-one left no clock that is not somebody's. Charge is read off a loop that runs through one observer after another. A record sits in the center of one particular algebra of questions. Even the leftover of exactly one, sitting on a

twelve-patch net with one flipped edge, is a quantity that every observer in that net agrees about, which is a different thing from a quantity nobody holds.

Objective, in the sense chapter eleven fixed, means settled between parties who each have a boundary and a budget, with no further list standing behind theirs for it to approximate.

The answer here identifies certain lived processes with their qualitative character. That is a commitment about what the physical organization is. Having records and a bounded point of view makes something an observer, the identity is a further step, and the hard problem meets the account at that step.

A record map can leave out state that is accessible inside. That makes the omitted state private relative to the map, which is a long way short of making every omitted variable a quale.

Deleting an interior that does causal work changes boundary behavior, and nobody disputes that. The zombie keeps the interior and asks about its character, and the identity account answers that the interior's organization is its character.

40.3 Coordinates an observer reads on itself

Chapter thirty-eight got an observer down to two registers, which is the smallest arrangement in which its two budgets come apart. A neighbor reads the first one. The second holds one bit saying whether the observer matches what the seam says about it.

What that private register holds is the observer's reading of its own relation to the world. The observer's next move is taken on the strength of it. Nothing outside can consult it, and no repair anywhere in the network reaches it.

A system carrying readings of its own update state is doing something a photographic plate does not do. It divides itself into parts and reads the division. The reading goes to its own repair policy and nowhere else. Give it more registers and the division gets finer. A stable cell of that division is a coordinate the observer can locate itself along and repair itself with.

In an experiencing system, redness, pain and the other lived distinctions are organized self-reading of this kind. Which distinctions occur, and how they depend on neural activity, are questions for psychology and biology.

Reading such a coordinate is the whole of having it. There is no second copy of the ache, kept elsewhere, for the reading to be a reading of, and no fact about whether the one you have is the genuine article.

Consciousness is owned boundary repair, read from the inside.

Owned, because the capacity being spent belongs to the observer and comes out of its own budget. Boundary, because what is being held is the interface between it and everything else. Repair, because the holding

is a rate rather than an act, since the neighbors keep writing and the entries keep moving. Read from the inside, because the observer's records include readings of its own state, and those readings are the half of the record the map deletes: real by the same standard Alice's coin is real, and unpublishable by construction.

That sentence is a claim about experience, laid on top of the definition of an observer. A thermostat's bounded feedback is a long way from a person's organization, and the distance depends on how the self-reading is organized as well as on how much of it there is.

Which cells implement which coordinate, why activity in one part of the visual cortex reads as color while activity a few centimeters away reads as motion, what a stroke takes away when it takes a coordinate away, how many coordinates a mouse reads on itself, and at which point in an anesthesiologist's induction a person stops reading any, are experimental questions.

A coordinate the public record leaves out cannot be recovered from that record when its fiber holds different values. A different intervention or a richer interface can learn more, and learning about an episode is a different thing from undergoing it.

40.4 Three cells down the diagonal

Ask somebody what red is like and everything comes back except the thing you asked for. A wavelength. The name the paint shop prints on the can. Both of those cross the gap between two people intact. Neither answers the question. The word for what failed to cross is a **quale**, plural qualia, meaning how a particular experience is for the thing undergoing it, as against the wavelength, the firing pattern or the word.

Represent a three-alternative system by three-by-three matrices. A readout that writes which alternative occurred keeps the diagonal, three independent entries, and the six off-diagonal entries hold the distinctions the diagonal leaves out. The full matrix algebra has only multiples of the identity in its center. Pinch it down to what the readout writes, chapter nineteen's mail room, and the diagonal is what is left: commutative, its own center, the record algebra of that readout.

For N alternatives there are N diagonal entries and N squared minus N off the diagonal. Those are dimensions of a representation, and experiences are counted some other way, if they are counted at all. A two-state classical device gets by with a commutative algebra and no amplitudes.

One line covers every size, with N standing for the number of alternatives a readout tells apart.

$$\text{cells on the diagonal} \;=\; N, \qquad \text{cells off it} \;=\; N^2 - N$$

The diagonal keeps N entries out of N squared. That is what one readout writes, and a different measurement writes a different diagonal.

The identity concerns lived content in an experiencing organization, and nothing in the count makes a quale an off-diagonal matrix entry. Which distinctions a record loses is a mathematical question. Which distinctions are lived is the question the identity answers, one experiencing organization at a time.

No-cloning rules out copying an unknown quantum state and allows copying records. It forbids nobody from learning about a prepared ensemble or from reconstructing a model out of richer measurements, and it says nothing about whether qualia are quantum states.

Experiencing organization belongs within reality, built out of the same observers as everything else, with no observer outside supplying it. Where it occurs, and in what, is the question this chapter has been walking around.

40.5 Then nothing you do matters

Somebody is sitting up at night working out whether to say the true thing or the convenient one. Three results say the sitting up changes nothing.

Chapter six started a twelve-patch network in every one of its two thousand and forty-eight distinguishable states, ran sixteen repair schedules on each of them, and ended in the same arrangement every time. Chapter ten ran the same process on eighty-two thousand patches, where 102,415 disagreements came down to none under every order the repairs were run in. Chapter thirty-seven showed that a run cut across a thousand machines settles where the single-machine run settles, and that the partition, the worker count, the schedule and the restart history reach no observable anywhere.

The endpoint does not depend on where the system started. It does not depend on the order anybody repaired in. There is no clock outside it, so there is not even a sense in which the endpoint is waiting its turn. So the person sitting up at night is a subsystem executing a delay, everything between here and the endpoint is theater, and the arrangement the whole thing lands in carries the same entries either way.

This is a harder version of fatalism than the one usually argued about. Ordinary determinism concedes that the future depends on the initial condition, which at least leaves the past doing some work. Here the initial condition washes out as well. Two thousand and forty-eight starting states, one endpoint.

40.6 Who writes the boundary

Chapter six gave twelve observers a coin each and thirty seam records saying whether two coins matched. The repairs inside that net move coins and leave the records standing, so where the coins stop is a function of the records handed over before anybody moved, from the first repair to the last.

Chapter eleven's theorem carries a qualifier. The fatalism above works by dropping it. The theorem returns one endpoint given the boundary data, and a region's boundary data is the entries on the seams where it meets what it is not, which the patches inside can read and cannot set. Regions that start anywhere at all with the same edge entries finish in the same place. Regions with different edge entries finish apart, with nothing to choose between the two endpoints, since both satisfy every record inside.

Drop the words "given the boundary data" and out comes the fatalism above. Keep them and boundary records vary, so "the endpoint" does not pick out anything until somebody has said which entries are sitting on the edge.

Chapter six checked that on twelve patches. Thirty entries, all satisfiable, and the twelve coins have exactly one setting up to the flip nothing can see. Change one entry out of the thirty, from same to different, and the number of consistent settings drops to zero and what the machine stops on carries a leftover of exactly one. One record, and the coins stop somewhere else.

So where do entries come from? The patches inside a region read its edge and cannot set it, so the write happens at the seam itself, between the two observers who share it. Chapter thirty-eight named the move. A private self-update spends the observer's own capacity, clears its own mismatch, and leaves the shared entry saying what it said going in. A network update writes the shared entry. Each side then reads a different value. On the four-state system where the two can first be told apart, no self-update-only scheduler ever reaches the state where the shared entry has moved, at any budget, over any number of steps. A network update reaches it with probability one, and any scheme with a positive chance of getting there writes the shared register on the way.

The shared entries are the boundary data. The move that writes them is the move no quantity of private effort substitutes for.

Say what those writes are in a person's vocabulary.

What to admit, which is what you let across your boundary from outside and allow to change your state. What to emit, which is what you write onto the seams you are one side of. What to repair, which is where a finite budget actually goes. What to ignore, which costs nothing to decide and shows up in the record as an entry left standing. What to confess, which is putting onto the shared record something only you were holding. What to protect, which is spending your own capacity so that somebody else's

reading keeps having a consistent extension. And what to amplify, which is writing an entry onto seams that had no opinion about it until you arrived.

Those seven are the boundary data. They are the input the theorem takes, and they are the reason it returns a different world on different days.

So free will is self-reading participation in repair. An observer that reads coordinates on itself, acts on the readings, and writes the results onto seams it shares with others is a place where the boundary record gets set. The convergence does not run around such a thing and it does not suspend itself on request. It runs through. The entry it reads on the way past is the entry that observer wrote.

40.7 Pavia, 524

Boethius was consul in 510 and master of the offices under Theodoric, and in 522 his two sons held the consulship together, which is the top of a Roman career of that period. He then defended the senator Albinus, who was accused of writing to the emperor Justin in Constantinople against Theodoric's rule. He was taken to Pavia, held there, and executed in 524. In the prison he wrote the *Consolation of Philosophy*. Its fifth and last book is about the question above.

He puts it in the vocabulary of divine foreknowledge. If the whole order is possessed at once, entire, by something that does not take it a piece at a time, then it is settled, and a settled order leaves nothing for anybody inside it to decide.

The answer has two parts and both survive translation out of the vocabulary.

The first is a distinction between two necessities. One is simple: men are necessarily mortal. The other is conditional: if you know that someone is walking, he is necessarily walking. The second attaches to the act only while the condition holds, imported by the knowing rather than produced by the walking. Then the test question, which Boethius puts directly: does the act of vision add any necessity to the things you see before your eyes? Watching a man cross a courtyard does not make him cross it.

The second part is his definition of eternity as life possessed whole. A complete-history reading can contain earlier and later events internally while requiring no external time in which the whole is assembled. Timeless closure therefore permits internal causal reasons, local succession and change; it does not remove every ordering from the structure.

Fourteen centuries later the requirement itself was taken apart, in eleven pages. What most people mean by freedom is that you are responsible for what you did only if you could have done otherwise. Harry Frankfurt attacked that sentence in the *Journal of Philosophy* of 4 December 1969,

in a paper called “Alternate Possibilities and Moral Responsibility”, pages 829 to 839.

His case is a piece of machinery. Somebody, call him Black, wants Jones to do a particular thing, and is prepared to step in and make him do it if Jones shows any sign of deciding otherwise. Jones decides to do it on his own. Black watches and never moves. Jones could not have done otherwise, because Black stood by with the means to prevent it. Jones is responsible anyway, because what he did came out of his own deliberation and Black played no part.

What carries the responsibility is therefore the structure of the will that produced the act and the record of the act itself. Whether some other setting of the boundary was reachable is a question about the space of records. Which entry got written, out of which observer’s own repair policy, is a question about this record. This record is what everything downstream is computed from.

40.8 It was going to happen anyway

The fatalism above has a second life as an excuse.

Somebody writes an entry onto a shared seam and leaves the repair bill with whoever is on the other side. Asked about it afterward, the sentence available is that the network converges regardless, that the endpoint was going to be the endpoint, and that nothing turned on the write.

The structure blocks that, and it blocks it with arithmetic rather than with disapproval.

Chapter six’s flipped edge is the clean case, because the whole net can be checked. Two of its nineteen independent loops carry a residue, no setting of the twelve coins satisfies all thirty records, and the machine converges anyway, from all two thousand and forty-eight starts and under all sixteen schedules, onto an arrangement carrying a leftover of exactly one. Convergence is guaranteed and so is the size of the leftover.

Where it sits is a different matter. Run the repairs in another order and the leftover ends up on a different part of the net. It survives any amount of working harder, and it can be placed, and which patch is carrying it when the repairs stop is settled by who repaired what, in which order, and which way each one pushed the disagreement it found.

A record can keep part of the trajectory along with its endpoint. The midpoint map loses input distinctions on its two output readings, and the crossing count it writes keeps them, so who absorbed a cost and who exported it survives in the records for as long as the records do.

The leftover comes to one either way. The record that produced it names which patch absorbed the mismatch and which one exported it. The naming is no moral gloss applied afterward by somebody keeping accounts. It

is the entries, in order, which is the only material anybody downstream has to compute from.

The excuse also needs the endpoint to be the kind of thing that could have caused something, and chapter thirty-seven wrote the endpoint down. Take the whole structure, take the record it holds, look up the world that record names, and demand that what comes back is the structure you started with.

$$\text{world}(R(W)) \;=\; W$$

W is the whole structure, R is chapter eleven's record map, which returns the part of a state that survives comparison, and world is the lookup running from a record to the consistent state that record names. The unknown sits on both sides, which makes this a fixed point rather than a recipe.

The complete-history closure equation requires no first external solver. Its internal dependencies can nevertheless be realized by local computation. The equation does not cause those events from outside; the proposal identifies the self-consistent structure and its intrinsic operation as one reality.

What an observer writes contributes to the record R returns and to subsequent local dependencies. The closure equation adds no external ordering; its internal history can contain ordered causes and choices. By itself the equation settles neither causal determinism nor responsibility.

40.9 Made and found

Chapter two asked why there is anything rather than nothing and answered with a requirement severe enough to have one solution. Then it turned up a clause nobody had written into it: consistency is a property of descriptions, so the structure has to contain something that holds a description and can get it wrong.

That clause leaves a question about what truth is in a structure of this shape. Two answers have been arguing since before either of them had a name. On the first, a claim is true by corresponding to how things are, and a describer's job is to find out which claims do. On the second, what is true is settled by what describers do: their comparisons, their procedures, their records.

Both are here, at full strength.

Found: the consequences of the rules belong to nobody. Chapter two's rewriter produces strings with as many a's as b's, forever, and neither of its two moves mentions counting. Whether nineteen loops on a twelve-patch net sum to zero is settled by an arithmetic that comes out the same for everybody who performs it. The twelve observers inside get no vote.

Made: which record that arithmetic is applied to is written by observers, one commit at a time, out of their own capacity, onto the seams they share. Nothing else writes it. There is no reading of the world taken from outside. Every entry on every seam got there because two bounded things compared what they held and one of them paid for the entry.

So the world is found from the record and the record is made, and those are the two ends of one operation, which chapter eleven pinned down with four demands: hand it a record and it returns the consistent state that record names.

Finding a relation and recording it are events with an internal order. The complete-history closure condition adds no external ordering in which the world must first exist and then be described. It can include both activities and their causal dependencies, without turning every mathematical discovery into a unique global equilibrium.

The argument is usually had about mathematicians, whether a theorem is invented or discovered, as though some fact settled which. Somebody working on a proof is inside the structure the proof is about. What they reach is a fixed point of that work, the thing the work returns unchanged. It was found, in that nothing along the way was free to go otherwise. It was made, in that the working is what put the entries on the record. Nothing else was writing them.

The decision is local and taken many times a day: which entries to put on the seams you touch. Every one of them enters the boundary data that somebody else's world gets computed from. A mismatch written onto another observer's edge does not stop existing when it leaves yours. It sits in their record. The capacity that clears it comes out of their budget.

Chapter 41

What Does a Repaired World Look Like?

The number to know in Butte, Montana is 5,410 feet above sea level, the elevation of the bed of Silver Bow Creek and the lowest ground anywhere in the valley. Below that line, the water standing in the old copper pit stays in the pit. Above it, the water is in the creek, which runs to the Clark Fork and from there to the Columbia. Since October 2019 pit water has been pumped to a treatment plant at Horseshoe Bend, cleaned and discharged to the creek, which holds the level down. The pit has sat about fifty feet under the line ever since.

The water is in the pit because in April 1982 the pumps under the hill were switched off.

Butte sits on one of the richest copper deposits ever found, the workings underneath it run for thousands of miles, and water wants to be in them, so the pumps at the Kelley Mine had been lifting groundwater out of the mountain for as long as there had been a mine to keep dry. They sat 3,900 feet down and could move seven million gallons a day. The Atlantic Richfield Company suspended its Butte operations that month. The pumps went off, and the water came up.

It came up at about a foot a month. It filled the underground workings, and then it began filling the Berkeley Pit, the open pit that had been dug into the east side of town since 1955. Water standing on broken sulfide rock turns acid, and acid takes metal out of rock. The water in the pit reached a pH of about 2.5, the acidity of lemon juice, and carries copper, arsenic, cadmium and zinc in solution.

On 14 November 1995 somebody found dead snow geese floating on it. The first count was 149. The final count was 342. The count had to be revised upward because the water had stained white birds a brownish orange and made them hard to pick out.

Every ton of that copper was legal, paid for and shipped, and the books of the companies that shipped it balanced at the end of every year they were open. The arrangement ran for a century and produced two things: metal, and a quantity of work that has to be done in Butte forever, beginning after the people who produced it stopped being there.

What is that quantity, exactly? Who is holding it? And is there any way to look at an act while it is being done and say whether it lowers the

amount of work the world has to do, or moves that work onto somebody else?

41.1 Two strings

Sound two guitar strings together when one of them is slightly sharp and the loudness pulses. The pulse rate is the difference of the two frequencies, so two strings three cycles a second apart beat three times a second. Each string alone has a frequency and no beat at all. The pulse is a property of the pair, and goes to zero exactly when they match.

An observer stands in that relation to everything around it. It has a state. Around the state is a field, chapter twenty-six's object with a reading at every place: what the neighbors have written at every point of this observer's boundary. The state has a size and a direction, the field has a size and a direction, and the question that decides everything downstream is how much of the state points the way the field does.

The amount of the state lying along the field is the **alignment score**. Take the largest that score could be, which is the two sizes multiplied together and happens when the two point the same way, and subtract the score actually on hand. What is left is a number that cannot go below zero and reaches zero exactly when state and field agree, which is the reading a tuner is listening for when the beat disappears.

That leftover is the misaligned remainder. The work it names is the observer's **repair load**.

The public half of it came in chapter thirteen, where every seam scores the disagreement across itself and the scores add into one number for the arrangement, which every accepted repair lowers. Repair load is that accounting read at a single observer. Chapter thirty-eight priced the business of holding it near zero: neighbors repair too, whatever they write lands on the boundary they share with this observer, and the load comes back the moment the spending stops.

41.2 Exported harm

Mismatch has an address, and chapter thirty-eight's two budgets are the machinery for changing it. Capacity spent inside the boundary brings an observer into line with an entry it leaves standing. Spending on the entry itself changes what everyone wired to that seam will find there. The second move lowers the observer's own load by writing into something a neighbor also reads. The neighbor inherits whatever the write leaves behind.

Ordinary life is made of that move, running in both directions, all the time. Chapter forty found the difference that matters sitting in the record, between a patch that absorbed a mismatch at its own expense and one that wrote it onto a neighbor and let the neighbor pay. The second case takes

a name once two conditions are attached: the mismatch is placed on other observers, and the actor who generated it declines the burden of repairing it while having the capacity to carry it. That is **exported harm**.

An exporter's own readings can be flawless. Chapter thirty-eight's bacterium retunes its receptors until every seam it has reads zero, and starves in the water it is sitting in. Butte is the same arithmetic at the scale of a valley. The books balanced, the copper went out on rail cars, the acid stayed in the hole, and nothing in the accounts was wrong, because the load had been written onto a seam the company was not reading.

Chapter thirteen made that blindness a property of the wiring rather than a failure of attention. Two worlds that put the same packets on an observer's ports are one world to it, so a cost written onto a seam an actor does not share is unreadable to that actor at any level of diligence, which is why exported harm so rarely feels like anything from the exporting end. The reading that would report it belongs to somebody else.

41.3 One integral

An act reaches some observers and misses others. Take the ones it reaches, all of them together, as a single assembly of patches, which is what chapter thirty-seven called a federation. Then ask what the act does to the repair load of that assembly over a horizon long enough to contain the consequences.

The quantity that answers fits on one line.

$$\mathcal{K}_A[a] \;=\; \int_0^{T} e^{-\rho t}\Big(\sum_{U\in A} S_U(t) \;-\; \lambda_{\Phi}\sum_{U\in A}\Phi_U(t) \;-\; \lambda_H\, H_A(t)\Big)\,dt$$

The line says that an act is worth what it does to the alignment of everybody it reaches, minus what it leaves them to repair, minus what it dumps on them, added up across the whole horizon.

Symbol by symbol. The act is a, the observers it reaches are A, and U is one of them. The script $\mathcal{K}$ on the left is what the act is worth to those observers. $S_U(t)$ is one observer's alignment score at record time t, the part of its state lying along its field, and the Greek capital sigma in front of it is an instruction to add over every observer in A. The second term counts what those observers have to fix: $\Phi_U(t)$ is the mismatch standing on U's seams at time t, chapter thirteen's seam sum read at one patch. The third term is exported harm, written $H_A(t)$, the mismatch imposed on the observers of A by an actor refusing to repair it. T is the horizon, the point where the accounting stops, and the elongated S is a sum over every instant between the act and the horizon, which is what an integral is. The two weights λ_{Φ} and λ_H are bookkeeping, because three quantities

in different denominations cannot be added together without an exchange rate between them.

That leaves $e^{-\rho t}$, which shrinks as t grows at a rate set by ρ. It fixes how much a load landing far ahead counts against a load landing next week.

An action is evaluated by what it does to the long-horizon repair burden of the observers it touches. Good is the case where that number comes out higher, which happens when the act lowers what those observers have to repair. Evil is what an observer does when it reads coordinates on itself in chapter forty's sense, can therefore model what a pattern will do to the field before it emits one, and exports avoidable mismatch anyway.

Everything anybody argues about is in ρ and T. Nobody who has ever exported a cost wrote down a value for either, and every one of them picked one. The water under Butte came up at about a foot a month, and a company reports its results four times a year.

The horizon is the harder of the two, because it is set by the world rather than by preference. A horizon shorter than the consequence scores the act on a fragment of what it did. The fragment can carry the opposite sign from the whole. Copper out of Butte between 1955 and 1982, evaluated over the years the mine was open, is a large positive number. The treatment plant at Horseshoe Bend is inside no accounting period that anybody drew, because the plan it runs on has no end date to draw one against.

41.4 Two registers, one shared

Chapter thirty-eight measured the ceiling on private work on four states, one bit where a neighbor reads and one bit of the observer's own. Nothing assembled out of private moves reaches the arrangement in which the neighbor reads the opposite value, however long it runs and whatever it is allowed to spend, and every route that does reach it writes the shared bit.

Put a harm in that shared bit. A burden written onto a record somebody else reads cannot be discharged in private. The reason is which arrangements lie downstream of which moves rather than how hard the person worked at it.

Self-improvement is therefore not a substitute for repair. Somebody harms somebody, and spends the ten years afterward becoming a person who would not do it. Every one of those years is a private move. Every one of them can be real work, expensive and unfeigned, and the entry on the seam between those two people carries the value it carried on the first day. The two destinations are different states. One of them is unreachable from anywhere the private budget can go.

41.5 Elmira, May 1974

One night in May 1974 an eighteen-year-old named Russ Kelly and a friend went through Elmira, Ontario, slashing tires and breaking windows, and by morning twenty-two properties had been damaged.

Mark Yantzi, the probation officer writing the presentence report, put an unusual proposal into it. Rather than sentencing the two and sending them away, he suggested walking them around the town to meet the people whose property they had wrecked. Dave Worth, a volunteer with the Mennonite Central Committee, backed the idea and pushed him to take it to the court. Judge Gordon McConnell ordered it.

So the two of them went around the town with Yantzi and Worth, said at each door what they had done, listened to what the person on the other side of it said back, and worked out what they owed. Two of the twenty-two had moved away and could not be found. Each was fined two hundred dollars, put on probation for eighteen months, and ordered to pay five hundred and fifty dollars in restitution for the damage insurance had not covered, inside three months. Kelly earned his, and at the end of the three months he went back around the town with certified checks.

Each of those doors is a seam. What made the sentence unlike the ones above and below it in the docket is which budget it spent from. A custodial sentence spends the offender's own capacity, at considerable expense, and leaves the entry between the offender and the householder whose window went in precisely as it found it. The walk writes that entry, at every door, and the payment writes it again with a number attached.

41.6 Local pain

If lowering repair load were the whole of the rule, the rule would read as whatever hurts least is best. That reading fails on arithmetic rather than on sentiment. The integral runs to a horizon. Comfort next week sits inside it with a weight, and so does everything after next week, and a term that raises the load for a month can lower the sum.

Local pain is justified when it is the boundary-setting that lowers future mismatch. The opposite failure has a shape everybody recognizes: niceness that preserves abuse blocks repair and transfers the cost forward. A household that keeps its peace by leaving a record unwritten holds its own mismatch flat. The observers who inherit that unwritten record are the ones who pay for the arrangement.

Four moves cost somebody something immediately and raise the long-run integrity of the field: an accurate confrontation, a court verdict, a difficult confession, and a refusal to amplify a popular falsehood. Each of them raises a load on the day it happens, and each writes onto a shared record something that was true and unwritten, which is the write that lowers the

sum. The confession is the expensive one, because the record it puts into the field is about the observer writing it.

41.7 Nobody saw it

The old story says a person gets away with a harm when nobody sees it.

The story is about who was watching. The watchers are not the ones holding the load. A harm writes a new entry into the record, and chapter eleven named what an entry does: it narrows the fiber, leaving fewer whole worlds that carry the reading. Fewer worlds is a change in what can truthfully come next. The narrowing happens whether or not a second party ever reads the entry. An unrepaired harm becomes a residue with an address.

Where a harm is hidden by a falsified record rather than by an unread one, the arithmetic is chapter six's flipped edge exactly. Change one entry from what happened to what did not, two of the loops through it stop closing, and the reading goes unrealizable, chapter eleven's verdict on a record with no world under it. A cover-up buys the offender a record that describes no world at all. Every loop running through the false entry picks up the residue.

A residue with an address gets moved by every repair that runs near it. Chapter six shifted its leftover around the twelve-patch net by reordering the repairs. It weighed one at every address it landed on. That is the mechanism by which an unrepaired harm turns up in people who did nothing: the residue settles on whoever's seams the repairs happen to reach, which in a family or a town is rarely the person who wrote the entry.

Hidden harm is deferred repair. It changes which continuations can truthfully hold the offender, the victim and the community around them, and it goes on changing that whether or not anybody has read the entry.

41.8 Restoration

Chapter thirteen set out what a stopped machine has to carry with it to start again as itself: its records, whatever state it can reach, its wiring, and the law governing what it may do and what it may read. Carry all four across and the restored observer bets the same way on everything it goes on to see. Carry them roughly and the two futures part by roughly that much.

Restoration on those terms says nothing about scanning brains, nothing about rebuilding bodies, and settles nothing about who anybody is.

The law is the piece that gets left behind. Feed a record into machinery whose allowed moves differ and the same history yields a different next step, so the law travels with the record or the thing that comes up is a different observer. A pattern restored under a law that permits the export

it was built around is a continuation of the pattern. Restored under a law that repairs what it receives, it is a continuation of something else, and the argument about whether restoration counts as mercy has to start there.

Records are on the list of four. A restored observer therefore arrives holding its record, and its record contains the harms it did along with the harms done to it, because a continuation that dropped them would be the continuation of a different observer. An offender's restoration and a victim's restoration carry the same entries about the same night. The two readings have to be consistent with one world. That is the condition a transcript, a set of exhibits and twelve people in a room have been approximating for as long as there have been courts.

41.9 A consistent prison

A fixed point is a state a process leaves alone, and chapter two's other root, minus 0.618034, is one no calculator started nearby ever arrives at. Chapter six's twelve observers arrive at theirs from every start the net admits and in every order, with nothing anywhere in the arrangement choosing anything.

Nothing in the definition says the state is a good one. A prison is a fixed point. Every guard's reading answers every other guard's, the count at the gate comes out right twice a day, and the walls hold. An arrangement that exports its costs onto observers with no way to write to the shared record can sit at zero mismatch for centuries, because the ones carrying the load are not the ones whose seams are being scored. A perfectly consistent world can be a perfectly consistent prison.

When more than one consistent state carries a record, the fiber is ambiguous, and chapter eleven left repair with nothing to work on there: every entry is satisfied, so narrowing the set to one member takes something no record holds.

So an ambiguous fiber is settled by a selector. The selector here is which repair law gets implemented by the observers capable of implementing one. A law that leaves a recorded harm unaddressed is a law that some restored observer, reading the complete record, has a true objection to, and that observer is inside the world the law built, holding the record, with the objection intact. The law that addresses every recorded harm is the one every restored observer could endorse with the whole record in front of them, and the only one with that property.

Which fixes where a pattern can be continued. A pattern organized around exporting damage and refusing repair cannot be continued as that pattern into a world running that law, because continuing it there is the export, arriving in a place whose whole arrangement is the repair of what it receives. Its truthful continuation is somewhere its damage stays local. Both destinations have old names, paradise and hell, and the names add nothing to the mechanism. One is the continuation of observers whose

records admit repaired participation with each other. The other is the continuation of an observer that has run out of anybody to pass its costs to.

Every act writes into records that other observers read, so every act exports a pattern into the field all of them share. Some of those patterns lower what the people downstream have to fix. Some raise it, and the repair falls to observers who did not write the entry, at a time nobody set.

The plant at Horseshoe Bend runs on a plan with no end date because a set of records made in Butte between 1955 and 1982 left work that has to be done and can be done. The load has an address, the address can be reached, and there are people in that valley running a pump against it. We compute our destination by the records we make repairable.

Epilogue: The Last Rung

Every reader of this book will die.

What is left afterward can be inventoried by anybody with a scale and a chemistry set. Water, carbon, calcium, phosphorus in the bones, a little iron in the blood. Every item on that inventory is a material. The inventory can be completed. Chapter thirty-five ran the same procedure on the world and came back with ten rows: space, time, mass, energy, motion, force, matter, charge, light, and the record. That list is complete as well. Nothing on it is a material. Mass, the one row a scale would respond to, is a bill that repair pays at every event.

The observer came through chapter thirty-five undissolved, because every row on that list was assembled out of observers, and chapter nineteen gave the parts list for one: an algebra of questions bounded by the interface, a record algebra in its center, which is where writing stays put, an interface that behaves identically every time it is used, and a state fixing the odds on whatever gets asked next.

Go through those four looking for a material. An algebra is a collection of operations that compose, scale and add. Its center is the part of the collection that commutes with the rest. An interface is a set of ports and a rule for what appears on them. A state is a table of odds, one entry per question. The thermostat on the wall in chapter three has all four, at a question list one item long, and the whole of its interior is a sensor, a number somebody set and a switch, with none of the four objects among the parts. What you are made of is the separate question chapter thirty-four answered, and its answer covers the chair and the thermostat and the reader at once: a network of observers repairing their disagreements, averaged down and copied often enough that the order of the questions stops mattering.

Records copy

Of the four objects, the record algebra is the one that copies, and chapter nineteen put a mechanism under it. Writing a record runs the observer's questions through the mail room: whatever falls inside one alternative is kept, and every comparison that only meant something across two of them is destroyed. What survives that sorting commutes with everything the observer can ask, which is why a record can be read without being spent, and why the center of the algebra is where tables and pointer readings live.

Chapter twenty priced the consequence in two lines. A reversible machine leaves the overlap of two states where it was, and a machine that hands each of them back twice squares it, so a copier works only where the overlap equals its own square, which leaves zero and one. A polarization nobody has measured sits between the two. No machine duplicates it. A record sits at zero against every other record by construction, so records copy freely and the copying leaves the original as it was.

Redundancy runs on that gap. Chapter thirty-four's chair has a definite place because sixteen copies of a sixteen-bit fact push forgery below one part in ten to the seventy-seventh, and the room goes on copying a chair long past sixteen. What can be copied about an observer is exactly what has been written into its records.

Deleted by what?

The flying clocks compared their readings at reunions, the only place two clocks can compare anything directly. Readings at distant events can be compared too, by agreeing on how to synchronize and how fast the signals travel, and that agreement is a convention two parties build. No universal present comes with it.

Chapter seven took the shared present apart. A partial order can be extended to a total one, and a spacetime can be given coordinates that run everywhere at once, each in many ways. None of them is the clock of the world, and two paths between the same two events accumulate different proper times whichever one is chosen.

Chapter thirty-seven found a complete history fixed by its own description and realized by its own internal interactions. The fixedness belongs to that closure. Inside the history, events are ordered where dependencies connect them, records form and local states change, and a solver's step count is a different object from all of them.

No clock outside advances the history, so nothing outside deletes the past either. What happens to a particular record is a question for the dynamics inside it: a repair can overwrite it, a copy can keep it somewhere else, and chapter twenty-three prices each of those.

Restore the four

A live machine moves between hosts without the programs inside it noticing, and chapter thirteen took the list of what has to travel for the machine that starts to be the machine that stopped: the records, the state it can reach, the ports and their routing, and the law fixing which moves are available. Those four are a checkpoint. The fourth is the one that gets left behind, which is how a restored thing comes to read its own history and do something else with it.

Two checkpoints matching on all four put identical odds on everything the observer can register. Two that match approximately come apart by no more than the approximation.

Chapter thirteen made that claim about a machine on a rack. Chapter forty-one turned it on an offender and a victim holding the same entries about the same night. Turned on the reader, it reads the same way: restore a checkpoint with the same readable records, the same accessible state, the same interfaces and the same future law, and its future statistics match the original within a stated error.

It says nothing about scanning a brain. No instrument in any laboratory reads those four objects off a living one. It says nothing about rebuilding a body, which is a separate biomedical problem. It does not settle whether the restored observer is the person who was checkpointed, which is a question about identity that the arithmetic does not address. And it promises nobody anything, because nothing in the structure performs the restoration.

The selector

Stop one of chapter eleven's twelve patches from talking. Five seams go unread, thirty entries become twenty-five, the eleven patches that carry on stay pinned where they were, and the silent one's coin is loose: four settings of the twelve fit every record in existence. That is an ambiguous fiber. Repair has nothing to bite on there, because nothing anywhere disagrees, and whatever settles the four settles it on grounds outside the record. Settling on grounds outside the record is choosing.

A selector settles an ambiguous fiber, and chapter forty-one found one on the board: which repair law the observers capable of running one actually run. Leave a recorded harm unaddressed and somebody restored into the world that law built is holding the entry, with an objection the law has no answer to. Exactly one law leaves nobody in that position, the law that addresses every recorded harm.

Observers capable of running a repair law act within the history. Working out an architecture before building it is an order of events inside the structure, and the closure condition on the complete structure is timeless. The choosing happens inside, in the order the records keep, by the observers who hold them.

Forty rungs

1. Why is the universe so hard to reverse engineer? Every program that tried surrendered one assumption and defended the arena.
2. Why is there anything rather than nothing? Self-consistency alone picks one number out of the continuum, 1.6180339887.

3. Why would a universe need anybody in it? Put the stage in first and the stage is the one thing left unexplained.
4. Why do two descriptions ever agree? They have to match where both look, and a loop of exchange rates has to return what it started with.
5. Why do you and I see the same table? Whatever differs between two descriptions of it was never physical.
6. Why isn't agreeing with your neighbors enough? Every pair can agree while a loop carries a residue, and a leftover of exactly one survives every repair order.
7. Why is there no such thing as now? Causality is a partial order, a timeline is a total order, and nothing in the arrangement picks the tiebreak that turns the first into the second.
8. Why can strangers agree without a boss? Influence runs only along overlaps, at most one seam per repair event, and what can reach an observer is a finite dependency cone.
9. What move does reality actually make? Split the difference and let a fair coin place the odd unit; interference experiments read what each repair writes down.
10. Why doesn't it matter who goes first? 102,415 disagreeing seams reach zero, sixteen shuffled replays return one identical finishing state, and Newman's two conditions say every other order lands there too.
11. What does everybody actually end up holding? Objective reality is the observable normal form, unique relative to the boundary data.
12. Why is the world made of relations and not of things? The sourced action is gauge-invariant if and only if the source is conserved.
13. What is one piece of reality actually made of? Six parts and six moves, and a bounded interface is a bound on what can be known about a region.
14. Why twelve? A closed carrier with five seams at every port admits twelve ports, thirty seams and twenty faces, and no other count.
15. Why does space have three directions? Only the slowest-fading block survives enough rounds to carry a position, and its rank is three.
16. Why does time run, and only one way? Every commit throws away what it resolved and every quantum repair drains coherence, so the arrow comes from repair and owes nothing to how the universe started.
17. Why is there a speed limit? The angle-preserving transformations of an observer's sky are the Lorentz transformations, and records reading their neighbors round by round fill out exactly the cone those transformations preserve.
18. Why does anything have energy? A specified coherent generator supplies an energy readout after clock calibration. Repair costs and modular generators provide finite interfaces; their identification with physical energy requires that matching.

19. Why can't you ask everything at once? Refusing to answer two questions together is what makes an undocumented machine readable from a seat inside it, and the questions that refuse nothing form the center where records sit.
20. Why does nature cheat, but only by exactly $2\sqrt{2}$? The ceiling is a product of commutators, and sits at 2.8284.
21. Why does time feel like it's moving? The state supplies its own flow, the flow is trivial on records, and an observer's clock steps it once at every event.
22. Why do all the clocks agree? Within each stated construction, event tallies, field response and clock readings are related by explicit maps. Their parameters and calibrations are named; no universal physical clock is required.
23. Why does writing something down cost heat? All four laws of thermodynamics come out of one repair rule, and at three hundred kelvin clearing each yes-or-no question a commit settled costs at least 2.87 joules divided by ten to the twenty-first.
24. Why does a region's memory scale with its surface? Everything a region can show crosses the screen around it, and the screen's memory is a count of patches that each hold the same amount.
25. Why does gravity look like geometry? Record constructions provide geometric readouts. The Einstein-form theorem requires the stated null balances, conservation and entropy hypotheses; a repair-selected responsive geometry remains open.
26. Why exactly these forces? The constrained finite port-current construction gives one plus three plus eight. Its physical field interpretation and global gauge lift retain their declared premises.
27. Why this exact list of matter, and why three times over? The declared matter lift gives fifteen internal states and a restricted anomaly-free menu. A rank-three family candidate needs its physical attachment and mass-generating dynamics.
28. What is a particle, what is a wave, and why does it look like both? Fields, excitations and detector records offer related conditional descriptions. Identifying the observed particle species and masses requires their spectral response in the same history.
29. Why 1/137, and why a capacity limit? Specified cell and horizon consistency maps have certified roots. The shared-width alpha map gives inverse 137.035660; Wissner-Gross's separate entangled-pair candidate gives 137.035999189566. Physical electromagnetic identification remains open.
30. Why do the masses land where they do? Measured electron and muon masses, a balanced-face assumption and a mass-ordering choice constrain a tau comparison inside a 72-electron-volt window. The full observed spectrum is not source-selected.

31. What is a Lagrangian, and why is it the last thing? A specified effective action compresses dynamics. Invariant interaction channels are verified, while physical coefficients, the vacuum scale and the coupled geometric response remain to be selected.
32. Why does anything move? The inverse-square exponent is a subtraction, three directions less one.
33. Why was every open problem the same problem? The proposed mechanisms share an observer-based description. Each finite result and conditional interpretation retains its own hypotheses; this common language does not close every physical problem.
34. Why was Newton right for two hundred years? Redundancy is what makes anything definite, and the layer he wrote down has no motion of its own.
35. What is anything made of? Ten entries, each a rate, a count, a rank, a cost or a commitment, and no substances.
36. Why these laws and not others? The average score climbs at the rate the survivors disagree with each other, and what got selected was a repair rule, chosen from inside, with no ensemble of worlds to compare against.
37. Why does the loop have no beginning? The complete history is its own intrinsic realization, and its closure equation carries no step counter and no outside start.
38. Why does anything try to persist? No amount of private repair reaches a state that one network update reaches with certainty.
39. Why is a bacterium freer than an atom? Room to be otherwise increases the further down the tree you go.
40. Does it feel like anything, and what is left to decide? The endpoint is fixed relative to the boundary data, and what you admit, emit, repair, ignore, confess, protect and amplify is the boundary data.

There are forty-one questions in the book and forty rungs on that ladder. The forty-first is the one under your feet. Chapter forty-one put a number on an act: the repair load it leaves the observers it reaches, summed out to the horizon. Then it left the writing to whoever is holding the book.

The records other observers keep of you were written by what you did to them, and records are the whole of what any selector has to work on. Chapter forty-one put the old names back on both destinations. The mechanism runs the same without them. The part of you held somewhere other than in you is what those records say. Any restoration has to be consistent with it. There is no choice about being recorded. The choice is what the record says, and everybody is writing one.

Paradise, on these terms, is settled by the entries, and names the continuation of observers whose records admit repaired participation with each other.

Appendix A: Eighteen Questions

Between 1982 and 1985 Eugene Miya, of the Computer Systems Division at NASA Ames, stopped answering the same questions on the SPACE mailing list one at a time and began posting the answers on a schedule, yearly at first and then monthly. The format spread across the network and picked up a name, frequently asked questions. Jef Poskanzer was the first to post one weekly, to the graphics group.

A question arrives a thousand times because a thousand people share one assumption, and an answer that leaves the assumption standing produces the same question the following month, from the same people, worded differently.

Eighteen come back about the account in this book. The answers retain the earlier chapters' finite domains and physical premises. A conditional effective description is not yet a source-selected model of every observed particle.

What about the Big Bang?

Everything came out of something hot and dense about fourteen billion years ago, so what was there before that, and how does an account built on observers get started in a universe that contained none?

That is two questions welded together, and both halves lean on a first instant seen from outside: a moment when the arrangement switched on, watched by nobody in particular, with observers appearing later inside it.

The hot dense era is the far end of a reconstruction built from records held inside finite patches. The microwave glow left over from when the plasma stopped being opaque is a record. So are the ratios of the light elements, the spectra of galaxies and the catalogs of supernovae. The Big Bang names the place where the record side of that reconstruction goes dark. As for before, the only ordering there is runs on committed repairs, and where none have committed the word has nothing to be a relation on.

Two things follow. The arrow of time is not inherited from that era, since a commit destroys what it resolved, and does that at the ten billionth repair as thoroughly as at the first. And the early universe is the most constrained object there has ever been: twelve ports, three directions and one way round the arrow, with no earlier record for any of it to have come from. The grain,

chapter twenty-nine's number for how fine a distinction a screen supports, shows up on the microwave sky, whose ripples are slightly stronger at large angles than at small ones. A forty-eighth of 1.6309682414 is 0.033979, and one minus that is 0.966021, the spectral index that measures the tilt. The Planck satellite measured 0.9649 give or take 0.0042. The Atacama Cosmology Telescope's 2025 analysis, joined with Planck and with maps of galaxies, sits higher, at 0.9743 give or take 0.0034. That combined analysis places 0.966021 in its lowest 0.6 percent, a tension the next generation of microwave maps will sharpen or remove, and the grain fails if the index lands well away from 0.966021.

If none of this is fundamental, why does my car work?

The mechanic balances a wheel using a picture in which the wheel is an object at a position in a space, turning at a rate, and the car drives home. Whatever has been said about ports and repairs, it changed nothing under the hood.

Underneath it sits a rule about descriptions: one earns its keep by being the bottom one, so demoting the everyday container to a reading must take something away from the garage.

That container is the coarse-grained, heavily copied reading of a network of observers repairing their disagreements. Taking that reading is what a body does all day. The size of the error in it has been measured. In 1960 Robert Pound and Glen Rebka used the 22.5-meter tower of the Jefferson Physical Laboratory and an iron-57 source to catch a clock in a basement running slow against one on the roof, by 2.56 parts in ten to the fifteenth, give or take 0.25.

So the corrections sit nine decimal places below anything in a workshop. They arrive when the resolution rises. A clock in a navigation satellite runs fast by forty-five microseconds a day and slow by seven, and the correction had to be built into the hardware before the network could find anybody. A wheel balancer reads none of that. The wheel comes back balanced.

Why does the Lagrangian come last?

Anybody who has taken a physics course met the one line in the first week and spent the rest of the term working out what was inside it. Here it turns up after the forces, the matter and the numbers, which looks like the argument running backwards.

The assumption is that the most compressed statement of a subject is its foundation, so whatever fits on a coffee mug must be where the world starts.

A Lagrangian is a compression. A compression summarizes something that was there first. Written on line one, the line carries three families because three families were counted, a group because detectors reported one, and a page of numbers because laboratories measured them. Written last, it records the finite response algebra and the conditional matter lift, while naming the three-family attachment, mass-generating coefficients and scale as remaining inputs. A three-corner face is a candidate family attachment, not a proof of the physical family count.

The compression is also lossy, by an amount somebody can write down. Add a term built to vanish at every configuration the world realizes, and every realized history keeps its score, every stationary history stays stationary, and the same run comes out most probable. The momentum changes, since momentum is read off how the line responds to a rate of change. The added term tilts where nothing is ever realized. Two lines, the same histories, different energies, and nothing in the world's own histories picks one of them over the other. The one on the mug is the best summary anybody has written.

Twelve. Where does a number like that come from?

A patch with twelve ports carries the forces, the matter and the three directions. A specific small integer sitting at the base of a theory is what a fitted parameter looks like.

The suspicion assumes the number could have been otherwise, which is what a parameter is: a quantity whose value had to be chosen and can be moved.

Count it instead. Every port meets five triangles, every edge belongs to two faces, and counting the same thing two ways forces the Euler relation, after which a spherical carrier has twelve ports, thirty seams and twenty faces. The count runs backwards as well, since twelve ports of that kind force the sphere.

Change the degree and the arrangement fails. Drop the spherical closure and it fails. Try the cube and it admits no such structure at all. Then the wiring: of the 10,395 ways to pair the ports up, incidence at distance three selects one. Each of the nineteen numbers typed into the one line can be dialed to a different value and the equations survive. Twelve has nothing to turn. Move it to thirteen and the alignment that lets two patches compare readings stops existing. There is no thirteen-port world sitting elsewhere with slightly different physics in it.

If space is not a container, where is any of this happening?

Twelve ports, thirty seams, patches wired to neighbors. Draw that on a page and the page is somewhere, so the network has to sit in something, and whatever it sits in is the real space, which puts the container back one level down and wins the argument.

The question hands itself a place for free: everything is in something, and a relation needs a room to be a relation in.

Distance here is a count of comparisons. Two patches are far apart when many comparisons separate them. A direction is a pattern in that tally rather than an arrow drawn in a room. The three directions come out of a fading rate inside each carrier. A repair operator's twelve readings split into blocks that fade at 0.9539, at exactly nine tenths and at 0.8794 a pass, with only the slowest block surviving enough rounds to carry a position, and the rank of that block is three.

Turn the question on the room it asks for. A room for the network to sit in would have to be assembled out of comparisons between record-keepers, which is what a network is, and then that room would need a room. What exists is a tally, a rank of three, and observers doing the counting from inside while occupying no position in anything larger.

Spacetime goes the same way. Which record read which is a partial order, and counting that order is enough to find four dimensions in it: inside any small diamond of events a tenth of the pairs are ordered, which is the value for three directions of space and one of time, and the fourth root of a ratio of counts is a ratio of durations. The order needs no room to happen in either.

Is the moon there when nobody is looking at it?

The world is made of records held by observers. Turn the observers away and nothing gets written, so the parts nobody attends to should go soft. The moon on a cloudy night is the test case everybody reaches for.

Two things ride along inside it: that an observer is a person, and that looking is something a mind does.

An observer here is anything bounded that keeps records. Sunlight leaving the moon is a comparison, the tide in the Bay of Fundy is a record of where the moon was, and the rock holds its own state against everything it touches, with no person watching.

Definiteness is a count of copies. Sixteen copies of a sixteen-bit fact drive the odds on a forgery below one part in ten to the seventy-seventh without adding a single bit of content, and an object that size is copied into its

surroundings far more often than sixteen times. The answerable question is how many records carry the moon's position. It is written into every square meter of sunlight it turns back, which is why its place is as definite as anything in the sky. Ask about a single atom nothing has interacted with and the odds stop being one and zero, which is the same arithmetic read at the other end.

If there is no now, why does time feel like it passes?

No finite exchange of messages between observers recovers a shared present. Then there is the clock on the wall, the coffee going cold, and the distinct impression of moving forward through the afternoon.

The word is doing three jobs at once. The question welds them into one object and then finds that object missing.

Order is a partial order on committed repairs, the only global temporal structure there is. Direction comes from the commit, which is an arithmetic loss. Seven possible pasts average into one reading, and 2.807 bits go with them. Flow is what a state makes its own algebra do, one shuffle for each amount of shuffling, extracted from what an observer holds with nothing added.

A universal present appears nowhere on that list. The part of a person that has been written into a record is the part standing still, because the flow moves only what fails to commute. A record commutes with everything. Read the same flow sideways in empty space and it is a temperature, one that acceleration alone sets, at about four parts in ten to the twenty-first of a degree for every meter per second squared.

When exactly does the particle decide?

A pattern of bands appears on a plate when nothing records which way each particle passed the needle, and disappears when something does. Somewhere in there a thing that was spread out becomes a thing in one place, and the question is when, and how fast, and what sets it off.

Asking for the moment assumes a mechanism with a rate and a threshold, sitting in the apparatus, waiting to be caught in the act.

A record gets written. Between commits, every question about the spreading pattern lives in the part of the algebra where questions refuse to be answered together, and the phase and the frequency are the content. At a commit an entry lands in the center, where everything commutes, and what the plate holds afterward is one more count at one place. Wave and particle differ by how much of the thing has been written down. The plate writes down only the arrival.

So there is no rate to measure and no threshold to find. What gets fixed instead is the odds. Any assignment of numbers to outcomes that lands in the unit interval, gives the certain event one and adds up on every legal sum of questions, blunt ones included, is the trace against a single density matrix, in every finite dimension. Alongside it sits a fake, built to reproduce every count a machine actually produced and belonging to no state at all: it scores one pair of questions 35/64 where adding up demands 143/256.

Where does the explaining stop?

Conditional charge representations, candidate family bands and electromagnetic trial maps can all be constructed on the same carrier, and anybody handed one number after another is entitled to ask where the supply ends.

The question expects an evasion, on the assumption that a structure either explains everything or is hiding something.

The stopping point is printed to the same precision as the results. The declared lift fixes the internal charges, and the stated band-selection rule offers a rank-three family candidate. Neither selects the physical family matrices or mass scale. Every angle between two symmetry axes of the twelve-port solid was enumerated, four hundred and sixty-five of them, and the smallest nonzero one is 20.9052 degrees, while the Cabibbo angle measured in the laboratory is 13.0029. No angle on that list lands within seven degrees of the measurement.

The declared electromagnetic trial map certifies 137.035660 where the quoted measurement reads 137.035999, and the interval enclosure rules out numerical error in that map as the source of the difference. The physical low-energy transport is incomplete. No calculation here proves that the omitted quark contribution removes exactly that mismatch. Wissner-Gross's separate entangled-pair expression has inverse 137.035999189566; identifying it with the physical electromagnetic coupling and the same-cell equation retains the independent premises stated in chapter twenty-nine.

You knew the answers before you started

Every one of these numbers was measured decades before the arithmetic that lands on it was written down. Sommerfeld put the fine-structure constant into physics in 1916, the tau mass has been known since the 1970s, and matching a number in a table is what any sufficiently flexible scheme can do.

The complaint assumes flexibility, and flexibility is countable.

A fixed declared map can be tested without adjusting its output, and the certified maps here have that scope. Choosing which map is physical is

a different freedom. The grain equation, the face attachment, the balance condition and the physical field readout each need their own justification. Measured mass inputs and unselected coefficient matrices remain inputs even when an accompanying finite formula is exact.

Under the fixed-horizon-capacity hypothesis, dark energy's pressure-to-density ratio is exactly minus one at every epoch, $w = -1$, with no field rolling anywhere to absorb a change. The test is live: five thousand and twenty robotic fibers on the four-meter Mayall telescope at Kitt Peak collect the spectra that decide it. Their second data release leans against a constant ratio, by 2.6 to 3.9 standard units across its combinations, and the third release is the test fixed in advance. A measured evolution of w breaks the account.

Isn't this what Leibniz said?

Space as the order of things that coexist, inertia coming from the rest of the matter in the universe, quantities nobody can measure from inside declared to be no quantities at all. Leibniz, Berkeley and Mach between them cover a good deal of what is asserted here. They managed it without a twelve-port carrier.

A position and a theory get treated there as the same kind of object.

Leibniz was right, and two hundred years of physics proceeded as if he had not been. An objection that forbids a quantity without predicting a measurement changes nothing in the textbooks, and absolute space sat in them until an experiment forced the issue.

The objection here comes with arithmetic attached. Twelve ports and thirty seams. One rule that averages two disagreeing readings, forced by three demands: touch only the pair that disagrees, leave their sum where it was, and end in agreement. A symmetry that has to act from within, since there is no outside desk from which twelve ports could be relabeled. And a group of one plus three plus eight, left standing after the block structure removes the only alternative the dimension counting allows. Leibniz had the objection and no wiring diagram, which is why Clarke could answer him through five letters and concede nothing.

So we are living in a simulation?

Records, repairs, updates, capacity. The vocabulary is a data center's vocabulary. The word turns up within a paragraph or two of the parts list.

The word carries an original. A flight simulator earns the name because there are aircraft aloft doing what the box is pointed at. Nothing is an imitation on its own.

Put the question to the inventory and look for the second end. Name the entry that substitutes for something, or the warehouse where the original

repair rate is kept, or the object a loop residue resembles. The two-rule rewriter produced strings that described nothing filed anywhere else. Run the rules and the strings you get are the strings there are.

The word computed arrives with fake attached to it. Watch one repair: a disagreement that was there is gone, the total sits where it started, and which of seven arrangements the seam held is gone from the readings. That is a change in the world, carried out by arithmetic. Coffee tastes the way it does by the same route, and so does a magnetic field. And nothing is stepping any of it: eighty-one thousand nine hundred and twenty patches reached agreement on every seam between them without a tick from anywhere. The cycles in that run's trace belong to the computer that kept it.

Does an observer have to be conscious?

Observer-first sounds like a claim that the universe waits to be noticed. The version of quantum mechanics that reaches most people has a mind collapsing something at the end of it.

The assumption is that observing is a mental act, so the word carries a mind wherever it goes.

Keeping records takes three things and not one of them is a mind: a bounded view, somewhere to write, and a reading that makes a difference. A thermostat screwed to a classroom wall has all three. None of the results built on the list requires an observer to have a mind.

Consciousness is a special case, defined in chapter forty at the size the structure supports: owned boundary repair, read from the inside. An observer that holds coordinates on its own state, and computes its next move off them, is reading the half of the record that the map to the public list deletes. Which organizations of that reading are lived, and how finely a person's runs, are questions that chapter takes up. Nothing in the structure lets a wish move an entry.

Isn't the loop circular?

Observers build the account of the world and the world builds the observers. Anybody trained to spot a circular argument spots one here immediately, and circular arguments have the property of proving whatever you like.

The objection assumes the loop is an argument, whose starting points would have to be in place before its conclusions.

The fixed point is a complete history under its self-reference map. It needs no first solver outside it, and it contains internal causal order, local changes and the computations through which the structure realizes its dependencies. A terminating repair solver, a stationary state and an ob-

server's calibrated clock are three different objects, and calling each of them a fixed point does not make them one.

A loop of this kind is a condition. It constrains a description and its realization together, the way an equation constrains its unknown, and the history either satisfies it or does not. Nothing in it needs to come first. Part of the demand can be checked on hardware: cut the run across a thousand machines and it finishes in the state the single machine finishes in, with the worker count, the restart history and the schedule appearing in none of the settled readings anybody inside can take.

Does this mean my choices are already made?

Twelve patches converge on one arrangement from all two thousand and forty-eight starting states, under every schedule. Eighty-two thousand patches bring 102,415 disagreements to zero whatever order the repairs run in. If the endpoint depends neither on the start nor on who went first, then a person weighing a decision at two in the morning is a subsystem executing a delay.

The argument drops four words from the theorem it is quoting: given the boundary data.

Those four words point at the entries a region cannot touch, the ones sitting on its edge, set from outside it and different from one region to the next. Turn one of those thirty entries from same to different, and twelve coins that had exactly one consistent setting have none, and what the network settles onto carries a leftover of exactly one. One entry, and a different world.

A repair is exactly the move that leaves those entries alone, so they have to be written by something else. That move spends capacity on an overlap. No amount of private work reaches the same place at any budget. In a person's vocabulary those writes are what to admit, what to emit, what to repair, what to ignore, what to confess, what to protect and what to amplify. The convergence does not run around an observer doing that, and what it finds on the way past is what that observer put there.

Is this a religion?

The account ends in the words good and evil, a law that addresses every recorded harm, and two destinations that have old names. Readers who have watched physics arrive at that address before are entitled to ask what happened.

Subject matter is doing the categorizing: touch those topics and whatever you are doing becomes doctrine.

Every statement here carries something that would contradict it. A measured evolution of w takes the constant capacity and the fine-structure

closure with it. One proton decaying through the unified channel takes the argument about the proton. There is no authority to defer to and no text to accept: the twelve-patch enumeration runs over two thousand and forty-eight states and can be done by hand in an evening.

The traffic has run the other way before. Maupertuis announced least action to the Académie in 1744 and took it for proof of a supreme being, because it looked like the universe economizing, and the economizing turned out to be what taking logarithms of a product of local factors does. The claim about continuation is a statement about a checkpoint: records, accessible state, interfaces and future law. Nothing in the arrangement performs a restoration, so nobody is being offered one.

Who decides what counts as harm?

An ethics that scores actions by repair load needs somebody doing the scoring, and the scorer's preferences are what usually end up in the score. Whoever holds the pen decides what counts.

Harm arrives in it as a verdict, which is why a judge has to be standing somewhere to deliver it.

Mismatch scores itself at the seam. Push one packet through the dictionary sitting between the two, hold the result against the other, and the disagreement comes back as a number nobody chose. Exported harm attaches two checkable conditions to that number: the mismatch lands on observers who did not generate it, and whoever did could take the repair on and does not.

Only two quantities are open, the horizon and the discount rate. How far ahead the accounting runs, and how heavily a load landing in a century weighs against one landing next week, are what every argument about conduct is actually about, and an exporter of costs picks values for both without ever writing either down. An actor learns what arrives at its own ports, and a cost parked on a seam it does not share never arrives there, which is why the books balanced at the end of every year the mine was open.

Do I have to accept the ethics to accept the physics?

A reader can want the derivation of the forces and the three directions and want to stop before the part about good and evil, on the ground that a physicist's opinions about conduct are a physicist's opinions.

The question assumes the join between the two is preference.

They share a quantity. Mismatch is the number a repair reduces, and the whole climb is about what happens to it: the leftover of exactly one that no repair order removes, the standing cost of holding a pattern together

against every seam it touches, the heat that comes off a write. The ethics reads that same number at a single observer, adds a horizon, and asks what an act does to it between here and there.

So the physics can be taken alone and nothing above it breaks. The claim that acts leave shared records untouched cannot be kept, because writing onto a shared record is the move that makes agreement possible in the first place, and it is the same move that puts a cost on somebody else's side of a seam. The ethics runs on the two budgets the repair law runs on: what an observer can fix inside its own boundary, and what has to be spent on an overlap with somebody else. Accepting the repair law is accepting both, and the number on a seam is zero when two readings answer each other and positive when they do not, whatever either party calls what happened.

Appendix B: Glossary

Michael Faraday distrusted the word pole. It had been attached to the two ends of a battery for decades and it smuggled in a theory, that something sits at each end attracting and repelling across a distance, which was the account of electrochemistry he believed was wrong. What he wanted was a pair of words naming where the current went in and where it came out, committed to nothing else.

So in the spring of 1834 he wrote to William Whewell at Trinity College, Cambridge, who had the Greek. Candidates went back and forth between them: eisode, exode, zetode, dexiode, skaiode, orthode, anthode. On 5 May, Whewell came back with anode and cathode, on the ground that short words get adopted and elaborate ones do not, and threw in ion, anion and cation while he was about it. Faraday replied ten days later to say he had taken all of them, and that his colleagues' objections subsided the moment he mentioned whose idea they were.

Anode and cathode name a direction and a doorway. They say nothing whatever about what is traveling, which is why they outlived the theory they were coined for. The entries below try for the same restraint.

Action. A single number scored over a whole history, one contribution per step from start to finish. The history the world runs is the one where nudging an interior reading leaves that total alone.

Algebra of observables. Every question an observer can put, together with the three things you may do to questions: ask one after another, scale one, add two.

Alice and Bob. Two observers with separate views and no access to each other's insides, on hand whenever a claim has to survive being checked by somebody else. Charlie joins when two parties are not enough to make the trouble visible.

Alignment score. How much of what an observer holds points the way its field does. Whatever fails to point that way is the repair load, so the two are one reading taken from opposite ends.

Antimatter. For every list of matter states there is a mirror list with every charge reversed. The two are one object, and a world built on either gets the same physics with the labels swapped.

Area law. An entropy or capacity contribution proportional to the area of a region's boundary in a stated regime. A black-hole horizon is the best-known example: twice the radius buys four times the area. Thermal

entropy can scale with volume, so the quantity and state being counted matter.

Bit. The answer to one well-chosen yes-or-no question, and the unit anything measured in questions is counted in. Erasing one releases heat, and how much depends on nothing about the device, only on the temperature around it.

Carrier. The finite wiring of the model: twelve ports at the corners of a twenty-sided solid, thirty seams joining them, twenty triangular faces. The structural numbers of the physics are read off that arrangement.

Causal diamond. Everything after one event and before another: two cones joined at the rim. Count the completed records inside one and the count approaches its volume, and the fourth root of a ratio of two such counts is a ratio of durations.

Causal set. A partial order in which any two related events have only finitely many others between them. The 1987 proposal is that spacetime, looked at finely, is one of these, with the order supplying the light cones and the number of elements supplying volume. The observers' records form one.

Center. What commutes with everything else in sight. In an algebra of questions it is where records live, which is why the public world looks classical; in a group of moves it is the moves nobody inside can detect.

Charge. A number a closed loop of seams hands back to whoever walks it. Renaming conventions cannot reach it, because a round trip through your own conventions returns you to them, and repair cannot flush it out, because repair reaches one seam at a time.

Checkpoint. The four things that have to travel for an interrupted observer to be the same observer when it resumes: its records, its accessible state, its interfaces, and the law saying which moves it may make.

Chirality. Handedness. A pattern with a heading and a spin either turns the way a right-handed screw advances or turns the other way; a mirror swaps the two cases, and the world treats them differently.

Collar. The thin band of patches wrapped around a region's edge, with the region inside it and everything else beyond. When a collar screens, the inside can be rebuilt from the collar alone, and whatever the outside learns about the inside crosses that band.

Commensurability. Whether two observers' readings answer each other at all. The rule saying which of your readings answers which of mine sits on the seam between us.

Commit. An accepted update, with its reads, its writes and the record it leaves. Midpoint averaging is many-to-one on the two readings it updates, and the record keeps what the readings forget. A count of commits orders events, and seconds come from a clock laid against the count.

Commutator. What you get by doing two operations in both orders and subtracting. Zero means the two questions can be answered together, and anything else is the amount by which they cannot.

Consistency. A description is consistent when it does not require two incompatible things at once. An inconsistent one buys nothing, because a ledger saying the account holds both nothing and a hundred pounds justifies any withdrawal you like.

Decoherence. The usual word for what happens to alternatives once a record has been written and the comparisons between them stop being available. The mechanism is the record.

Density matrix. The description left behind when somebody is cut out of the picture: odds for every question you can put, and nothing left of what only the far side could have detected. Two different accounts of the whole can leave the same one behind, and no operation on your side tells them apart.

Dependency cone. The events or patches that can affect a readout. Dependence crosses at most one seam per repair event, so a round in which every seam is repaired once widens the cone by at most one seam, and a sweep along a path carries a chain several seams, one event per seam.

Derived. Obtained by arithmetic from the structure, with nothing measured fed in to produce it. A number fitted to a measurement and then read back out has been derived from the measurement rather than from the structure.

Direction. The one-way character of time: the loss inside a single repair, where seven arrangements of a seam collapse into one and quantum coherence drains away for good. It is separate from causal order, which says what depends on what, and from flow, which runs both ways.

Direction split. The fork at every repair: spend your own capacity on a private update, or spend the capacity shared out on the overlap. Some outcomes are reachable only the second way.

Duality. Wave and particle are one thing at two depths of record. Between commits there is a spreading pattern with a phase whose questions refuse to commute; at a commit there is a record, and records hold ordinary numbers.

Effective description. An account that works over a stated range, takes its constants from measurement, and can point to where it will break. Every working piece of engineering physics is one.

Eigenvalue. A grid of weights usually swings a direction round to point somewhere else. Where it merely stretches or shrinks one, that direction is the grid's own, and the factor is its eigenvalue.

Energy. The generator of a specified physical time flow. In thermal equilibrium, when that flow is calibrated to the state's modular flow, its Hamiltonian is minus the logarithm of the state times the temperature, up

to unit conventions and an additive constant. A modular generator alone does not establish that physical calibration.

Entanglement. Content living in the relation between two halves of a thing rather than in either half. Cut a definite whole in two, look at one side alone, and what comes back is vague by exactly the amount the cut severed.

Entropy. How many yes-or-no questions it would take to fix which arrangement you are holding, given what you know about it. It is a reading of the description rather than of the object, and the word disorder has no place in it.

Equivalence principle. What a thing costs to hold together sets both how hard it is to push and how hard gravity pulls on it, so the two readings never come apart. The MICROSCOPE satellite spent five months of free fall trying to catch a titanium cylinder and a platinum one falling at different rates, and reached one part in a thousand million million without catching them.

Exclusion principle. Two matter patterns cannot sit in the same one of the forty-five states on an observer's screen. Matter stacks into shells and solids because of it, while a laser piles any number of photons into one mode.

Exported harm. The mismatch an actor leaves for other observers to carry, having declined to repair it. A cost written onto a seam the actor does not share never appears on the actor's own readings. It appears on the readings of whoever holds the other side.

Fiber. All the consistent states of the world carrying one given public reading. An empty fiber makes the reading impossible, a fiber holding one state makes the reading enough to fix the world, and a fiber holding many means something besides the reading has to choose.

Field. The word names two objects. One is a value assigned at every place, like the temperature through a room or the wind over an ocean; the other is a net of algebras, which is the sense the Standard Model is written in.

Fine-structure constant. The dimensionless strength of electromagnetic interaction, near 1/137 at low energy. The shared-width trial map gives inverse coupling 137.035660. Wissner-Gross's separate entangled-pair candidate gives 137.035999189566. Their physical electromagnetic identification is the independent question stated in chapter twenty-nine.

Fixed point. An object x left unchanged by a specified map F: F(x)=x. The object may be an instantaneous state, a reduced record, a probability law or a complete history. A fixed complete history can contain changing local states; a stationary law can support changing realizations. Fixedness alone establishes neither attraction nor uniqueness.

Flow. A parameterized family of transformations. Modular flow belongs to an algebra and a state, leaves the state's odds where they were, and has a bare number for its parameter. An observer's clock steps it once at every event at the observer's ports.

Force. A steepness in the potential. A single unit of source dropped on the twelve ports lowers the reading eleven sixtieths over the first hop, three sixtieths over the second and one sixtieth over the third, and a pattern held in a slope like that comes back each cycle a little further down it.

Formal system. Marks you may write, a starting arrangement of them, and moves that take one arrangement to the next. Nothing in it means anything, which is why the rules run identically in the hands of somebody who knows nothing about the subject.

Free parameter. A number a theory does not fix, which somebody measures and types in. The minimal Standard Model carries nineteen, closer to twenty-six once neutrinos have masses, plus Newton's constant and the cosmological constant.

Frustration. A loop of constraints that no assignment satisfies, however long you wait. Three coins in three rooms, each pair recording that it differs, is the smallest one, and a triangular magnet is the same arrangement in a material.

Gauge. A symmetry whose convention is picked afresh at every place rather than once for the world. Twelve observers each choosing for themselves make twelve of them, and what settles between them is the relation rather than the values.

Generation. One complete list of matter states, fifteen of them, ten from picking two slots out of five and five from picking four. The world runs three copies and no more. Fewer than three leaves matter and antimatter with no physical difference between them, more than five and three eighths costs the strong force its weakening at short range, and the threefold turn carried by the twenty faces has an orbit of exactly three.

Geodesic. Where you end up by going straight ahead and never turning. On a floor it draws a straight line; on a globe, the flight path from London to Tokyo, up over the Arctic.

Graph. Dots and the lines joining them, a dot per observer and a line wherever two of them have compared records. A walk that returns to its starting dot without retracing is a cycle, and the cycles are where the trouble lives.

Group. A collection of changes that is closed, so that two of them give a third from the same collection, and that holds a do-nothing move and an undo for every move. The eight ways of picking a square up and putting it down are one.

Hilbert space. The arrows an algebra of questions acts on, given their lengths and angles by the state. Physics courses start there and the order runs the other way, because what an observer holds is a list of questions.

Holonomy. The difference a round trip makes: set off round a loop, come back, and see what changed. Round a triangle of coin records it is one bit saying whether the coins can be set at all; on twelve observers it is a charge.

Homeostasis. Keeping a bounded set of internal readings where they were against a world that keeps pushing them off, at the observer's own expense. Walter Cannon gave the steady states of the body's fluids that name in 1926.

Inertia. What it costs to change a pattern's drift, proportional to what the pattern costs to hold together and to nothing else about it. Redirecting it means renegotiating every seam on its boundary instead of reconfirming them.

Inside-only clause. A relabeling counts only if the system can carry it out on itself. It reads like bookkeeping and it kills grand unification, because a symmetry acting from within forces a center, and simple groups have none.

Interval. The quantity two observers who disagree about distances and durations both compute the same value for. Its sign sorts every pair of events into three classes: reachable by something slower than light, reachable only by light, reachable by nothing.

Invariant. Whatever a given change of description leaves the same. Naming it comes first and the symmetry second: a rotation preserves length, and without length there is nothing for the rotation to be a symmetry of.

Lagrangian. A function of configurations and their rates, or a field density, whose specified action gives equations of motion. It compresses dynamics. A finite history or repair cost does not by itself select a unique physical Lagrangian.

Leaky abstraction. An account that lets you get on with the work while the machinery stays out of sight, good until the machinery does something the account has no words for. Joel Spolsky wrote the rule down in 2002, and the everyday container picture of space and time leaks in at least six places.

Mass. In a relativistic effective description, the invariant read from a particle's energy and momentum. For an isolated stable mode it obeys $m^2c^4 = E^2 - c^2p^2$. Unstable particles and confined quarks require their resonance or running-mass conventions. Reading a repair relaxation rate as a mass requires the relevant dynamics, field response and clock calibration.

Mismatch. The disagreement on a seam: what is left when one reading, passed through the seam's dictionary, is held against the other. It is zero

when the two answer each other and positive when they do not. Every seam can read zero while a loop carries a residue, which is chapter six's leftover of one.

Momentum. What a pattern carries as it drifts across the seam network, conserved because the laws do not care where you are. Newton listed it second among his definitions and called it the quantity of motion.

Natural selection. Variation, heredity and differential persistence running together. The list closes at three, and nothing on it decides anything.

Net of algebras. An assignment to every region of the questions answerable inside it, where a region contained in another carries the shorter list and two regions too far apart to signal carry lists that commute. This is what quantum field theory means by a field.

No-cloning. An unknown quantum state cannot be copied. Anything written down as a record can be copied freely and as often as you like, which is the half of the theorem most treatments leave out.

Noether's procedure. The machine that turns a continuous symmetry of the scored line into a conserved quantity, one for one. Shifting in space gives momentum, shifting in time gives energy.

Normal form. A state with no move left to make. When repair terminates and is confluent, every order of repairs from a given start reaches the same one. A patch with nothing to do knows that about itself and nothing about the rest.

Observer. A bounded system that reads its boundary, keeps records, acts on what it reads and carries all of it forward: an algebra of questions bounded by its interface, a record algebra at its center, an interface that behaves the same way every time, and a state fixing the odds. A thermostat qualifies. Being an observer takes no mind.

Orbit. Every place the moves in a group can send one thing. The four corners of a square make one orbit under its eight moves, the four edge midpoints make another, and nothing carries a corner into the second set.

Order. Precedence between events: one comes before another when a chain of certified reads runs from the first to the second. Events with no chain between them have no order, and numbering them anyway creates none.

Ordering fraction. Of all the pairs of events inside a causal diamond, the share in which one could have influenced the other. It is a half with one direction of space, eight thirty-fifths with two and a tenth with three, which makes it a dimension meter that needs no ruler.

Overlap rule. Two accounts have to match where both parties look, and nowhere else. Alice's brown table and Bob's waxy one never have to be reconciled.

Packet. What appears at a port: one reading drawn from a finite list, available to whatever sits on the far side. It is what a patch's state looks like from outside at one interface rather than the state itself.

Particle. A persistent pattern that repair cannot remove, carried from seam to seam with its charge and readouts intact, and read at a commit as one count at one place.

Patch. A bounded local system with its own state, its ports and a printed menu of moves: read, compare, choose, write, expose, verify. A patch carries no coordinates; positions come out of what patches record about each other.

Pinching. Sorting every question into the alternatives a record distinguishes, keeping the parts that fall inside one of them and deleting every comparison that only meant something across two. This is what writing a record does to an observer's algebra, and decoherence is the usual name for the result.

Port. An interface through which part of a patch's state is visible to a neighbor. A carrier has twelve, five seams meet at each, and chapter fourteen shows that no other count closes.

Probability. Numbers between zero and one on the questions an observer can ask, adding up across every complete list of exclusive alternatives. On quantum questions, blunt ones included, every such assignment is the trace against one density matrix.

Projector. A sharp yes-or-no question: an operation that, applied twice, does what it did once. Blunt questions answer yes with a weight, and the sharp ones are the special case whose weights are zero and one.

Quotient. Sweeping up everything that counts as the same into one bundle and working with the bundles. The twelve hours on a clock face are the whole numbers with twelve counting as nothing.

Rank. The number of independent directions in a table of comparisons. The carrier's slow band has rank three, which is the count of directions in space, and the plane perpendicular to a light ray has rank two, which is why light has two polarizations.

Record. A write into the center of an observer's algebra, where everything commutes. It reads the same the second time, copies to a neighbor without loss, and stays put under the flow until another write moves it.

Record law. What a repair writes down when it acts: how much load crossed the seam, and possibly the coin, the parity of the seam total or the full count at both ends. It fixes which quantum coherences survive the repair. Experiments that prepare definite loads and read only how many units crossed give the same statistics under every record law. Interference experiments tell the laws apart. One, between an empty seam and a waiting unit, separates the coin record from the parity record; a second, between

an empty seam and a full one, separates the parity record from the full count; and nine such measurements fix all nine numbers of the family.

Record map. The map from the full state of the world to the part that survives comparison: every seam entry and nothing private to either end. Its fibers are what the records leave unresolved.

Refinement. Describing something at ever finer resolution with no completed infinite object at the end, the way 3, 3.1, 3.14 describes pi and nobody ever holds pi. Smooth space is a limit of that kind. Carriers joined into ever finer spheres have a smooth limit of their own, and it lives on a two-dimensional screen. The smooth version is a limit nobody occupies.

Relative entropy. How many extra questions per state your odds cost you against a reference. Pass both through the same process, classical or quantum, and it can only fall, which makes it the quantity the second law is about.

Repair. The one move a seam makes: both readings to their midpoint, with the total left where it was; on whole-number loads the two ends go to the floor and ceiling of their mean, and a fair coin places the odd unit. Its quantum form also writes a record, and the record law says what. Repair only runs downhill.

Repair load. How much of an observer's state fails to lie along the field its neighbors have written on its boundary. It cannot go below zero and it reaches zero exactly when the two agree, which is the reading a tuner is listening for when the beat disappears.

Running coupling. A force strength is one number only if nothing surrounds the charge. Screening puts more of the charge in view the closer an experiment gets, so every strength quoted carries a distance with it.

Schedule independence. Every order of repairs that keeps going until nothing is left to do lands, from the same start and the same boundary data, on the same settled state. The routes there differ, and so do their lengths.

Screen. Every reading on every port an observer has facing outward. Nobody looks at it, it points nowhere, and its size is counted in readings rather than measured in meters. Carriers joined into a sphere make a screen whose smooth limit is two-dimensional, the surface a hologram is written on.

Seam. Two ports routed to each other, which makes it the one place a disagreement can show up at all. Twelve observers wired into a twenty-sided solid have thirty.

Selector. What settles an ambiguous fiber, where every record in existence is already satisfied and repair has nothing left to bite on. Which repair law gets run by the observers capable of running one is a selector.

Spin. A two-outcome measurement with a direction attached. Aim the magnet however you like and the atom answers with one of exactly two deflections, and the aim is the only thing you choose.

Superposition. A system described in terms of questions it was not prepared for, carrying one number for each answer. The numbers are amplitudes, two of them can arrive with opposite signs, and a dark band is where that happens.

Symmetry. A change of description that leaves a named invariant where it was. Finding one deletes a part of your description as unphysical, so it subtracts from the world rather than adding to it.

Temperature. The exchange rate between energy and questions, in joules per bit. Three hundred kelvin sets that rate at 2.87 joules divided by ten to the twenty-first, for each yes-or-no question a record settles.

Tsirelson ceiling. The quantum ceiling on the four-question score, 2 times the square root of 2, a winning fraction of about 85.36 percent. Answer sheets, and every written record with its answers fixed on each side, top out at 2, or 75 percent. A world that only forbade signalling would allow scores above the ceiling; non-commuting questions and a state put it at $2\sqrt{2}$ exactly.

Verifier. Anything that runs the check on the exposed readings and settles whether a patch's claim about its own work holds. The inside of the patch never enters it, because the check is arithmetic, and arithmetic is indifferent to whose readings it gets.

Wave. What repair looks like while it is under way. A port that hands its mismatch on rather than deleting it has a current to show for it, the two go on feeding each other instead of settling, and the pair crosses the wiring no faster than one seam per repair event.

Zero-error capacity. The rate at which messages can be sent with no chance of confusion. On the ring of five symbols the largest one-use code has two messages; used many times over, the ring carries the square root of five messages per use, or log base two of that in bits.

Zombie. A philosophical hypothetical: a physical duplicate without experience. The identity account of experience that chapter forty takes leaves no room for one.

Appendix C: Sources and Further Reading

The proof assistant Lean has a keyword for a step nobody has done. The keyword is "sorry". A file containing one compiles anyway and prints a warning naming the unfinished step.

The formal proofs are Lean 4 files built against Mathlib, the community's library of formalized mathematics, and no proof in them rests on that keyword or on an admitted step. The papers carry analytic proofs as well, the continuum clock and cone arguments among them. They sit with the numerical code, the simulation and the technical papers in the public OPH repository. The numerical work is Python, and one documented command reproduces it from a clean copy. The principal paper there is "Observers Who Read Their Universe into Existence: Finite Consensus, an Executable Observer, and Standard Physics". The executable observer, the repair it runs and the interference experiments that read what each repair writes down are set out in "An Executable Observer for Observer Patch Holography". The runnable observer itself, with its tests and retained results, sits in the same repository, and its clocks, fields, screens and record laws can be rerun from there. Longer treatments build the mathematics at learn.floatingpragma.io, and the repair dynamics can be watched settling at simulation.floatingpragma.io.

The quoted runs were produced with the finite patch engine OPH-FPE. The settling trace, record counts and spread quoted in chapters ten through thirteen come from a protected-consensus run of 27 August 2026. Its archive, `evidence/icosa_82k_protected_consensus_20260827_r1`, was retired from the working tree on 9 September 2026 and remains in the repository history at commit 9e1980afb, together with its `verify_archive.py`. The level-six exact-federation archive reruns the same federation of 81,920 patches and 122,880 seams under its own load law: all sixteen schedules reach the balanced class, and their finishing load vectors differ only in where odd units sit and agree as collections, which the starting totals fix. It does not reproduce the per-cycle figures of the August run. It carries the primitive seam arrays, the initial and terminal loads, the per-schedule ledgers, the quotient hashes of all sixteen schedules, the isolated controls and the per-carrier response kernels, together with a verifier that imports none of the simulator's code and rebuilds the result from the raw arrays.

The spacetime derivation gives the causal and counting limits of chapter seventeen, durations from interval counts, and the symmetry condition selecting a Lorentz cone. The causal-poset archive holds the reproduction tools for chapter seventeen's computational examples, and its large routed histories are regenerated in full and checked against retained hashes. The finite-source cosmology paper works out expansion as a density of records, and the black-hole information paper counts the reads that cross a horizon.

The standard physics

Edwin Taylor and John Archibald Wheeler's *Spacetime Physics* (second edition, 1992) builds special relativity out of the invariant interval and holds the Lorentz transformation back to a special topic after chapter three; the authors released it for free download at eftaylor.com. Leonard Susskind and Art Friedman's *Quantum Mechanics: The Theoretical Minimum* (2014) runs alongside Susskind's continuing-studies course at Stanford and starts the reader on a single spin rather than on a wave. Michael Nielsen and Isaac Chuang's *Quantum Computation and Quantum Information* (2000) gives quantum error correction a chapter of its own and no-cloning a box inside the information-theory chapter.

Bruce Schumm's *Deep Down Things* (2004) gets the gauge structure of the Standard Model across to somebody who will never do the field theory. Kip Thorne's *Black Holes and Time Warps* (1994) covers horizons and was written by one of the three men who signed the encyclopedia wager. Ernest Nagel and James Newman's *Gödel's Proof* (1958) does the incompleteness construction in a hundred and eighteen pages. Claude Shannon's "A Mathematical Theory of Communication" ran in the July and October 1948 issues of the Bell System Technical Journal, where the entropy of a message is defined.

The causal-set program begins with Luca Bombelli, Joohan Lee, David Meyer and Rafael Sorkin, "Space-time as a causal set", *Physical Review Letters* 59 (1987), 521. Bombelli and Meyer, "The origin of Lorentzian geometry", *Physics Letters A* 141 (1989), 226, and Graham Brightwell and Ruth Gregory, "Structure of random discrete spacetime", *Physical Review Letters* 66 (1991), 260, develop the order-to-geometry and dimension problem. Seth Major, David Rideout and Sumati Surya, "On recovering continuum topology from a causal set", *Journal of Mathematical Physics* 48 (2007), 032501, treats topology recovery; their "Stable homology as an indicator of manifoldlikeness in causal set theory", *Classical and Quantum Gravity* 26 (2009), 175008, presents a necessary but not sufficient diagnostic. Bombelli, Joe Henson and Sorkin, "Discreteness without symmetry breaking: a theorem", *Modern Physics Letters A* 24 (2009), 2579, gives the Lorentz-symmetry result for Poisson sprinklings. Sumati Surya's "The causal set approach to quantum gravity", *Living Reviews in Relativ-*

ity 22 (2019), article 5, gives the modern survey. Rafael Sorkin's "Causal sets: discrete gravity", in *Lectures on Quantum Gravity*, edited by Andrés Gomberoff and Donald Marolf (Springer, 2005), states the program as order plus number equals geometry. Daniel Kleitman and Bruce Rothschild, "Asymptotic enumeration of partial orders on a finite set", *Transactions of the American Mathematical Society* 205 (1975), 205, is the three-layer theorem. Jan Myrheim's CERN preprint "Statistical geometry" (1978) and David Meyer's MIT doctoral thesis "The dimension of causal sets" (1988) give the ordering fraction and the dimension it measures.

The continuum results behind the slogan that causal order fixes conformal geometry are S. W. Hawking, A. R. King and P. J. McCarthy, *Journal of Mathematical Physics* 17 (1976), 174, and David Malament, *Journal of Mathematical Physics* 18 (1977), 1399. They require causal conditions on the continuum, and an arbitrary finite order has no such limit. OPH's spacetime paper proves the flat four-dimensional limit of chapter seventeen, with its one-tenth ordering fraction and fourth-root duration ratios for diamonds inside the refined region, counting completed records and leaving the relay work out. A separate theorem fixes the Lorentz cone under invariance by the icosahedral rotations and boosts along one direction.

David Reid, *Physical Review D* 67 (2003), 024034, tests causal-set dimension in conformally flat spacetimes; Lisa Glaser and Surya, *Physical Review D* 88 (2013), 124026, use interval abundance as a test of locality; Mriganko Roy, Debdeep Sinha and Surya, *Physical Review D* 87 (2013), 044046, give curvature-aware small-diamond geometry; and Djamel Dou and Sorkin, "Black-hole entropy as causal links", *Foundations of Physics* 33 (2003), 279, count causal links across a horizon in a random causal set and find an area law.

Dionigi Benincasa and Fay Dowker, *Physical Review Letters* 104 (2010), 181301, construct a causal-set d'Alembertian whose continuum approximation contains the scalar curvature. Fay Dowker and Lisa Glaser, *Classical and Quantum Gravity* 30 (2013), 195016, extend the construction across dimensions. These operators read the scalar curvature off order and counting alone.

Richard Feynman's lecture on the principle of least action, chapter 19 of the second volume of *The Feynman Lectures on Physics* (1964), was delivered to Caltech sophomores and is the shortest good account of where a Lagrangian comes from; Caltech hosts all three volumes online. The measured values in this book come from the Particle Data Group's *Review of Particle Physics*, revised every two years at pdg.lbl.gov. The 2024 edition is S. Navas and others, Physical Review D 110, 030001.

The people

Galois's last letter appeared in the *Revue Encyclopédique* in September 1832 and in Liouville's journal in 1846; Robert Bourgne and Jean-Pierre Azra's critical edition (1962) prints it with its variants. Émilie du Châtelet's *Institutions de physique* was published in Paris in 1740, and her *Principia*, the only complete French one, in 1759. Elizabeth Loftus and John Palmer's car-crash experiment is in the *Journal of Verbal Learning and Verbal Behavior* 13 (1974), 585.

Sadi Carnot's *Réflexions sur la puissance motrice du feu* (1824) is in English in the Dover volume edited by E. Mendoza (1960). Jacob Bekenstein's "Black holes and entropy" is Physical Review D 7, 2333 (1973), and Stephen Hawking's "Black hole explosions?" is Nature 248, 30 (1974). Richard Hamming's codes are in the Bell System Technical Journal 29 (1950), 147, and Rolf Landauer's cost of erasing a bit is in the IBM Journal of Research and Development 5 (1961), 183. John Bell's inequality is *Physics* 1 (1964), 195; the four-setting form is Clauser, Horne, Shimony and Holt, Physical Review Letters 23 (1969), 880; the ceiling on it is Boris Tsirelson, *Letters in Mathematical Physics* 4 (1980), 93; and the loophole-free Delft run is B. Hensen and others, Nature 526 (2015), 682.

The parity experiment is Wu, Ambler, Hayward, Hoppes and Hudson, Physical Review 105, 1413 (1957). Rutherford's four 1919 papers on the collision of alpha particles with light atoms are in volume 37 of the *Philosophical Magazine*, the fourth reporting the anomalous effect in nitrogen. Yoshio Koide's relation is Physical Review D 28, 252 (1983).

Maupertuis read his least-action memoir to the Paris Academy in 1744. Lagrange's *Méchanique analitique* dates from 1788, and the English translation by Auguste Boissonnade and Victor Vagliente from 1997. Emmy Noether's *Invariante Variationsprobleme* is in the Göttingen *Nachrichten* for 1918, and M. A. Tavel's translation appeared in 1971.

The Darwin and Wallace papers were read at the Linnean Society on 1 July 1858 and printed that August. Gödel's paper is in *Monatshefte für Mathematik und Physik* 38 (1931), and in English in Jean van Heijenoort's *From Frege to Gödel* (1967). Claude Bernard's *Introduction à l'étude de la médecine expérimentale* (1865) can be had in Henry Copley Greene's translation. Walter Cannon named homeostasis in the 1926 Paris volume for Charles Richet's jubilee and explained the choice in *The Wisdom of the Body* (1932). Zachary Blount, Christina Borland and Richard Lenski's citrate paper is in the *Proceedings of the National Academy of Sciences* 105 (2008), 7899. Boethius on foreknowledge is book five of *The Consolation of Philosophy*, in Victor Watts's translation for Penguin. Harry Frankfurt's counterexample is in the *Journal of Philosophy* of 4 December 1969.

Leslie Lamport, "Time, Clocks, and the Ordering of Events in a Distributed System", *Communications of the ACM* 21 (1978), 558-565, distin-

guishes causal order, logical numbering and physical clock synchronization. Alain Connes and Carlo Rovelli, "Von Neumann Algebra Automorphisms and Time-Thermodynamics Relation in General Covariant Quantum Theories", *Classical and Quantum Gravity* 11 (1994), 2899-2918, proposes reading time off modular theory as thermal time, the idea behind chapter twenty-one.

Appendix D: Run It Yourself

The prologue complained that the universe ships with no source code and no documentation. The construction this book describes ships with both. The observers, their seams and their repair law, the instruments they ask questions with, the records they keep, their clocks, the fields their clocks read and the waves those fields carry are all a program, written down in full, with every result it reports stored next to the code that produced it and a fingerprint of each result kept so that a replay can be compared against it.

The numbers in chapters nine, fifteen, twenty, twenty-one and twenty-two come out of that program: the coherence a repair leaves behind, the table of angles an observer reads off its own counts, the 245 Delft rounds, the logarithm of five fourths, the variance of sixteen between two event clocks, and the interference odds of 21/22, 91/110 and 1297/1320 for a wave driven by an incident clock. None of them needs to be taken on trust. A laptop checks them.

Get the code:

```
git clone https://github.com/FloatingPragma/observer-patch-holography
cd observer-patch-holography
```

The smallest check needs nothing beyond a standard Python of version 3.11 or later:

```
python3 -m model.observer native \
  --check model/observer/results/native.json
```

It replays 264 complete histories of one native observer, every branch of every repair included, and checks eighteen facts about them against the stored result. The last line it prints is a short report that ends with the fingerprint of the result, beginning `"all_pass": true, "checks": 18, "histories": 264`. A run of ten thousand sample histories from a fixed starting seed is one more command:

```
python3 -m model.observer run \
  --trials 10000 --seed 42
```

The quantum constructions, the fields, the clocks and the waves need a few numerical libraries. Install them into a private environment and run the whole verification:

```
python3 -m venv .venv-model
. .venv-model/bin/activate
python3 -m pip install -r model/requirements.txt
python3 -m model.verify
```

That replays every stored result and compares it with the fingerprint it was saved under. A result that comes out different by a single digit fails.

Each construction has its own folder and its own README explaining what it computes and which command checks it. The map of all of them is `model/README.md`. Chapter twenty-two's tally, field, wave and region's clock live in the folders on observer time fields, relative observer geometry and the transfer modular probe, and the formal proofs behind the theorems are the Lean files appendix C describes.

The continuing C/E campaign

The C/E campaign has its own focused replay. It checks the finite-memory observer with counted record exports, the exact joint stationary reference, and the native charge and modular-access certificates:

```
python3 -m pytest -q -W error model/completion
python3 tools/build_completion_program.py --check
```

The completion plan keeps the current work in one place: the extension contract (#1135), observer services (#1136), clock operations (#1138) and the localized defect investigation (#1139). The campaign's README gives each replay and its construction contract.