

Nexivarin 10 mg Tablets — eCTD Submission
Module 5 — Clinical Study Reports

Submission Type	NDA — 505(b)(1)	Sequence Number	0000
Submission Date	2021-03-15	Section	5.2
Prepared by	PharmaCorp Inc.	Confidentiality	Proprietary and Confidential

Table of Contents — [5.2](#) Tabular Listing of All Clinical Studies

Section	Title	Description
5.2	Tabular Listing of All Clinical Studies	Complete inventory of all clinical studies supporting NDA 123456
Phase I Studies	First-in-Human and PK Studies	SAD, MAD, hepatic/renal impairment, DDI studies (NXV-PK-01 to -05)
Phase II Studies	Dose-Ranging Studies	Study NXV-201: Phase II dose-ranging (5, 10, 20 mg)
Phase III Studies	Pivotal Efficacy and Safety Studies	NXV-301 (placebo-controlled) and NXV-302 (active-controlled)

PharmaCorp Inc.
NDA 123456

Nexivarin 10 mg Tablets

Section 5.2
Tabular Listing of Clinical Studies

Submission Type	NDA — 505(b)(1)
Sequence Number	0000
Submission Date	2021-03-15
Prepared by	PharmaCorp Inc.
Confidentiality	Proprietary and Confidential

5.2 Tabular Listing of All Clinical Studies

Study No.	Phase	Design	Population	Dose	Duration	N	Primary Endpoint	Status
NXV-PK-01	I	Open-label, SAD	Healthy volunteers	1–80 mg single dose	1 day	32	PK (C _{max} , AUC, t _{1/2})	Complete
NXV-PK-02	I	Open-label, MAD	Healthy volunteers	5–40 mg/day	14 days	24	Steady-state PK; safety	Complete
NXV-BA-01	I	Crossover, open	Healthy volunteers	10 mg (tablet vs. IV)	Single dose	16	Absolute BA (F%)	Complete
NXV-BA-02	I	Crossover, open	Healthy volunteers	10 mg (fed/fasted)	Single dose	24	Food effect on AUC, C _{max}	Complete
NXV-ADME-01	I	Mass balance, open	Healthy volunteers	10 mg [14C]-labeled	Single dose	8	Total recovery, metabolite profile	Complete
NXV-PK-03	I	Open, parallel	Hepatic impairment (Child-Pugh A,B,C) vs. matched healthy	10 mg single	1 day	32	PK comparison	Complete
NXV-PK-04	I	Open, parallel	Renal impairment (GFR stages) vs. healthy	10 mg single	1 day	32	PK comparison	Complete
NXV-DDI-01	I	Crossover, open	Healthy volunteers	NXV 10mg ± ketoconazole 200mg	10 days	20	CYP3A4 DDI (AUC ratio)	Complete
NXV-DDI-02	I	Crossover, open	Healthy volunteers	NXV 10mg ± simvastatin 40mg	14 days	22	Simvastatin PK change	Complete
NXV-201	II	DB, RCT, parallel, PLC	Hypertension patients	2.5, 5, 10 mg/day	8 weeks	320	SeDBP at Week 8	Complete
NXV-301	III	DB, RCT, parallel, PLC	Hypertension	10 mg/day	12 weeks	892	ΔSeDBP at Week 12	Complete
NXV-302	III	DB, RCT, parallel, PLC + active	Hypertension	10 mg/day vs. amlodipine 10mg vs. PLC	24 weeks	908	NI vs. amlodipine (ΔSeDBP)	Complete
NXV-PED-01	III	DB, RCT	Pediatric hypertension (6–17y)	2.5–5 mg/day	8 weeks	154	SeDBP	Ongoing