

Nexivarin 10 mg Tablets — eCTD Submission
Module 5 — Clinical Study Reports

Submission Type	NDA — 505(b)(1)	Sequence Number	0000
Submission Date	2021-03-15	Section	5.3.4.2
Prepared by	PharmaCorp Inc.	Confidentiality	Proprietary and Confidential

Table of Contents — [5.3.4.2](#) Clinical Study Report: Study NXV-301

Section	Title	Description
5.3.4.2	Clinical Study Report — Study NXV-301	Phase III randomized, double-blind, placebo-controlled trial
Synopsis	Study Synopsis	Multicenter; 902 randomized; 12-week treatment; primary endpoint SeDBP
Patient Demographics	Patient Demographics (ITT Population, N=892)	Balanced baseline characteristics across treatment arms
Primary Efficacy Results	Primary Efficacy Results (ITT, LOCF)	SeDBP reduction: -12.8 vs -4.2 mmHg; p<0.0001; responder rate 68.4% vs 31.7%
Safety Results	Safety Results (Safety Population, N=902)	Edema 13.9% vs 1.8% ; no deaths related to study drug

PharmaCorp Inc.
NDA 123456

Nexivarin 10 mg Tablets
Section 5.3.4.2
Phase III Pivotal Study NXV-301

Submission Type	NDA — 505(b)(1)
Sequence Number	0000
Submission Date	2021-03-15
Prepared by	PharmaCorp Inc.
Confidentiality	Proprietary and Confidential

5.3.4.2 Clinical Study Report: Study NXV-301

Synopsis

Item	Details
Study Title	A Multicenter, Randomized, Double-Blind, Placebo-Controlled Study to Evaluate the Efficacy and Safety of Nexivarin Hydrochloride 10 mg in Patients with Mild-to-Moderate Essential Hypertension
Study Number	NXV-301
Phase	III
Study Design	Parallel-group, double-blind, placebo-controlled, multicenter
Number of Sites	48 centers (USA, Canada, EU)
Duration	12-week double-blind treatment + 2-week follow-up
Enrolled	902 randomized; 892 ITT; 874 PP
Primary Endpoint	Mean change from baseline in seated diastolic blood pressure (SeDBP) at Week 12
Secondary Endpoints	SeSBP change; responder rate (SeDBP <90 mmHg); normalization rate (SeDBP <85 mmHg); quality of life (SF-36)
Randomization Ratio	1:1 (Nexivarin 10 mg : Placebo)
Blinding	Double-blind; identical appearance tablets
Statistical Analysis	ANCOVA with treatment, center, and baseline as covariates; LOCF imputation; alpha=0.05 (two-sided)

Patient Demographics (ITT Population, N=892)

Characteristic	Nexivarin 10 mg (N=446)	Placebo (N=446)
Age, mean ± SD (y)	52.4 ± 10.8	51.9 ± 11.2
Age range (y)	21 – 78	23 – 76
Male sex (%)	54.3%	52.7%
White (%)	68.2%	67.5%
Black (%)	18.6%	19.1%
Hispanic/Latino (%)	9.4%	9.8%
BMI, mean ± SD (kg/m2)	28.9 ± 4.6	29.1 ± 4.8
Baseline SeDBP (mmHg)	97.8 ± 5.4	98.1 ± 5.6
Baseline SeSBP (mmHg)	154.6 ± 11.2	155.1 ± 10.8
Duration of hypertension (y, mean)	5.2 ± 4.1	5.4 ± 4.3
Prior antihypertensive use (%)	42.8%	44.2%
Diabetes mellitus (%)	14.6%	15.2%
Current smoker (%)	18.4%	17.9%

Primary Efficacy Results (ITT, LOCF)

Endpoint	Nexivarin 10 mg	Placebo	LS Mean Difference (95% CI)	p-value
Baseline SeDBP (mmHg)	97.8 ± 5.4	98.1 ± 5.6	—	—
Week 12 SeDBP (mmHg)	84.9 ± 7.2	93.9 ± 6.8	—	—
Change from Baseline SeDBP	-12.8 ± 0.6	-4.2 ± 0.6	-8.6 (-10.1 to -7.1)	<0.001
Baseline SeSBP (mmHg)	154.6 ± 11.2	155.1 ± 10.8	—	—
Change from Baseline SeSBP	-18.4 ± 1.2	-7.3 ± 1.1	-11.1 (-13.8 to -8.4)	<0.001
Responder rate (SeDBP <90 mmHg)	68.4%	31.7%	OR 4.38 (3.10–6.18)	<0.001
Normalization (SeDBP <85 mmHg)	49.6%	18.4%	OR 4.26 (2.92–6.22)	<0.001

Safety Results (Safety Population, N=902)

Parameter	Nexivarin 10 mg (N=451)	Placebo (N=451)
Any TEAE	58.3%	50.1%
Severe TEAE	1.8%	1.6%
Serious AE (SAE)	1.6%	1.3%
Discontinuation due to AE	2.9%	1.1%
Edema	13.9%	1.8%
Headache	7.1%	7.6%
Dizziness	3.1%	1.3%
Flushing	2.4%	0.4%
Fatigue	4.2%	2.9%
Palpitations	1.2%	0.4%
Deaths	0	1 (unrelated to study drug)

Investigator's Conclusion

Study NXV-301 met its primary endpoint, demonstrating statistically significant and clinically meaningful reductions in SeDBP and SeSBP with nexivarin 10 mg compared to placebo in patients with mild-to-moderate essential hypertension over 12 weeks. The safety profile was consistent with the calcium channel blocker class. These results support the benefit/risk assessment and regulatory approval of nexivarin 10 mg for the treatment of hypertension.