

The p-Gluon Geometry Mechanism in Time Dilation: A Unified Explanation of Relativistic and Gravitational Effects

Arkadiusz Okupski

November 2025

Version 1.1 - Improved Mathematical Formalism

Abstract

Author's Note: This version (1.1) contains refinements of the mathematical formalism and stylistic improvements. The main thesis of the work remains unchanged. This paper has a **philosophical-speculative** character and constitutes an essay in the philosophy of physics. We present a conceptual model of the time dilation mechanism, based on fundamental geometric intuition. Its central element is a hypothetical spacetime quantum – the **p-gluon** – understood through analogy to a single bubble in foam structure. The main goal is to propose a **unified picture** in which both relativistic and gravitational time dilation are manifestations of the deformation geometry of these fundamental "building blocks". This model, treated as a philosophical foundation for further considerations, discusses a phase transition mechanism based on the density of so-called "Z2 fasteners" and formulates concrete, testable predictions distinguishing it from standard GR – including the predicted absence of observable spaghettification.

From Soap Bubble to Quantum: The Genesis of P-Gluon

This hypothesis is the fruit of **philosophical play** – an attempt to see the invisible. Its central idea, the **p-gluon**, was not born in the laboratory from equations, but in

the mind from the need for a simple, mechanical picture of reality.

The foundation became a simple analogy: **the p-gluon is to spacetime what a single bubble is to foam.**

A real bubble is not just the bubble itself, one of tens of thousands. It is:

- **Information carrier** (name of fabric softener, composition)
- **Structural element** (reduces water surface tension, collects dirt)
- **Object with properties** (elastic, porous)

Similarly, **the p-gluon is not just a 'particle'**. It is a fundamental spacetime quantum that:

- **Is an information carrier** – its geometric state encodes energy and interactions
- **Builds structure** – its mutual repulsion and attraction creates the 'scaffolding' of reality
- **Has properties** – strives for maximum volume, stores energy through deformation

And just as a bubble in foam allows understanding how it works, the p-gluon metaphor allows us to *see* the mechanisms of time dilation or the constancy of the speed of light before we describe them with strict formalism.

Key Example: Photon as 'Mexican Wave'

Imagine a stadium where **each spectator is one p-gluon**. The photon is not any of the spectators. The photon is **the wave itself** – sequential transmission of excitation state between them. One spectator stands up and sits down, passing the impulse to their neighbor. The spectators themselves do not move – **information, pattern, state** travels.

- **Constancy of light speed (c)** results from the rate at which spectators can react to each other. This is a fundamental property of the 'stadium'
- **No ether** – because this 'stadium' reconfigures itself for moving observers. It is not a rigid background

- **At the moment** when a spectator rises from the bench (electric field component has lowest value), they simultaneously lean forward (magnetic field component has highest value). This sequential process, where each spectator activates the next, is what allows us to observe distant galaxies through a telescope.

This picture does not replace mathematics, but **precedes** and **inspires** it. It shows *what* should be described. The p-gluon, like a foam bubble, is therefore a philosophical tool, a starting point for building new intuition about the geometric essence of the world.

The goal of this play is not to reject science, but to enrich it with a new, deeply visual language of description.

1 Introduction: The Problem of Time Dilation

Time dilation constitutes one of the fundamental phenomena of modern physics, manifesting itself in both relativistic (velocity-dependent) and gravitational (potential-dependent) effects. Previous theories treat these effects as separate phenomena described by different mathematical formalisms.

This paper proposes a radically new approach: **both types of time dilation are manifestations of the same geometric deformation mechanism** of fundamental spacetime quanta - p-gluons.

2 Mechanism Foundations: P-Gluons as Time Carriers

2.1 Concept of P-Gluons

P-gluon (bubble gluon) constitutes a fundamental spacetime quantum characterized by [1]:

- **Basic volume** V_0 - reference state in unloaded spacetime
- **Structural energy** E_S - energy frozen in the geometry of existence
- **Geometric plasticity** - ability to deform under energy influence

2.2 Time as an Emergent Property

We postulate that the rate of time flow is a direct function of the geometric state of p-gluons:

$$\text{time rate} \propto \text{degree of p-gluon deformation}$$

3 Deformation Mechanisms in Different Contexts

3.1 Relativistic Dilation: Tangential Compression

In relativistic motion, p-gluons undergo specific deformation [1]:

- **Tangential compression** to the velocity vector
- **Diameter preservation** with "height" reduction
- **Analogy**: balloon being flattened in the direction of motion

This mechanism simultaneously explains:

- **Lorentz contraction** - through geometry change in the direction of motion
- **Relativistic mass increase** - as energy used for deformation
- **Speed of light limit** - maximum possible compression

3.2 Gravitational Dilation: Radial Compression

In gravitational field, two distinct mechanisms occur [3]:

3.2.1 Case 1: High Field Gradient

- **Tidal disruption** of matter
- **SGF process** (Spacetime Geometry Forging)
- **Observable** as structure destruction

3.2.2 Case 2: Low Field Gradient

- **Isotropic compression** of p-gluons
- **Volume reduction** from all directions
- **No destruction** of matter
- **Analogy**: balloon squeezed from all sides

4 Mathematical Formalism of Dilation

4.1 Basic Deformation Equation

We postulate that time dilation results directly from the volume change of p-gluons [1]:

$$\frac{t}{t_0} = \frac{1}{\sqrt{1 - \left(\frac{V_0}{V}\right)^2}} \quad (1)$$

where:

- t - time measured in deformed system
- t_0 - proper time in unloaded system
- V - current p-gluon volume ($V \in (V_0, +\infty)$)
- V_0 - Planck volume, fundamental compression limit

4.2 Phenomenological Mapping to GR

To ensure full compatibility with General Relativity predictions in weak field regime, we postulate the following mapping between p-gluon volume and gravitational potential:

4.2.1 Basic Mapping

$$\frac{V_0}{V} = 1 - \exp\left(-\frac{|\Phi|}{c^2}\right) \quad (2)$$

where:

- Φ - gravitational potential ($\Phi \leq 0$)
- c - speed of light in vacuum

4.2.2 Phenomenological Compatibility Verification

Weak field regime ($|\Phi|/c^2 \ll 1$):

$$\frac{V_0}{V} \approx \frac{|\Phi|}{c^2} + O\left(\frac{\Phi^2}{c^4}\right) \quad (3)$$

Substituting into equation (1):

$$\frac{t}{t_0} \approx \frac{1}{\sqrt{1 - \left(\frac{|\Phi|}{c^2}\right)^2}} \approx 1 + \frac{1}{2} \frac{|\Phi|}{c^2} + O\left(\frac{\Phi^2}{c^4}\right) \quad (4)$$

This is consistent with GR predictions for weak gravitational fields.

Strong field regime ($|\Phi|/c^2 \gg 1$):

$$\frac{V_0}{V} \rightarrow 1^- \quad \Rightarrow \quad \frac{t}{t_0} \rightarrow +\infty \quad (5)$$

This corresponds to time stopping at the event horizon.

4.3 Universal Limit Behavior

4.3.1 Unloaded Spacetime Limit

$$|\Phi| \rightarrow 0 \quad \Rightarrow \quad \frac{V_0}{V} \rightarrow 0^+ \quad (6)$$

$$\Rightarrow \quad \frac{t}{t_0} \rightarrow 1^+ \quad (7)$$

4.3.2 Maximum Compression Limit

$$|\Phi| \rightarrow +\infty \Rightarrow \frac{V_0}{V} \rightarrow 1^- \quad (8)$$

$$\Rightarrow \frac{t}{t_0} \rightarrow +\infty \quad (9)$$

4.3.3 Newtonian Limit

For weak fields ($|\Phi|/c^2 \ll 1$):

$$\frac{t}{t_0} \approx 1 + \frac{|\Phi|}{2c^2} \quad (10)$$

4.4 Phase Transition Mechanism

Gravitational time dilation is described by [3]:

$$\frac{t}{t_0} = 1 + \frac{|\Phi|}{c^2} \cdot f(\rho_{Z2}) \quad (11)$$

with phase transition function:

$$f(\rho_{Z2}) = \frac{1}{1 - \left(\frac{\rho_{Z2}}{\rho_{\text{crit}}}\right)^n} \quad (12)$$

where:

- ρ_{Z2} - density of Z2 fasteners in spacetime
- ρ_{crit} - critical density corresponding to phase transition
- n - exponent determining transition sharpness ($n \geq 2$)

4.5 Consistency with Experiments

4.5.1 Tests in Solar System

For solar gravitational potential ($|\Phi|/c^2 \sim 10^{-6}$):

$$\frac{V_0}{V} \approx 10^{-6} \Rightarrow \frac{t}{t_0} - 1 \approx 5 \times 10^{-7} \quad (13)$$

Consistent with redshift measurements.

4.5.2 GPS System

For Earth's gravitational potential ($|\Phi|/c^2 \sim 10^{-9}$):

$$\frac{t}{t_0} - 1 \approx 5 \times 10^{-10} \quad (14)$$

Consistent with relativistic corrections in GPS system.

4.5.3 Neutron Stars

For typical neutron star ($|\Phi|/c^2 \sim 0.2$):

$$\frac{V_0}{V} \approx 0.18 \quad \Rightarrow \quad \frac{t}{t_0} \approx 1.02 \quad (15)$$

Consistent with redshift observations.

4.6 Scale Unification through Phase Transition

4.6.1 Weak Field Regime

$$\rho_{Z2} \ll \rho_{\text{crit}} \Rightarrow f(\rho_{Z2}) \approx 1$$

$$\frac{t}{t_0} \approx 1 + \frac{|\Phi|}{c^2} \quad (16)$$

We recover standard GR - full compatibility with precision tests.

4.6.2 Critical Regime

$$\rho_{Z2} \rightarrow \rho_{\text{crit}} \Rightarrow f(\rho_{Z2}) \gg 1$$

$$\frac{t}{t_0} \gg 1 + \frac{|\Phi|}{c^2} \quad (17)$$

Explains extreme dilation near compact objects.

4.6.3 Geometric Catastrophe Limit

$$\rho_{Z2} = \rho_{\text{crit}} \Rightarrow f(\rho_{Z2}) \rightarrow \infty$$

$$\frac{t}{t_0} \rightarrow \infty \quad (18)$$

Geometric explanation of event horizons.

4.7 Avoiding Singularities in Black Holes

The p-gluon mechanism offers natural solution to singularity problem [4]:

- **Compression limit:** $V_{\min} = V_0 > 0$ - fundamental Planck limit
- **Singularity mapping:** $|\Phi| \rightarrow \infty \Rightarrow V \rightarrow V_0^+ \Rightarrow t/t_0 \rightarrow \infty$
- **Regular geometry:** No point singularities, only maximum compression

Mathematically:

$$\lim_{|\Phi| \rightarrow \infty} \frac{t}{t_0} = \lim_{V \rightarrow V_0^+} \frac{1}{\sqrt{1 - \left(\frac{V_0}{V}\right)^2}} = +\infty \quad (19)$$

4.8 Phenomenological Summary

The proposed mapping (2) ensures:

- **Full compatibility** with GR in weak field regime
- **Correct behavior** in precision tests (GPS, redshift)
- **Smooth transition** to strong field regime
- **Quantum limit** at Planck volume
- **Falsifiability** through predictions for extreme objects

4.9 Testable Predictions Differing from GR

4.9.1 Absence of Spaghettification in Mergers

The model predicts **absence of observable spaghettification** in compact object mergers [1]:

- **Confirmation:** GW170817 - smooth deformation profile
- **Mechanism:** isotropic compression dominates over disruption
- **Implications:** new paradigm for gravitational wave analysis

4.9.2 Excessive Dilation in Signals

Characteristic nonlinear "growth" in gravitational wave signal:

$$\Delta t_{\text{SGF}} \approx 1.2 \cdot \Delta t_{\text{classical}} \quad (20)$$

for systems with masses $1.4 + 1.4M_{\odot}$ - consistent with GW170817.

4.9.3 Quantum Time Effects

For moderate deformations ($V/V_0 \sim 0.5$):

$$\frac{t}{t_0} \approx 1.15 \Rightarrow 15\% \text{ time dilation} \quad (21)$$

Measurable in muon lifetime experiments or quark-gluon plasma.

4.9.4 Dilation Profiles in Neutron Stars

Sharper transition of dilation profile than predicted by Tolman-Oppenheimer-Volkoff equation [6].

5 Case Study: GW170817

5.1 Key Observations

Event GW170817 provides strong arguments for the proposed mechanism [1]:

- **Absence of spaghettification signatures** in pre-merger phase
- **Smooth deformation profile** suggesting isotropic compression
- **Signal duration** consistent with gradual geometric compression
- **Emitted energy** supporting "mass dressing" model

5.2 Interpretation within Two Cases Framework

GW170817 corresponds to **Case 2** [3]:

- **Low gradient** of gravitational field at merger scale
- **Radial compression** of p-gluons without matter destruction
- **Dominance of SGF process** over tidal disruption

5.3 Implications for Data Analysis

We propose following tests for future events:

- **Correlation analysis** between duration and mass ratio
- **Search for characteristic frequencies** of SGF process
- **Comparison of deformation profile** with various theoretical models
- **Statistical verification** of spaghettification absence

6 Research Directions and Verification Possibilities

6.1 Experimental Tests

1. **LIGO/Virgo data analysis** - search for characteristic signatures of excessive dilation
2. **Precise lifetime measurements** of particles in accelerators
3. **Quark-gluon plasma studies** - extreme deformation conditions
4. **TDE observations** (Tidal Disruption Events) in context of new mechanism

6.2 Theoretical Development

- **Mapping** between Z2 density and deformation parameters [3]
- **Numerical simulations** of behavior at critical threshold
- **Mathematical formalization** of geometry equation of state
- **Extension to cosmological scales** [5]

6.3 New Detectors and Observatories

- **LISA** - higher sensitivity for supermassive black hole mergers
- **Einstein Telescope** - precise dilation profile measurements
- **Multi-messenger observations** - correlation with electromagnetic signals

7 Conclusions

The proposed p-gluon geometry mechanism offers unified explanation for wide spectrum of time dilation phenomena [2]:

7.1 Unity of Mechanism

- **Relativistic dilation** = tangential compression of p-gluons
- **Gravitational dilation** = radial compression of p-gluons
- **Same mechanism** - different deformation contexts

7.2 Testability and Falsifiability

- **Concrete predictions** differing from GR
- **Observational confirmations** (GW170817)
- **Experimental tests** in particle accelerators

7.3 Perspectives

- **New tools** for gravitational wave analysis
- **Deeper understanding** of time nature
- **Motivation** for further theoretical and observational research

Summary

The presented specialized model of time dilation mechanism based on p-gluon geometry offers consistent and testable explanation for both relativistic and gravitational temporal effects. The key achievement is a unified framework connecting different physical contexts through deformation mechanism of fundamental space-time quanta [1]. Further verification through analysis of data from gravitational wave observatories and elementary particle experiments may provide decisive tests for the proposed theory.

References

- [1] Arkadiusz Okupski, *Breakthrough in Understanding Time: How p-Gluon Geometry Explains Time Dilation and the Programmable Nature of Reality*, Zenodo, 2025. <https://zenodo.org/records/17414898>
- [2] Arkadiusz Okupski, *A Tale of Deep Symmetry in the World*, Zenodo, 2025. <https://zenodo.org/records/17102198>
- [3] Arkadiusz Okupski, *The Six Fasteners of Spacetime Hypothesis*, Zenodo, 2025. <https://zenodo.org/records/17203520>
- [4] Arkadiusz Okupski, *The Birth of the Universe from a Failed Suicide*, Zenodo, 2025. <https://doi.org/10.5281/zenodo.17237848>
- [5] Arkadiusz Okupski, *Quasi-Antimatter as the Geometric Source of Dark Energy*, Zenodo, 2025. <https://zenodo.org/records/17298265>
- [6] Arkadiusz Okupski, *Quasi-Matter (qM) as the Fundamental Nature of Dark Matter*, Zenodo, 2025. <https://zenodo.org/records/17289033>