

Expected Value and Inequalities

Expectation is Lebesgue integration with respect to a probability measure.

Let (Ω, \mathcal{F}, P) be a probability space. If $X \geq 0$,

$$\mathbb{E}[X] := \int_{\Omega} X \, dP \in [0, \infty].$$

Expectation may be infinite.

For any real-valued X ,

$$X^+ = \max(X, 0), \quad X^- = \max(-X, 0).$$

Define

$$\mathbb{E}[X] = \mathbb{E}[X^+] - \mathbb{E}[X^-]$$

whenever at least one term is finite.

Expectation inherits properties of the integral.

If expectations exist:


- ▶ Linearity: $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$
- ▶ Affine invariance: $\mathbb{E}[aX + b] = a\mathbb{E}[X] + b$
- ▶ Monotonicity: $X \leq Y \Rightarrow \mathbb{E}[X] \leq \mathbb{E}[Y]$

Theorem. If φ is convex and expectations exist, then

$$\varphi(\mathbb{E}[X]) \leq \mathbb{E}[\varphi(X)].$$

Expectation interacts predictably with convex functions.

Key geometric fact (supporting line)



A convex function admits a supporting (tangent) line at every point.

Proof of Jensen's Inequality

Let

$$x_0 := \mathbb{E}[X].$$

From the supporting-line inequality,

$$\varphi(X) \geq aX + b \quad \text{a.s.}$$

Taking expectations and using linearity,

$$\mathbb{E}[\varphi(X)] \geq a \mathbb{E}[X] + b = ax_0 + b = \varphi(x_0) = \varphi(\mathbb{E}[X]).$$

This proves the result.

Jensen example: $g(x) = x^2 - 3x + 3$

Let X satisfy $P(X = 1) = P(X = 3) = \frac{1}{2}$.

Compute the mean:

$$\mathbb{E}[X] = \frac{1}{2} \cdot 1 + \frac{1}{2} \cdot 3 = 2.$$

Compute both sides of Jensen:

$$g(1) = 1 - 3 + 3 = 1, \quad g(3) = 9 - 9 + 3 = 3,$$

so

$$\mathbb{E}[g(X)] = \frac{1}{2}g(1) + \frac{1}{2}g(3) = \frac{1}{2}(1 + 3) = 2.$$

Also

$$g(\mathbb{E}[X]) = g(2) = 4 - 6 + 3 = 1.$$

Thus

$$g(\mathbb{E}[X]) = 1 < 2 = \mathbb{E}[g(X)].$$

Theorem. Let $p, q \geq 1$ satisfy $\frac{1}{p} + \frac{1}{q} = 1$. If $\mathbb{E}|X|^p < \infty$ and $\mathbb{E}|Y|^q < \infty$, then

$$\mathbb{E}|XY| \leq \|X\|_p \|Y\|_q, \quad \|X\|_p := (\mathbb{E}|X|^p)^{1/p}.$$

Young's inequality for $a, b \geq 0$:

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}.$$

Apply pointwise to

$$a = \frac{|X|}{(\mathbb{E}|X|^p)^{1/p}}, \quad b = \frac{|Y|}{(\mathbb{E}|Y|^q)^{1/q}},$$

then integrate.

Proof of Hölder (via Young + normalization)

If $(\mathbb{E}|X|^p)^{1/p} = 0$ or $(\mathbb{E}|Y|^q)^{1/q} = 0$, then $XY = 0$ a.s. and the claim is trivial. Assume both are positive and define

$$U := \frac{|X|}{(\mathbb{E}|X|^p)^{1/p}}, \quad V := \frac{|Y|}{(\mathbb{E}|Y|^q)^{1/q}}.$$

Then $\mathbb{E}[U^p] = 1$ and $\mathbb{E}[V^q] = 1$.

Apply Young pointwise:

$$UV \leq \frac{U^p}{p} + \frac{V^q}{q} \quad \text{a.s.}$$

Taking expectations,

$$\mathbb{E}[UV] \leq \frac{1}{p} \mathbb{E}[U^p] + \frac{1}{q} \mathbb{E}[V^q] = \frac{1}{p} + \frac{1}{q} = 1.$$

Finally,

$$\mathbb{E}|XY| = (\mathbb{E}|X|^p)^{1/p} (\mathbb{E}|Y|^q)^{1/q} \mathbb{E}[UV] \leq (\mathbb{E}|X|^p)^{1/p} (\mathbb{E}|Y|^q)^{1/q}. \quad \square$$

Corollary (Hölder, $p = q = 2$). If $\mathbb{E}[X^2] < \infty$ and $\mathbb{E}[Y^2] < \infty$, then

$$\mathbb{E}|XY| \leq (\mathbb{E}[X^2])^{1/2}(\mathbb{E}[Y^2])^{1/2}.$$

Cauchy–Schwarz is the L^2 case of Hölder.

There is also a direct proof using

$$\mathbb{E}[(X + tY)^2] \geq 0 \quad \text{for all } t \in \mathbb{R},$$

which shows that expectation behaves like an inner product.
We will rely on the Hölder formulation.

Proof. For any $t \in \mathbb{R}$,

$$0 \leq \mathbb{E}[(X + tY)^2] = \mathbb{E}[X^2] + 2t \mathbb{E}[XY] + t^2 \mathbb{E}[Y^2].$$

Thus the quadratic

$$Q(t) := \mathbb{E}[X^2] + 2t \mathbb{E}[XY] + t^2 \mathbb{E}[Y^2]$$

satisfies $Q(t) \geq 0$ for all $t \in \mathbb{R}$. Hence its discriminant must be nonpositive:

$$(2\mathbb{E}[XY])^2 - 4\mathbb{E}[X^2]\mathbb{E}[Y^2] \leq 0.$$

Therefore,

$$|\mathbb{E}[XY]| \leq (\mathbb{E}[X^2])^{1/2} (\mathbb{E}[Y^2])^{1/2}. \quad \square$$

Theorem (Markov). If $X \geq 0$ and $a > 0$, then

$$P(X \geq a) \leq \frac{\mathbb{E}[X]}{a}.$$

Proof. On the event $\{X \geq a\}$ we have $X \geq a$, so

$$X \geq a \mathbf{1}_{\{X \geq a\}} \quad \text{a.s.}$$

Taking expectations and using linearity and monotonicity,

$$\mathbb{E}[X] \geq a \mathbb{E}[\mathbf{1}_{\{X \geq a\}}] = a P(X \geq a).$$

Divide by a to obtain the result. □

Apply Markov to X^2 .

If $\mathbb{E}[X^2] < \infty$ and $a > 0$,

$$P(|X| \geq a) = P(X^2 \geq a^2) \leq \frac{\mathbb{E}[X^2]}{a^2}.$$

Chebyshev is a corollary of Markov.