



UNIVERSITY OF GENEVA

Faculty of Science

My Course

Digital Forensics

14x065

Michel Jean Joseph Donnet

2026-02-23



Contents

Exercise 1 3

Exercise 2 4

Exercise 3 6

Exercise 4 6

Exercise 5 7

Exercise 6 8

Exercise 7 8

Exercise 8 8

Exercise 9 8

Exercise 1 8



Exercise 1

Let $|\psi\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}$. Write out $|\psi\rangle^{\otimes 2}$ and $|\psi\rangle^{\otimes 3}$ explicitly, both in terms of tensor products like $|0\rangle |1\rangle$, and using the Kronecker product.

- As we often use the abbreviated notations $|v\rangle \otimes |w\rangle = |vw\rangle$ for the tensor product, we have in term of tensor products:

$$\begin{aligned} |\psi\rangle^{\otimes 2} &= |\psi\rangle \otimes |\psi\rangle \\ &= \frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes \frac{|0\rangle + |1\rangle}{\sqrt{2}} \\ &= \frac{1}{\sqrt{2}} \cdot \frac{1}{\sqrt{2}} (|0\rangle \otimes |0\rangle + |0\rangle \otimes |1\rangle + |1\rangle \otimes |0\rangle + |1\rangle \otimes |1\rangle)^1 \\ &= \frac{|00\rangle + |01\rangle + |10\rangle + |11\rangle}{2} \end{aligned}$$

- As $|0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $|1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, we have by using the Kronecker² product:

$$\begin{aligned} |\psi\rangle^{\otimes 2} &= |\psi\rangle \otimes |\psi\rangle \\ &= \frac{1}{\sqrt{2}} \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) \otimes \frac{1}{\sqrt{2}} \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) \\ &= \frac{1}{2} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 1 \end{pmatrix} \\ &= \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \end{aligned}$$

- For $|\psi\rangle^{\otimes 3}$, we have in term of tensor products:

$$\begin{aligned} |\psi\rangle^{\otimes 3} &= |\psi\rangle \otimes |\psi\rangle^{\otimes 2} \\ &= \frac{|0\rangle + |1\rangle}{\sqrt{2}} \otimes \frac{|00\rangle + |01\rangle + |10\rangle + |11\rangle}{2} \\ &= \frac{1}{2\sqrt{2}} (|0\rangle \otimes (|00\rangle + |01\rangle + |10\rangle + |11\rangle) + |1\rangle \otimes (|00\rangle + |01\rangle + |10\rangle + |11\rangle)) \\ &= \frac{1}{2\sqrt{2}} (|000\rangle + |001\rangle + |010\rangle + |011\rangle + |100\rangle + |101\rangle + |110\rangle + |111\rangle) \end{aligned}$$

¹ Distributivity of tensorial product ² Kronecker product: Given vectors $\mathbf{a} \in \mathbb{C}^m$ and $\mathbf{b} \in \mathbb{C}^n$, their Kronecker product is defined as: $\mathbf{a} \otimes \mathbf{b} = (a_1\mathbf{b}, a_2\mathbf{b}, \dots, a_m\mathbf{b})^T \in \mathbb{C}^{mn}$



- And by using the Kronecker product:

$$\begin{aligned}
 |\psi\rangle^{\otimes 3} &= |\psi\rangle \otimes |\psi\rangle^{\otimes 2} \\
 &= \frac{1}{\sqrt{2}} \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) \otimes \frac{1}{2} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \\
 &= \frac{1}{2\sqrt{2}} \left(\begin{pmatrix} 1 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \right) = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}
 \end{aligned}$$

Exercise 2

The Hadamard operator on one qubit may be written as

$$H = \frac{1}{\sqrt{2}} [(|0\rangle + |1\rangle)\langle 0| + (|0\rangle - |1\rangle)\langle 1|].$$

Show explicitly that the Hadamard transform on n qubits, $H^{\otimes n}$, may be written as

$$H^{\otimes n} = \frac{1}{\sqrt{2^n}} \sum_{x,y} (-1)^{x \cdot y} |x\rangle\langle y|.$$

Write out an explicit matrix representation for $H^{\otimes 2}$.

- The notation $\langle \varphi|$ is used to represent the vector dual to $|\varphi\rangle$, so the vector φ conjugate and transposed.
- The notation $|\varphi\rangle\langle \psi|$ denote the matrix product between the vectors φ and ψ .

So we have:

$$\begin{aligned}
 H &= \frac{1}{\sqrt{2}} [(|0\rangle + |1\rangle)\langle 0| + (|0\rangle - |1\rangle)\langle 1|] \\
 &= \frac{1}{\sqrt{2}} \left[\begin{pmatrix} 1 \\ 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \end{pmatrix} + \begin{pmatrix} 1 \\ -1 \end{pmatrix} \begin{pmatrix} 0 & 1 \end{pmatrix} \right] \\
 &= \frac{1}{\sqrt{2}} \left[\begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ 0 & -1 \end{pmatrix} \right] \\
 &= \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}
 \end{aligned}$$

For $H^{\otimes 2}$, we have:



$$\begin{aligned}
 H^{\otimes 2} &= H \otimes H \\
 &= \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \otimes \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \\
 &= \frac{1}{2} \begin{pmatrix} 1 \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} & 1 \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \\ 1 \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} & -1 \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \end{pmatrix} \\
 &= \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix}
 \end{aligned}$$

We want to prove that $H^{\otimes n} = \frac{1}{\sqrt{2^n}} \sum_{x,y} (-1)^{x \cdot y} |x\rangle\langle y|$ with x and y that represent n qubits, so with $|x\rangle = |x_1 x_2 \dots x_n\rangle$ and $|y\rangle = |y_1 y_2 \dots y_n\rangle$. So x and y are two sequences of bits.

We can define $(x \cdot y)$ as the bitwise inner product modulo 2:

$$x \cdot y = \sum_{i=1}^n x_i y_i \bmod 2$$

where $x_i, y_i \in \{0, 1\}$.

For the initialization, we have:

$$\begin{aligned}
 H^{\otimes 1} &= H = \frac{1}{\sqrt{2}} [(|0\rangle + |1\rangle)\langle 0| + (|0\rangle - |1\rangle)\langle 1|] \\
 &= \frac{1}{\sqrt{2}} (|0\rangle\langle 0| + |1\rangle\langle 0| + |0\rangle\langle 1| - |1\rangle\langle 1|) \\
 &= \frac{1}{\sqrt{2}} [(-1)^{0 \cdot 0} |0\rangle\langle 0| + (-1)^{1 \cdot 0} |1\rangle\langle 0| + (-1)^{0 \cdot 1} |0\rangle\langle 1| + (-1)^{1 \cdot 1} |1\rangle\langle 1|] \\
 &= \frac{1}{\sqrt{2}} \sum_{a,b \in \{0,1\}} (-1)^{a \cdot b} |a\rangle\langle b|
 \end{aligned}$$

Suppose we have

$$H^{\otimes(n-1)} = \frac{1}{\sqrt{2^{n-1}}} \sum_{x', y' \in \{0,1\}^{n-1}} (-1)^{x' \cdot y'} |x'\rangle\langle y'|$$

By induction,



$$\begin{aligned}
H^{\otimes n} &= H^{\otimes(n-1)} \otimes H \\
&= \frac{1}{\sqrt{2^{n-1}}} \sum_{x', y' \in \{0,1\}^{n-1}} (-1)^{x' \cdot y'} |x'\rangle \langle y'| \otimes \frac{1}{\sqrt{2}} \sum_{a, b \in \{0,1\}} (-1)^{a \cdot b} |a\rangle \langle b| \\
&= \frac{1}{\sqrt{2^{n-1}} \times 2} \left(\sum_{x', y' \in \{0,1\}^{n-1}} (-1)^{x' \cdot y'} |x'\rangle \langle y'| \otimes \sum_{a, b \in \{0,1\}} (-1)^{a \cdot b} |a\rangle \langle b| \right) \\
&= \frac{1}{\sqrt{2^n}} \left(\sum_{x', y' \in \{0,1\}^{n-1}} \sum_{a, b \in \{0,1\}} (-1)^{x' \cdot y'} (-1)^{a \cdot b} |x'\rangle \langle y'| \otimes |a\rangle \langle b| \right)^1 \\
&= \frac{1}{\sqrt{2^n}} \left(\sum_{x', y' \in \{0,1\}^{n-1}} \sum_{a, b \in \{0,1\}} (-1)^{x' \cdot y'} (-1)^{a \cdot b} |x'\rangle \otimes |a\rangle \langle y'| \otimes \langle b| \right) \\
&= \frac{1}{\sqrt{2^n}} \left(\sum_{x', y' \in \{0,1\}^{n-1}} \sum_{a, b \in \{0,1\}} (-1)^{x' \cdot y'} (-1)^{a \cdot b} |x'a\rangle \langle y'b| \right) \\
&= \frac{1}{\sqrt{2^n}} \left(\sum_{x, y \in \{0,1\}^n} (-1)^{x \cdot y} |x\rangle \langle y| \right)
\end{aligned}$$

So we have proved by induction that

$$H^{\otimes n} = \frac{1}{\sqrt{2^n}} \sum_{x, y} (-1)^{x \cdot y} |x\rangle \langle y|.$$

Exercise 3

(Commutation relations for the Pauli matrices) Verify the commutation relations

$$[X, Y] = 2iZ; \quad [Y, Z] = 2iX; \quad [Z, X] = 2iY.$$

There is an elegant way of writing this using ε_{jkl} , the antisymmetric tensor on three indices, for which $\varepsilon_{jkl} = 0$ except for $\varepsilon_{123} = \varepsilon_{231} = \varepsilon_{312} = 1$, and $\varepsilon_{321} = \varepsilon_{213} = \varepsilon_{132} = -1$:

$$[\sigma_j, \sigma_k] = 2i \sum_{l=1}^3 \varepsilon_{jkl} \sigma_l.$$

Exercise 4

Suppose $[A, B] = 0$, $\{A, B\} = 0$, and A is invertible. Show that B must be 0.

¹ We can write this thanks to the bilinearity of the tensor product: $(X_1 + X_2) \otimes (Y_1 + Y_2) = \sum_{i,j} X_i \otimes Y_j$



Exercise 5

Suppose A and B are commuting Hermitian operators. Prove that $\exp(A)\exp(B) = \exp(A + B)$.



Exercise 6

Suppose we have a qubit in the state $|0\rangle$, and we measure the observable X . What is the average value of X ? What is the standard deviation of X ?

Exercise 7

Calculate the probability of obtaining the result $+1$ for a measurement of $\vec{v} \cdot \vec{\sigma}$, given that the state prior to measurement is $|0\rangle$. What is the state of the system after the measurement if $+1$ is obtained?

Exercise 8

Show that the average value of the observable $X_1 Z_2$ for a two qubit system measured in the state $(|00\rangle + |11\rangle)/\sqrt{2}$ is zero.

Exercise 9

Verify that the Bell basis forms an orthonormal basis for the two qubit state space.

Exercise 1

Suppose E is any positive operator acting on Alice's qubit. Show that $\langle\psi|E \otimes I|\psi\rangle$ takes the same value when $|\psi\rangle$ is any of the four Bell states. Suppose some malevolent third party ('Eve') intercepts Alice's qubit on the way to Bob in the superdense coding protocol. Can Eve infer anything about which of the four possible bit strings 00, 01, 10, 11 Alice is trying to send? If so, how, or if not, why not?

To archive all results (solutions, report together with your code, etc.) and submit them using Quantum computing course moodle.