Session 17 · Thursday 22 October 2026 · Four to work. Everything on this sheet is about the same thing: the number you measured on a part may not belong to the part.

What a part is, and what it costs to make one

The claim the whole hour tests. A parts list promises you this:

$$\text{protein out} \;=\; p \times r$$

where p is what the promoter does and r is what the RBS does. Two numbers, each belonging to one part, multiplied. Every abstraction in this course so far has assumed something of this shape: **composing two things combines their behaviours and leaves the things alone.**

The three ways a promoter refuses to be a thing.

no boundary	5′-RACE tells you where transcription *started*. There is no second edge — not operationally, not in the annotation, not in any model.
not contained	strength depends on flanking DNA, on the host, on ribosome availability. None of that is in the tube, and we have no model for how it moves the number.
joining changes both	the transcript's 5′UTR is encoded at the junction, and the 5′UTR sets RNA stability. Attaching a promoter to a gene changes the mRNA's lifetime — and so the RNA level you were about to call "promoter strength".

Three classes of context, after Cardinale & Arkin (2012). Use these names; they have different fixes.

compositional	what the neighbours do. *Physical*: parts on one DNA, RNA or polypeptide interacting. *Functional*: two consumers drawing on one shared output — retroactivity.
host	parasitic interaction with the cell's own machinery; competition for ribosomes, tRNAs, proteases; coupling to growth and replication.
environmental	temperature, pH, medium — directly, and through the host's response.

1 · Move a promoter, and say what kind of trouble you are in

You characterised promoter **P** carefully: high-copy plasmid, GFP downstream, DH5α, LB, 37 °C. You now want it on the chromosome, in front of a different gene, in a production strain, in minimal medium.

(a) Name **four** things that could move the number, and classify each one as compositional, host or environmental.

(b) Exactly one of your four is different in kind from the other three. Which, and what is the difference? *One sentence.*

(c) For each of the other three, say whether you would **model it**, or **hold it fixed and report it**. There is no third option, and the choice is not the same for all three.

---

2 · How much of the answer do the parts own?

What they built. Kosuri *et al.* (2013) designed all 114 × 111 promoter × RBS combinations — 12,654, less one that created an incompatible restriction site, so **12,653** — synthesised them as an oligo library, and cloned them into a single vector carrying **sfGFP** as the output and a constitutive **mCherry on the same plasmid** as an internal control. Protein levels came from **FlowSeq**: sort the population into twelve log-spaced bins by the sfGFP/mCherry ratio, sequence each bin, and reconstruct every construct's level from where its reads landed. RNA came from RNA-seq, reported as an RNA/DNA ratio.

Note what the mCherry is for: it divides out the cell-to-cell and plasmid-to-plasmid variation that would otherwise be mistaken for part behaviour. **The instrument needs insulating before the parts do.**

With that, the variance decomposes as:

	PROMOTER	RBS	UNEXPLAINED
RNA level	92%	4%	4%
protein level	54%	30%	16.7%

(a) The RBS is a *translational* element. Give a mechanism by which it takes 30% of the variance in protein but only 4% in RNA.

(b) In the same study, a 10-fold increase in translation efficiency **correlates with** about a 3-fold increase in RNA abundance — though it saturates: the weakest-to-strongest RBS span of 87-fold corresponds to only about 4.3-fold in mRNA.

Which of the three failures on the front page is this? And why does it mean the $p \times r$ model is not merely inaccurate but has **no true p to estimate** — so that collecting more data cannot fix it?

(c) 64% of protein levels land within twofold of the $p \times r$ prediction; the largest 5% of deviations are off by **an average of 13-fold**. You are designing a ten-gene pathway and need every gene within twofold. Assuming independence, what fraction of ten-gene builds has every gene inside tolerance? Is independence a safe assumption here, and does it make your estimate optimistic or pessimistic?

---

3 · Where prediction works, and what that tells you

What they built. You cannot measure a terminator with one reporter — you would be measuring the promoter too. So Cambray *et al.* (2013) built a measurement device: **green and red fluorescent reporters bracketing the terminator test site**, and read-through estimated as the ratio of downstream to upstream fluorescence. Termination efficiency is what is left; 100% means every arriving polymerase was stopped.

A ratio is still contaminated, because mRNA stability differs on either side. So the device also carries **RNase sites flanking the test site, to standardise the ends of the processed mRNA** — the same move as item 2's

mCherry, one layer down.

They then fitted a **linear model**: four sequence features chosen by stepwise selection out of twelve candidates, fitted on **31 curated terminators**, and then used to predict **23 that were held out**. The features are biophysically motivated — hairpin stability normalised by length, the U-tail, the closing base pair.

The paper reports **r = 0.9 (cross-validated 0.85) on the 31 it trained on**, and r = 0.67 (CV 0.61) across all 54. Below, recomputed from the supplementary table, is the median **signed** error per class — positive means the model predicted more termination than was measured:

CLASS	N	MEDIAN PREDICTED − MEASURED (TE POINTS, OUT OF 100)
`terminator` — the curated training set	31	+0.7
extended (ET)	6	**−8.3**
low folding frequency (LFFT)	9	+5.3
low efficiency (LET)	7	**+44.4**
both (LFFT/ET)	1	+26.1

(a) A colleague reports the pooled correlation over all 54 and concludes the model is weak. Two things are wrong with stopping there. Name both: one is about the **classes**, and one is about **which terminators the model was fitted on**.

(b) Look at the sign column. The model does not miss at random: LET and LFFT are **over**predicted, ET is **under**predicted, and the paper says exactly this. Explain why a model that fails in named, *directional* ways is more useful than one with the same average error scattered at random.

(c) Complete the sentence in your own words, in **no more than fifteen words**: *"Prediction from sequence works when `_____`."*

---

4 · Joining them, and the arithmetic of overhangs

Golden Gate uses a Type IIS enzyme, which cuts at a fixed offset *outside* its own recognition site, so the site leaves with the scrap and the junction is **four bases you chose** — which neither restriction cloning nor Gibson gives you.

(a) There are $4^4 = 256$ four-base overhangs. A **palindromic** overhang equals its own reverse complement, so it anneals to a copy of itself and the fragment can go in either way round — directionality is lost. How many palindromic four-base overhangs are there? *Show the counting*.

(b) An overhang and its reverse complement are the same junction identity. Using your answer to (a), how many **distinct usable** junction identities are there?

(c) Place the junctions. One transcriptional unit from three parts in a backbone — **backbone → promoter → 5′UTR/RBS → CDS → backbone** — is four junctions, each leaving four chosen bases behind *in the product*.

For each, say in a few words **what sequence the scar sits inside**, and name the one you would **most want measured** before trusting the construct. Cambray *et al.* (2018) put **83%** of what designed properties explain in the structure within 30 nt either side of the start codon; use that. (An *E. coli* result; (d)'s standard is a plant one.)

The common syntax puts a junction at the start codon anyway, with the overhang fixed at **AATG**. **What does fixing it buy, and what does it not?** *One line.*

(d) Check your rule against the field's set. A reasonable rule: every pair of junctions must differ in at least two bases, compared directly **and** against the other's reverse complement. The eleven fusion sites analysed by Pryor *et al.* are

> *GGAG TGAC TCCC TACT CCAT AATG AGCC TTCG GCTT GGTA CGCT*

Apply it to **TACT** and **GGTA**, reverse-complementing GGTA first. Over all 55 pairs the rule flags **three**.

Potapov *et al.* (2018) measured ligation for all 256 possible overhangs; Pryor *et al.* (2020) used that data to predict this set at **81%** error-free, about **92%** without the GGTA/TACT pair. **Note the dates. So what is your rule good for, and what is it not?** *Two lines.*

Stretch — the one nobody has answered

Manufacturing works — item 3's 47% against 93%, same question, same paper. The same group, the same year, also proposed a **measurement standard** for part collections: an ANOVA framework reporting a part's *activity* and a second statistic, its *quality*, tracking how much that activity moves across contexts. The standard did not take.

Write one paragraph. What would a part-measurement standard have to offer a working lab for that lab to pay the cost of adopting it? Be concrete about who pays and who benefits, and name one reason the 2013 proposal might have failed that is *not* "people are lazy".

Sources: Cardinale & Arkin, Biotechnol J (2012); Kosuri et al., PNAS (2013); Cambray et al., NAR (2013); Mutalik et al., Nat Methods (2013), two papers; Cambray, Guimaraes & Arkin, Nat Biotechnol (2018); Pryor et al., PLOS ONE (2020); plant common syntax (Phytobricks, 2015). The 81% and 92% are predicted, not measured.