

# Matrix Theory (EE5609) Assignment 20

Arkadipta De  
MTech Artificial Intelligence  
AI20MTECH14002

**Abstract**—This document solves a problem on a functional and linear transformation. Where,

All the codes for the figure in this document can be found at

[https://github.com/Arko98/EE5609/blob/master/Assignment\\_20](https://github.com/Arko98/EE5609/blob/master/Assignment_20)

$$\mathbf{A} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad (2.0.6)$$

$$(2.0.7)$$

Consider the following mapping,

$$g = T^t f \quad (2.0.8)$$

Then,  $\forall (x_1, x_2) \in \mathbb{F}^2$  we have,

$$g(x_1, x_2) = T^t f(x_1, x_2) \quad [\text{From (2.0.8)}] \quad (2.0.9)$$

$$= f(T(x_1, x_2)) \quad [\text{From (2.0.4)}] \quad (2.0.10)$$

$$= \mathbf{a}^T \mathbf{A} \mathbf{x} \quad (2.0.11)$$

$$= \mathbf{a}^T \begin{pmatrix} x_1 \\ 0 \end{pmatrix} \quad [\text{From (2.0.5)}] \quad (2.0.12)$$

$$= ax_1 \quad [\text{From (2.0.1)}] \quad (2.0.13)$$

Hence, (2.0.13) is the required answer.

## 2 SOLUTION

The linear functional  $f$  on  $\mathbb{F}^2$  is defined by,

$$f(x_1, x_2) = \mathbf{a}^T \mathbf{x} \quad \forall (x_1, x_2) \in \mathbb{F}^2 \quad (2.0.1)$$

where,

$$\mathbf{a} = \begin{pmatrix} a \\ b \end{pmatrix} \quad (2.0.2)$$

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \quad (2.0.3)$$

We use the following theorem,

Let  $\mathbb{V}$  and  $\mathbb{W}$  be vector spaces, over the field  $F$ . For each linear transformation  $T : \mathbb{V} \rightarrow \mathbb{W}$ , there is a unique linear transformation  $T^t : \mathbb{W}^* \rightarrow \mathbb{V}^*$  such that,

$$(T^t g)(\alpha) = g(T\alpha) \quad (2.0.4)$$

The given linear operator  $T$  defined as,

$$T(x_1, x_2) = \mathbf{A} \mathbf{x} = \begin{pmatrix} x_1 \\ 0 \end{pmatrix} \quad \forall (x_1, x_2) \in \mathbb{F}^2 \quad (2.0.5)$$