

Matrix Theory (EE5609) Assignment 14

Arkadipta De
MTech Artificial Intelligence
AI20MTECH14002

Abstract—This document proves the existence of inverse of Hilbert Matrix.

All the codes for the figure in this document can be found at

https://github.com/Arko98/EE5609/blob/master/Assignment_14

1 PROBLEM

Prove that the following matrix is invertible and \mathbf{A}^{-1} has integer entries.

$$\mathbf{A} = \begin{pmatrix} 1 & \frac{1}{2} & \cdots & \frac{1}{n} \\ \frac{1}{2} & \frac{1}{3} & \cdots & \frac{1}{n+1} \\ \vdots & \vdots & \cdots & \vdots \\ \frac{1}{n} & \frac{1}{n+1} & \cdots & \frac{1}{2n-1} \end{pmatrix}$$

2 SOLUTION

Let \mathbf{A}_3 be 3×3 matrix i.e

$$\mathbf{A}_3 = \begin{pmatrix} 1 & \frac{1}{2} & \frac{1}{3} \\ \frac{1}{2} & \frac{1}{3} & \frac{1}{4} \\ \frac{1}{3} & \frac{1}{4} & \frac{1}{5} \end{pmatrix} \quad (2.0.1)$$

Now we find the inverse of the matrix \mathbf{A}_3 as follows,

$$\begin{pmatrix} 1 & \frac{1}{2} & \frac{1}{3} & 1 & 0 & 0 \\ \frac{1}{2} & \frac{1}{3} & \frac{1}{4} & 0 & 1 & 0 \\ \frac{1}{3} & \frac{1}{4} & \frac{1}{5} & 0 & 0 & 1 \end{pmatrix} \quad (2.0.2)$$

$$\begin{matrix} R_2 = R_2 - \frac{1}{2}R_1 \\ R_3 = R_3 - \frac{1}{3}R_1 \end{matrix} \begin{pmatrix} 1 & \frac{1}{2} & \frac{1}{3} & 1 & 0 & 0 \\ 0 & \frac{1}{12} & \frac{1}{12} & -\frac{1}{2} & 1 & 0 \\ 0 & \frac{1}{12} & \frac{1}{45} & -\frac{1}{3} & 0 & 1 \end{pmatrix} \quad (2.0.3)$$

$$\begin{matrix} R_3 = R_3 - R_2 \end{matrix} \begin{pmatrix} 1 & \frac{1}{2} & \frac{1}{3} & 1 & 0 & 0 \\ 0 & \frac{1}{12} & \frac{1}{12} & -\frac{1}{2} & 1 & 0 \\ 0 & 0 & \frac{1}{180} & \frac{1}{6} & -1 & 1 \end{pmatrix} \quad (2.0.4)$$

$$\begin{matrix} R_2 = 12R_2 \\ R_3 = 180R_3 \end{matrix} \begin{pmatrix} 1 & \frac{1}{2} & \frac{1}{3} & 1 & 0 & 0 \\ 0 & 1 & 1 & -6 & 12 & 0 \\ 0 & 0 & 1 & 30 & -180 & 180 \end{pmatrix} \quad (2.0.5)$$

$$\begin{matrix} R_2 = R_2 - R_3 \\ R_1 = R_1 - R_3 \end{matrix} \begin{pmatrix} 1 & \frac{1}{2} & 0 & -9 & 60 & -60 \\ 0 & 1 & 0 & -36 & 192 & -180 \\ 0 & 0 & 1 & 30 & -180 & 180 \end{pmatrix} \quad (2.0.6)$$

$$\begin{matrix} R_1 = R_1 - \frac{1}{2}R_2 \end{matrix} \begin{pmatrix} 1 & 0 & 0 & 9 & -36 & 30 \\ 0 & 1 & 0 & -36 & 192 & -180 \\ 0 & 0 & 1 & 30 & -180 & 180 \end{pmatrix} \quad (2.0.7)$$

Hence we see that \mathbf{A}_3 is invertible and the inverse contains integer entries and \mathbf{A}_3^{-1} is given by,

$$\mathbf{A}_3^{-1} = \begin{pmatrix} 9 & -36 & 30 \\ -36 & 192 & -180 \\ 30 & -180 & 180 \end{pmatrix} \quad (2.0.8)$$

Now, expressing \mathbf{A}_4 using \mathbf{A}_2 we get,

$$\mathbf{A}_4 = \begin{pmatrix} \mathbf{A}_3 & \mathbf{u} \\ \mathbf{u}^T & \mathbf{d} \end{pmatrix} \quad (2.0.9)$$

where,

$$\mathbf{u} = \begin{pmatrix} \frac{1}{4} \\ \frac{1}{5} \\ \frac{1}{6} \end{pmatrix} \quad (2.0.10)$$

$$\mathbf{d} = \left(\frac{1}{7} \right) \quad (2.0.11)$$

Now assuming \mathbf{A}_4 has an inverse, then from (2.0.9), the inverse of \mathbf{A}_4 can be written as follows,

$$\mathbf{A}_4^{-1} = \begin{pmatrix} \mathbf{A}_3^{-1} + \mathbf{A}_3^{-1}\mathbf{u}\mathbf{X}^{-1}\mathbf{u}^T\mathbf{A}_3^{-1} & -\mathbf{A}_3^{-1}\mathbf{u}\mathbf{X}^{-1} \\ -\mathbf{X}^{-1}\mathbf{u}^T\mathbf{A}_3^{-1} & \mathbf{X}^{-1} \end{pmatrix} \quad (2.0.12)$$

where,

$$\mathbf{X} = \mathbf{d} - \mathbf{u}^T\mathbf{A}_3^{-1}\mathbf{u} \quad (2.0.13)$$

Now, the assumption of \mathbf{A}_4 being invertible will hold if and only if \mathbf{A}_3 is invertible, which has been proved in (2.0.8) and \mathbf{X} from (2.0.13) is invertible. We now prove that \mathbf{X} is invertible as follows,

$$\mathbf{X} = \left(\frac{1}{7}\right) - \begin{pmatrix} \frac{1}{4} & \frac{1}{5} & \frac{1}{6} \end{pmatrix} \begin{pmatrix} 9 & -36 & 30 \\ -36 & 192 & -180 \\ 30 & -180 & 180 \end{pmatrix} \begin{pmatrix} \frac{1}{4} \\ \frac{1}{5} \\ \frac{1}{6} \end{pmatrix} \quad (2.0.14)$$

$$\Rightarrow \mathbf{X} = \left(\frac{1}{2800}\right) \quad (2.0.15)$$

Hence, \mathbf{X} is a square invertible matrix as it has linearly independent columns, hence \mathbf{X}^{-1} is given by,

$$\mathbf{X}^{-1} = (2800) \quad (2.0.16)$$

Hence, \mathbf{A}_4 is invertible. Now putting the values of \mathbf{A}_3^{-1} , \mathbf{X}^{-1} and \mathbf{u} we get,

$$\mathbf{A}_3^{-1} + \mathbf{A}_3^{-1}\mathbf{u}\mathbf{X}^{-1}\mathbf{u}^T\mathbf{A}_3^{-1} = \begin{pmatrix} 16 & -120 & 240 \\ -120 & 1200 & -2700 \\ 240 & -2700 & 6480 \end{pmatrix} \quad (2.0.17)$$

$$-\mathbf{A}_3^{-1}\mathbf{u}\mathbf{X}^{-1} = \begin{pmatrix} -140 \\ 1680 \\ -4200 \end{pmatrix} \quad (2.0.18)$$

$$-\mathbf{X}^{-1}\mathbf{u}^T\mathbf{A}_3^{-1} = \begin{pmatrix} -140 & 1680 & -4200 \end{pmatrix} \quad (2.0.19)$$

$$\mathbf{X}^{-1} = (2800) \quad (2.0.20)$$

Putting values from (2.0.17), (2.0.18), (2.0.19) and (2.0.20) into (2.0.12) we get,

$$\mathbf{A}_4^{-1} = \begin{pmatrix} 16 & -120 & 240 & -140 \\ -120 & 1200 & -2700 & 1680 \\ 240 & -2700 & 6480 & -4200 \\ -140 & 1680 & -4200 & 2800 \end{pmatrix} \quad (2.0.21)$$

Hence, from (2.0.21) we prove that, \mathbf{A}_4 is invertible and has integer entries. Hence by induction we conclude that as \mathbf{A}_{n-1} will always have inverse with

integer entries hence, \mathbf{A}_n will also have inverse with integer entries for all integer n .

Hence Proved.